

H. Sugimoto, Sugimoto Clinic, Dr. H. Tanaka, Tanaka Clinic, Dr. M. Haji, Kyushu Rosai Hospital, Dr. I. Shiroma, Nakagami Hospital, Dr. T. Nakachi, Chatan Hospital, Dr. H. Yoshida, Shonan Hospital, Dr. K. Ono, Takagi Hospital, Dr. E. Higashi, Social Insurance Chikuhō Hospital, Dr. H. Shinozaki, Social Insurance Inatsuki Hospital, Dr. M. Masakado, Nakatsu Municipal Hospital, Dr. T. Yamashita, Yamashita Tsukasa Clinic.

Conflict of interest

There are no conflicts of interest.

REFERENCES

- [1] J.M. Murabito, J.C. Evans, M.G. Larson, K. Nieto, D. Levy, P.W. Wilson, The ankle-brachial index in the elderly and risk of stroke, coronary disease, and death: the Framingham Study, *Arch. Intern. Med.* 163 (2003) 1939-1942.
- [2] A.B. Newman, L. Shemanski, T.A. Manolio, M. Cushman, M. Mittelmark, J.F. Polak, et al., Ankle-arm index as a predictor of cardiovascular disease and mortality in the Cardiovascular Health Study. The Cardiovascular Health Study Group, *Arterioscler. Thromb. Vasc. Biol.* 19 (1999) 538-545.
- [3] M.H. Criqui, R.D. Langer, A. Fronek, H.S. Feigelson, M.R. Klauber, T.J. McCann, et al., Mortality over a period of 10 years in patients with peripheral arterial disease, *N. Engl. J. Med.* 326 (1992) 381-386.
- [4] M.H. Criqui, V. Vargas, J.O. Denenberg, E. Ho, M. Allison, R.D. Langer, et al., Ethnicity and peripheral arterial disease: the San Diego Population Study, *Circulation* 112 (2005) 2703-2707.
- [5] A. Matsuda, T. Kuzuya, Arteriosclerosis obliterans in Japanese diabetic patients, *Tohoku J. Exp. Med.* 141 (Suppl.) (1983) 569-573.
- [6] D.P. Faxon, M.A. Creager, S.C. Smith Jr., R.G. Pasternak, J.W. Olin, M.A. Bettmann, et al., Atherosclerotic Vascular Disease Conference: Executive summary: Atherosclerotic Vascular Disease Conference proceeding for healthcare professionals from a special writing group of the American Heart Association, *Circulation* 109 (2004) 2595-2604.
- [7] A.T. Hirsch, M.H. Criqui, D. Treat-Jacobson, J.G. Regensteiner, M.A. Creager, J.W. Olin, et al., Peripheral arterial disease detection, awareness, and treatment in primary care, *JAMA* 286 (2001) 1317-1324.
- [8] P. Greenland, J. Abrams, G.P. Aurigemma, M.G. Bond, L.T. Clark, M.H. Criqui, et al., Prevention Conference V: Beyond secondary prevention: identifying the high-risk patient for primary prevention: noninvasive tests of atherosclerotic burden: Writing Group III, *Circulation* 101 (2000) E16-E22.
- [9] J.A. Dormandy, R.B. Rutherford, Management of peripheral arterial disease (PAD). TASC Working Group. TransAtlantic Inter-Society Consensus (TASC), *J. Vasc. Surg.* 31 (2000) S1-S296.
- [10] T. Fujiwara, S. Saitoh, S. Takagi, H. Ohnishi, J. Ohata, H. Takeuchi, et al., Prevalence of asymptomatic arteriosclerosis obliterans and its relationship with risk factors in inhabitants of rural communities in Japan: Tanno-Sobetsu study, *Atherosclerosis* 177 (2004) 83-88.
- [11] T.C. Collins, M. Suarez-Almazor, N.J. Peterson, An absent pulse is not sensitive for the early detection of peripheral arterial disease, *Fam. Med.* 38 (2006) 38-42.
- [12] S. Laing, R.M. Greenhalgh, The detection and progression of asymptomatic peripheral arterial disease, *Br. J. Surg.* 70 (1983) 628-630.
- [13] S.L. Eason, N.J. Petersen, M. Suarez-Almazor, B. Davis, T.C. Collins, Diabetes mellitus, smoking, and the risk for asymptomatic peripheral arterial disease: whom should we screen? *J. Am. Board Fam. Pract.* 18 (2005) 355-361.
- [14] P.E. Norman, W.A. Davis, D.G. Bruce, T.M. Davis, Peripheral arterial disease and risk of cardiac death in type 2 diabetes: the Fremantle Diabetes Study, *Diabetes Care* 29 (2006) 575-580.
- [15] M. McKenna, S. Wolfson, L. Kuller, The ratio of ankle and arm arterial pressure as an independent predictor of mortality, *Atherosclerosis* 87 (1991) 119-128.
- [16] M.M. McDermott, J. Feinglass, R. Slavensky, W.H. Pearce, The ankle-brachial index as a predictor of survival in patients with peripheral vascular disease, *J. Gen. Intern. Med.* 9 (1994) 445-449.
- [17] C.J. Sikkink, W.N. van Asten, M.A. van't Hof, H. van Langen, J.A. van der Vliet, Decreased ankle/brachial indices in relation to morbidity and mortality in patients with peripheral arterial disease, *Vasc. Med.* 2 (1997) 169-173.
- [18] H.E. Resnick, R.S. Lindsay, M.M. McDermott, R.B. Devereux, K.L. Jones, R.R. Fabsitz, et al., Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: the Strong Heart Study, *Circulation* 109 (2004) 733-739.
- [19] T.C. Collins, M. Suarez-Almazor, R.L. Bush, N.J. Petersen, Gender and peripheral arterial disease, *J. Am. Board Fam. Med.* 19 (2006) 132-140.
- [20] K. Wattanakit, A.R. Folsom, E. Selvin, B.D. Weatherley, J.S. Pankow, F.L. Brancati, et al., Risk factors for peripheral arterial disease incidence in persons with diabetes: the Atherosclerosis Risk in Communities (ARIC) Study, *Atherosclerosis* 180 (2005) 389-397.
- [21] A.R. Yataco, A.W. Gardner, Acute reduction in ankle/brachial index following smoking in chronic smokers with peripheral arterial occlusive disease, *Angiology* 50 (1999) 355-360.

Naoki Nakashima, MD, PhD

Associate Professor
Department of Medical Informatics
Kyushu University Hospital

3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, JAPAN

TEL 092-642-5881

FAX 092-642-5889

mail to: nnaoki@info.med.kyushu-u.ac.jp

March, 2009