

- (1999) *Nature*, 399, 700-704.
- 6) Yang, W. & Woodgate, R. (2007) *Proc. Natl. Acad. Sci. U.S.A.*, 104, 15591-15598.
- 7) Masuda, Y. & Kamiya, K. (2002) *FEBS Lett.*, 520, 88-92.
- 8) Auerbach, P., Bennett, R.A., Bailey, E.A., Krokan, H.E., & Demple, B. (2005) *Proc. Natl. Acad. Sci. U.S.A.*, 102, 17711-17716.
- 9) Acharya, N., Johnson, R.E., Prakash, S., & Prakash, L. (2006) *Mol. Cell. Biol.*, 26, 9555-9563.
- 10) Murakumo, Y., Roth, T., Ishii, H., Rasio, D., Numata, S., Croce, C.M., & Fishel, R. (2000) *J. Biol. Chem.*, 275, 4391-4397.
- 11) Masuda, Y., Ohmae, M., Masuda, K., & Kamiya, K. (2003) *J. Biol. Chem.*, 278, 12356-12360.
- 12) Haracska, L., Unk, I., Prakash, L., & Prakash, S. (2006) *Proc. Natl. Acad. Sci. U.S.A.*, 103, 6477-6482.
- 13) Wood, A., Garg, P., & Burgers, P.M. (2007) *J. Biol. Chem.*, 282, 20256-20263.
- 14) Guo, C., Sonoda, E., Tang, T.S., Parker, J.L., Bielen, A.B., Takeda, S., Ulrich, H.D., & Friedberg, E.C. (2006) *Mol. Cell.*, 23, 265-271.
- 15) Masuda, Y. & Kamiya, K. (2006) *J. Biol. Chem.*, 281, 24314-24321.

増田 雄司, 神谷 研二  
 (広島大学原爆放射線医科学研究所  
 分子発がん制御研究分野)

Induced mutagenesis and translesion DNA synthesis—structure and function of REV1—  
 Yuji Masuda and Kenji Kamiya (Department of Experimental Oncology, Research Institute for Radiation Biology and Medicine, Hiroshima University, Minami-ku, Hiroshima 734-8553, Japan)

## [Fe]-ヒドロゲナーゼ (Hmd) の構造解析から見てきたヒドロゲナーゼ活性中心の収斂進化

### はじめに

ヒドロゲナーゼは水素ガス ( $H_2$ ) を活性化し、ヒドリドとプロトンに分解する反応を触媒する酵素である。多くのヒドロゲナーゼはヒドリドをさらに分解し、二つの電子とプロトンにする。その触媒反応は可逆であり、逆反応では  $H_2$  が生産される<sup>1)</sup>。ヒドロゲナーゼのバイオテクノロジーへの応用として、燃料電池電極や水素生産の触媒としての活用が目ざされている<sup>2)</sup>。

自然界において、ヒドロゲナーゼは微生物生態系の水素


代謝で重要な役割を担っている。嫌気条件下、有機物は細菌や原生動物などによって分解され、有機酸や  $H_2$  になる。ここで大量の  $H_2$  が発生するが ( $\sim 10^6$  トン/年)<sup>3)</sup>、 $H_2$  はメタンや酢酸の生産および硫酸塩の還元に使われるため、環境中に存在する  $H_2$  濃度は低く保たれている。一部の  $H_2$  は好気環境に分散し、そこで好気性細菌によって酸化される<sup>4)</sup>。ヒドロゲナーゼはこれらの水素代謝系で、 $H_2$  の生産と酸化を触媒する。

ヒドロゲナーゼはおよそ 80 年前に発見され、これまでに [NiFe] ヒドロゲナーゼと [FeFe] ヒドロゲナーゼという 2 種類がよく研究されてきた。名称からも明らかなように、活性中心に含まれる金属の種類が異なり、前者はニッケルと鉄を、後者は 2 原子の鉄を含んでいる (図 1)<sup>5-7)</sup>。20 年ほど前、メタン菌から新しいタイプのヒドロゲナーゼが発見された。この新規酵素はひとつの鉄のみを活性中心に含んでいることから、[Fe] ヒドロゲナーゼ (Hmd) と呼ばれている (図 1)<sup>8)</sup>。

本稿では、Hmd の活性中心に関する知見を概説した後、ヒドロゲナーゼの特徴的な鉄錯体構造から、3 種類のヒドロゲナーゼの進化について述べる。

### 1. [Fe]-ヒドロゲナーゼ (Hmd) の機能と性質

[Fe] ヒドロゲナーゼの分類名は  $H_2$ -forming methylenetetrahydromethanopterin dehydrogenase で、略称は Hmd である。その触媒反応では、 $H_2$  を活性化してヒドリドとプロトンに分解し、ヒドリドをメチルテトラヒドロメタンプロテリン (メチル-H.MPT<sup>+</sup>) に転移し、メチレンテトラヒドロメタンプロテリン (メチレン-H.MPT) をつくる。逆反応では、メチレン-H.MPT とプロトンから水素ガスを発生する (式 1)<sup>9)</sup>。



テトラヒドロメタンプロテリンはメタン菌の C1 キャリヤーであり、Hmd の触媒する反応はメタン生成代謝に含まれる。Hmd は水素利用性のメタン菌のいくつかに見出されているが、*Methanothermobacter marburgensis* 由来の酵素がよく研究されている。このメタン菌では、培地中のニッケルが制限された条件 (50nM 以下) で Hmd が大量に生産され、全タンパク質の 5% を超えることが知られている<sup>8)</sup>。

Hmd は分子量 38,000 のサブユニットのホモ二量体からなり、2 分子の鉄コファクター (FeGP コファクター) を含んでいる。FeGP コファクターはピリジノール環が GMP

# Biochemical analysis of human PIF1 helicase and functions of its N-terminal domain

Yongqing Gu<sup>1,2</sup>, Yuji Masuda<sup>1</sup> and Kenji Kamiya<sup>1\*</sup><sup>1</sup>Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima 734-8553, Japan and<sup>2</sup>School of Medicine, Shihezi University, Shihezi, Xinjiang 832002, China

Received May 1, 2008; Revised September 8, 2008; Accepted September 9, 2008

## ABSTRACT

The evolutionary conserved PIF1 DNA helicase family appears to have largely nonoverlapping cellular functions. To better understand the functions of human PIF1, we investigated biochemical properties of this protein. Analysis of single-stranded (ss) DNA-dependent ATPase activity revealed nonstructural ssDNA to greatly stimulate ATPase activity due to a high affinity for PIF1, even though PIF1 preferentially unwinds forked substrates. This suggests that PIF1 needs a ssDNA region for loading and a forked structure for translocation entrance into a double strand region. Deletion analysis demonstrated novel functions of a unique N-terminal portion, named the PIF1 N-terminal (PINT) domain. When the PINT domain was truncated, apparent affinity for ssDNA and unwinding activity were much reduced, even though the maximum velocity of ATPase activity and  $K_m$  value for ATP were not affected. We suggest that the PINT domain contributes to enhancing the interaction with ssDNA through intrinsic binding activity. In addition, we found DNA strand-annealing activity, also residing in the PINT domain. Notably, the unwinding and annealing activities were inhibited by replication protein A. These results suggest that the functions of PIF1 might be restricted with particular situations and DNA structures.

## INTRODUCTION

Helicases are ubiquitous enzymes that catalyze the unwinding of DNA duplexes using ATP as their energy source. They therefore play vital roles in nearly all DNA metabolic processes, including DNA replication, recombination and repair. The PIF1 subfamily of 5'–3' DNA helicases (1–10) belongs to the SFI superfamily, conserved in diverse organisms (11).

In *Saccharomyces cerevisiae* (*Sc*), ScPif1 was originally identified because of its involvement in recombination

of mitochondrial DNA (mtDNA) (12,13). Dysfunction of ScPif1 leads to mitochondrial genetic instability due to spontaneous oxidative damage (14–16), and induces mtDNA damage (17). ScPif1 has not only mitochondrial targeting but also nuclear targeting signals and therefore is localized in both the mitochondria and nucleus (18). Nuclear ScPif1 has multiple functions. When it is overproduced, telomeres become shorter, while they elongate when it is eliminated (18,19). In the absence of nuclear ScPif1, gross chromosome rearrangement is also increased, and healing of double-stranded broken ends via telomere addition increases ~200- to 1000-fold. These data suggest a negative regulatory role of ScPif1 in telomere metabolism (18–24). Indeed, ScPif1 catalytically inhibits telomerase activity *in vitro* (25). Other genetic data suggest that ScPif1 plays roles in Okazaki fragment processing (26), pausing of replication progression at ribosomal DNA loci (27) and unwinding of hemicatenans (28).

*Schizosaccharomyces pombe* encodes a PIF1 homolog, Pfh1 (8,10) which is required for cell cycle progression in late S-phase and for appropriate responses to DNA damage agents (8,10). It is also implicated in lagging strand DNA processing (7).

Previous biochemical studies of human PIF1 were performed using N-terminal-truncated forms of PIF1, containing only the conserved helicase motifs, located in the C-terminus, since it earlier proved impossible to obtain purified full-length human PIF1 protein (3,4,9). However, we found that the N-terminal region of human PIF1, named here as the PIF1 N-terminal (PINT) domain, is well conserved in the PIF1 family, suggesting a possible functional role. Here, we established a method to purify full-length human PIF1 protein and provided the first evidence that the PINT domain has crucial functions in this enzyme.

## MATERIALS AND METHODS

### Plasmid construction

The nucleotide sequence of the 5'-end of human PIF1 cDNA was obtained by 5'-rapid amplification of cDNA ends. Then the full-length human PIF1 cDNA was

\*To whom correspondence should be addressed. Tel: +81 82 257 5842; Fax: +81 82 257 5844; Email: kkamiya@hiroshima-u.ac.jp

amplified from a HeLa cDNA library using primers, CATATGCTCTCGGGCATAGAGGGCGGCGGAGG GGAATATGAGGACTCG and TCAGAGGATTGGG TCCATGTT by PCR, then the nucleotide sequences of the clones were verified and submitted to the database with the accession number, EU084033. The entire coding region with a histidine tag at the N-terminus was inserted into the pET20b(+) vector to yield pET20b-PIF1. The truncated forms, PIF<sup>167-641</sup> and PIF<sup>1-180</sup> consisting of the numbered amino acid residues were also cloned into pET20b(+) and pET15b, respectively, to produce his-tagged fusion proteins. The structures of the resultant plasmids, pET20b-PIF1, pET20b-PIF<sup>167-641</sup> and pET15b-PIF<sup>1-180</sup> are shown in Supplementary Figure S1. In this article, PIF<sup>167-641</sup> and PIF<sup>1-180</sup> are referred to as C-terminal region of PIF1 (PIF1C) and N-terminal region of PIF1 (PIF1N), respectively.

### Protein purification

RPA was purified as described (29) from over producing *Escherichia coli* cells (30). PIF1 and its deletion derivatives were purified as his-tagged fusion proteins at the N-termini. During all the purification steps, induced proteins were monitored by SDS-PAGE followed by staining with Coomassie Brilliant Blue R-250, or western blotting using Penta-His antibody (#34660, QIAGEN, Tokyo, Japan) or anti-PIF1 antibodies. Protein concentrations were determined by Bio-Rad protein assay using BSA (Bio-Rad, Tokyo, Japan) as the standard.

His-tagged full-length PIF1 and PIF1C were purified from overexpressing *E. coli* cells, BL21 (DE3) (31). The strain harboring a plasmid pMStRNA1, in which tRNAs for rare codons were cloned into a R6K derived kanamycin resistant plasmid (32), and pET20b-PIF1 was grown in 3 l of LB supplemented with ampicillin (250 µg/ml) and kanamycin (30 µg/ml) at 15°C, with aeration until the culture reached an  $A_{600}$  value of 0.6. Isopropyl β-D-thiogalactopyranoside (IPTG) was added to 0.2 mM, and the incubation was continued for 14 h. The resultant cell paste (9 g) was resuspended in 18 ml of buffer I (50 mM HEPES NaOH pH 7.5, 0.1 mM EDTA, 10 mM β-mercaptoethanol, 1 M NaCl) and frozen in liquid nitrogen. The cells were thawed in ice water and lysed by addition of 3 ml buffer I containing 100 mM spermidine and 4 mg/ml lysozyme. After incubation on ice for 30 min, heating in a 37°C water bath for 2 min and further incubation on ice for 30 min, the lysate was clarified by centrifugation twice at 85000g for 30 min at 4°C. Subsequent column chromatography was carried out at 4°C using a fast protein liquid chromatography (FPLC) system (GE Healthcare, Tokyo, Japan). After adding imidazole to 50 mM, the lysate was applied at 0.2 ml/min to a 1-ml HiTrap chelating column (GE Healthcare), which had been treated with 0.1 M NiSO<sub>4</sub> and then equilibrated with buffer A (50 mM HEPES NaOH pH 7.5, 10% glycerol, 10 mM β-mercaptoethanol, 1 M NaCl) containing 50 mM imidazole. The column was washed with 10 ml of equilibration buffer at 0.2 ml/min and his-tagged PIF1 was eluted with 10 ml of buffer A containing 100 mM imidazole. Fractions eluted with 100 mM imidazole were pooled

and diluted to 50 mM imidazole with buffer A, then loaded again onto a 1-ml HiTrap chelating column at 0.2 ml/min. The column was washed, and PIF1 was eluted with buffer A containing 300 mM imidazole, then loaded at 0.1 ml/min onto a Superdex 200 10/300 GL column (GE Healthcare) equilibrated with buffer A. PIF1 peak fractions were pooled, frozen in liquid nitrogen, and stored at -80°C. His-tagged PIF1C was purified under the same conditions as described for his-tagged PIF1.

His-tagged human PIF1N was purified from overexpressing *E. coli* cells, Rosetta 2 (DE3) (Novagen, Tokyo, Japan). The strain harboring pET15-PIF1N was grown in 3 l of LB supplemented with ampicillin (250 µg/ml) and chloramphenicol (30 µg/ml) at 15°C with aeration until the culture reached an  $A_{600}$  value of 0.6. IPTG was added to 0.2 mM, the incubation was continued for 14 h, and the cells were lysated as described. After adding imidazole to 50 mM, the lysate was applied at 0.2 ml/min to a 1-ml HiTrap chelating column, which had been treated with 0.1 M NiSO<sub>4</sub> and then equilibrated with buffer A containing 50 mM imidazole. The column was washed with 10 ml of equilibration buffer at 0.2 ml/min and then with 10 ml of buffer A containing 100 mM imidazole. His-tagged PIF1N was eluted with 10 ml of 300 mM imidazole in buffer A. Fractions containing PIF1N were pooled, diluted with buffer B (50 mM HEPES NaOH pH 7.5, 10 mM β-mercaptoethanol) to 100 mM of NaCl, and applied at 0.5 ml/min to a 1-ml HiTrap SP HP column (GE Healthcare) equilibrated with buffer B containing 100 mM NaCl. The column was washed with 10 ml of equilibration buffer at 0.1 ml/min, and the PIF1N was eluted with 20 ml of a linear gradient of 100–1000 mM NaCl in buffer B. Fractions containing PIF1N were pooled, frozen in liquid nitrogen and stored at -80°C.

### Antibodies

To obtain polyclonal antibodies against PIF1, truncated his-tagged PIF1 proteins (1–180 and 338–641 amino acids) were expressed in Rosetta 2 (DE3), purified and used to immunize rabbits.

### DNA substrates

The oligonucleotides employed for the preparation of DNA substrates are listed in Table 1. Oligonucleotides were 5'-end labeled using [ $\gamma$ -<sup>32</sup>P]ATP (GE Healthcare) and polynucleotide kinase (New England BioLabs, Tokyo, Japan). The schematic structures of substrates are shown in figures, and the labeled oligonucleotides are indicated with asterisks. Annealing reaction mixtures (30 µl) containing the 5'-<sup>32</sup>P-labeled oligonucleotides at 1 µM, all unlabeled oligonucleotides at 3 µM, 10 mM Tris-HCl (pH 7.5), 7 mM MgCl<sub>2</sub> and 200 mM NaCl were heated at 95°C for 10 min, transferred directly to 65°C and held at that temperature for 1 h, slow-cooled to 25°C over a period of 2 h and held at that temperature for 30 min and then cooled to 4°C. Substrates were then purified by electrophoresis through 15–25% polyacrylamide using 0.5× TBE (33) as the electrophoresis buffer. Substrates were eluted from the gel by crushing the gel

slice in TE buffer and incubating overnight at 4°C. The slurry was then filtered through Micro Bio-Spin (Bio-Rad) columns, and the DNA was recovered by ethanol precipitation and resuspended in TE buffer.

#### ATPase assays

ATPase activity was measured in a standard reaction mixture (20  $\mu$ l) containing 50 mM Tris-HCl (pH 8.0), 2 mM DTT, 1.2 mM MgCl<sub>2</sub>, 0.25 mg/ml BSA, 2 mM [ $\gamma$ -<sup>32</sup>P]ATP, indicated DNA and 1  $\mu$ l of protein sample diluted with buffer D (50 mM Tris-HCl pH 8.0, 1 M NaCl, 2 mM DTT, 10% glycerol, 0.1 mg/ml BSA) to obtain indicated final concentrations. After preincubation for 30 s at 30°C, reactions were initiated by the addition of PIF1 proteins and further incubated for 10 min. After the reaction was stopped with 4  $\mu$ l of 20 mM EDTA (pH 8.0), an aliquot (2  $\mu$ l) was spotted onto a polyethyleneimine-cellulose plate (Merck, Tokyo, Japan) and developed in 0.3 M LiCl/0.9 M formic acid. The products were analyzed using a Bio-Imaging Analyzer BAS2000 (Fuji Photo Film Co., Ltd., Tokyo, Japan). The extents of ATP hydrolysis were measured with reference to the relative ratios of radioactivity of inorganic phosphate to uncleaved ATP.

Kinetic assays to determine  $K_m$  values for ATP were performed for 10 min in 20  $\mu$ l reaction mixtures using 14 nM of PIF1 and PIF1C with 3.8  $\mu$ M and 150  $\mu$ M (in nucleotides) of M13 mp7 single-stranded (ss) DNA, respectively. Concentrations of ATP ranged from 25 to 400  $\mu$ M.  $K_m$  values were evaluated from the plot of the initial velocity versus the ATP concentration using a hyperbolic curve-fitting program with correlation coefficients ( $R^2$ ) >0.99.

#### DNA helicase assays

DNA helicase activity was measured under ATPase assay conditions (20  $\mu$ l) with the indicated DNA substrate (0.35 nM) and 1  $\mu$ l of protein sample diluted with buffer D to obtain the indicated final concentrations. After preincubation for 30 s at 30°C, reactions were initiated by the addition of PIF1 proteins and incubated for 10 min at 30°C. Helicase reactions were terminated with 10  $\mu$ l of stop solution (150 mM EDTA, 30% glycerol, 2% SDS, 0.1% bromophenol blue). In kinetic experiments, a 160- $\mu$ l reaction mixture was incubated at 30°C and 10- $\mu$ l aliquots were withdrawn at the indicated times. The reaction products were subjected to electrophoresis through 15–25% polyacrylamide gels in Tris-glycine buffer (33). The gels were dried on DE81 paper (Whatman, Tokyo, Japan) and autoradiographed. DNA products were quantified using a Bio-Imaging Analyzer.

#### Electrophoretic mobility shift assays

Oligonucleotides were labeled with polynucleotide kinase (New England BioLabs) and [ $\gamma$ -<sup>32</sup>P]ATP. Assays of DNA binding were performed with a modification of a method described previously (34). The reactions (20  $\mu$ l) were carried out under ATPase assay conditions, sometimes omitting MgCl<sub>2</sub> or ATP, with 25 pM oligonucleotides and 1  $\mu$ l of protein sample diluted with buffer D to obtain the

indicated final concentrations. Incubation was carried out on ice for 10 min followed by loading on prerunning 5% polyacrylamide gels (79:1 acrylamide/bis-acrylamide). The electrophoresis buffer contained 6 mM Tris-HCl (pH 8.0), 5 mM sodium acetate and 1 mM EDTA, and the gels were subjected to a constant voltage of 8 V/cm for 100 min at 4°C. Following gel electrophoresis, the products were analyzed as described for the helicase assay. For quantification, fractions of free DNA were measured, and the binding fractions were determined by subtraction from the amount of the free DNA at 0 nM of the protein (35).

For RPA binding experiments, the reactions (20  $\mu$ l) were carried out under the ATPase assay conditions, sometimes omitting ATP. The substrate 3F:4L (0.35 nM) was incubated, directly or after heating at 100°C for 5 min, with 1  $\mu$ l of RPA sample diluted with buffer (50 mM HEPES NaOH pH 7.5, 250 mM NaCl, 10 mM  $\beta$ -mercaptoethanol, 10% glycerol) to obtain indicated final concentrations. Incubation was carried out on ice for 10 min, and the products were analyzed as in the PIF1 binding experiments.

#### DNA strand annealing assays

Strand annealing reactions (20  $\mu$ l) were carried out under the ATPase assay conditions, but in the presence of the indicated concentrations of ATP, with the 5'-end labeled substrate DNA for helicase assay (0.35 nM), which had been boiled at 100°C for 5 min and quickly chilled on ice before adding to the reaction mixture, and 1  $\mu$ l of protein sample diluted with buffer D to obtain the indicated final concentrations. After preincubation for 30 s at 30°C, reactions were initiated by the addition of PIF1 proteins, with incubation for 10 min at 30°C. After terminating the reactions with 10  $\mu$ l of stop solution, the products were analyzed as described for the helicase assay. Kinetic experiments were also carried out as described for the helicase assay.

## RESULTS

#### Purification of PIF1 protein and its deletion derivatives

Previously, biochemical studies of human PIF1 were performed using N-terminal-truncated forms of PIF1 containing only the conserved helicase motifs, located in the C-terminus (Supplementary Figure S2A). However, we found that the PINT domain is well conserved in the PIF1 family (Supplementary Figure S2B), suggesting that it plays a functional role. To examine biochemical activity, we established a procedure to purify full-length PIF1 with a 6 $\times$  histidine tag at the N-terminus at quantities sufficient for detailed biochemical studies from overproducing *E. coli* cells (Supplementary Figure S2C). We also purified a C-terminal truncated form (PIF1N) and a N-terminal truncated form (PIF1C) (Supplementary Figure S2A) to address biochemical functions of individual domains. PIF1C and PIF1N consist of only the seven helicase motifs and only the PINT domain, respectively (Supplementary Figure S2A and Materials and methods section). Elution profiles from gel filtration chromatography

suggested PIF1 and PIF1C to be monomers (data not shown) as described in yeast homologs (1,5,7). The purified proteins were analyzed by SDS-PAGE followed by CBB staining and western blotting, showing PIF1, PIF1C and PIF1N to have molecular sizes of 71, 54 and 22 kDa, respectively (Supplementary Figure S2C).

#### Unwinding activities of PIF1 and PIF1C, a mutant lacking PINT domain but containing the helicase domain

We first measured DNA helicase activity of the purified proteins using the indicated DNA substrate (Figure 1A). When 5' overhang (1F:2S) and 3' overhang (1R:2S) DNA were used as substrates, we detected helicase activity only on the 5'-overhang substrate, consistent with previous reports (3,9), although the activity was very low (Figure 1B). In the titration experiment, activity was only detected clearly at the maximum concentration of the purified sample (Figure 1B). Since yeast PIF1 homologs preferentially unwind forked structures (1,5,7), we tested a forked substrate (Figure 1C). The titration experiment demonstrated that human PIF1 efficiently unwound the forked substrate with about 10 times higher activity than that for the 5'-overhang substrate (Figure 1C), suggesting that the property was conserved in evolution. Time course experiments using the forked substrate demonstrated that PIF1 could unwind up to 50% of the substrate in a 15 min reaction, although longer incubation did not increase the products (Figure 1D).


To compare the unwinding activity of PIF1C, titration experiments were performed with the forked substrate.

We found that PIF1C exhibited helicase activity, but it required more than 100 times more protein to obtain equivalent activity to that of PIF1 (Figure 1C). As a control, we showed that PINT domain itself could not unwind the substrate (Figure 1C). These results suggested that the function of the PINT domain could be enhancement of the unwinding activity of the helicase domain.


#### Nonstructural ssDNA, but not forked-structural DNA, preferentially stimulates ATPase activity of PIF1

DNA helicases are enzymes with an associated DNA-dependent ATPase activity. They are presumed to use the hydrolysis of ATP to translocate to ssDNA and to subsequently break the hydrogen bonds of duplex DNA. Characterization of ATPase activity could provide important information as a helicase. To analyze ssDNA dependent ATPase activity of PIF1, titration of M13 mp7 ssDNA was performed in reactions with optimal concentrations of ATP and  $MgCl_2$  (Figure 2A). The result showed that the ATPase activity was increased depending on the concentrations of ssDNA and reached a plateau between 3 and 50  $\mu M$  (in nucleotide equivalents) of ssDNA (Figure 2A). The maximum rate of ATP hydrolysis was calculated to be about 1 000  $min^{-1}$ , which was equivalent to 5000  $min^{-1}$  at 37°C of *ScPi1* (5) and 4000  $min^{-1}$  at 30°C of the fission yeast homolog, *Pfh1* (7), which have been determined under reaction conditions with nearly saturated concentrations of ssDNA.

To describe precisely the concentration of ssDNA required for ATPase activity, a kinetic parameter,



**Figure 1.** Helicase activity of PIF1. (A) Schematic structures of the DNA substrates. The asterisks indicate  $^{32}P$ -labeled 5' phosphate. (B) Helicase activity of PIF1. Increasing levels of PIF1 were incubated with substrates (0.35 nM) with the 5' overhang, 1F:2S (upper panel), or 3' overhang, 1R:2S (lower panel) under standard reaction mixtures at 30°C for 10 min. Reaction products were separated on a 15–25% polyacrylamide gel. (C) Helicase activity of PIF1, PIF1N and PIF1C proteins. The forked structure partial duplex DNA substrate, 3F:4L (0.35 nM), was incubated with the indicated concentrations of PIF1, PIF1N and PIF1C under standard reaction conditions at 30°C for 10 min. The quantified data are shown graphically. The errors in the experiments were <10%. (D) Time course of unwinding reactions with PIF1. The forked structure partial duplex DNA substrate, 3F:4L (0.35 nM), was incubated at 30°C for the indicated time under standard reaction conditions with PIF1 (33 nM). The quantified data are shown graphically. The errors in the experiments were <10%.



**Figure 2.** Effects of various DNA molecules on stimulation of ATPase activity of PIF1. (A) Titration of M13 mp7 ssDNA for its stimulation of ATPase activity of PIF1, PIF1N and PIF1C. ATPase activities were measured under standard reaction conditions with PIF1 proteins (13.9 nM) at 30°C for 10 min. An aliquot (2  $\mu\text{l}$ ) of the reaction products was then analyzed by thin layer chromatography. The extent of ATP hydrolysis was measured by determining the relative radioactivity of the released inorganic phosphate. Inset is a plot of the data at lower concentrations of ssDNA. The errors in the experiments were <10%. (B) Effects of the various DNA molecules on stimulation of ATPase activity of PIF1. Assays were performed with the indicated DNA (3.8  $\mu\text{M}$  in nucleotides) and PIF1 (13.9 nM). (C) Effects of oligonucleotides with different compositions on stimulation of ATPase activity by PIF1. Assays were performed with indicated DNA (7.5  $\mu\text{M}$  in nucleotides) and PIF1 (14.6 nM). (D) Effects of oligonucleotides with different compositions on stimulation of ATPase activity by PIF1C. Assays were performed with indicated DNA (150  $\mu\text{M}$  in nucleotides), and PIF1C (13.9 nM).

$K_{\text{eff}}$ , defined as the concentration of ssDNA required to achieve half-maximal ATP hydrolysis (5,36,37) was calculated from the titration curve (Figure 2A) using a hyperbolic curve-fitting program (Table 2). The calculated  $K_{\text{eff}}$  value, 0.35  $\mu\text{M}$  (in nucleotides) for M13 mp18 ssDNA, agreed with the 0.6  $\mu\text{M}$  (in nucleotides) of *ScPif1* reported for the same M13 ssDNA (5).

Since PIF1 preferentially unwound the forked substrate (Figure 1), we consider that substrate specificity for unwinding activity could be correlated with stimulation of ATPase activity. To seek preferential structures for stimulation for ATPase of PIF1, we tested various DNAs, including a fork-structure, a primer-template-structure, a hairpin-structure and a linear oligonucleotide (Table 1), as well as M13 mp18 ssDNA and double stranded (ds) DNA (Figure 2B). The reactions were carried out with 3.8  $\mu\text{M}$  (in nucleotides) of each oligonucleotide, which was nearly saturated concentration for M13 mp7 ssDNA (Figure 2A inset). Among these, M13 mp18 ssDNA proved to strongly and dsDNA to poorly stimulate ATPase activity (Figure 2B). Moreover, we found surprisingly that a oligonucleotide, 1F, which would not be expected to form secondary structures (Table 1), stimulated ATPase activity to the highest level, equivalent to that of M13 mp18 ssDNA (Figure 2B). The result suggested that an ssDNA region itself, rather than the structure of the DNA, is crucial for stimulation of ATPase activity of PIF1. Consequently, we compared ATPase activity stimulated by seven different 60-mer oligonucleotides composed of one or two nucleotides, guaranteed not to form secondary structures. The reactions were performed in the presence of a nearly saturated concentration of the oligonucleotides (7.5  $\mu\text{M}$  in nucleotides) (Supplementary Figure S3A).

The results revealed general features of ssDNA for stimulation of ATPase (Figure 2C), with the order of stimulation being poly(purine-pyrimidine)  $\geq$  polypyrimidine  $\gg$  polypurine.

Since these experiments were performed with nearly saturated concentrations of oligonucleotides, the results do not reflect affinities of the respective ssDNAs. We therefore determined a kinetic parameter,  $K_{\text{eff}}$ , calculated from data of ssDNA-titration experiments (Supplementary Figure S3A, data not shown) using a hyperbolic curve-fitting program (Table 2). The  $K_{\text{eff}}$  value for ssDNA was more than 20 times higher than those for dsDNA, showing a preference to ssDNA. For ssDNA, the  $K_{\text{eff}}$  values, except with polypurine, were essentially the same ( $\sim 0.1 \mu\text{M}$  in nucleotides), and slightly lower than with M13 ssDNA and structured DNAs, which was probably due to over estimation of the ssDNA region because of the presence of local double-stranded regions generated by the secondary structure, since the  $K_{\text{eff}}$  values were expressed in nucleotide equivalents. These results supported our conclusion that the predominant requirement for stimulation of ATPase is nonstructural ssDNA.

#### Evaluation of ATPase activity of PIF1C, a mutant lacking the PINT domain

We noted that the  $K_{\text{eff}}$  value for ssDNA of PIF1 was in line with one report for *ScPif1* (5), but significantly lower than that for human PIF1 with a N-terminal truncated form (4). We consider that the difference could be attributed to the missing function of the PINT domain. To test this possibility, we examined ATPase activity of PIF1C (Figure 2A), and also tested PIF1N as a control. First, we confirmed no ATPase activity of PIF1N (Figure 2A),




suggested again that the helicase domain is sufficient to express ssDNA-dependent ATPase activity.

### Characterization of ssDNA binding activities of PIF1 and PIF1C

It seemed that the apparent affinity to ssDNA for the helicase domain (PIF1C) was much lower than that for PIF1 in ATPase reactions. We considered that the defect might be attributed to binding ability to ssDNA. To measure DNA binding of PIF1 and PIF1C directly, we performed electrophoretic mobility shift assays (EMSA) using a oligonucleotide, d(AC)60 (Table 1) as a model substrate. It has been reported that this assay detects DNA-protein complexes in yeast and human PIF1 in an ATP-independent manner (7,9). Binding reactions were carried out under the optimal ATPase assay conditions, then products were loaded on gels as described in the

Materials and methods section. The titration experiment displayed PIF1-DNA complexes, which were increased depending on the concentration of PIF1 (Figure 3A). The apparent  $K_d$ , which is approximately equal to the protein concentration at which half the free DNA has become bound (35), was determined to be about 3 nM. This was in good agreement with the  $K_{eff}$  value of d(AC)60 for ATPase activity, because the 0.13  $\mu$ M (in nucleotides) (Table 2) corrected for the concentration of the 60-mer oligonucleotide became 2 nM.

When PIF1C was tested, we could detect PIF1C-DNA complexes, but the affinity seemed much lower than that with PIF1, demonstrating a defect in ssDNA binding activity. When the point at which half the free DNA has become bound was extrapolated from the binding curve, the apparent  $K_d$  value was estimated at about 10 nM. However, the  $K_d$  value was not directly correlated with the  $K_{eff}$  value, since it was still 6 times lower than the



**Figure 3.** EMSA of PIF1 and PIF1C for binding to ssDNA. (A) EMSA of ssDNA binding to PIF1 and PIF1C under standard reaction conditions. The  $5'$ - $^{32}$ P labeled oligonucleotide, d(AC)60, was incubated with the indicated concentrations of PIF1. Arrowheads indicate the positions of free DNA. The quantified data are shown graphically. (B) EMSA in the absence of ATP. Experiments were performed as described in (A). The quantified data are shown graphically. (C) EMSA in the absence of  $MgCl_2$ . Experiments were performed as described in (A). The quantified data are shown graphically together with data from (A) and (B). The errors in the experiments were  $<10\%$ .



apparent  $K_{\text{eff}}$  value of 60 nM (in oligonucleotides), with the  $K_{\text{eff}}$  value of 3.7  $\mu\text{M}$  (in nucleotides) (Table 2), corrected for the concentration of the 60-mer oligonucleotide.


Since it was very likely that ATP could influence the binding reaction, we performed the same experiments without ATP. With the full length PIF1, the titration curve was not affected (Figure 3B). In contrast that for PIF1C in the absence of ATP was significantly changed. The apparent affinity was increased to about 0.3 nM (Figure 3B). To determine whether the effect of ATP is caused by binding or hydrolysis, titration experiments were carried out omitting  $\text{MgCl}_2$  to prevent hydrolysis of ATP (Figure 3C). Under this condition, the ATPase activity was under the background level (data not shown), suggesting that, even though trace amounts of Mg ions were present as contaminants of chemicals, the contribution was negligible. The result clearly demonstrated the

binding curves to be identical to that under standard assay conditions containing ATP and  $\text{MgCl}_2$  (Figure 3C), suggesting that ATP binding itself affected the interaction between ssDNA and the helicase domain.

#### ssDNA binding activity of the PIF1N

Our results suggested that the PINT domain plays a role in modulating the ssDNA binding activity of the helicase domain. Therefore, we carried out the same binding assays with PIF1N (Figure 4A). The assays detected PIF1N-DNA complexes, and the titration curves were not affected by ATP and  $\text{MgCl}_2$  (Figure 4A), with an apparent  $K_d$  value of about 10 nM. These results suggested that PIF1N itself possesses ATP-independent ssDNA binding activity.

In contrast to the binding reactions with PIF1 and PIF1C shown in Figure 3, further shifts of the mobility



**Figure 4.** EMSA of PIF1N for binding to ssDNA. (A) Titration of PIF1N under different conditions. The 5'-<sup>32</sup>P labeled oligonucleotide, d(AC)60, was incubated with the shown concentrations of PIF1N, omitting ATP or  $\text{MgCl}_2$  as indicated. Arrowheads indicate the positions of free DNA. The quantified data are shown graphically. (B) Binding ability of PIF1N to different sizes of oligonucleotides. Experiments were performed using indicated oligonucleotides as substrates under standard reaction conditions in the absence of ATP. The quantified data are shown graphically. The errors in the experiments were <10%.

of the complexes with PIF1N in higher concentrations were observed (Figure 4A). This could be due to more than one protein molecule binding to the 60-mer oligonucleotide. To determine the minimal ssDNA size for binding of PIF1N, six different oligonucleotides in lengths ranging from 55 to 15 bases were subjected to binding assays. When the 55-mer was used as the substrate, the result was essentially identical to that for a 60-mer. However, further reduction of the length to 45-mer and 35-mer decreased the extent of multiple binding. With the 30-mer oligonucleotide, but not the 20-mer and 15-mer, a distinct band of the complex was observed. These results demonstrated that the minimal ssDNA size could be between 30 and 20 bases.

#### DNA strand annealing activity of PIF1, residing in the PINT domain

During helicase assays, we unexpectedly found that PIF1 possessed robust annealing activity. For detailed analysis, a forked substrate (1F:2L) (Figure 5A) was denatured by heating at 100°C for 5 min, and then incubated with PIF1 under the conditions for the standard ATPase assay omitting ATP to avoid unwinding reactions. Titration of PIF1 showed about 50% of ssDNA could be annealed within 10 min when 46 nM PIF1 was present (Figure 5B). To determine the region of PIF1 responsible for the annealing activity, PIF1C and PIF1N were tested for their ability to promote the reaction. The results demonstrated that PIF1N, but not PIF1C, efficiently annealed ssDNA, with activity only 3-fold lower than that of PIF1 when 1F and 2L substrates were tested (Figure 5B). This result indicated that the annealing activity of PIF1 resides in the PINT domain.

Since ATP is essential for helicase activity, we tested effects of ATP on the annealing reaction. To avoid the effect of unwinding, titration of ATP was carried out in the absence of MgCl<sub>2</sub>. The result demonstrated both PIF1 and PIF1N to be inhibited by ATP (Figure 5C). Time course experiments demonstrated that, in the absence of ATP, PIF1 annealed up to 70% of ssDNA, whereas, in the presence of the optimal concentration of ATP (2 mM) for helicase activity, the annealed fraction reached only 20% (Figure 5D). The properties of PIF1 were same as those of PIF1N, 2 mM ATP also inhibiting annealing activity from 50% to 17% (Figure 5E). Importantly, the ATP-titration curve and time course with PIF1 were essentially identical to those for PIF1N (Figure 5C–E), suggesting that the properties are intrinsic to the PINT domain.

Further confirming this conclusion, we used completely complementary strands as a substrate. The annealing product has no ssDNA region, preventing product unwinding under the standard reaction conditions containing ATP and MgCl<sub>2</sub>. Titration curves of ATP for PIF1 and PIF1N demonstrated again inhibitory effects, although significant fractions of the substrates were spontaneously annealed under the assay conditions (Figure 5F).

#### Effects of RPA on unwinding and annealing reactions of PIF1

Elucidation of whether PIF1 has the potential to promote unwinding and annealing reactions on RPA-coated

substrates is valuable for understanding cellular functions of PIF1. To do this, first, we determined the optimal concentration of RPA for the substrate DNA binding by EMSA. The substrate, 3F:4L was used for the binding reactions at the final concentration of 0.35 nM. The results of titration of RPA are shown in Figure 6A. When the concentration of RPA was increased to 0.9 nM, one molecule of RPA bound to at least one strand of the forked substrate. At >1.8 nM, each strand of the substrate was probably occupied by one RPA molecule (Figure 6A). The levels of apparent affinity of RPA for ssDNA were in good agreement with previous reports (38).


Then we examined unwinding activity of PIF1. RPA was mixed with the forked substrate, 3F:4L, under the standard reaction conditions on ice, then PIF1 was introduced and incubation was performed at 30°C for 10 min. We found that RPA did not affect unwinding reactions at low concentrations <0.4 nM (Figure 6B, left panel), in which majority of the substrate was RPA-free (Figure 6A). When the concentration of RPA reached 0.9 nM, at which almost all the oligonucleotides were occupied with RPA (Figure 6A), severe inhibition was observed (Figure 6B, left panel). At much higher concentrations of RPA, we observed unwinding products (Figure 6B, left panel). However, a control experiment without PIF1 revealed that the products detected at higher concentrations >1.8 nM of RPA were PIF1 independent (Figure 6B, right panel). The quantified results shown in Figure 6C demonstrate no difference in the two reactions at higher concentrations of RPA (>0.9 nM). These results suggest that RPA does not enhance, but rather inhibits, the helicase activity of PIF1.

To examine annealing activity in the presence of RPA, we also determined the optimal concentration of RPA for ssDNA binding. In the reactions, we used the same substrate as for the unwinding assay (3F:4L) at the final concentration of 0.35 nM, but after denaturation by heating. The assay detected only complexes with the labeled oligonucleotide 3F (55-mer), although another fragment, 4L (51-mer), was present. The results of titration of RPA are shown in Figure 6D. When the concentration of RPA was increased to 0.9 nM, almost all the oligonucleotides 3F and probably 4L were occupied by at least one molecule of RPA. At >1.8 nM, the oligonucleotides were occupied by two molecules of RPA (Figure 6D).

Next, we carried out annealing assays under the same conditions. RPA was mixed with heat denatured substrate (3F and 4L) on ice, then PIF1 was introduced with incubation at 30°C for 10 min. The result clearly demonstrated an inhibitory effect of RPA (Figure 6E). The quantified results for inhibition of annealing reactions well correlated inversely with the amount of RPA binding (Figure 6F).

## DISCUSSION

In this article, we document for the first time the biochemical properties of full-length human PIF1 together with those of truncated forms consisting of individual domains. We could establish intrinsic properties of the helicase domain and functional roles of the PINT domain.



**Figure 5.** DNA strand annealing activity. (A) Schematic structures of the DNA substrates. The asterisks indicate  $^{32}\text{P}$ -labeled 5'-phosphate. (B) Annealing activities of PIF1, PIF1C and PIF1N. The fork-structural partial duplex DNA substrate, 1F:2L (0.35 nM), was boiled for 5 min, then incubated with increasing levels of PIF1, PIF1C and PIF1N as indicated under standard reaction conditions omitting ATP at 30°C for 10 min. The quantified data are shown graphically. (C) Titration of ATP on annealing reactions mediated by PIF1 and PIF1N. The fork-structural partial duplex DNA substrate, 3F:4L (0.35 nM), was boiled for 5 min, then incubated with PIF1 (33 nM) or PIF1N (195 nM) and increasing levels of ATP under standard reaction conditions omitting  $\text{MgCl}_2$  at 30°C for 10 min. The quantified data are shown graphically. (D) Time course of annealing reactions mediated by PIF1 under standard reaction conditions omitting  $\text{MgCl}_2$ . Heat denatured substrate DNA, 1F:2L (0.35 nM), was incubated with PIF1 (46 nM). The quantified data are shown graphically. (E) Time course of annealing reactions mediated by PIF1N under standard reaction conditions omitting  $\text{MgCl}_2$ . Heat denatured substrate DNA, 1F:2L (0.35 nM), was incubated with PIF1N (195 nM). The quantified data are shown graphically. (F) Titration of ATP on annealing reactions mediated by PIF1 and PIF1N under standard reaction conditions. A complete double stranded oligonucleotide, 3S:4S (0.35 nM) was boiled for 5 min, then incubated with PIF1 (30 nM) or PIF1N (195 nM) and increasing levels of ATP under standard reaction conditions at 30°C for 10 min. The quantified data are shown graphically. The errors in the experiments were <10%.





Figure 5. Continued.



**Figure 6.** Effects of RPA on the unwinding and annealing reactions. (A) EMSA of RPA binding. The fork-structural partial duplex DNA substrate, 3F:4L (0.35 nM) was incubated with RPA at indicated concentrations on ice for 10 min under the standard reaction conditions. (B) Unwinding assay in the presence of RPA. The fork-structural partial duplex DNA substrate, 3F:4L (0.35 nM), was incubated with RPA at indicated concentrations on ice for 10 min under standard reaction conditions. Then, PIF1 (16 nM) (left panel) and buffer (right panel) were introduced and incubated at 30°C for 10 min. Reaction products were separated on a 15–25% polyacrylamide gel. (C) The quantified data of unwinding reactions in (B) are shown graphically with the RPA binding curve in (A). (D) EMSA of RPA binding. The fork-structural partial duplex DNA substrate, 3F:4L (0.35 nM), was boiled for 5 min, then incubated with RPA at indicated concentrations on ice for 10 min under the standard reaction conditions omitting ATP. (E) Annealing assay in the presence of RPA. The fork-structural partial duplex DNA substrate, 3F:4L (0.35 nM) (Figure 5A), was boiled for 5 min, then incubated with RPA at indicated concentrations on ice for 10 min under standard reaction conditions omitting ATP. Then PIF1 (30 nM) was introduced and incubated at 30°C for 10 min. Reaction products were separated on a 15–25% polyacrylamide gel. (F) The quantified data of annealing reactions in (E) are shown graphically with the RPA binding curve in (D). The errors in the experiments were <10%.

Biochemical analysis of yeast PIF1 homologs has demonstrated that they preferentially unwind forked substrates (1,5,7). We showed that the property is conserved in human PIF1. However, we found that forked substrates were not optimal with respect to stimulation of ATPase activity. Rather nonstructural ssDNA greatly stimulated ATPase activity. The finding that the  $K_{\text{eff}}$  value for non-structural ssDNA was lower than for other DNA molecules, including forked structures, suggested preferential binding of PIF1 to ssDNA. From these results, we suggest that PIF1 needs an ssDNA region for loading and a forked structure for entrance to the double strand region by translocation.

We present several lines of evidence that the enzymatic characters of PIF1C reflect the intrinsic properties of the helicase domain. First, PIF1C expressed ATPase activity to the level equivalent to full-length PIF1 (about  $1000 \text{ min}^{-1}$ ) and also equivalent to that for yeast homologs (5,7). Second, full-length PIF1 and PIF1C both showed a similar preference for poly(purine–pyrimidine) and polypyrimidine, but not polypurine, for stimulation of ATPase. This property is also conserved in yeast homologs (5,7). Third, the  $K_m$  values for ATP of full-length PIF1 and PIF1C were essentially identical. This is also in agreement with a previous report for N-terminal truncated PIF1, purified as a C-terminus GST-fusion protein (4).

These results suggest that the determined properties of PIF1C are intrinsic to the helicase domain, and could also exclude the possibility that the his-tag at the N-terminal of PIF1C interferes with activities of the helicase domain.

Interestingly, we noted a significant difference between PIF1 and PIF1C with regard to the required concentration of ssDNA for stimulation of ATPase activity. PIF1C needed a 20 times higher concentration and also exhibited much lower unwinding activity. The results suggest that the defects could be attributed to missing functions of the PINT domain. The difference in the  $K_{\text{eff}}$  values could be due to lower binding affinity of the helicase domain for ssDNA. Consequently, we demonstrated ssDNA binding of PIF1 directly by EMSA. The apparent  $K_d$  value, 3 nM was in good agreement with the  $K_{\text{eff}}$  value of 2 nM when expressed with reference to the oligonucleotide concentration, suggesting that this assay well reflected the functional interaction between ssDNA and PIF1. In this assay, as expected, we demonstrated lower affinity of PIF1C to ssDNA. We suggested that the defect in PIF1C could be due, at least in part, to lower binding affinity for ssDNA, and the PINT domain plays a role for increasing this affinity of the helicase domain. Notably, the  $K_d$  value (10 nM) of PIF1C determined by EMSA was still 6 times lower than the  $K_{\text{eff}}$  value (60 nM in oligonucleotides) for the ATPase assay. We consider the following possible explanations for the discrepancy. We found that PIF1C exhibited a much higher affinity for ssDNA without binding of ATP. With the EMSA, an ATP-free fraction of PIF1C could exist, even in the presence of ATP. Therefore, the results could be an overestimation, due to high affinity binding of the ATP-free fraction of PIF1C. Alternatively, the PINT domain could possess another function for enhancing activity of the helicase domain by modulating the mode of interaction with ssDNA. The higher affinity of PIF1C for ssDNA without binding of ATP could be intrinsic to the helicase domain. The results suggested that ATP modulated the binding mode with ssDNA. The higher affinity to ssDNA before binding of ATP is reduced by binding of ATP. The alteration must be alternatively repeated during turnover of reactions of ATPase. With the full-length PIF1, such alteration due to ATP binding was not detected, suggesting that the PINT domain somehow could suppress the alteration during turnover of ATPase reactions. In this study, we demonstrated that the PINT domain also possesses ssDNA binding activity. We suggest that its enhancement of the activities of the helicase domain is due, at least in part, to this ssDNA binding activity.

We unexpectedly found the PINT domain to further possess ssDNA annealing activity. Among the proteins handled in this study having ssDNA binding activity, including PIF1C, PIF1N and RPA, we could detect annealing activity only with PIF1N. We consider that the annealing activity could be mediated by ssDNA binding, although not attributable to the general effect of high affinity ssDNA binding. Annealing activity has been reported to be associated with the RecQ family helicase in general (39–45). While the RecQ family is distinct from PIF1 family, annealing activity shares similar properties

in common. It is located outside of the conserved helicase domains (39,40), is inhibited by RPA, and is ATP-independent or rather inhibited by ATP (39–42,44). We demonstrated that inhibition by ATP is not a consequence of the unwinding reaction, suggesting that it is an intrinsic property of the PINT domain.

At the present time, the biological significance of our findings cannot be readily assessed. Importantly, we showed that RPA inhibited unwinding and annealing reactions, suggesting that these functions of PIF1 might be restricted under particular situations in DNA metabolism. There is a marked difference from RecQ helicases, whose unwinding activity is proficient on RPA-coated ssDNA and stimulated by RPA, although annealing activity is suppressed (39–41,46–48). Notably, unwinding activity of *ScPif1* was stimulated by RPA (1), but that of the fission yeast homolog, *Pfh1*, was inhibited (7) like human PIF1, suggesting that the outcomes in cells would differ. This could be related to the fact that budding yeast has another member of the PIF1 superfamily, *Rrm3*, but human and fission yeast have only one. Further analysis of the precise cellular roles of PIF1 should shed light on functions in maintenance of genetic stability.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

## ACKNOWLEDGEMENTS

We thank Dr Marc S. Wold (University of Iowa College of Medicine, Iowa City, Iowa, USA), Dr Hisaji Maki (Nara Institute of Science and Technology, Nara, Japan) and Dr Tadashi Shimamoto (Hiroshima University, Hiroshima, Japan) for, respectively, providing an RPA-expression plasmid, the M13 mp7 plasmid and the *E. coli* strain to produce ss M13 DNA. Several cloning vectors were obtained from National BioResource Project (NIG, Japan). We are grateful to Kumiko Mizuno, Tomoka Nakashima, Masako Okii, Fumie Okubo, Kazumi Shimamoto, Hatsue Wakayama and Mai Yoshida for their laboratory assistance.

## FUNDING

Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to Y.M. and K.K.); 21st Century Center of Excellence Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to K.K.); Health and Labour Science Research Grants (to K.K.); Grants-in-Aid for Cancer Research from the Ministry of Health, Labour and Welfare (to K.K.). Funding to pay the Open Access publication charges for this article was provided by Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to Y.M.).

*Conflict of interest statement.* None declared.

## REFERENCES

- Boulé, J.B. and Zakian, V.A. (2007) The yeast Pif1p DNA helicase preferentially unwinds RNA DNA substrates. *Nucleic Acids Res.*, **35**, 5809–5818.
- Foury, F. and Lahaye, A. (1987) Cloning and sequencing of the PIF gene involved in repair and recombination of yeast mitochondrial DNA. *EMBO J.*, **6**, 1441–1449.
- Futami, K., Shimamoto, A. and Furuichi, Y. (2007) Mitochondrial and nuclear localization of human Pif1 helicase. *Biol. Pharm. Bull.*, **30**, 1685–1692.
- Huang, Y., Zhang, D.H. and Zhou, J.Q. (2006) Characterization of ATPase activity of recombinant human Pif1. *Acta. Biochim. Biophys. Sin. (Shanghai)*, **38**, 335–341.
- Lahaye, A., Leterme, S. and Foury, F. (1993) PIF1 DNA helicase from *Saccharomyces cerevisiae*. Biochemical characterization of the enzyme. *J. Biol. Chem.*, **268**, 26155–26161.
- Lahaye, A., Stahl, H., Thines-Sempoux, D. and Foury, F. (1991) PIF1: a DNA helicase in yeast mitochondria. *EMBO J.*, **10**, 997–1007.
- Ryu, G.H., Tanaka, H., Kim, D.H., Kim, J.H., Bae, S.H., Kwon, Y.N., Rhee, J.S., MacNeill, S.A. and Seo, Y.S. (2004) Genetic and biochemical analyses of Pif1 DNA helicase function in fission yeast. *Nucleic Acids Res.*, **32**, 4205–4216.
- Tanaka, H., Ryu, G.H., Seo, Y.S., Tanaka, K., Okayama, H., MacNeill, S.A. and Yuasa, Y. (2002) The fission yeast *pif1+* gene encodes an essential 5' to 3' DNA helicase required for the completion of S-phase. *Nucleic Acids Res.*, **30**, 4728–4739.
- Zhang, D.H., Zhou, B., Huang, Y., Xu, L.X. and Zhou, J.Q. (2006) The human Pif1 helicase, a potential *Escherichia coli* RecD homologue, inhibits telomerase activity. *Nucleic Acids Res.*, **34**, 1393–1404.
- Zhou, J.Q., Qi, H., Schulz, V.P., Mateyak, M.K., Monson, E.K. and Zakian, V.A. (2002) *Schizosaccharomyces pombe pif1+* encodes an essential 5' to 3' DNA helicase that is a member of the PIF1 subfamily of DNA helicases. *Mol. Biol. Cell.*, **13**, 2180–2191.
- Bessler, J.B., Torredagger, J.Z. and Zakian, V.A. (2001) The Pif1p subfamily of helicases: region-specific DNA helicases? *Trends Cell Biol.*, **11**, 60–65.
- Foury, F. and Dyck, E.V. (1985) A PIF-dependent recombinogenic signal in the mitochondrial DNA of yeast. *EMBO J.*, **4**, 3525–3530.
- Foury, F. and Kolodny, J. (1983) *pif* mutation blocks recombination between mitochondrial  $\rho^+$  and  $\rho^-$  genomes having tandemly arrayed repeat units in *Saccharomyces cerevisiae*. *Proc. Natl Acad. Sci. USA*, **80**, 5345–5349.
- Doudican, N.A., Song, B., Shadel, G.S. and Doetsch, P.W. (2005) Oxidative DNA damage causes mitochondrial genomic instability in *Saccharomyces cerevisiae*. *Mol. Cell Biol.*, **25**, 5196–5204.
- O'Rourke, T.W., Doudican, N.A., Mackereth, M.D., Doetsch, P.W. and Shadel, G.S. (2002) Mitochondrial dysfunction due to oxidative mitochondrial DNA damage is reduced through cooperative actions of diverse proteins. *Mol. Cell Biol.*, **22**, 4086–4093.
- O'Rourke, T.W., Doudican, N.A., Zhang, H., Eaton, J.S., Doetsch, P.W. and Shadel, G.S. (2005) Differential involvement of the related DNA helicases Pif1p and Rrm3p in mtDNA point mutagenesis and stability. *Gene*, **354**, 86–92.
- Cheng, X., Dunaway, S. and Ivessa, A.S. (2007) The role of Pif1p, a DNA helicase in *Saccharomyces cerevisiae*, in maintaining mitochondrial DNA. *Mitochondrion*, **7**, 211–222.
- Zhou, J., Monson, E.K., Teng, S.C., Schulz, V.P. and Zakian, V.A. (2000) Pif1p helicase, a catalytic inhibitor of telomerase in yeast. *Science*, **289**, 771–774.
- Schulz, V.P. and Zakian, V.A. (1994) The *Saccharomyces PIF1* DNA helicase inhibits telomere elongation and *de novo* telomere formation. *Cell*, **76**, 145–155.
- Kanaar, R., Wyman, C. and Rothstein, R. (2008) Quality control of DNA break metabolism: in the 'end', it's a good thing. *EMBO J.*, **27**, 581–588.
- Mangahas, J.L., Alexander, M.K., Sandell, L.L. and Zakian, V.A. (2001) Repair of chromosome ends after telomere loss in *Saccharomyces*. *Mol. Biol. Cell.*, **12**, 4078–4089.
- Mateyak, M.K. and Zakian, V.A. (2006) Human PIF helicase is cell cycle regulated and associates with telomerase. *Cell Cycle*, **5**, 2796–2804.
- Myung, K., Chen, C. and Kolodner, R.D. (2001) Multiple pathways cooperate in the suppression of genome instability in *Saccharomyces cerevisiae*. *Nature*, **411**, 1073–1076.
- Pennaneach, V., Putnam, C.D. and Kolodner, R.D. (2006) Chromosome healing by *de novo* telomere addition in *Saccharomyces cerevisiae*. *Mol. Microbiol.*, **59**, 1357–1368.
- Boulé, J.B., Vega, L.R. and Zakian, V.A. (2005) The yeast Pif1p helicase removes telomerase from telomeric DNA. *Nature*, **438**, 57–61.
- Budd, M.E., Reis, C.C., Smith, S., Myung, K. and Campbell, J.L. (2006) Evidence suggesting that Pif1 helicase functions in DNA replication with the Dna2 helicase/nuclease and DNA polymerase  $\delta$ . *Mol. Cell Biol.*, **26**, 2490–2500.
- Ivessa, A.S., Zhou, J.Q. and Zakian, V.A. (2000) The *Saccharomyces* Pif1p DNA helicase and the highly related Rrm3p have opposite effects on replication fork progression in ribosomal DNA. *Cell*, **100**, 479–489.
- Wagner, M., Price, G. and Rothstein, R. (2006) The absence of Top3 reveals an interaction between the Sgs1 and Pif1 DNA helicases in *Saccharomyces cerevisiae*. *Genetics*, **174**, 555–573.
- Masuda, Y., Suzuki, M., Piao, J., Gu, Y., Tsurimoto, T. and Kamiya, K. (2007) Dynamics of human replication factors in the elongation phase of DNA replication. *Nucleic Acids Res.*, **35**, 6904–6916.
- Henriksen, L.A., Umbricht, C.B. and Wold, M.S. (1994) Recombinant replication protein A: expression, complex formation, and functional characterization. *J. Biol. Chem.*, **269**, 11121–11132.
- Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.*, **185**, 60–89.
- Snaitch, M.R., Kilby, N.J. and Murray, J.A. (1996) An *Escherichia coli* system for assay of F1p site-specific recombination on substrate plasmids. *Gene*, **180**, 225–227.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Masuda, Y., Takahashi, M., Tsunekuni, N., Minami, T., Sumii, M., Miyagawa, K. and Kamiya, K. (2001) Deoxycytidyl transferase activity of the human REV1 protein is closely associated with the conserved polymerase domain. *J. Biol. Chem.*, **276**, 15051–15058.
- Carey, J. (1991) Gel retardation. *Methods Enzymol.*, **208**, 103–117.
- Kornberg, A., Scott, J.F. and Bertsch, L.L. (1978) ATP utilization by rep protein in the catalytic separation of DNA strands at a replicating fork. *J. Biol. Chem.*, **253**, 3298–3304.
- Matson, S.W. and George, J.W. (1987) DNA helicase II of *Escherichia coli*. Characterization of the single-stranded DNA-dependent NTPase and helicase activities. *J. Biol. Chem.*, **262**, 2066–2076.
- Wold, M.S. (1997) Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu. Rev. Biochem.*, **66**, 61–92.
- Cheok, C.F., Wu, L., Garcia, P.L., Janscak, P. and Hickson, I.D. (2005) The Bloom's syndrome helicase promotes the annealing of complementary single-stranded DNA. *Nucleic Acids Res.*, **33**, 3932–3941.
- Garcia, P.L., Liu, Y., Jiricny, J., West, S.C. and Janscak, P. (2004) Human RECQ5 $\beta$ , a protein with DNA helicase and strand-annealing activities in a single polypeptide. *EMBO J.*, **23**, 2882–2891.
- Kanagaraj, R., Saydam, N., Garcia, P.L., Zheng, L. and Janscak, P. (2006) Human RECQ5 $\beta$  helicase promotes strand exchange on synthetic DNA structures resembling a stalled replication fork. *Nucleic Acids Res.*, **34**, 5217–5231.
- Machwe, A., Xiao, L., Groden, J., Matson, S.W. and Orren, D.K. (2005) RecQ family members combine strand pairing and unwinding activities to catalyze strand exchange. *J. Biol. Chem.*, **280**, 23397–23407.
- Macris, M.A., Krejci, L., Bussen, W., Shimamoto, A. and Sung, P. (2006) Biochemical characterization of the RECQ4 protein, mutated in Rothmund-Thomson syndrome. *DNA Repair (Amst.)*, **5**, 172–180.

44. Sharma,S., Sommers,J.A., Choudhary,S., Faulkner,J.K., Cui,S., Andreoli,L., Muzzolini,L., Vindigni,A. and Brosh,R.M. Jr (2005) Biochemical analysis of the DNA unwinding and strand annealing activities catalyzed by human RECQ1. *J. Biol. Chem.*, **280**, 28072–28084.
45. Machwe,A., Lozada,E.M., Xiao,L. and Orren,D.K. (2006) Competition between the DNA unwinding and strand pairing activities of the Werner and Bloom syndrome proteins. *BMC Mol. Biol.*, **7**, 1.
46. Brosh,R.M. Jr, Orren,D.K., Nehlin,J.O., Ravn,P.H., Kenny,M.K., Machwe,A. and Bohr,V.A. (1999) Functional and physical interaction between WRN helicase and human replication protein A. *J. Biol. Chem.*, **274**, 18341–18350.
47. Brosh,R.M. Jr, Li,J.L., Kenny,M.K., Karow,J.K., Cooper,M.P., Kurekattil,R.P., Hickson,I.D. and Bohr,V.A. (2000) Replication protein A physically interacts with the Bloom's syndrome protein and stimulates its helicase activity. *J. Biol. Chem.*, **275**, 23500–23508.
48. Cui,S., Arosio,D., Doherty,K.M., Brosh,R.M. Jr, Falaschi,A. and Vindigni,A. (2004) Analysis of the unwinding activity of the dimeric RECQ1 helicase in the presence of human replication protein A. *Nucleic Acids Res.*, **32**, 2158–2170.