

- 4 Jemec B. Studies of the goitrogenic and tumorigenic effect of two goitrogens in combination with hypophysectomy or thyroid hormone treatment. *Cancer* 1980; **45**: 2138–48.
- 5 Hiasa Y, Ohshima M, Kitabori Y, Yuasa T, Fujita T, Iwata C. Promoting effects of 3-amino-1,2,4-triazole on the development of thyroid tumors in rats treated with *N*-bis (2-hydroxypropyl) nitrosamine. *Carcinogenesis* 1982; **3**: 381–4.
- 6 Hard GC. Recent developments in the investigation of thyroid regulation and thyroid carcinogenesis. *Environ Health Perspect* 1998; **106**: 427–36.
- 7 Smith P, Williams ED, Wynford-Thomas D. *In vitro* demonstration of a TSH-specific growth desensitizing mechanism in rat thyroid epithelium. *Mol Cell Endocrinol* 1987; **51**: 51–8.
- 8 Brewer C, Yeager N, Di Cristofano A. Thyroid-stimulating hormone initiated proliferative signals converge *in vivo* on the mTOR kinase without activating AKT. *Cancer Res* 2007; **67**: 8002–6.
- 9 Mitsumori K, Onodera H, Takahashi M *et al*. Effect of thyroid stimulating hormone on the development and progression of rat thyroid follicular cell tumors. *Cancer Lett* 1995; **92**: 193–202.
- 10 Roy G, Mugesh G. Bioinorganic chemistry in thyroid gland: effect of antithyroid drugs on peroxidase-catalyzed oxidation and iodination reactions. *Bioinorg Chem Appl* 2006; **2006**: 23214.
- 11 Shibutani M, Uneyama C, Miyazaki K, Toyoda K, Hirose M. Methacarn fixation: a novel tool for analysis of gene expressions in paraffin-embedded tissue specimens. *Lab Invest* 2000; **80**: 199–208.
- 12 Uneyama C, Shibutani M, Masutomi N, Takagi H, Hirose M. Methacarn fixation for genomic DNA analysis in microdissected, paraffin-embedded tissue specimens. *J Histochim Cytochem* 2002; **50**: 1237–45.
- 13 Takagi H, Shibutani M, Kato N *et al*. Microdissected region-specific gene expression analysis with methacarn-fixed, paraffin-embedded tissues by real-time RT-PCR. *J Histochim Cytochem* 2004; **S2**: 903–13.
- 14 Takagi H, Shibutani M, Lee KY *et al*. Impact of maternal dietary exposure to endocrine-active chemicals on progesterone receptor expression in microdissected hypothalamic medial preoptic areas of rat offspring. *Toxicol Appl Pharmacol* 2005; **208**: 127–36.
- 15 Shibutani M, Lee KY, Igashira K *et al*. Hypothalamus region-specific global gene expression profiling in early stages of central endocrine disruption in rat neonates injected with estradiol benzoate or flutamide. *Dev Neurobiol* 2007; **67**: 253–69.
- 16 Lee KY, Shibutani M, Inoue K, Kuroiwa KUM, Woo GH, Hirose M. Methacarn fixation – effects of tissue processing and storage conditions on detection of mRNAs and proteins in paraffin-embedded tissues. *Anal Biochem* 2006; **351**: 36–43.
- 17 Hardisty JF, Boorman GA. Thyroid gland. In: Boorman GA, Eustis SL, Ellwell MR, Montgomery CA Jr, MacKenzie WF, eds. *Pathology of the Fischer Rat. Reference and Atlas*. San Diego: Academic Press, 1990: 519–36.
- 18 Imai T, Onose J, Hasumura M, Ueda M, Takizawa T, Hirose M. Sequential analysis of development of invasive thyroid follicular cell carcinomas in inflamed capsular regions of rats treated with sulfadimethoxine after *N*-bis (2-hydroxypropyl) nitrosamine-initiation. *Toxicol Pathol* 2004; **32**: 229–36.
- 19 Goldstein IM, Kaplan HB, Edelson HS, Weissmann G. Ceruloplasmin: A scavenger of superoxide anion radicals. *J Biol Chem* 1979; **254**: 4040–5.
- 20 Lin JD. Thyroglobulin and human thyroid cancer. *Clin Chim Acta* 2008; **388**: 15–21.
- 21 Zlak M, Meier M, Novak-Hofer I, Roth J. Ceruloplasmin carries the anionic glycan oligo/poly o2,8 deaminonurameric acid. *Biochem Biophys Res Commun* 2002; **295**: 597–602.
- 22 Scott IS, Morris LS, Bird K *et al*. A novel immunohistochemical method to estimate cell-cycle phase distribution in archival tissue: implications for the prediction of outcome in colorectal cancer. *J Pathol* 2003; **201**: 187–97.
- 23 Yamamoto H, Monden T, Ikeda K *et al*. Coexpression of cdk2/cdc2 and retinoblastoma gene products in colorectal cancer. *Br J Cancer* 1995; **71**: 1231–6.
- 24 Faggiano A, Caillou B, Lacroix L *et al*. Functional characterization of human thyroid tissue with immunohistochemistry. *Thyroid* 2007; **17**: 203–11.
- 25 Kuramitsu K, Ikeda W, Inoue N, Tamaru Y, Takai Y. Novel role of neitin: implication in the co-localization of JAM-A and claudin-1 at the same cell-cell adhesion membrane domain. *Genes Cells* 2008; **13**: 797–805.
- 26 Carroll M, Hamzeh M, Robaire B. Expression, localization, and regulation of inhibitor of DNA binding (Id) proteins in the rat epididymis. *J Androl* 2006; **27**: 212–24.
- 27 Leikin PL, Hahn KL, Sukovich DA, Karmiol S, Schmidt DH. On the possible role of reactive oxygen species in angiogenesis. *Adv Exp Med Biol* 1998; **454**: 295–310.
- 28 Sub YA, Arnold RS, Lassegue B *et al*. Cell transformation by the superoxide-generating oxidase Mox1. *Nature* 1999; **401**: 79–82.
- 29 Stassar MJ, Devitt G, Brosius M *et al*. Identification of human renal cell carcinoma associated genes by suppression subtractive hybridization. *Br J Cancer* 2001; **85**: 1372–82.
- 30 Hough CD, Cho KR, Zonderman AB, Schwartz DR, Morin PJ. Coordinate up-regulated genes in ovarian cancer. *Cancer Res* 2001; **61**: 3869–76.
- 31 Wang KK, Liu N, Radulovic N *et al*. Novel candidate tumor marker genes for lung adenocarcinoma. *Oncogene* 2002; **21**: 7598–604.
- 32 Tuccari G, Barresi G. Immunohistochemical demonstration of ceruloplasmin in follicular adenomas and thyroid carcinomas. *Histopathology* 1987; **11**: 723–31.
- 33 Song B. Immunohistochemical demonstration of epidermal growth factor receptor and ceruloplasmin in thyroid diseases. *Acta Pathol Jpn* 1991; **41**: 336–43.
- 34 Kondi-Pafiti A, Smyniotis V, Frangou M, Papayannopoulos A, Englezou M, Deligeorgi H. Immunohistochemical study of ceruloplasmin, lactoferrin and secretory component expression in neoplastic and non-neoplastic thyroid gland diseases. *Acta Oncol* 2000; **39**: 753–6.
- 35 Draetta G, Luca F, Westendorf J, Brizuela L, Ruderman J, Beach D. cdc2 protein kinase is complexed with both cyclin A and B. Evidence for proteolytic inactivation of MPF. *Cell* 1989; **56**: 829–38.
- 36 Chen H, Huang Q, Dong J, Zhai DZ, Wang AD, Lan Q. Overexpression of CDC2/CyclinB1 in gliomas, and CDC2 depletion inhibits proliferation of human glioma cells *in vitro* and *in vivo*. *BMC Cancer* 2008; **8**: 29.
- 37 Wiseman SM, Griffith OL, Deen S *et al*. Identification of molecular markers altered during transformation of differentiated into anaplastic thyroid carcinoma. *Arch Surg* 2007; **142**: 717–29.
- 38 Simmons DL. Dissecting the modes of interactions amongst cell adhesion molecules. *Dev Suppl* 1993; **3**: 193–203.
- 39 Geraghty RJ, Krummenacher C, Cohen GH, Eisenberg RJ, Spear PG. Entry of alphaherpesviruses mediated by poliovirus receptor-related protein 1 and poliovirus receptor. *Science* 1998; **280**: 1618–20.
- 40 Nelson GM, Padra TP, Garkavtsev I, Shiota T, Jain RK. Differential gene expression of primary cultured lymphatic and blood vascular endothelial cells. *Neoplasia* 2007; **9**: 1038–45.
- 41 Takai Y, Miyoshi J, Ikeda W, Ogita H. Neitin and neitin-like molecules: roles in contact inhibition of cell movement and proliferation. *Nat Rev Mol Cell Biol* 2008; **9**: 603–15.
- 42 Murakami Y. Involvement of a cell adhesion molecule, TSLC1/IGSF4, in human oncogenesis. *Cancer Sci* 2005; **96**: 543–52.
- 43 Wakayama T, Sai Y, Ito A *et al*. Heterophilic binding of the adhesion molecules poliovirus receptor and immunoglobulin superfamily 4A in the interaction between mouse spermatogenic and Sertoli cells. *Biol Reprod* 2007; **76**: 1081–90.
- 44 Ben Ezra R, Davis RL, Lockshon D, Turner DL, Weintraub H. The protein Id: A negative regulator of helix-loop-helix DNA binding proteins. *Cell* 1990; **61**: 49–59.
- 45 Ellis HM, Spann DR, Posakony JW. *Extramacrociliae*, a negative regulator of sensory organ development in *Drosophila*, defines a new class of helix-loop-helix proteins. *Cell* 1990; **61**: 27–38.
- 46 Sikder HA, Devlin MK, Dunlap S, Ryu B, Alani RM. Id proteins in cell growth and tumorigenesis. *Cancer Cell* 2003; **3**: 525–30.
- 47 Arnold JM, Mok SC, Purdie D, Chenevix-Trench G. Decreased expression of the *Id3* gene at 1p36.1 in ovarian adenocarcinomas. *Br J Cancer* 2001; **84**: 352–9.
- 48 Nikolova DN, Zembutsu H, Sechanov T *et al*. Genome-wide gene expression profiles of thyroid carcinoma: Identification of molecular targets for treatment of thyroid carcinoma. *Oncol Rep* 2008; **20**: 105–21.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. List of genes showing altered expression in microdissected focal follicular cell hyperplasias (FFCH) + adeomas (Ad) induced in the thyroids of rats using a two-stage thyroid carcinogenesis model (≥ 2 -fold, ≤ 0.5 -fold).

Table S2. List of genes showing altered expression in microdissected carcinomas (Ca) induced in the thyroid of rats using a two-stage thyroid carcinogenesis model (≥ 2 -fold, ≤ 0.5 -fold).

Table S3. List of genes showing altered expression in common in all types of thyroid proliferative lesion induced in rats using a two-stage thyroid carcinogenesis model (≥ 2 -fold, ≤ 0.5 -fold).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.