

17. Honda, K., H. Nakano, H. Yoshida, S. Nishikawa, P. Rennert, K. Ikuta, M. Tamechika, K. Yamaguchi, T. Fukumoto, T. Chiba, and S. I. Nishikawa. 2001. Molecular basis for hematopoietic/mesenchymal interaction during initiation of Peyer's patch organogenesis. *J. Exp. Med.* 193: 621–630.
18. Finke, D., H. Acha-Orbea, A. Mattis, M. Lipp, and J. Krahenbuhl. 2002. CD4+CD3+ cells induce Peyer's patch development: role of $\alpha_4\beta_1$ integrin activation by CXCR5. *Immunity* 17: 363–373.
19. Weil, F., and J. Caamano. 2003. Regulation of secondary lymphoid organ development by the nuclear factor- κ B signal transduction pathway. *Immunol. Rev.* 195: 91–105.
20. Yokota, Y., A. Mansouri, S. Mori, S. Sugawara, S. Adachi, S. Nishikawa, and P. Gruss. 1999. Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature* 397: 702–706.
21. Sun, Z., D. Unutmac, Y. R. Zou, M. J. Sunshine, A. Pierani, S. Brenner-Morton, R. E. Mebius, and D. R. Littman. 2000. Requirement for ROR γ t in thymocyte survival and lymphoid organ development. *Science* 288: 2369–2373.
22. Eberl, G., S. Marmon, M. J. Sunshine, P. D. Rennert, Y. Choi, and D. R. Littman. 2004. An essential function for the nuclear receptor ROR γ t in the generation of fetal lymphoid tissue inducer cells. *Nat. Immunol.* 5: 64–73.
23. Kim, D., R. E. Mebius, J. D. MacMicking, S. Jung, T. Cupedo, Y. Castellanos, J. Rho, B. R. Wong, R. Josien, N. Kim, et al. 2000. Regulation of peripheral lymph node genesis by the tumor necrosis factor family member TRANCE. *J. Exp. Med.* 192: 1467–1478.
24. Kong, Y. Y., H. Yoshida, I. Sarosi, H. L. Tan, E. Timms, C. Capparelli, S. Momoy, A. J. Oliveira-dos-Santos, G. Van, A. Itie, et al. 1999. OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature* 397: 315–323.
25. Dougall, W. C., M. Glaccum, K. Charrier, K. Rohrbach, K. Brasel, T. De Smedt, E. Daro, J. Smith, M. E. Tometsko, C. R. Maliszewski, et al. 1999. RANK is essential for osteoclast and lymph node development. *Genes Dev.* 13: 2412–2424.
26. Yoshida, H., A. Naito, J. Inoue, M. Satoh, S. M. Santee-Cooper, C. F. Ware, A. Togawa, S. Nishikawa, and S. Nishikawa. 2002. Different cytokines induce surface lymphotactin- α on IL-7 receptor- α cells that differentially engender lymph nodes and Peyer's patches. *Immunity* 17: 823–833.
27. Farr, A. G., M. L. Berry, A. Kim, A. J. Nelson, M. P. Welch, and A. Aruffo. 1992. Characterization and cloning of a novel glycoprotein expressed by stromal cells in T-dependent areas of peripheral lymphoid tissues. *J. Exp. Med.* 176: 1477–1482.
28. Berney, C., S. Herren, C. A. Power, S. Gordon, L. Martinez-Pomares, and M. H. Kosco-Vilbois. 1999. A member of the dendritic cell family that enters B cell follicles and stimulates primary antibody responses identified by a mannose receptor fusion protein. *J. Exp. Med.* 190: 851–860.
29. McNagny, K. M., R. P. Bucy, and M. D. Cooper. 1991. Reticular cells in peripheral lymphoid tissues express the phosphatidylinositol-linked BP-3 antigen. *Eur. J. Immunol.* 21: 509–515.
30. Katakai, T., T. Nomura, H. Gonda, M. Sugai, Y. Agata, A. Nishio, T. Masuda, S. Sakaguchi, and A. Shimizu. 2006. Spontaneous large-scale lymphoid neogenesis and balanced autoimmunity versus tolerance in the stomach of H $+$ /K $^{+}$ -ATPase-reactive TCR transgenic mouse. *J. Immunol.* 177: 7858–7867.
31. Katagiri, K., A. Maeda, M. Shimonaka, and T. Kinashi. 2003. RAPL, a Rap1-binding molecule that mediates Rap1-induced adhesion through spatial regulator of LFA-1. *Nat. Immunol.* 4: 741–748.
32. Sukumar, S., A. K. Srakal, and J. G. Tew. 2006. Isolation of functionally active murine follicular dendritic cells. *J. Immunol. Methods* 313: 81–95.
33. Katakai, T., T. Hara, M. Sugai, H. Gonda, and A. Shimizu. 2003. Th1-biased tertiary lymphoid tissue supported by CXC chemokine ligand 13-producing stromal network in chronic lesions of autoimmune gastritis. *J. Immunol.* 171: 4359–4368.
34. Cupedo, T., M. F. Vondenhoff, E. J. Heeregrave, A. E. De Weerd, W. Jansen, D. G. Jackson, G. Kraal, and R. E. Mebius. 2004. Presumptive lymph node organizers are differentially represented in developing mesenteric and peripheral nodes. *J. Immunol.* 173: 2968–2975.
35. Rennert, P. D., J. L. Browning, R. Mebius, F. Mackay, and P. S. Hochman. 1996. Surface lymphotactin- $\alpha\beta$ complex is required for the development of peripheral lymphoid organs. *J. Exp. Med.* 184: 1999–2006.
36. Rennert, P. D., D. James, F. Mackay, J. L. Browning, and P. S. Hochman. 1998. Lymph node genesis is induced by signaling through the lymphotactin β receptor. *Immunity* 9: 71–79.
37. Shinkura, R., K. Kitada, F. Matsuda, K. Tashiro, K. Ikuta, M. Suzuki, K. Kogishi, T. Serikawa, and T. Honjo. 1999. Alymphoplasia is caused by a point mutation in the mouse gene encoding NF- κ B-inducing kinase. *Nat. Genet.* 22: 74–77.
38. Koike, R., T. Nishimura, R. Yasumizu, H. Tanaka, Y. Hataba, T. Watanabe, S. Miyawaki, and M. Miyasaka. 1996. The splenic marginal zone is absent in alymphoplastic aly mutant mice. *Eur. J. Immunol.* 26: 669–675.
39. Bajenoff, M., J. G. Egen, L. Y. Koo, J. P. Laugier, F. Brau, N. Glaichenhaus, and R. N. Germain. 2006. Stromal cell networks regulate lymphocyte entry, migration, and territoriality in lymph nodes. *Immunity* 25: 989–1001.
40. Luther, S. A., T. Lopez, W. Bai, D. Hanahan, and J. G. Cyster. 2000. BLCl expression in pancreatic islets cause B cell recruitment and lymphotxin-dependent lymphoid neogenesis. *Immunity* 12: 471–481.
41. Aloisi, F., and R. Pujo-Borrell. 2006. Lymphoid neogenesis in chronic inflammatory diseases. *Nat. Rev. Immunol.* 6: 205–217.
42. Gonzalez, M., F. Mackay, J. L. Browning, M. H. Kosco-Vilbois, and R. J. Noelle. 1998. The sequential role of lymphotactin and B cells in the development of splenic follicles. *J. Exp. Med.* 187: 997–1007.
43. Hashi, H., H. Yoshida, K. Honda, S. Fraser, H. Kubo, M. Awane, A. Takabayashi, H. Nakano, Y. Yamaoka, and S. Nishikawa. 2001. Compartmentalization of Peyer's patch anlagen before lymphocyte entry. *J. Immunol.* 166: 3702–3709.
44. Cupedo, T., F. E. Lund, V. N. Ngo, T. D. Randall, W. Jansen, M. J. Greuter, R. de Waal-Malefyt, G. Kraal, J. G. Cyster, and R. E. Mebius. 2004. Initiation of cellular organization in lymph nodes is regulated by non-B cell-derived signals and is not dependent on CXC chemokine ligand 13. *J. Immunol.* 173: 4889–4896.
45. Taylor, R. T., S. R. Patel, E. Lin, B. R. Butler, J. G. Lake, R. D. Newberry, and I. R. Williams. 2007. Lymphotactin-independent expression of TNF-related activation-induced cytokine by stromal cells in cryptopatches, isolated lymphoid follicles, and Peyer's patches. *J. Immunol.* 178: 5659–5667.
46. Kanamori, Y., K. Ishimaru, M. Nanno, K. Maki, K. Ikuta, H. Nariuchi, and H. Ishikawa. 1996. Identification of novel lymphoid tissues in murine intestinal mucosa where clusters of e-kit⁺IL-7R⁺Thy1⁺ lympho-hemopoietic progenitors develop. *J. Exp. Med.* 184: 1449–1459.
47. Cupedo, T., and R. E. Mebius. 2005. Cellular interactions in lymph node development. *J. Immunol.* 174: 21–25.
48. Itano, A. A., and M.K. Jenkins. 2003. Antigen presentation to naive CD4 T cells in the lymph node. *Nat. Immunol.* 4: 733–739.
49. Iwasaki, A., and B. L. Kelsall. 2000. Localization of distinct Peyer's patch dendritic cell subsets and their recruitment by chemokines macrophage inflammatory protein (MIP)-3 α , MIP-3 β , and secondary lymphoid organ chemokine. *J. Exp. Med.* 191: 1381–1394.
50. Mowat, A. M. 2003. Anatomical basis of tolerance and immunity to intestinal antigens. *Nat. Rev. Immunol.* 3: 331–341.
51. Pape, K. A., D. M. Catron, A. A. Itano, and M. K. Jenkins. 2007. The humoral immune response is initiated in lymph nodes by B cells that acquire soluble antigen directly in the follicles. *Immunity* 26: 491–502.
52. Carrasco, Y. R., and F. D. Batista. 2007. B cells acquire particulate antigen in a macrophage-rich area at the boundary between the follicle and the subcapsular sinus of the lymph node. *Immunity* 27: 160–171.
53. Phan, T. G., I. Grigorova, T. Okada, and J. G. Cyster. 2007. Subcapsular encounter and complement-dependent transport of immune complexes by lymph node B cells. *Nat. Immunol.* 8: 992–1000.
54. Junt, T., E. A. Moseman, M. Iannaccone, S. Massberg, P. A. Lang, M. Boes, K. Fink, S. E. Henrickson, D. M. Shayakhmetov, N. C. Di Paolo, et al. 2007. Subcapsular sinus macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells. *Nature* 450: 110–114.
55. Lo, C. G., T. T. Lu, and J. G. Cyster. 2003. Integrin-dependence of lymphocyte entry into the spleen white pulp. *J. Exp. Med.* 197: 356–361.
56. Woolf, E., I. Grigorova, A. Sagiv, V. Grabovsky, S. W. Feigelson, Z. Shulman, T. Hartmann, M. Sixt, J. G. Cyster, and R. Alon. 2007. Lymph node chemokines promote sustained T lymphocyte motility without triggering stable integrin adhesiveness in the absence of shear forces. *Nat. Immunol.* 8: 1076–1085.
57. Lämmermann, T., B. L. Bader, S. J. Monkey, T. Worbs, R. Wedlich-Söldner, K. Hirsch, M. Keller, R. Förster, D. R. Critchley, R. Fässler, and M. Sixt. 2008. Rapid leukocyte migration by integrin-independent flowing and squeezing. *Nature* 453: 51–55.