

Fig. 5. Cytotoxicity (RS) and MN induction in TK6 cells treated with AA or DBN for 4 h in the presence (open symbol) or absence (closed symbol) of rat liver S9. Asterisk (*) statistically significant experiments in both pair-wise comparison and trend test ($P < 0.05$).

the bacterial and mammalian *HPRT* gene mutation assays detect only point mutations and small intragenic deletions [13]. AA was positive only in the *TK* mutation assay, suggesting that AA causes predominantly large-scale chromosomal changes. Our molecular analysis of the *TK* mutants supported this hypothesis. The majority of the AA-induced *TK* mutants showed hemi-LOH, which is the result of a deletion, although the other types were also induced (Fig. 3). Deletions are thought to result from the repair of double strand breaks by non-homologous end-joining [13]. Radiation-induced double strand breaks are repaired by non-homologous end-joining, which leads to hemi-LOH. LOH-mapping analysis, however, revealed that AA frequently induces intermediate-sized deletions (100–3000 kb); the deletions encompass exons 4 and 7 of the *TK* locus but do not extend to the microsatellites loci of the vicinity. This type of deletion is rarely observed in radiation-inducing *TK* mutants [13]. Because the COM assay indicated that AA did not induce DNA damage, the deletion may not be caused by DNA damage directly. Mechanisms associated with global genomic instability should also be considered [10] because the LOH patterns, except for the intermediate-sized deletions, are generally similar to those observed in spontaneous mutants. Most GA-induced *TK* mutants, on the other hand, were the non-LOH type, as were most spontaneous ones, strongly

supporting the positive results in bacterial gene mutation assay [29]. In contrast to AA, GA is a mutagen, inducing primarily point mutations.

AA is known to be metabolized to GA by CYP2E1 [9]. GA, an epoxide, forms adducts directly with DNA and protein, causing cytotoxicity and genotoxicity. GA forms mainly *N7*-(2-carbamoyl-2-hydroxyethyl) guanine and *N3*-(2-carbamoyl-2-hydroxyethyl) adenine and reacts with hemoglobin and cytoskeletal proteins [31–33]. Rat S9, however, did not affect AA cytotoxicity or genotoxicity, although it did enhance the cytotoxicity and genotoxicity of DBN, which is also metabolized by CYP2E1. This suggests that rat S9 does not work for activating AA. AA and GA are detoxified through glutathione conjugation, and GA is also detoxified by epoxy hydrolase (EH), which catalyzes the hydrolysis of GA to dihydroxy propionamide [34,35]. Other *in vitro* studies also failed to demonstrate the enhancement of AA genotoxicity by rat S9 [36,37]. Our results do not mean that AA is always detoxified rather than activated because DNA adducts are found in mice and rats given oral AA, and the genotoxicity of AA is consistently observed in *in vivo* studies [8,31,36,37]. Recently, Manjanatha et al. demonstrated in transgenic Big BlueTM mice that AA as well as GA induces endogenous *Hprt* and transgenic *cII* mutation at same level, and both chemicals cause predominantly base substitutions and frameshift mutations.

This result may indicate that AA is metabolized to GA in vivo [38]. Tests that use rat liver S9 for metabolic activation may not be appropriate for in vitro investigations of AA genotoxicity and metabolism. Transgenic cells expressing CYP2E1, however, would be useful for demonstrating the in vitro genotoxicity of AA [39].

In conclusion, AA is weakly genotoxic, causing chromosome aberrations and a type of genomic instability. GA, its epoxide metabolite, is highly reactive with DNA. GA is a strong mutagen, inducing predominantly point mutations, and it may contribute to human cancers.

Acknowledgment

This study was supported by Health and Labor Sciences Research Grants in Japan.

References

- [1] IARC, Acrylamide, IARC Monographs on the Evaluation of Carcinogen Risk to Human: Some Industrial Chemicals, vol. 60, International Agency for Research on Cancer, Lyon, 1994, pp. 389–433.
- [2] E. Tareke, P. Rydberg, P. Karlsson, S. Eriksson, M. Tornqvist, Acrylamide: a cooking carcinogen? *Chem. Res. Toxicol.* 13 (2000) 517–522.
- [3] R.H. Stadler, I. Blank, N. Varga, F. Robert, J. Hau, P.A. Guy, M.C. Robert, S. Riediker, Acrylamide from Maillard reaction products, *Nature* 419 (2002) 449–450.
- [4] d.S. Mottram, B.L. Wedzicha, A.T. Dodson, Acrylamide is formed in the Maillard reaction, *Nature* 419 (2002) 448–449.
- [5] J.M. Rice, The carcinogenicity of acrylamide, *Mutat. Res.* 580 (2005) 3–20.
- [6] E.J. Lehnig, C.D. Balaban, J.F. Ross, M.A. Reid, R.M. LoPachin, Acrylamide neuropathy. I. Spatiotemporal characteristics of nerve cell damage in rat cerebellum, *Neurotoxicology* 23 (2002) 397–414.
- [7] R.M. LoPachin, C.D. Balaban, J.F. Ross, Acrylamide axonopathy revisited, *Toxicol. Appl. Pharmacol.* 188 (2003) 135–153.
- [8] K.L. Dearfield, G.R. Douglas, U.H. Ehling, M.M. Moore, G.A. Sega, D.J. Brusick, Acrylamide: a review of its genotoxicity and an assessment of heritable genetic risk, *Mutat. Res.* 330 (1995) 71–99.
- [9] S.C. Sumner, T.R. Fennell, T.A. Moore, B. Chanas, F. Gonzalez, B.I. Ghanayem, Role of cytochrome P450 2E1 in the metabolism of acrylamide and acrylonitrile in mice, *Chem. Res. Toxicol.* 12 (1999) 1110–1116.
- [10] M. Friedman, Chemistry, biochemistry, and safety of acrylamide. A review, *J. Agric. Food Chem.* 51 (2003) 4504–4526.
- [11] H.M. Bolt, Genotoxicity—threshold or not? Introduction of cases of industrial chemicals, *Toxicol. Lett.* 140–141 (2003) 43–51.
- [12] H.L. Liber, W.G. Thilly, Mutation assay at the thymidine kinase locus in diploid human lymphoblasts, *Mutat. Res.* 94 (1982) 467–485.
- [13] M. Honma, Generation of loss of heterozygosity and its dependency on p53 status in human lymphoblastoid cells, *Environ. Mol. Mutagen.* 45 (2005) 162–176.
- [14] M. Honma, M. Hayashi, T. Sofuni, Cytotoxic and mutagenic responses to X-rays and chemical mutagens in normal and p53-mutated human lymphoblastoid cells, *Mutat. Res.* 374 (1997) 89–98.
- [15] K. Sekihashi, H. Saitoh, Y. Sasaki, Genotoxicity studies of stevia extract and steviol by the comet assay, *J. Toxicol. Sci.* 27 (Suppl. 1) (2002) 1–8.
- [16] M. Watanabe-Akanuma, T. Ohta, Y.F. Sasaki, A novel aspect of thiabendazole as a photomutagen in bacteria and cultured human cells, *Mutat. Res.* 158 (2005) 213–219.
- [17] L. Zhan, H. Sakamoto, M. Sakuraba, D.S. Wu, L.S. Zhang, T. Suzuki, M. Hayashi, M. Honma, Genotoxicity of microcystin-LR in human lymphoblastoid TK6 cells, *Mutat. Res.* 557 (2004) 1–6.
- [18] T. Matsushima, M. Hayashi, A. Matsuoka, M. Ishidate Jr., K.F. Miura, H. Shimizu, Y. Suzuki, K. Morimoto, H. Ogura, K. Mure, K. Koshi, T. Sofuni, Validation study of the in vitro micronuclei test in a Chinese hamster lung cell line (CHL/IU), *Mutagenesis* 14 (1999) 569–580.
- [19] S.A. Amundson, H.L. Liber, A comparison of induced mutation at homologous alleles of the tk locus in human cells, *Mutat. Res.* 247 (1991) 19–27.
- [20] E.E. Furth, W.G. Thilly, B.W. Penman, H.L. Liber, W.M. Rand, Quantitative assay for mutation in diploid human lymphoblasts using microtiter plates, *Anal. Biochem.* 110 (1981) 1–8.
- [21] T. Omori, M. Honma, M. Hayashi, Y. Honda, I. Yoshimura, A new statistical method for evaluating of L5178Y± mammalian cell data using microwell method, *Mutat. Res.* 517 (2002) 199–208.
- [22] M. Honma, M. Momose, H. Tanabe, H. Sakamoto, Y. Yu, J.B. Little, T. Sofuni, M. Hayashi, Requirement of wild-type p53 protein for maintenance of chromosomal integrity, *Mol. Carcinog.* 28 (2000) 203–214.
- [23] E. Zeiger, B. Anderson, S. Haworth, T. Lawlor, K. Mortelmans, W. Speck, Salmonella mutagenicity tests: III. Results from the testing of 255 chemicals, *Environ. Mutagen.* 9 (Suppl. 9) (1987) 1–109.
- [24] H. Tsuda, C.S. Shimizu, M.K. Taketomi, M.M. Hasegawa, A. Hamada, K.M. Kawata, N. Inui, Acrylamide; induction of DNA damage, chromosomal aberrations and cell transformation without gene mutations, *Mutagenesis* 8 (1993) 23–29.
- [25] A.G. Knaap, P.G. Kramers, C.E. Voogd, W.G. Bergkamp, M.G. Groot, P.G. Langebroek, H.C. Mout, J.J. van der Stel, H.W. Verharen, Mutagenic activity of acrylamide in eukaryotic systems but not in bacteria, *Mutagenesis* 3 (1988) 263–268.
- [26] T. Sofuni, M. Hayashi, A. Matsuoka, M. Sawada, Mutagenicity tests on organic chemical concomitants in city water and related compounds. II. Chromosome aberration tests in cultured mammalian cells, *Eisei Shiken. Hok.* 103 (1985) 64–75.
- [27] M. Baum, E. Fauth, S. Fritzen, A. Herrmann, P. Mertes, K. Merz, M. Rudolphi, H. Zankl, G. Eisenbrand, Acrylamide and glycidamide: genotoxic effects in V79-cells and human blood, *Mutat. Res.* 580 (2005) 61–69.
- [28] M.M. Moore, A. Amtower, C. Doerr, K.H. Brock, K.L. Dearfield, Mutagenicity and clastogenicity of acrylamide in L5178Y mouse lymphoma cells, *Environ. Mutagen.* 9 (1987) 261–267.
- [29] K. Hashimoto, H. Tani, Mutagenicity of acrylamide and its analogues in *Salmonella typhimurium*, *Mutat. Res.* 158 (1985) 129–133.
- [30] C.S. Aaron, G. Bolcsfoldi, H.R. Glatt, M. Moore, Y. Nishi, L. Stankowski, Theiss F J., E. Thompson, Mammalian cell gene mutation assays working group report, *Mutat. Res.* 312 (1994) 235–239.

- [31] J.C. Gamboa, M.I. Churchwell, L.P. Hamilton, L.S. Von Tungeln, F.A. Beland, M.M. Marques, D.R. Doerge, DNA adduct formation from acrylamide via conversion to glycidamide in adult and neonatal mice, *Chem. Res. Toxicol.* 16 (2003) 1328–1337.
- [32] D.M. Lapadula, M. Bowe, C.D. Carrington, L. Dulak, M. Friedman, M.B. Abou-Donia, In vitro binding of [¹⁴C]acrylamide to neurofilament and microtubule proteins of rats, *Brain Res.* 481 (1989) 157–161.
- [33] S.C. Sumner, C.C. Williams, R.W. Snyder, W.L. Krol, B. Asgharian, T.R. Fennell, Acrylamide: a comparison of metabolism and hemoglobin adducts in rodents following dermal, intraperitoneal, oral, or inhalation exposure, *Toxicol. Sci.* 75 (2003) 260–270.
- [34] B. Paulsson, A. Rannug, A.P. Henderson, B.T. Golding, M. Tornqvist, M. Warholm, In vitro studies of the influence of glutathione transferases and epoxide hydrolase on the detoxification of acrylamide and glycidamide in blood, *Mutat. Res.* 580 (2005) 53–59.
- [35] S.C. Sumner, L. Selvaraj, S.K. Nauhaus, T.R. Fennell, Urinary metabolites from F344 rats and B6C3F1 mice coadministered acrylamide and acrylonitrile for 1 or 5 days, *Chem. Res. Toxicol.* 10 (1997) 1152–1160.
- [36] A. Besaratinia, G.P. Pfeifer, Genotoxicity of acrylamide and glycidamide, *J. Natl. Cancer Inst.* 96 (2004) 1023–1029.
- [37] A. Besaratinia, G.P. Pfeifer, DNA adduction and mutagenic properties of acrylamide, *Mutat. Res.* 580 (2005) 31–40.
- [38] M.G. Manjanatha, A. Aidoo, S.D. Shelton, M.E. Bishop, L.P. MacDaniel, D.R. Doerge, Genotoxicity of acrylamide and its metabolite glycidamide administered in drinking water to male and female Big Blue mice, *Environ. Mol. Mutagen.*, in press (Epub ahead of prints).
- [39] H. Glatt, H. Schneider, Y. Liu, V79-hCYP2E1-hSULT1A1, a cell line for the sensitive detection of genotoxic effects induced by carbohydrate pyrolysis products and other food-borne chemicals, *Mutat. Res.* 580 (2005) 41–52.



Evaluation of the ability of a battery of three *in vitro* genotoxicity tests to discriminate rodent carcinogens and non-carcinogens II. Further analysis of mammalian cell results, relative predictivity and tumour profiles

David Kirkland^{a,*}, Marilyn Aardema^b, Lutz Müller^c, Makoto Hayashi^d

^a Covance Laboratories Limited, Otley Road, Harrogate HG3 1PY, United Kingdom

^b The Procter and Gamble Company, Miami Valley Laboratories, P.O. Box 538707, Cincinnati, OH 45253-8707, USA

^c Non-Clinical Drug Safety, F Hoffmann-La Roche Ltd., PRBN-S, Bldg 73311b, CH-4070 Basel, Switzerland

^d Division of Genetics and Mutagenesis, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

Received 2 February 2006; received in revised form 10 April 2006; accepted 30 April 2006

Available online 12 June 2006

Abstract

One of the consequences of the low specificity of the *in vitro* mammalian cell genotoxicity assays reported in our previous paper [D. Kirkland, M. Aardema, L. Henderson, L. Müller, Evaluation of the ability of a battery of three *in vitro* genotoxicity tests to discriminate rodent carcinogens and non-carcinogens. I. Sensitivity, specificity and relative predictivity, *Mutat. Res.* 584 (2005) 1–256] is industry and regulatory agencies dealing with a large number of false-positive results during the safety assessment of new chemicals and drugs. Addressing positive results from *in vitro* genotoxicity assays to determine which are “false” requires extensive resources, including the conduct of additional animal studies. In order to reduce animal usage, and to conserve industry and regulatory agency resources, we thought it was important to raise the question as to whether the protocol requirements for a valid *in vitro* assay or the criteria for a positive result could be changed in order to increase specificity without a significant loss in sensitivity of these tests. We therefore analysed some results of the mouse lymphoma assay (MLA) and the chromosomal aberration (CA) test obtained for rodent carcinogens and non-carcinogens in more detail. For a number of chemicals that are positive only in either of these mammalian cell tests (i.e. negative in the Ames test) there was no correlation between rodent carcinogenicity and level of toxicity (we could not analyse this for the CA test as insufficient data were available in publications), magnitude of response or lowest effective positive concentration. On the basis of very limited *in vitro* and *in vivo* data, we could also find no correlation between the above parameters and formation of DNA adducts. Therefore, a change to the current criteria for required level of toxicity in the MLA, to limit positive calls to certain magnitudes of response, or to certain concentration ranges would not improve the specificity of the tests without significantly reducing the sensitivity.

We also investigated a possible correlation between tumour profile (trans-species, trans-sex and multi-site *versus* single-species, single-sex and single-site) and pattern of genotoxicity results. Carcinogens showing the combination of trans-species, trans-sex and multi-site tumour profile were much more prevalent (70% more) in the group of chemicals giving positive results in all three *in vitro* assays than amongst those giving all negative results. However, single-species, single-sex, single-site carcinogens were not very prevalent even amongst those chemicals giving three negative results *in vitro*. Surprisingly, when mixed positive and negative results were compared, multi-site carcinogens were highly prevalent amongst chemicals giving only a single positive result in the battery of three *in vitro* tests.

* Corresponding author. Tel.: +44 1423 848401; fax: +44 1423 848983.

E-mail address: david.kirkland@covance.com (D. Kirkland).

Finally we extended our relative predictivity (RP) calculations to combinations of positive and negative results in the genotoxicity battery. For two out of three tests positive, the RP for carcinogenicity was no higher than 1.0 and for 2/3 tests negative the RP for non-carcinogenicity was either zero (for Ames + MLA + MN) or 1.7 (for Ames + MLA + CA). Thus, all values were less than a meaningful RP of two, and indicate that it is not possible to predict outcome of the rodent carcinogenicity study when only 2/3 genotoxicity results are in agreement.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Carcinogens; Mouse lymphoma assay; Chromosomal aberration; Genotoxicity; Tumour profiles

1. Introduction

Current regulatory guidelines for genotoxicity testing require chemicals to be evaluated at high concentrations/molarity and/or cellular toxicity *in vitro* for a valid test. The low specificity in the mammalian cell assays reported in our previous paper [1] raises the question as to whether such protocol requirements for a valid assay or the criteria for a positive result (according to ICH [2] and OECD [3] guidelines) could be changed in order to increase specificity, thereby reducing the number of false-positive results obtained. Since the standard approach to determine the biological relevance of positive *in vitro* genotoxicity assays involves conduct of additional *in vivo* genotoxicity assays, efforts to reduce this high false-positive rate could lead to substantial reductions in animal testing, as well as the cost and time to run these tests. Therefore, we examined such questions as whether the sensitivity of a particular mammalian cell assay could be maintained, but the specificity improved, if lower levels of toxicity were required for a valid assay? Did the carcinogens induce genotoxic responses of greater magnitude than the non-carcinogens, such that by requiring a higher level of response for a positive result, specificity might be improved? Also, were the non-carcinogens positive at much higher concentrations than the carcinogens, such that by requiring a lower top concentration than the current 10 mM, specificity might be improved? In this paper we have analysed some of the mouse lymphoma assay (MLA) and chromosomal aberration (CA) test results with rodent carcinogens and non-carcinogens (as defined in our previous paper [1]) in light of these questions.

We also examined whether a consistent pattern of genotoxicity results e.g. positive in Ames, MLA and either *in vitro* micronucleus (MN) or CA tests correlated better with trans-species, trans-sex, multi-site carcinogens than with single-species, single-sex, single-site carcinogens.

Finally, the new concept of relative predictivity (RP) that we introduced in our previous paper [1] and that had been calculated and reported only for those chemicals giving all three results positive or all three results

negative in the *in vitro* battery, was examined for other combinations of test results. Previously we reported that a reasonable correlation (RP of two or more) with rodent carcinogenicity was obtained when all three *in vitro* tests gave positive results (>3× more likely that a chemical would be a rodent carcinogen if all three tests were positive), and a reasonable correlation with non-carcinogenicity was obtained when all three tests were negative (>2× more likely that a chemical would be a non-carcinogen if all three tests were negative). As most compounds tested by industry give mixtures of positive and negative results across a battery of three tests (2 positive + 1 negative, or 1 positive + 2 negative), and not all regulatory guidelines request the conduct of an extensive *in vitro* battery, we calculated the RP for rodent carcinogenesis and non-carcinogenesis from these combinations of results.

2. Quantitative analysis of various aspects of the mammalian cell tests

An important source of poor specificity in our previous analysis [1] was for non-carcinogens where a single positive result was obtained in one of the mammalian cell tests in the battery. We therefore decided to examine whether changing the criteria for a valid assay or the criteria for a positive result would eliminate some of the isolated positives with non-carcinogens, whilst preserving the single positive results with carcinogens, thereby improving the specificity without decreasing the sensitivity. We decided to look at three parameters:

- Level of toxicity
- Magnitude of positive response
- Lowest clearly positive concentration

As indicated above, we focussed this analysis on those chemicals where the only positive response was in either the MLA or CA test, accompanied by negative results in the Ames test (we included one chemical with no Ames result), and either negative or no results in the other mammalian cell test.

Analysing these three parameters was relatively easy to do for the MLA because most published data (and many results came from NTP tests [4]) contained sufficient detail. However, very few CA publications, including the many NTP tests, provided detail on toxicity. We did have access to the laboratory notebooks of Dr. Ishidate, but were able to obtain toxicity information on too few clastogens to allow a meaningful analysis to be undertaken. For the MN test, where there are many fewer publications, there were so few situations where the MN was the only positive in the battery that this was not worthy of further analysis at this time.

As is the case in a retrospective analysis of this sort, it is important to recognise that the data from some studies may not meet current guideline requirements. Whilst it would be ideal to review each published result in terms of compliance with current guidelines, and focus our analysis only on those that complied, this goes beyond the scope of the project. As indicated previously [1], we did re-evaluate a significant number of the MLA studies from Mitchell et al. [6], and a number of published studies that we considered to be “technically compromised” were excluded from the analysis. However, there was insufficient information to do this for other study types, in particular the CA test. Nonetheless, we believe the trends observed here would largely still hold true if the analysis was limited to studies compliant with current guidelines.

2.1. Mouse lymphoma assay (MLA) results

The criteria for conclusions of positive, negative, equivocal or technically compromised have been described in our previous paper [1], as have the reasons why some previously reported MLA results were reassessed in order to fit our categorisation. The carcinogens and non-carcinogens where only the MLA gave a positive result (i.e. either both Ames and MN/CA were negative or results were not available) were identified from our Carcinogenicity Genotoxicity eXperience (CGX) database (as detailed in [1]; it can be viewed at <http://www.lhasalimited.org/cgx>). This amounted to only 45 chemicals, which is a small number, and analysis of other categories of chemicals in our database may provide more useful information. We re-examined the original reports and recorded the following:

- The highest relative total growth (RTG—i.e. lowest level of toxicity) at which a significant increase in mutant frequency occurred. Increases in MF above concurrent control levels of 90×10^{-6} for the agar method and 126×10^{-6} for the microwell method are

considered significant according to the latest recommendations [5]. Most of the published studies in our database used the agar method.

- The maximum increase in MF at any concentration giving acceptable RTG (i.e. RTG >0.1).
- The lowest effective concentration at which a significant increase in MF (as defined above) was observed.

For the MLA-positive chemicals shown in Tables 1 and 2, all except one were negative in the Ames test—for pyrilamine maleate there was no Ames test result. Most were also negative for CA and/or MN but for a small number there was no result for either CA or MN, as noted in the Table. For pyrilamine maleate there was also no result in either CA or MN, so the MLA was the only result. For two carcinogens (*o*-benzyl-*p*-chlorophenol and C.I. Direct Blue 218) the MLA data were referred to by Mitchell et al. [6] in their revised EPA Gene-Tox review, and therefore included in our CGX database, but the original data were not available for review.

Tables 1 and 2 show the highest RTG levels (i.e. lowest levels of toxicity) at which significant positive responses were seen for carcinogens and non-carcinogens, respectively. As can be seen, there were no differences between carcinogens and non-carcinogens in the levels of toxicity at which the minimum positive response was observed. Carcinogens and non-carcinogens are distributed across the whole range of toxicity for the point at which the first positive response is seen. The lowest level of toxicity for a biologically relevant positive response ranged from 23 to 89% (RTG from 0.77 to 0.11) for carcinogens, and from 16 to 91% (RTG 0.84–0.09) for non-carcinogens.

Thus, if the upper limit of toxicity required for a valid assay were lowered, several carcinogens would be missed. Also, sufficient non-carcinogens were positive at low levels of toxicity that the specificity would not be improved. For instance, if the toxicity limit were lowered to 50% (consistent with the chromosomal aberration test), seven non-carcinogens would no longer be positive (see Table 2A) but 15 carcinogens would also no longer be positive (see Table 1A).

Tables 1 and 2 also show the maximum induced MF (IMF) at any concentration that produced acceptable (non-excessive) toxicity (i.e. RTG not lower than 0.1). It should be noted that, at the time most of the MLA studies were performed, there was limited appreciation of the importance of small-colony mutants, and therefore some tests may have achieved conditions that favoured growth of small-colony mutants, whereas others would not. Clearly this could have an impact on

Table 1
Toxicity, maximum mutagenic response and lowest mutagenic concentrations in the MLA for Ames-negative carcinogens

Chemical name	CAS number	RTG for minimum positive response ^a	Maximum induced MF ($\times 10^{-6}$) at any treatment with RTG of 0.1 or higher	Lowest effective concentration ($\mu\text{mol/l}$) for minimum positive or highest non-significant response ^b	Induction of DNA adducts [reference]
A: Carcinogens positive only in MLA (i.e. negative or equivocal in Ames and CA/MN)					
11-Aminoundecanoic acid	2432-99-7	0.51	149	99.4	No data
Benzofuran	271-89-6	0.22	172	1270	No data
<i>o</i> -Benzyl- <i>p</i> -chlorophenol	120-32-1	Data not publicly available			
Butylated hydroxytoluene	128-37-0	0.26	145	45	No data
Chlorobenzilate	510-15-6	0.30	244	277	No data
Chloroform	67-66-3	0.34	92	0.4	+ Calf thymus DNA ^b [9] and hepatocytes [10] <i>in vitro</i>
4-Chloro- <i>o</i> -toluidine	3165-93-3	0.17	71 ^c	848	+ Liver <i>in vivo</i> [11]
C.I. Direct blue 218	28407-37-6	Data not publicly available			
Cinnamyl anthranilate	87-29-6	0.24	205	30.8	No data
Dicofol	115-32-2	0.30	36 ^c	135	No data
<i>N,N'</i> -Diethyl-2-thiourea	105-55-5	0.42	2307	11344	No data
Ethylbenzene	100-41-4	0.34	529	754	No data
Kepon (AKA chlordecone)	143-50-0	0.38	62 ^c	71.3	No data
Malonaldehyde sodium salt	24382-04-5	0.67	239	5553	+ Liver <i>in vivo</i> [12]
Methyl <i>tert</i> -butyl ether	1634-04-4	0.61	181	22.7	+ Hepatocytes <i>in vitro</i> ^d [13]
Piperonyl butoxide	51-03-6	0.17	443	148	No data
Piperonyl sulphoxide	120-62-7	0.70	85 ^c	7.7	No data
Toluene	108-88-3	0.21	216	2442	+ HL-60 cells <i>in vitro</i> [14]
2,4,6-Trichlorophenol	88-06-2	0.11	144	608	No data
Trimethylthiourea	2489-77-2	0.21	377	35533	No data
4-Vinylcyclohexene	100-40-3	0.40	79 ^c	740	+ Skin <i>in vivo</i> [15]
B: Carcinogens positive in MLA, negative Ames but no test (or TC) in CA/MN					
1,4-Benzoquinone	106-51-4	0.34	200	3.15	+ Plasmid DNA <i>in vitro</i> [16]
Dimethyl methylphosphonate	756-79-6	0.73	54 ^c	40.3	No data
FD&C Red 1 (Ponceau 3R)	3564-09-8	0.36	100	3440	No data
Procarbazine HCl (Natulan)	366-70-1	0.77	356	49.7	+ Mammary tissue [17], liver and bone marrow [18] <i>in vivo</i>
Thioacetamide	62-55-5	0.69	177	10240	+ Liver <i>in vivo</i> [19]
Trichloroacetic acid	76-03-9	0.28	121	12241	+ Liver <i>in vivo</i> ^b [20]
Vinylidene chloride	75-35-4	0.70	212	64990	No data
C: Carcinogens positive in MLA but no test (or TC) in Ames and CA/MN					
Pyrimidine maleate	59-33-6	0.55	155	1619	No data

NB: There were no chemicals positive in MLA, negative in CA/MN but no test (or TC) in Ames.

^a Minimum clear positive response considered where mutant frequency (MF) increased at least 90×10^{-6} (all studies used agar method) over concurrent control as recommended by Moore et al. [5]. In some cases, due to spacing of doses, the lowest significant increase in MF was much greater than the minimum, which would therefore have occurred at a higher RTG and lower concentration.

^b Oxidative damage to DNA.

^c Minimum increase in MF not achieved. These chemicals should be reclassified as equivocal rather than positive.

^d DNA-protein crosslinks.

^e Although induced MF was $<90 \times 10^{-6}$, the dose response was very steep and it is expected that a significant response would have been achieved at higher concentrations.

Table 2
Toxicity, maximum mutagenic response and lowest effective mutagenic concentrations in the MLA Ames-negative non-carcinogens

Chemical name	CAS number	RTG for minimum positive response ^a	Maximum induced MF ($\times 10^{-6}$) at any treatment with RTG of 0.1 or higher	Lowest effective concentration ($\mu\text{mol/l}$) for minimum positive or highest non-significant response ^a	Induction of DNA adducts [reference]
A: Non-carcinogens positive only in MLA (i.e. Ames and CA/MN negative or equivocal)					
Aldicarb	116-06-3	0.84	111	21024	No data
Anilazine	101-05-3	0.72	298	10.5	No data
Barium chloride dihydrate	10326-27-9	0.40	27 ^b	4094	No data
1,2-Dichlorobenzene	95-50-1	0.81	357	68	+ Calf thymus DNA <i>in vitro</i> [21]
FD&C yellow number 6	2783-94-0	0.79	62 ^b	8842	No data
4-Hexylresorcinol	136-77-6	0.29	110	154	No data
Hydrochlorothiazide	58-93-5	0.51	264	2519	No data
Isopropyl-N-(3-chlorophenyl)carbamate	101-21-3	0.13	68 ^b	147	No data
Malaoxon	1634-78-2	0.22	75 ^b	477	No data
Oxytetracycline HCl	2058-46-0	0.30	4329	616	No data
Phenyl- β -naphthylamine	135-88-6	0.36	140	59.3	No data
Sodium diethylcarbamate trihydrate	148-18-5	0.55	326	0.39	No data
Sulfisoxazole	127-69-5	0.39	96	3741	No data
2,3,5,6-Tetrachloro-4-nitroanisole	2438-88-2	0.55	151	24.1	No data
B: Non-carcinogens positive in MLA, negative in Ames but no test (or TC) in CA/MN					
Alpha-methyl-dopa sesquihydrate	41372-08-1	0.27	164	168	No data
Rhodamine 6G	989-38-8	0.09	94	16.7	No data

NB: There were no chemicals positive in MLA, negative in CA/MN but no test (or TC) in Ames.

^a Minimum clear positive response considered where mutant frequency (MF) increased at least 90×10^{-6} (all studies used agar method) over concurrent control as recommended by Moore et al. [5]. In some cases, due to spacing of doses, the lowest significant increase in MF was much greater than the minimum, which would therefore have occurred at a higher RTG and lower concentration.

^b Minimum increase in MF not achieved. Chemical should be reclassified as equivocal rather than positive.

the IMF, and highlights some of the difficulties of trying to apply current standards to old studies. Non-carcinogens (Table 2) were just as likely to produce very high IMF as were carcinogens (Table 1), and some carcinogens produced very low IMF. In fact, several carcinogens (as well as four non-carcinogens) failed to achieve the minimum IMF (90×10^{-6} for the agar method) currently recommended for a biologically relevant response [5] and should be re-classified as equivocal. These changes will be made in the web version of the CGX database (<http://www.lhasalimited.org/cgx>). Thus, if we were to consider increasing the required IMF for a positive response, several carcinogens would be missed.

The lowest concentrations of chemicals at which the minimum positive responses were seen (or in the case of those where minimum IMF was not reached, the highest non-significant concentration) are also shown in Tables 1 and 2. The lowest-effective concentrations clearly vary widely from <1 to $>10,000 \mu\text{M}$ for both carcinogens (Table 1) and non-carcinogens (Table 2). It will be noted that some chemicals are listed where

the concentration for a minimum positive response was $>10 \text{ mM}$. If tested according to current standards such high concentrations would not be used. Thus five carcinogens and two non-carcinogens would drop out of the analysis if positive results were only included when obtained below 10 mM . However, the range of lowest effective concentrations would still be large, both for carcinogens and non-carcinogens. Thus, changing the maximum concentration requirement for a valid test might improve the specificity of the MLA but it would lower the sensitivity.

We investigated whether formation (or not) of DNA adducts might discriminate between MLA positives of different potency. The *in vitro* and *in vivo* DNA-adduct data that could be found in the literature are also included in Tables 1 and 2. The following comments can be made on these limited data:

- There were nine carcinogens positive only in the MLA that had adduct data, and all formed DNA adducts. Within this sub-group, the level of toxicity varied widely (from 23 to 83% toxicity for a minimum pos-

itive response), the maximum IMF varied from 71 to 356×10^{-6} , and the lowest effective concentration varied over six orders of magnitude from 0.4 to 64,990 μM .

- There was only one non-carcinogen giving an isolated positive result in the MLA for which adduct data could be found, and this compound formed DNA adducts. This was positive at low toxicity (19% toxicity for minimum positive response).

Since all the MLA-positive carcinogens and the one MLA-positive non-carcinogen induced DNA adducts, it is not possible to use these data to draw conclusions regarding DNA reactivity and the other parameters we studied (level of toxicity required, minimum effective

concentration or magnitude of response). However, it does not appear that any changes to currently recommended toxicity levels or maximum concentration of test chemical, or requiring a higher induced mutant frequency for a positive "call" would improve the specificity of the MLA without causing serious deterioration in the sensitivity.

2.2. Chromosomal aberration (CA) test results

As mentioned earlier, there were very few published examples where CA results were accompanied by toxicity data. Although one of us (MH) was able to look at the laboratory notebooks of Dr. Ishidate, who published many of the CA results in our previous analysis [1], we

Table 3
Analysis of CA test responses for Ames-negative carcinogens

Chemical name	CAS number	Survival (cell count, confluence or mitotic index as a % of control) for minimum positive response ^a	Maximum % of aberrant cells (excluding gaps) at any treatment with survival 50% or higher	Lowest effective concentration ($\mu\text{mol/l}$) for minimum positive or highest non-significant response ^a	Induction of DNA adducts [reference]
A: Carcinogens positive only in CA ^b (i.e. negative in Ames and MLA) ^c					
Pentachloronitrobenzene	82-68-8	Not given	19	8.1	No data
Naphthalene	91-20-3	Not given	16	234	No data
Zearalenone	17924-92-4	Not given	13	47.1	+ Kidneys, liver and ovaries <i>in vivo</i> [22]
B: Carcinogens positive in CA ^b , negative in Ames but no test (or TC) in MLA ^d					
Atrazine	1912-24-9	64	28.7	5.0	- Liver <i>in vivo</i> [23]
Heptachlor	76-44-8	Not given	14	67.0	+ Hop plants <i>in vivo</i> [24]
Hexanamide	628-02-4	Not given	10	39071	No data
4-Methoxyphenol	150-76-5	Not given	6.8	250	No data
N-Methyloacrylamide	924-42-5	Not given	26	2473	No data
Methylphenidate HCl	298-59-9	Not given	9	3707	No data
Nafepopin	3771-19-5	100	Cannot be established	30.0	No data
Nitrobenzene	98-95-3	Not given	33.2	61.3	+ Liver <i>in vivo</i> [25]
C: Carcinogens positive in CA ^b but no test (or TC) in Ames and MLA ^e					
Haloperidol	52-86-8	Not given	Only one concentration used	Only one concentration used	No data
Retinol acetate ^f	127-47-9	Not given	13	200	No data

NB: There were no chemicals positive in CA, negative in MLA but no test (or TC) in Ames.

^a Minimum positive response was considered as >5% cells with aberrations (excluding gaps) as this would usually be considered a positive response for CHO, CHL cells and human lymphocytes.

^b Chemicals that were also positive in MN were excluded from these tables.

^c Another chemical in this category, but for which there were insufficient data in the publications to complete any of the columns, was sodium saccharin.

^d Other chemicals in this category, but for which there were insufficient data in the publications to complete any of the columns, were aldrin, aniline HCl, asbestos, carboxymethylnitrosourea, clofibrate, lead acetate, methimazole, styrene and 12-O-tetradecanoylphorbol 13-acetate.

^e Other chemicals in this category, but for which there were insufficient data in the publications to complete any of the columns, were 1-amy-1-nitrosourea, sodium barbital, manganese ethylenebisthiocarbamate and petastinenine.

^f Equivocal in MN.

were only able to find toxicity data on very few additional clastogens. Therefore, it was not possible to analyse (as we did above for the MLA) whether any change to the required level of toxicity (currently at least 50% according to OECD [3]) would provide improved specificity whilst retaining sensitivity.

For a small number of chemicals that were positive only in this test, we could identify the maximum induced CA response (although we could not know whether this occurred at levels of toxicity much higher than 50%) and the lowest effective concentration required for a minimum positive response. For the latter, since most published data were from Chinese hamster cells, we accepted that CA frequencies (excluding gaps) of >5% would probably be statistically significant and exceed historical control ranges. These data are tabulated in Tables 3 and 4 for carcinogens and non-carcinogens, respectively. Unfortunately there were a number of chemicals for which the published reports contained insufficient details to establish the maximum frequency of aberrant cells or the minimum concentration for a positive effect.

From Tables 3 and 4 it can be seen that the maximum percentage of aberrant cells ranged from 9 to 33% for carcinogens and from 7 to 23% for non-carcinogens.

The minimum positive concentration ranged from 5 to 39,000 μM for carcinogens and from 24 to 5000 μM for non-carcinogens. In other words, for this small sample of Ames-negative clastogens there were no clear differences in maximum response or minimum positive concentration between carcinogens and non-carcinogens. It will be noted that one chemical (hexanamide, Table 3) is listed where the concentration for a minimum positive response was >10 mM. If tested according to current standards such high concentrations would not be used. Thus this one carcinogen would drop out of the analysis if positive results were only included when obtained below 10 mM. However, the range of lowest effective concentrations would still be large, both for carcinogens and non-carcinogens. Thus it does not appear that any change to the maximum concentration, or the introduction of a requirement for the induced frequency of aberrant cells to reach a certain level would improve the specificity of the CA test without causing serious deterioration in the sensitivity.

As for the analysis of MLA data, we investigated whether DNA reactivity may correlate better with potency. Any *in vitro* or *in vivo* DNA-adduct data that could be found in the literature for these chemicals positive only in the CA test are shown in Tables 3 and 4.

Table 4
Analysis of CA test responses for Ames-negative non-carcinogens

Chemical name	CAS number	Survival (cell count, confluence or mitotic index as a % of control) for minimum positive response ^a	Maximum % of aberrant cells (excluding gaps) at any treatment with survival 50% or higher	Lowest effective concentration ($\mu\text{mol/l}$) for minimum positive or highest non-significant response ^a	Induction of DNA adducts [reference]
A: Non-carcinogens positive only in CA ^b (i.e. Ames and MLA negative) ^c					
Diphenhydramine HCl	147-24-0	Not given	17	343	No data
4,4'-Isopropylidenediphenol	80-05-7	45	23	399	+ Liver <i>in vivo</i> [26] and Syrian hamster cells <i>in vitro</i> [27] No data
Tin (II) chloride	7772-99-8	Not given	22	132	No data
B: Non-carcinogens positive in CA, negative in Ames but no test (or TC) in MLA ^d					
Carbromal	77-65-6	Not given	7	5061	No data
Chlorpheniramine maleate	113-92-8	Not given	18	1279	No data
Fenvalerate	51630-58-1	Not given	20	23.8	No data
Hexachlorocyclopentadiene	77-47-4	Not given	9.5	27.5	No data

NB: There were no chemicals positive in CA, negative in MLA but no test (or TC) in Ames.

^a Minimum positive response was considered as >5% cells with aberrations (excluding gaps) as this would usually be considered a positive response for CHO, CHL cells and human lymphocytes.

^b Chemicals that were also positive in MN were excluded from these tables.

^c Other chemicals in this category, but for which there were insufficient data in the publications to complete any of the columns, were acetohexamide, benzoin, FD&C Red number 3 and tetracycline HCl.

^d Other chemicals in this category, but for which there were insufficient data in the publications to complete any of the columns, were sodium benzoate, caffeine, chlorpropamide, FD&C Yellow number 5 and vinyl toluene.

Adduct data were found for only a small number of chemicals. However, the following comments can be made:

- Of the carcinogens that were positive only for CA, there were only four with adduct data, and one of these (atrazine) did not form DNA adducts. Whereas we might have expected DNA-reactive chemicals to be positive at lower concentrations, the non-DNA-reactive atrazine was positive at the lowest concentration of all the 13 carcinogens studied in this set.
- Of the carcinogens with adduct data the maximum CA response was quite variable (13–33%) as was the lowest effective concentration (5–67 μM).
- Only one non-carcinogen had adduct data (which was positive).

Since all except one CA-positive carcinogen and the one CA-positive non-carcinogen induced DNA adducts, it is not possible to draw conclusions regarding DNA reactivity and the other parameters we studied (magnitude of response, or lowest effective concentration). Thus it does not appear that the specificity of the CA test can be improved, without serious deterioration in sensitivity, by changing the maximum required test concentration or setting a limit for magnitude of response.

3. Tumour profiles

We reviewed the tumour profiles as detailed in the Carcinogenic Potency DataBase (CPDB) of Gold [7] for those carcinogens giving positive results across all three *in vitro* tests (Table 5) or giving negative results across all three *in vitro* tests (Table 6). The prevalence of the trans-species, trans-sex and multi-site carcinogens amongst the two categories of results we examined (positive or negative across all three *in vitro* tests) is shown in Table 7A. The prevalence of chemicals showing “all positive” or “all negative” amongst the single-species, single-sex, single-site carcinogens is shown in Table 7B. The following conclusions can be drawn from these analyses:

- There are no differences between “all positive for genotoxicity” and “all negative for genotoxicity” amongst the carcinogens as far as inducing tumours in both males and females. In other words, trans-sex carcinogens are equally likely to be genotoxic in all *in vitro* tests as to be non-genotoxic in all *in vitro* tests.
- Trans-species carcinogens are 39% more prevalent in the “all positive for genotoxicity” group than in the “all negative for genotoxicity” group.

- Multi-site carcinogens are 31% more prevalent in the “all positive for genotoxicity” group than in the “all negative for genotoxicity” group.
- Carcinogens showing the combination of trans-species, trans-sex and multi-site tumour profile are much more prevalent (70% more) in the “all positive for genotoxicity” group than in the “all negative for genotoxicity” group.
- Single-species, single-sex, single-site carcinogens are not very prevalent even amongst those chemicals giving three negative results *in vitro*.

We also reviewed the tumour profiles for carcinogens giving positive results in only one of the *in vitro* genotoxicity tests whilst being negative in the other two tests in the battery. The results are shown in Table 8A for Ames-positive only, Table 8B for MLA-positive only, and Table 8C for MN- or CA-positive only. The prevalence of trans-species, trans-sex or multi-site carcinogens amongst these compounds positive in a single genotoxicity test is shown in Table 9. The following conclusions can be drawn:

- Multi-site carcinogens are surprisingly prevalent amongst those chemicals inducing only single positive genotoxicity results.
- Multi-site carcinogens are more prevalent amongst Ames-positive genotoxins, but the database is very small.
- The prevalence of trans-species or trans-sex or multi-site carcinogens amongst single genotoxicity test positive compounds is intermediate between those in “all positive” and “all negative” groups.

4. Relative predictivity for batteries with mixed positive and negative results

Because we did not observe any obvious changes to protocols for the mammalian cell tests that would improve specificity without impairing sensitivity, we revisited the topic of relative predictivity described in our previous paper [1] and we calculated relative predictivity (RP) for rodent carcinogenicity from different combinations of test results. As described previously, positive RP for predicting carcinogenicity was acceptable (>2) only for the Ames test of the single tests, but was most informative when all three tests were positive. We also calculated the RP for non-carcinogenicity from one, two or three *in vitro* tests all giving negative results. Only a few combinations of all negative results gave acceptable RP for non-carcinogenicity.

Table 5

Tumour profiles for carcinogens (tested in rats and mice, males and females) positive in Ames plus MLA plus either MN or CA

Carcinogen	CAS number	Tumour profile		
		Rats and mice	Males and females	Multi-site
N-Acetoxy-2-acetylaminofluorene	6098-44-8	Yes	Yes	Yes
2-Acetylaminofluorene	53-96-3	Yes	Yes	Yes
2-Aminoanthracene ^a	613-13-8	Yes	Yes	Yes
2-Aminofluorene ^a	153-78-6	Yes	Yes	Yes
2-Amino-4-nitrophenol	99-57-0	No	No	No
4-Amino-2-nitrophenol	119-34-6	No	No	No
2-Amino-5-nitrothiazole	121-66-4	No	No	Yes
5-Azacytidine	320-67-2	Yes	Yes	Yes
Benzo[a]pyrene	50-32-8	Yes	Yes	No ^b
Benzyl chloride	100-44-7	Yes	Yes	Yes
bis(2-Chloro-1-methylethyl)ether, technical grade	108-60-1	No	Yes	Yes
Calcium chromate	13765-19-0	Yes	Yes	Yes
Captan	133-06-2	Yes	Yes	Yes
Chlorodibromomethane	124-48-1	No	Yes	No
3-(Chloromethyl)pyridine HCl	6959-48-4	Yes	Yes	No
C.L. Disperse blue 1	2475-45-8	No	Yes	No
C.L. Disperse orange 2 (1-amino-2-methyl-anthraquinone)	82-28-0	Yes	Yes	Yes
Cyclophosphamide monohydrate	6055-19-2	Yes	Yes	Yes
Cytembena	21739-91-3	No	Yes	Yes
2,4-Diaminoanisole sulphate	39156-41-7	Yes	Yes	Yes
2,4-Diaminotoluene	95-80-7	Yes	Yes	Yes
1,2-Dibromo-3-chloropropane	96-12-8	Yes	Yes	Yes
1,2-Dibromoethane	106-93-4	Yes	Yes	Yes
Dichloroacetic acid	79-43-6	Yes	Yes	No
Dichloromethane	75-09-2	Yes	Yes	Yes
2,6-Dichloro-p-phenylenediamine	609-20-1	No	Yes	No
1,2-Dichloropropane	78-87-5	No	Yes	No
Dichlorvos	62-73-7	Yes	Yes	Yes
Diglycidyl resorcinol ether, technical grade	101-90-6	Yes	Yes	No
3,3'-Dimethylbenzidine	119-93-7	Yes	Yes	Yes
Dimethyl hydrogen phosphite	868-85-9	No	No	Yes
Epichlorohydrin	106-89-8	No	Yes	No
1,2-Epoxybutane	106-88-7	No	No	Yes
Ethyl methanesulphonate	62-50-0	Yes	Yes	Yes
Formaldehyde	50-00-0	Yes	Yes	Yes
Furylfuramide (AF-2)	3688-53-7	Yes	Yes	No ^b
Glycidol	556-52-5	Yes	Yes	Yes
Hydrazine sulphate	10034-93-2	Yes	Yes	Yes
Methylazoxymethanol acetate	592-62-1	No	Yes	Yes
4,4'-Methylenedianiline 2HCl	13552-44-8	Yes	Yes	Yes
N-Methyl-N'-nitro-N-nitrosoguanidine	70-25-7	Yes	Yes	Yes
2-Naphthylamine	91-59-8	Yes	Yes	No ^b
o-Nitroanisole	91-23-6	Yes	Yes	Yes
5-Nitro-2-furaldehyde semicarbazone (AKA Nitrofurazone)	59-87-0	Yes	No	Yes
1-[(5-Nitrofurfurylidene)amino]hydantoin (AKA Nitrofurantoin)	67-20-9	Yes	No	No ^b
2-Nitro-p-phenylenediamine	5307-14-2	No	No	No
4-Nitroquinoline-N-oxide	56-57-5	Yes	Yes	Yes
N-Nitrosodimethylamine (dimethylnitrosamine)	62-75-9	Yes	Yes	Yes
p-Nitrosodiphenylamine	156-10-5	Yes	No	No
4,4'-Oxydianiline	101-80-4	Yes	Yes	Yes
Phenobarbital	50-06-6	No	Yes	No
Beta-Propiolactone	57-57-8	Yes	Yes	No
1,2-Propylene oxide	75-56-9	Yes	Yes	Yes
Quercetin	117-39-5	No	Yes	Yes
p-Quinone dioxime	105-11-3	No	No	No
Selenium sulphide	7446-34-6	Yes	Yes	Yes
Styrene oxide	96-09-3	Yes	Yes	No
1,2,3-Trichloropropane	96-18-4	Yes	Yes	Yes
Zinc dimethyldithiocarbamate (Ziram)	137-30-4	No	Yes	Yes

NT: not tested in that species or sex.

^a Carcinogens positive in Ames + MLA + MN but not tested or equivocal in CA. All others positive in Ames + MLA + CA.^b Single-site tumours only within a species. However, different tumours may have arisen in rats and mice.

Table 6
 Tumour profiles for carcinogens negative in Ames plus MLA plus either MN or CA

Carcinogen	CAS number	Tumour profile		
		Rats and mice	Males and females	Multi-site
3-Amino-1,2,4-triazole (Amitrole)	61-82-5	Yes	Yes	Yes
<i>tert</i> -Butyl alcohol	75-65-0	Yes	No ^a	No ^b
5-Chloro- <i>o</i> -toluidine	95-79-4	No	Yes	Yes
Decabromodiphenyl oxide	1163-19-5	No	Yes	No
Diethanolamine	111-42-2	No	Yes	Yes
Di(2-ethylhexyl)phthalate	117-81-7	Yes	Yes	No
1,4-Dioxane	123-91-1	Yes	Yes	Yes
<i>DL</i> -Ethionine	67-21-0	Yes	Yes	No
Melamine	108-78-1	No	No	No
Methyl carbamate	598-55-0	No	Yes	No
Nitriiotriacetic acid, trisodium salt, monohydrate	18662-53-8	No	Yes	Yes
<i>N</i> -Nitrosodiphenylamine	86-30-6	No	Yes	No
Progesterone	57-83-0	Yes	Yes	Yes
Pyridine	110-86-1	Yes	Yes	No ^b
Reserpine	50-55-5	Yes	Yes	Yes
2,3,7,8-Tetrachlorodibenzo- <i>p</i> -dioxin	1746-01-6	Yes	Yes	Yes
1,1,2,2-Tetrachloroethane	79-34-5	No	Yes	No
Tris(2-ethylhexyl)phosphate	78-42-2	No	No	No

By inhalation.

^a Tumours in male rats but female mice.

^b Single-site tumours only within a species. However, different tumours may have arisen in rats and mice.

Table 7A
 Prevalence of trans-species, trans-sex or multi-site carcinogens amongst consistent positive or negative genotoxicity responses

	Trans-species carcinogens (%)	Trans-sex carcinogens (%)	Multi-site carcinogens (%) ^a	Trans-species, multiple site and trans-sex carcinogens (%)
Positive in Ames plus MLA plus either MN or CA	69.5 (41/59)	83.1 (49/59)	72.9 (43/59)	56 (33/59)
Negative in Ames plus MLA plus either MN or CA	50.0 (9/18)	83.3 (15/18)	55.6 (10/18)	33 (6/18)

^a Including those carcinogens producing single-site tumours within one species but tumours at different sites in different species.

Table 7B
 Prevalence of single-species, single-sex and single-site carcinogens amongst consistent positive or negative genotoxicity responses

	Single-species, single-sex, single-site carcinogens (%)
Positive in Ames plus MLA plus either MN or CA	6.8 (4/59)
Negative in Ames plus MLA plus either MN or CA	11.1 (2/18)

Table 8A
 Tumour profile for carcinogens positive in Ames but negative in MLA, MN or CA

Carcinogen	CAS number	Tumour profile		
		Rats and mice	Males and females	Multi-site
D&C red 9	5160-02-1	No	No	Yes
Trifluralin, technical grade	1582-09-8	No	No	Yes
Urethane ^a	51-79-6	Yes	Yes	Yes

^a There are a number of reports that urethane is negative in the Ames test. As noted in our previous paper [1] we decided to accept the deliberations of Zeiger [28] for this test with urethane.

Table 8B

Tumour profile for carcinogens (with results in rats and mice, males and females) positive in MLA but negative in Ames, MN or CA

Carcinogen	CAS number	Tumour profile		
		Rats and mice	Males and females	Multi-site
11-Aminoundecanoic acid	2432-99-7	No	No	Yes
Benzofuran	271-89-6	Yes	Yes	Yes
<i>o</i> -Benzyl- <i>p</i> -chlorophenol	120-32-1	No	No	No
Butylated hydroxytoluene	128-37-0	No	No	Yes
Chlorobenzilate	510-15-6	No	Yes	No
Chloroform	67-66-3	Yes	Yes	Yes
C.I. Direct blue 218	28407-37-6	Yes	Yes	No ^a
Cinnamyl anthranilate	87-29-6	Yes	Yes	Yes
Dicofol	115-32-2	No	No	No
<i>N,N</i> -Diethyl-2-thiourea	105-55-5	No	Yes	No
Ethylbenzene	100-41-4	Yes	Yes	Yes
Kepon (AKA Chlordecone)	143-50-0	Yes	Yes	No
Malonaldehyde sodium salt	24382-04-5	Yes	Yes	Yes
Methyl <i>tert</i> -butyl ether	1634-04-4	Yes	Yes	Yes
Piperonyl butoxide	51-03-6	Yes	Yes	No
Piperonyl sulphoxide	120-62-7	No	No	No
Toluene	108-88-3	No	Yes	Yes
Trichloroethylene (with and without epichlorhydrin)	79-01-6	Yes	Yes	Yes
2,4,6-Trichlorophenol;	88-06-2	Yes	Yes	No ^a
Trimethylthiourea	2489-77-2	No	No	No

^a Single-site tumours only within a species. However, different tumours may have arisen in rats and mice.

Table 8C

Tumour profile for carcinogens (with results in rats and mice, males and females) positive in MN or CA but negative in Ames and MLA

Carcinogen	CAS number	Tumour profile		
		Rats and mice	Males and females	Multi-site
DDT	50-29-3	Yes	Yes	Yes
17- β -Estradiol	50-28-2	Yes	Yes	Yes
Nitrotriacetic acid	139-13-9	Yes	Yes	Yes
Oxazepam	604-75-1	No	Yes	Yes
Pentachloronitrobenzene	82-68-8	No	No	No
Saccharin, sodium ^a	128-44-9	No	No	No
Zeralenone	17924-92-4	No	Yes	Yes

^a Sodium saccharin induces bladder tumours via chronic irritation from the formation of precipitates at high doses, and the CA positive response was only obtained at concentrations >10 mM, so it is unlikely these effects are connected.

Table 9

Prevalence of trans-species, trans-sex or multi-site carcinogens amongst single-positive genotoxicity responses

	Trans-species carcinogens (%)	Trans-sex carcinogens (%)	Multi-site carcinogens (%) ^a	Trans-species, multiple site and trans-sex carcinogens (%)
Positive Ames only	33.3 (1/3)	33.3 (1/3)	100 (3/3)	33.3 (1/3)
Positive MLA only	55.0 (11/20)	70.0 (14/20)	60.0 (12/20)	45.0 (9/20)
Positive MN or CA only	42.9 (3/7)	71.4 (5/7)	71.4 (5/7)	42.9 (3/7)
Positive in any single test but Negative in the other two	50.0 (15/30)	66.7 (20/30)	66.7 (20/30)	43.3 (13/30)

^a Including those carcinogens inducing single-site tumours within one species, but inducing tumours at different sites in different species.

Table 10A
Numbers of carcinogens and non-carcinogens with 2/3 results positive or negative^a

	Ames plus MLA plus MN (%)	Ames plus MLA plus CA (%)
Carcinogens		
2/3 Tests positive	16/50 (32.0)	51/176 (29.0)
2/3 Tests negative	6/50 (12.0)	30/176 (17.0)
Non-carcinogens		
2/3 Tests positive	8/12 (66.7)	22/76 (28.9)
2/3 Tests negative	0/12 (0)	22/76 (28.9)

^a E and TC results excluded from analysis (counted as blank = no result).

Table 10B
Relative predictivity of carcinogen/non-carcinogen status when 2/3 results indicative (data from Table 10A)

Combination of three tests	Chemicals positive in 2/3 tests (%)		Relative predictivity that 2/3 positive results indicate a carcinogen A/B	Chemicals negative in 2/3 tests (%)		Relative predictivity that 2/3 negative results indicate a non-carcinogen C/D
	Carcinogens (A)	Non-carcinogens (B)		Non-carcinogens (C)	Carcinogens (D)	
Ames plus MLA plus MN	32.0	66.7	0.48	0	12.0	0
Ames plus MLA plus CA	29.0	28.9	1.00	28.9	17.0	1.70

As a battery of three *in vitro* tests most often gives a mixture of positive and negative results, we decided to calculate RP for predicting carcinogenicity when two tests were positive but one negative, and to calculate RP for predicting non-carcinogenicity when two tests were negative but one positive. The numbers of results falling into each category are shown in Table 10A and the RP values in Table 10B. It can be seen that in all cases the RP values are less than the meaningful value of two, and in some cases are <1, which means there would be a better chance of predicting the outcome of the carcinogenicity studies by flipping a coin.

As all of the RP values were worse for 2/3 tests than for 3/3 tests, we decided not to analyse further the RP from 1/3 tests.

5. Discussion and conclusions

Although limited data were available, our analyses of chemicals giving isolated positive genotoxicity results in one of the mammalian cell tests, suggest that revising the requirements for toxicity limits or highest test concentration, or to apply some "threshold" response that must be exceeded, would improve the poor specificity of the MLA and CA tests, but would at the same time significantly impair sensitivity. An examination of DNA reactivity within these small groups did not suggest any associations between DNA reactivity and level of toxicity or concentration needed for a minimum positive

response, or the magnitude of the genotoxic response. It is possible that if we examine all MLA and CA positives, and view separately those carcinogens known to interact with DNA, and/or known to have structural alerts, we may find that the "profile" of the genotoxicity results (concentration, level of toxicity, magnitude of response) would be more useful in predicting carcinogenic outcome. This is currently being explored and will be the subject of a future manuscript.

An analysis of tumour profiles indicated that trans-species, trans-sex and multi-site carcinogens are much more prevalent amongst those chemicals giving positive results in all three *in vitro* genotoxicity tests than amongst those giving negative results in all three tests. This is consistent with the relative predictivity (RP) analysis we reported previously [1] where chemicals giving positive results in all three *in vitro* tests would be at least 3 × more likely to be rodent carcinogens than non-carcinogens. However, unexpectedly, single-species, single-sex and single-site carcinogens were not prevalent amongst those chemicals giving negative results in all three *in vitro* tests. This suggests that non-genotoxic carcinogens are just as likely to produce single-site tumours in one sex of one species as to produce multi-site tumours in both sexes of rats and mice. It was also surprising that multi-site carcinogens were highly prevalent amongst those chemicals positive in one but negative in the other two genotoxicity tests. Multi-site carcinogens were particularly prevalent amongst Ames-positive chemicals, but the

database was small and further examples are needed for a rigorous evaluation. Thus, it appears that the profile of genotoxicity results cannot be used with any confidence to predict the tumour profile in subsequent carcinogenicity studies. This is consistent with a more modern view that mutational events and chromosomal rearrangements are, although required for tumourigenesis, not a major driving factor for prevalence and incidence of tumours in lifetime rodent carcinogenicity tests with chemical compounds [8]. In such long-term experiments, hormonal disturbances, genetic strain background, general tumour promotional effects, etc. are very important. Such effects may equally be exerted by many of the investigated chemicals that also possess genotoxic properties.

Although our previous RP analysis [1] indicated that chemicals giving positive results in all three *in vitro* tests were >3× more likely to be rodent carcinogens than non-carcinogens, and that chemicals giving all negative results were >2× more likely to be non-carcinogens than carcinogens, in industry mixed positive and negative results are most common. As might be expected, RP analysis of “mixed” genotoxicity results revealed that these cannot be used to predict carcinogenic outcome. It should be noted that predictivity analyses are dependent on the choice of compounds analysed and influenced by the prevalence of carcinogens and non-carcinogens in the data set. Different predictivities from those presented here could be obtained with different numbers of carcinogens and non-carcinogens in the data set. Also, only some of the mechanisms (e.g. genotoxic) giving rise to cancer are the same as those leading to mutations and chromosomal aberrations, and non-genotoxic mechanisms would not be expected to be predicted by genotoxicity assays. Therefore, the predictivity of genotoxicity tests for cancer will depend on the mechanism.

As indicated previously [1], the disappointing findings from these analyses suggest it may be time for a complete rethink with regard to *in vitro* genotoxicity testing. It does not appear that a redefinition of the conditions (cytotoxicity, solubility, etc.) for *in vitro* testing will improve the low specificity for prediction of rodent tumourigenesis, and therefore new, more robust assays may be needed. We should also consider, however, that the rodent bioassay is not relevant for predicting human carcinogenicity in many cases, and therefore new, more robust carcinogenicity assays may also be needed.

Regarding the availability of complete data sets, or of critical information (e.g. cytotoxicity) related to single-assay results, the authors are astonished that only small subsets of chemicals have full data sets. Hence, despite more than 30 years of genotoxicity testing, the published literature still contains large data gaps. These gaps cer-

tainly do not facilitate far-reaching and important conclusions on correlations between important parameters. Thus in many cases our only recourse is to some kind of patchwork analysis when working with published data. As expected, most adduct data have been reported for the carcinogens and there are very few studies of adducts formed by non-carcinogens.

References

- [1] D. Kirkland, M. Aardema, L. Henderson, L. Müller, Evaluation of the ability of a battery of three *in vitro* genotoxicity tests to discriminate rodent carcinogens and non-carcinogens. I. Sensitivity, specificity and relative predictivity, *Mutat. Res.* 584 (2005) 1–256.
- [2] ICH Topic S2A, Genotoxicity: guidance on specific aspects of regulatory genotoxicity tests for pharmaceuticals. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use, Harmonised tripartite guideline CPMP/ICH/141/95, approved September 1995.
- [3] OECD, Guidelines for the Testing of Chemicals: Genotoxicity. Organisation for Economic Co-operation and Development, Paris, Revised and new guidelines adopted, 1997.
- [4] NTP website at <http://ntp-server.niehs.nih.gov>.
- [5] M.M. Moore, M. Honma, J. Clements, G. Bolcsfoldi, B. Burlinson, M. Cifone, J. Clarke, R. Delongchamp, R. Durward, M. Fellows, B. Gollapudi, S. Hou, P. Jenkinson, M. Lloyd, J. Majeska, B. Myhr, M. O'Donovan, T. Omori, C. Riach, R. San, L.F. Stankowski Jr., A.K. Thakur, F. Van Goethem, S. Wakuri, I. Yoshimura, Mouse lymphoma thymidine kinase gene mutation assay: follow-up meeting of the international workshop on genotoxicity tests—Aberdeen, Scotland, 2003, *Environ. Mol. Mutagen.* 47 (2006) 1–5.
- [6] A.D. Mitchell, A.E. Auletta, D. Clive, P.E. Kirby, M.M. Moore, B.C. Myhr, The L5178Y/*tk*^{+/−} mouse lymphoma specific gene and chromosomal mutation assay. A phase III report of the U.S. Environmental protection Agency Gene-Tox Program, *Mutat. Res.* 394 (1997) 177–303.
- [7] L.S. Gold, The carcinogenic potency project. <http://potency.berkeley.edu/cpdb.html> (2004).
- [8] R.A. Beckman, L.A. Loeb, Genetic instability in cancer: theory and experiment, *Semin. Cancer Biol.* 15 (2005) 423–435.
- [9] A.B. Drenzo, A.J. Gandolfi, I.G. Sipes, Microsomal bioactivation and covalent binding of aliphatic halides to DNA, *Toxicol. Lett.* 11 (1982) 243–252.
- [10] E.J. Beddowes, S.P. Faux, J.K. Chipman, Chloroform, carbon tetrachloride and glutathione depletion induce secondary genotoxicity in liver cells via oxidative stress, *Toxicology* 187 (2003) 101–115.
- [11] P. Bentley, F. Waechter, F. Bieri, W. Staubli, W. Muecke, Species differences in the covalent binding of *p*-chloro-*o*-toluidine to DNA, *Arch. Toxicol. Suppl.* 9 (1986) 163–166.
- [12] A. Kautiainen, C.E. Vaca, F. Granath, Studies on the relationship between hemoglobin and DNA adducts of malonaldehyde and their stability *in vivo*, *Carcinogenesis* 14 (1993) 705–708.
- [13] M. Casanova, H.A. Heck, Lack of evidence for the involvement of formaldehyde in the hepatocarcinogenicity of methyl tertiary-butyl ether in CD-1 mice, *Chem. Biol. Interact.* 105 (1997) 131–143.

- [14] N.W. Gaikwad, W.J. Bodell, Formation of DNA adducts in HL-60 cells treated with the toluene metabolite *p*-cresol: a potential biomarker for toluene exposure, *Chem. Biol. Interact.* 145 (2003) 149–158.
- [15] N. Mabon, K. Randerath, ³²P-postlabeling of 1,3-butadiene and 4-vinyl-1-cyclohexene metabolite-DNA adducts: *in vitro* and *in vivo* applications, *Toxicology* 113 (1996) 341–344.
- [16] M. Gaskell, K.I. McLuckie, P.B. Farmer, Genotoxicity of benzene metabolites *para*-benzoquinone and hydroquinone, *Chem. Biol. Interact.* 153/154 (2005) 267–270.
- [17] L.Y. Fong, D.E. Jensen, P.N. Magee, DNA methyl-adduct dosimetry and O⁶-alkylguanine-DNA alkyl transferase activity determinations in rat mammary carcinogenesis by procarbazine and *N*-methylnitrosourea, *Carcinogenesis* 11 (1990) 411–417.
- [18] V. Plesta, C. Valavanis, J.H. van Delft, M.J. Steenwinkel, S.A. Kryptopoulos, DNA damage and mutagenesis induced by procarbazine in lambda lacZ transgenic mice: evidence that bone marrow mutations do not arise primarily through miscoding by O⁶-methylguanine, *Carcinogenesis* 18 (1997) 2191–2196.
- [19] A.E. Pegg, M.E. Dolan, D. Scicchitano, K. Morimoto, Studies of the repair of O⁶-alkylguanine and O⁴-alkylthymine in DNA by alkyltransferases from mammalian cells and bacteria, *Environ. Health Perspect.* 62 (1985) 109–114.
- [20] E.W. Austin, J.M. Parrish, D.H. Kindler, R.J. Bull, Lipid peroxidation and formation of 8-hydroxydeoxyguanosine from acute doses of halogenated acetic acids, *Fund. Appl. Toxicol.* 31 (1996) 77–82.
- [21] M. Paolini, L. Pozzetti, P. Silingardi, C. Della Croce, G. Bronzetti, G. Cantelli-Forti, Isolation of a novel metabolising system enriched in phase-II enzymes for short-term genotoxicity assays, *Mutat. Res.* 413 (1998) 205–217.
- [22] A. Pfohl-Leszkowicz, L. Chekir-Ghedira, H. Bacha, Genotoxicity of zearalenone, an estrogenic mycotoxin: DNA adduct formation in female mouse tissues, *Carcinogenesis* 16 (1995) 2315–2320.
- [23] S.E. George, R.W. Chadwick, M.J. Kohan, J.C. Allison, S.H. Warren, R.W. Williams, Atrazine treatment potentiates excretion of mutagenic urine in 2,6-dinitrotoluene-treated Fischer 344 rats, *Environ. Mol. Mutagen.* 26 (1995) 178–184.
- [24] J.A. Laouedj, C. Schenk, A. Pfohl-Leszkowicz, G. Keith, D. Schontz, P. Guillemat, B. Rether, Detection of DNA adducts in declining hop plants grown on fields formerly treated with heptachlor, a persistent insecticide, *Environ. Pollut.* 90 (1995) 409–414.
- [25] H. Li, H. Wang, H. Sun, Y. Liu, K. Liu, S. Peng, Binding of nitrobenzene to hepatic DNA and hemoglobin at low doses in mice, *Toxicol. Lett.* 139 (2003) 25–32.
- [26] A. Atkinson, D. Roy, *In vivo* DNA adduct formation by Bisphenol A, *Environ. Mol. Mutagen.* 26 (1995) 60–66.
- [27] T. Tsutsui, Y. Tamura, E. Yagi, K. Hasegawa, M. Takahashi, N. Maizumi, F. Yamaguchi, J.C. Barrett, Bisphenol-A induces cellular transformation, aneuploidy and DNA adduct formation in cultured Syrian hamster embryo cells, *Int. J. Cancer* 75 (1998) 290–294.
- [28] E. Zeiger, Genotoxicity database, in: L.S. Gold, E. Zeiger (Eds.), *Handbook of Carcinogenic Potency and Genotoxicity Databases*, CRC Press Inc., Boca Raton, 1997, pp. 687–729.

Flow Cytometric Analysis of Micronuclei in Peripheral Blood Reticulocytes: I. Intra- and Interlaboratory Comparison with Microscopic Scoring

Stephen D. Dertinger,* Michelle E. Bishop,† James P. McNamee,‡ Makoto Hayashi,§ Takayoshi Suzuki,§ Norihide Asano,¶ Madoka Nakajima,|| Junichiro Saito,||| Martha Moore,† Dorothea K. Torous,* and James T. MacGregor|||¹

*Litron Laboratories, Rochester, New York 14623; †U.S. Food and Drug Administration, National Center for Toxicological Research, Jefferson, Arkansas 72079; ‡Health Canada, Ottawa, Ontario, Canada K1A 0L2; §National Institute of Health Sciences, Tokyo 158-8501, Japan; ¶Nitto Denko Corporation, Osaka 567-8680, Japan; ||An-Pyo Center, Shizuoka 437-1213, Japan; |||Astellas Pharma Inc., Tokyo 174-8511, Japan; |||U.S. Food and Drug Administration, National Center for Toxicological Research, Rockville, Maryland 21012

Received February 2, 2006; accepted July 31, 2006

Accumulating evidence suggests that reticulocytes (RETs) in the peripheral blood of rats may represent a suitable cell population for use in the micronucleus assay, despite the ability of the rat spleen to selectively remove micronucleated erythrocytes from the peripheral circulation. To evaluate the analytical performance of a previously described flow cytometric method (Torous *et al.*, 2003, *Toxicol. Sci.* 74, 309–314) that may allow this assay to be conducted using peripheral blood *in lieu* of bone marrow sampling, we compared the sensitivity and performance characteristics of the flow cytometric technique with two established microscopy-based scoring methods. Peripheral blood samples from single Sprague-Dawley rats treated for 6 days with either vehicle or cyclophosphamide were prepared in replicate for scoring by the three methods at different laboratories. These blood-based measurements were compared to those derived from bone marrow specimens from the same animals, stained with acridine orange, and scored by microscopy. Through the analysis of replicate specimens, inter- and intralaboratory variability were evaluated for each method. Scoring reproducibility over time was also evaluated. These data support the premise that rat RETs harvested from peripheral blood are a suitable cell population to assess genotoxicant-induced micronucleus formation. The interlaboratory comparison provides evidence of the general robustness of the micronucleus endpoint using different analytical approaches. Furthermore, data presented herein demonstrate a clear advantage of flow cytometry-based scoring over microscopy—significantly lower inter- and intralaboratory variation and higher statistical sensitivity.

Key Words: flow cytometric analysis; reticulocytes; micronucleus test; CD71.

The *in vivo* rodent erythrocyte micronucleus (MN) test is widely used in research and regulatory safety assessment to evaluate the potential of chemical and physical agents to cause chromosomal damage. Historically, MN studies based on rat peripheral blood have been avoided as it has been assumed that the efficiency by which the rat spleen filters out erythrocytes with intracellular inclusions would reduce assay sensitivity (Hayashi *et al.*, 2000; Wakata *et al.*, 1998). However, accumulated data suggest that peripheral blood from intact rats can be used effectively to detect chemical-induced genotoxicity (Abramsson-Zetterberg *et al.*, 1999; Asanami *et al.*, 1995; Hamada *et al.*, 2001; Hayashi *et al.*, 1992; Hynes *et al.*, 2002; Romagna and Staniforth, 1989; Torous *et al.*, 2000, 2003; Wakata *et al.*, 1998). Thus, it appears that MN studies using peripheral blood sampling in the rat have the potential to substitute for labor-intensive, bone marrow-based tests. In addition, the ability to use low-volume blood samples will facilitate integration of the assay into routine toxicology and/or pharmacokinetic studies and may make it unnecessary to conduct separate assays for the evaluation of chromosomal damage (Asanami *et al.*, 1995; Hamada *et al.*, 2001; MacGregor *et al.*, 1995; Wakata *et al.*, 1998).

Before rat blood-based MN assays gain wider acceptance, especially in the context of regulatory testing requirements, additional information that allows direct comparisons between bone marrow and blood data is needed. Furthermore, the performance characteristics of the most widely utilized scoring techniques require further study. The experiments described herein were designed to address these issues of analytical performance by directly comparing values in blood and bone marrow obtained at different laboratories with three widely used methodologies, comparing values derived from two microscopy-based methods with a flow cytometry-based method that incorporates a calibration standard.

For each of the three scoring techniques, at least three proficient laboratories received replicate, coded samples for reticulocyte

¹ To whom correspondence should be sent at the present address: Toxicology Consulting Services, 201 Nomini Drive, Arnold, MD 21012. Fax: 410-975-0481. E-mail: jtmacgregor@earthlink.net.

TABLE 1
Participating Laboratories

Laboratory	Code	Specimens analyzed	Scoring method	Instrumentation (magnification)
U.S. FDA-NCTR, Rockville, MD and Jefferson, AR	L1	BM, PB	MeOH-AO	Zeiss Axioskop 50 ($\times 630$), Zeiss PlanApomat $\times 63$ oil objective
			FCM	BD-FACSort, BD-FACScan
Litron Laboratories, Rochester, NY	L2	PB	FCM	BD-FACSCalibur
Health Canada, Ottawa, Ontario, Canada	L3	PB	FCM	BD-FACSCalibur
National Institute of Health Sciences, Tokyo, Japan	L4	Coordinated SV-AO laboratories		
Nitto Denko Corporation, Osaka, Japan	L5	PB	SV-AO	Olympus AHB3-RFC ($\times 600$)
An-Pyo Center, Shizuoka, Japan	L6	PB	SV-AO	Olympus BX50-RFL ($\times 800$)
Astellas Pharma Inc., Tokyo, Japan	L7	PB	SV-AO	Olympus BH-RFL ($\times 600$)
N/A to this study	L8			
Contract testing laboratory 1*	L9	BM, PB	MeOH-AO	Leitz Laborlux 12 ($\times 1000$)
Contract testing laboratory 2*	L10	BM	MeOH-AO	Olympus BH2 ($\times 1000$)
Contract testing laboratory 3*	L11	BM, PB	MeOH-AO	Zeiss STD 14 ($\times 1000$)

Note. Abbreviations: FDA-NCTR = U.S. Food and Drug Administration, National Center of Toxicological Research; BM = bone marrow; PB = peripheral blood; MeOH-AO = acridine orange staining of methanol-fixed smears; FCM = flow cytometry; SV-AO = supravital staining using acridine orange-coated slides; N/A = not applicable. *The three contract testing laboratories are BioReliance, Covance, and SR1 International, but their identities as L9, L10, or L11 is confidential.

(RET) and MN-RET scoring. Proficiency was assumed based on the high level of training that has occurred at these laboratories (L1, L2, L3, L5, L6, and L7) and/or the frequency with which they contribute *in vivo* rodent MN data for regulatory submission purposes (L9, L10, and L11). See Table 1 for more detailed information regarding collaborating laboratories.

Data presented herein describe the performance characteristics of the three scoring methods evaluated, address the sensitivity of the rat peripheral blood compartment for detecting genotoxicant-induced micronuclei, and support recommendations concerning the minimum number of rat blood RET that should be evaluated for micronuclei.

MATERIALS AND METHODS

Chemicals and other reagents. Cyclophosphamide (CP) (CAS No. 6055-19-2) was purchased from Sigma, St Louis, MO. Acridine orange (AO)-coated slides used for supravital staining, prepared according to the method of Hayashi *et al.* (1990), were provided by the National Institute of Health Sciences, Japan. Flow cytometry reagents, including fixed malaria-infected rat blood (malaria biostandard) were from Rat MicroFlow^{PLUS} Kits contributed by Litron Laboratories (available from Litron Laboratories, Rochester, NY and BD Biosciences Pharmingen, San Diego, CA).

Animals and treatment regimens. Animal studies were conducted in compliance with guidelines of the National Research Council (1996) "Guide for the Care and Use of Laboratory Animals" and were approved by the appropriate Institutional Animal Care and Use Committees. Two female Sprague-Dawley rats, 4- to 5-weeks old, were purchased from Taconic, Germantown, NY. Animals were housed singly and were assigned randomly to treatment groups. The animals were acclimated for approximately 2 weeks before the experiment was initiated. Food and water were available *ad libitum* throughout the acclimation and experimentation periods. One rat was treated via oral gavage with distilled water, and the other rat was treated by the same route with 10 mg CP/kg/day for 6 consecutive days.

Blood/bone marrow sample collection and storage. Each day, before vehicle or CP treatment, low-volume blood samples (approximately 100 μ l) were collected from the tail vein using a 26.5-gauge needle and syringe after a brief warming period under a heat lamp. These samples were fixed for flow cytometric analysis of RET and MN-RET frequencies according to procedures described in the Rat MicroFlow^{PLUS} manual (v020213). Fixed samples were stored at -85°C until analysis. Approximately 24 h after the last administration of vehicle or CP, blood samples were collected into tubes containing heparin solution (500 USP units heparin per milliliter of phosphate buffered saline) as follows: into a small tube containing 75 μ l heparin solution, blood was collected until a final volume of approximately 750 μ l was obtained; into a second tube containing 5 ml heparin solution, approximately 1 ml blood was collected. To tubes with the 750 μ l blood suspension, an equal volume of heat-inactivated fetal bovine serum (FBS) was added. These FBS-diluted suspensions were used to prepare replicate AO-supravital (SV) slides (8 μ l per slide) according to the method of Hayashi *et al.* (1990, 1992). These slides were frozen, shipped to collaborating SV-AO laboratories on dry ice, and stored frozen until analysis. FBS-diluted blood suspensions were also used to prepare slides for conventional acridine orange staining of methanol-fixed smears (MeOH-AO) staining (5 μ l per slide). These blood smears were prepared by drawing the cell suspensions behind a second slide with smoothed edges (a "spreader slide"). These smears were allowed to air dry and were then fixed with absolute methanol for 10 min. The slides were stored in a slide box until they were shipped to collaborating MeOH-AO laboratories for MN scoring according to their standard operating procedures. Replicate bone marrow slides were prepared as smears, air dried, methanol fixed, and shipped similarly. These bone marrow cells were harvested from two femurs per rat, whereby both ends of each femur were cut and its contents flushed with 1 ml FBS. The cells were centrifuged at approximately 1100 rpm for 5 min and then resuspended with approximately 600 μ l FBS. As with the peripheral blood, 5 μ l of cell suspension was applied to each slide. The 6 ml heparinized peripheral blood suspensions were fixed with ultracold methanol according to procedures described in the Rat MicroFlow^{PLUS} manual (v020213) in order to preserve cells for flow cytometric analysis. These cell suspensions were stored at -85°C until analysis or shipment on dry ice to collaborating flow cytometry laboratories.

The samples obtained were divided into three identical pools, and replicate samples of each pool were provided to participating laboratories with three separate codes. Thus, laboratories received triplicate samples of each condition, but were not aware that they were from an identical pool. Thus, the analyses

conducted allow assessment of both intralaboratory variability of replicate analysis of identical samples and interlaboratory variability of the same analysis. Each laboratory also conducted analysis of each of these pools on three separate occasions, allowing assessment of variability of analysis over time.

Standard acridine orange slide scoring (MeOH-AO). Blood and bone marrow smears were scored using the MeOH-AO scoring technique at the Food and Drug Administration-National Center for Toxicological Research Laboratory (L1) and three contract testing laboratories (L9, L10, and L11). Methanol fixation leads to a diffuse distribution of RNA, and erythrocytes are classified as normochromatic or as RETs based on the presence or absence of RNA-associated fluorescence. This technique is not well suited for visually classifying subpopulations of RETs. RET frequencies were determined by inspecting 500 or 1000 total erythrocytes per bone marrow or blood sample, respectively. MN-RET incidence was determined by inspecting 2000 RETs per sample. At L1, micronuclei were defined by the criteria of Schmid (1976) with the added requirements that they exhibit the characteristic yellow to yellow-green fluorescence characteristic of AO staining and that they exhibit the smooth boundary expected from a membrane-bound body. Laboratories L9, L10, and L11 were instructed to follow the standard operating procedures they use for regulatory submissions to support new drug or food additive development. Thus, the acquisition of data by these facilities allows for comparisons with three highly experienced contract laboratories under conditions associated with regulatory testing.

Supravital acridine orange slide scoring (SV-AO). Laboratories L5, L6, and L7 scored peripheral blood samples using the SV-AO scoring technique. This staining procedure aggregates RNA, leading to punctate staining patterns. These staining characteristics allow RET to be classified into four age cohorts: Type I (youngest) through Type IV (oldest) RETs as described by Hayashi *et al.* (1990, 1992). The frequency of MN-RETs was determined by analyzing 2000 Type I and Type II RETs (L5 and L7) or 2000 Type I RETs (L6). An index of cytotoxicity was obtained by inspecting at least 400 RETs and calculating the percentage of Type I and Type II RET among total RETs (L5 and L7) or the percentage of Type I RETs among Type I and Type II RETs (L6). AO-coated slides were purchased from TOYOBO (Osaka, Japan). Supravital stained triplicate slides were frozen and sent to the Japanese reference laboratory (Nitto Denko) with dry ice. Each set of slides was also sent to two other laboratories for replicate scoring by fluorescence microscopy.

Flow cytometry-based scoring. Methanol-fixed blood samples were washed and labeled for flow cytometric analysis by L1, L2, and L3 according to procedures described in the Rat MicroFlow^{PLUS} Kit (v020213). Samples were analyzed with 488-nm capable instruments (FACSCalibur, FACSort, and FACScan, all from Becton Dickinson, San Jose, CA). Anti-CD71-FITC and propidium iodide fluorescence signals were detected in the FL1 and FL3 channels, respectively (stock filter sets). Calibration of the flow cytometers for the MN scoring application, across laboratories and between experiments within each laboratory, was accomplished by staining *Plasmodium berghei*-infected rat blood (malaria biostandards) in parallel with test samples on each day of analysis (Dertinger *et al.*, 2000; Tometsko *et al.*, 1993; Torous *et al.*, 2001). By adjusting voltages applied to the photomultiplier tube, it was possible to standardize the FL3 fluorescence channel into which erythrocytes with single (MN like) parasites fell. In this manner, analysis regions were consistent across laboratories and between experiments. Flow cytometry-based MN-RET measurements reported herein are based on an immature fraction of peripheral blood RETs (approximately the youngest 30–50% of propidium iodide-positive erythrocytes, based on CD71 expression level; Torous *et al.*, 2001, 2003). This is thought to be analogous to scoring the youngest (Types I and II) RETs using the SV-AO method, which may be beneficial in view of reports which have suggested that the influence of rat spleen filtration function can be minimized by scoring the younger RETs (Abramsson-Zetterberg *et al.*, 1999; Hayashi *et al.*, 1992; Hynes *et al.*, 2002; Torous *et al.*, 2000, 2003). Data were acquired with CellQuest software (v3.3, BD-Immunocytometry Systems, San Jose, CA), with the stop mode set so that 20,000 high CD71-expressing RETs were analyzed per blood sample. The number of mature (CD71 negative) erythrocytes was determined concurrently, providing an index of cytotoxicity (%RETs).

Calculations. All calculations were performed with Excel (Office X for Mac or Microsoft Office Excel 2002 for XP Windows Professional, Microsoft Corp., Seattle, WA). The incidences of MN-RETs are expressed as frequency percent. The percentage of RETs among total erythrocytes was measured by the flow cytometric and MeOH-AO laboratories and served as an index of bone marrow cytotoxicity. The three SV-AO laboratories used percentage of RETs in different stages of maturity as an index of toxicity; therefore, these indices are not directly comparable to those obtained by the flow cytometric and MeOH-AO microscopy laboratories. Percent coefficient of variance values (%CV, i.e., standard deviation (SD) as percent of the mean) were used to describe intralaboratory variability associated with multiple readings of replicate samples and also interlaboratory variation of vehicle control and CP-induced MN-RET measurements that were pooled according to scoring method.

RESULTS AND DISCUSSION

Confirmation of Steady State

RET and MN-RET measurements obtained from the daily low-volume blood specimens were analyzed to confirm that the




FIG. 1. The frequency of peripheral blood RETs (%RET, panel a) and peripheral blood micronucleated RETs (%MN-RET, panel b) as a function of time in the individual rats used to generate reference samples for analytical comparison. These data were acquired by flow cytometric analysis (laboratory L2) and demonstrate the attainment of a steady-state MN-RET frequency, facilitating subsequent comparisons between bone marrow and peripheral blood compartments.