

FIG. 2. Panels (a-c): Bivariate graphs illustrate the staining characteristics of rat blood specimens over the course of several days of CP treatment. Note the appearance of micronuclei at Day 1 in the very youngest (highest anti-CD71-FITC fluorescence) RETs (Panel b) and the more uniform distribution among RETs after a steady state has been reached on Day 3 of treatment (Panel c). Panel (d) illustrates the use of the malaria standard, with distinct fluorescence intensities corresponding to inclusion of one, two, or three parasites. This allows the instrument settings to be standardized to the DNA content of the parasite, which is controlled biologically to a quantity similar to that in an average MN.

MN-RET frequency of the vehicle-treated animal was stable over the duration of the experiment and that CP treatment caused the MN-RET frequency to increase to a steady-state level of approximately 10-fold the control frequency (Fig. 1). Since the frequency of MN-RETs was at steady state in both cases, the values in bone marrow and peripheral blood should be directly comparable—that is, expected to be equal in the absence of selective removal of MN-RETs from blood or methodological differences in measurement. Thus, the samples collected in this manner allow the direct comparisons between measurements in the bone marrow and blood compartments that follow. The use of large samples from a single treated and a single control rat allows differences in methodology and scoring laboratory to be assessed independently of sample variation.

The dose of CP (10 mg/kg/day) had a moderate effect on erythropoiesis, as indicated by the decline in RET frequency (terminal day specimen showed a greater than 50% decrease from pretreatment value; see Fig. 1, panel a). This level of bone marrow cytotoxicity is well within the range of target toxicity recommended by current regulatory guidances (i.e., $\leq 80\%$, see Organisation for Economic Cooperation and Development, 1997, Guideline 474; U.S. Food and Drug Administration, 2000).

To illustrate the nature and source of the flow cytometry-based data described above, bivariate fluorescence intensity plots are provided (Fig. 2). Note the appearance of micronuclei on Day 1 in the very youngest (highest anti-CD71-FITC fluorescence) RETs (Panel b) and the more uniform

TABLE 2
Reticulocyte Data (cytotoxicity determinations)

Laboratory	Method	Compartment	Treatment	Cytotoxicity index ^a	Average %RET ^b ± SEM	%CV	%Change
L1	MeOH-AO	BM	Vehicle	%RET	81.0 ± 0.70	1.5	
			CP		67.8 ± 2.36	6.0	- 16
L9	MeOH-AO	BM	Vehicle	%RET	65.4 ± 1.03	2.7	
			CP		51.9 ± 1.83	6.1	- 21
L10	MeOH-AO	BM	Vehicle	%RET	58.2 ± 1.65	4.9	
			CP		60.0 ± 0.66	1.9	+ 3
L11	MeOH-AO	BM	Vehicle	%RET	63.1 ± 1.67	4.6	
			CP		57.1 ± 1.14	3.5	- 10
<i>Pooled^b L1, 9, 10, 11</i>			Vehicle		66.9 ± 2.63	13.6	
			CP		59.2 ± 1.87	10.9	- 12
L1	MeOH-AO	PB	Vehicle	%RET	7.7 ± 0.19	4.2	
			CP		5.7 ± 0.27	8.2	- 26
L9	MeOH-AO	PB	Vehicle	%RET	6.2 ± 0.27	7.4	
			CP		5.6 ± 0.90	27.8	- 9
L11	MeOH-AO	PB	Vehicle	%RET	6.6 ± 0.52	13.5	
			CP		4.9 ± 0.33	11.7	- 26
<i>Pooled L1, 9, 11</i>			Vehicle		6.9 ± 0.29	12.5	
			CP		5.4 ± 0.32	17.6	- 21.7
L5	SV-AO	PB	Vehicle	%Type I + II/III-IV	55.2 ± 1.95	6.1	
			CP		42.2 ± 0.12	0.5	- 24
L6	SV-AO	PB	Vehicle	%Type I/II + III	42.4 ± 2.8	11.5	
			CP		29.1 ± 2.1	12.4	- 31
L7	SV-AO	PB	Vehicle	%Type I + II/III-IV	52.3 ± 2.1	6.8	
			CP		34.8 ± 3.1	15.6	- 34
<i>Pooled^c L5, 7</i>			Vehicle		53.7 ± 1.4	6.5	
			CP		38.5 ± 2.2	13.8	- 28
L1	FCM	PB	Vehicle	%RET ^{High} CD71+	3.40 ± 0.02	1.18	
			CP		1.53 ± 0.01	0.75	- 55
L2	FCM	PB	Vehicle	%RET ^{High} CD71+	3.32 ± 0.02	1.26	
			CP		1.44 ± 0.01	1.44	- 57
L3	FCM	PB	Vehicle	%RET ^{High} CD71+	3.32 ± 0.05	2.42	
			CP		1.40 ± 0.08	9.88	- 58
<i>Pooled L1, 2, 3</i>			Vehicle		3.34 ± 0.02	1.93	
			CP		1.46 ± 0.03	6.33	- 56

Note. Abbreviations: RET = reticulocyte; MeOH-AO = acridine orange staining of methanol-fixed smears; SV-AO = supravital staining using acridine orange-coated slides; FCM = flow cytometry; BM = bone marrow; PB = peripheral blood; CP = cyclophosphamide; SEM = standard error of the mean.

^aEach laboratory evaluated cytotoxicity based on immature erythrocyte parameters. This was accomplished in several different manners: %RET = percentage of RETs relative to total erythrocytes; %Type I + II/III-IV = percentage of Type I and Type II RETs relative to total RETs; %Type I/II + III = percentage of Type I RETs relative to Type I and Type II RETs; and %RET^{High} CD71+ = percentage of RETs that express high levels of CD71 relative to total erythrocytes.

^bValues are the mean of three separately coded, but identical, samples. By "Pooled" it is meant that like-method data from two, three, or four laboratories were combined for these calculations.

^cOnly data from the two SV-AO laboratories that measured toxicity similarly (%Type I + II/III-IV) were combined for these calculations.

among RETs after a steady state has been reached on Day 3 of treatment (Panel c). Panel (d) illustrates the use of the malaria biostandard, with distinct fluorescence intensities corresponding to inclusion of one, two, or three parasites. This allows the instrument settings to be standardized to the DNA content of the parasite, which is controlled biologically to a quantity similar to that in an average MN. For research purposes, the regions may be adjusted to allow measurements in different age populations of RETs and/or micronuclei with different DNA contents. For analytical purposes, the standard can be used to achieve comparable instrument performance across time within a laboratory or across different instruments in different laboratories.

Intra- and Interlaboratory Variability

Replicate bone marrow and/or peripheral blood specimens obtained after 6 consecutive days of treatment were provided to each collaborating laboratory. As noted above, the frequencies of MN-RETs were at steady state and therefore not changing as a function of time. Each laboratory received three separately coded samples from each of the high and low MN-RET frequency pools but were not aware that the three separately coded samples were identical. Tabular values are the means of the values of the three separately coded samples.

TABLE 3
Intra- and Interlaboratory Micronucleated Reticulocyte Data

Laboratory	Method	Compartment	Treatment	%MN-RET		
				Average ^a ± SEM	%CV	Fold difference
L1	MeOH-AO	BM	Vehicle	0.15 ± 0.03	33.3	
			CP	3.35 ± 0.10	5.4	22.3
L9	MeOH-AO	BM	Vehicle	0.05 ± 0.05	173.2	
			CP	1.63 ± 0.27	28.4	32.6
L10	MeOH-AO	BM	Vehicle	0.03 ± 0.02	86.6	
			CP	2.33 ± 0.23	17.3	77.7
L11	MeOH-AO	BM	Vehicle	0.18 ± 0.03	31.5	
			CP	2.44 ± 0.20	13.9	13.6
<i>Pooled^a L1, 9, 10, 11</i>				<i>Vehicle</i>	<i>0.10 ± 0.02</i>	<i>80.5</i>
			<i>CP</i>	<i>2.44 ± 0.21</i>	<i>29.1</i>	<i>24.4</i>
L1	MeOH-AO	PB	Vehicle	0.05 ± 0.03	100.0	
			CP	1.77 ± 0.17	16.6	35.4
L9	MeOH-AO	PB	Vehicle	0.05 ± 0.00	0.0	
			CP	0.50 ± 0.03	10.0	10.0
L11	MeOH-AO	PB	Vehicle	0.18 ± 0.04	41.7	
			CP	1.42 ± 0.10	12.3	7.9
<i>Pooled L1, 9, 11</i>				<i>Vehicle</i>	<i>0.09 ± 0.03</i>	<i>85.6</i>
			<i>CP</i>	<i>1.23 ± 0.20</i>	<i>48.2</i>	<i>13.7</i>
L5	SV-AO	PB	Vehicle	0.13 ± 0.03	43.3	
			CP	1.83 ± 0.15	13.7	14.1
L6	SV-AO	PB	Vehicle	0.12 ± 0.07	99.0	
			CP	1.77 ± 0.32	31.2	14.8
L7	SV-AO	PB	Vehicle	0.22 ± 0.14	113.8	
			CP	1.47 ± 0.27	31.7	6.7
<i>Pooled L5, 6, 7</i>				<i>Vehicle</i>	<i>0.16 ± 0.05</i>	<i>94.3</i>
			<i>CP</i>	<i>1.69 ± 0.14</i>	<i>24.7</i>	<i>10.6</i>
L1	FCM	PB	Vehicle	0.12 ± 0.02	24.8	
			CP	0.99 ± 0.04	6.5	8.3
L2	FCM	PB	Vehicle	0.11 ± 0.02	31.5	
			CP	1.04 ± 0.04	6.7	9.5
L3	FCM	PB	Vehicle	0.11 ± 0.02	32.9	
			CP	1.11 ± 0.04	6.8	10.1
<i>Pooled L1, 2, 3</i>				<i>Vehicle</i>	<i>0.11 ± 0.01</i>	<i>26.5</i>
			<i>CP</i>	<i>1.05 ± 0.03</i>	<i>7.6</i>	<i>9.5</i>

Note. Abbreviations: MN-RET = micronucleated reticulocyte; MeOH-AO = acridine orange staining of methanol-fixed smears; SV-AO = supravital staining using acridine orange-coated slides; FCM = flow cytometry; BM = bone marrow; PB = peripheral blood; CP = cyclophosphamide; SEM = standard error of the mean; %CV = percent coefficient of variance.

^aValues are the mean of three separately coded, but identical, samples. By "Pooled" it is meant that like-method data from three or four laboratories were combined for these calculations.

Most laboratories detected a reduction in %RET for the CP-treated rat (see Table 2). However, this was somewhat variable across microscopy-based laboratories, especially when the MeOH-AO technique was used to evaluate bone marrow specimens. In two of the three laboratories that scored both bone marrow and peripheral blood, peripheral blood measurements demonstrated greater CP-associated reduction of %RETs than in bone marrow. Intra- and interlaboratory %CV values for the replicate RET analyses are presented in Table 2. Flow cytometric measurements were more consistent within and across laboratories than microscopic scoring. For instance, vehicle control specimens' %CV for pooled laboratory MeOH-AO/bone marrow data was 13.6%, while the corresponding

blood-based analyses for flow cytometric, SV-AO, and MeOH-AO techniques were 1.93, 6.5, and 12.5%, respectively.

The interlaboratory %CV values for MN-RET determinations and the intralaboratory %CV values for the triplicate blinded analyses conducted within each laboratory are provided in Table 3. The flow cytometric analyses demonstrate superior intra- and interlaboratory consistency relative to both microscopy-based methods. %CV values for MN-RET measurements performed on vehicle control blood specimens pooled across like-method laboratories were 26.5, 94.3, and 85.6% for the flow cytometric, SV-AO, and MeOH-AO methods, respectively, and 80.5% for MeOH-AO scored bone marrow. Analogous %CV values for CP blood samples were 7.6, 24.7,




FIG. 3. Values are the mean of three identical, but separately coded, samples. Panel (a): The average frequency of bone marrow RETs (%RET) as measured by the standard MeOH-AO microscopy technique are graphed (with standard error of mean [SEM] bars). These data were collected on two separate occasions at each laboratory. Panel (b): The average frequency of peripheral blood RET as measured by the flow cytometric (FCM) technique are graphed (with SEM bars). These data were collected on three or four separate occasions.




FIG. 4. Values are the mean of three identical, but separately coded, samples. Panel (a): The average frequency of bone marrow micronucleated RETs (%MN-RET) as measured by the standard MeOH-AO microscopy technique are graphed (with standard error of mean [SEM] bars). These data were collected on two separate occasions at each laboratory. Panel (b): The average frequency of peripheral blood MN-RET as measured by the flow cytometric (FCM) technique are graphed (with SEM bars). These data were collected on three or four separate occasions.

and 48.2%, respectively, and 29.1% for MeOH-AO scored bone marrow.

Fold difference values based on each laboratory's average MN-RET frequencies, as well as for like-method pooled data, are also presented (Table 3). It was somewhat surprising that the fold difference in MN-RETs between vehicle and CP-associated blood specimens, as well as absolute MN-RET frequencies, were no higher with the flow cytometric or SV-AO techniques than with the conventional MeOH-AO method as it has been reported that restriction of MN analysis to an immature RET cohort based on RNA content or CD71 expression levels could reduce, if not eliminate, the influence of the spleen's erythrophagocytotic activity (Abramsson-Zetterberg *et al.*, 1999; Hayashi *et al.*, 1992). Splenic activity and its effects on assay sensitivity for blood-based analyses have been investigated thoroughly, and these data are discussed in a companion paper that appears in this issue (MacGregor *et al.*).

Intralaboratory Variability Across Time

In addition to the inter- and intralaboratory analyses, an evaluation of scoring reproducibility over time was studied. This was accomplished by having flow cytometry laboratories analyze coded peripheral blood specimens on three or four different occasions, while triplicate vehicle and CP bone marrow slides were submitted to L9, L10, and L11 laboratories for analysis on two separate occasions. Reagents were prepared separately for each day of analysis. The resulting repeat-analysis RET data are presented in Figure 3 and demonstrate higher reproducibility for the multiple flow cytometric analyses compared to MeOH-AO.

As with RET enumeration, repeat-analysis MN-RET microscopy data were also quite variable. For instance, laboratories using the MeOH-AO method reported average CP-induced values that differed from their original mean reading by 19.4,

50.9, 58.4, and 8.6% (L1, L9, L10, and L11, respectively; see Fig. 4). Repeat-analysis MN-RET data generated by the flow cytometric technique were considerably more reproducible as average values were all within 11% of the originally reported mean frequencies. It should also be noted that the fourth flow cytometric analyses by L2 was performed more than 2 years after blood fixation, demonstrating this procedure's compatibility with long-term storage of fixed blood specimens.

CONCLUSIONS

Distribution of replicate bone marrow and blood specimens obtained from single rats that were first shown to exhibit steady-state spontaneous or genotoxicant-induced MN-RET frequencies were used to assess inter- and intralaboratory scoring variability using two widely used microscopic and one flow cytometric procedure. These results demonstrate that the quantification of MN-RETs benefits from an objective flow cytometry-based method of data acquisition. The flow cytometric method provides better reproducibility, and the high throughput capability allows interrogation of tens of thousands of RETs per specimen. Enhanced scoring precision is important as it is necessary to offset the spleen-dependent loss of dynamic range observed in peripheral blood relative to bone marrow—a phenomenon that was observed in this as well as other reports (MacGregor *et al.*, this issue; Wakata *et al.*, 1998). A recent report by Torous *et al.* (2006) delineates the consequential improvements to assay power when the number of cells scored per specimen is increased and supports this view.

Beyond overcoming lower genotoxicant-induced MN-RET frequencies in blood relative to bone marrow, further incentive for automating rat peripheral blood MN-RET measurements comes from a recent recommendation of the *In Vivo* MN Assay Expert Group of the International Working Group on Genetic Toxicology Testing (IWGT; Hayashi *et al.*, in press). Specifically, IWGT has recommended that a sufficient number of RETs should be scored to ensure that the MN-RET counting error is kept below the level of interanimal variability. This allows the sensitivity of the experiment to be limited by the variability of spontaneous MN-RET frequency among animals, rather than being limited by the statistical variation of count. Based on the flow cytometric scoring of 20,000 peripheral blood RETs from each of the 15 control animals from the three experiments reported in the MacGregor *et al.* companion paper in this issue (laboratory L2, the reference laboratory), we find that the mean incidence of MN-RET ± 1 SD is $0.11\% \pm 0.045$. This is a 41% CV. Poisson distribution theory allows us to calculate that 6 MN-RETs per animal must be scored to limit counting error to this level of variation (SD of the Poisson count = $\sqrt{\text{absolute count}}$). At a spontaneous MN-RET frequency of approximately 0.1%, this means that an average of 6000 RETs per individual need to be scored for

micronuclei in order to achieve a CV that is at or below the interanimal variance. This is a significantly higher number of RETs per animal than required to be scored under the current OECD MN assay guideline (which recommends scoring 2000 RETs per animal) and is difficult to achieve by manual microscopic scoring.

In conclusion, the data presented herein and in the companion paper that follows support the growing consensus that rat peripheral blood can be used to perform *in vivo* MN tests more effectively than the standard bone marrow-based assay. The ability of the described automated scoring procedure to greatly enhance the precision of MN-RET measurements overcomes the somewhat attenuated genotoxicant-induced frequencies observed in peripheral blood relative to bone marrow. This conclusion is supported by experiments described in the accompanying paper whereby intact and splenectomized rats were treated with diverse genotoxicants (MacGregor *et al.*, this issue).

ACKNOWLEDGMENTS

This work was supported by an intercenter research award from the Food and Drug Administration Office of Science and Health Coordination to J.T.M. The contents are the sole responsibility of the authors and do not necessarily represent the official views of the U.S. Food and Drug Administration, the National Institute of Health Sciences, Japan, or Health Canada. We would like to thank Nikki Hall for her expert assistance with bleeding and dosing procedures. Thanks also to Puntipa Kwanyuen and Dr Neal Cariello, two researchers who were early adopters of the flow cytometric technique described herein. Their experiences with an early version of the method led to the development of malaria-infected erythrocytes as daily calibration standards. Conflict of interest disclosure: Litron Laboratories holds patents pertaining to flow cytometric enumeration of micronucleated erythrocyte populations and also sells kits that facilitate these analyses. The authors express their appreciation for the cooperation and contributions of the staff of the 3 contract laboratories that participated in these studies: Drs Robert Young and Rama Gudi of BioReliance, Rockville, MD; Dr Gregory Erexson of Covance, Inc., Vienna, VA; and Dr John Mirsalis and Edward Riccio of SRI International, Menlo Park, CA.

REFERENCES

- Abramsson-Zetterberg, L., Grawe, J., and Zetterberg, G. (1999). The micronucleus test in rat erythrocytes from bone marrow, spleen and peripheral blood: The response to low doses of ionizing radiation, cyclophosphamide and vincristine determined by flow cytometry. *Mutat. Res.* **423**, 113–124.
- Asanami, S., Shimono, K., Sawamoto, O., Kurisu, K., and Uejima, M. (1995). The suitability of rat peripheral blood in subchronic studies for the micronucleus assay. *Mutat. Res.* **347**, 73–78.
- Dertinger, S. D., Camphausen, K., MacGregor, J. T., Bishop, M. E., Torous, D. K., Avlasevich, S., Cairns, S., Tometsko, C. R., Menard, C., Muanza, T., *et al.* (2004). Three-color labeling method for flow cytometric measurement of cytogenetic damage in rodent and human blood. *Environ. Mol. Mutagen.* **44**, 427–435.
- Dertinger, S. D., Torous, D. K., Hall, N. E., Tometsko, C. R., and Gasiewicz, T. A. (2000). Malaria-infected erythrocytes serve as biological standards to ensure reliable and consistent scoring of micronucleated erythrocytes by flow cytometry. *Mutat. Res.* **464**, 195–200.

- Hamada, S., Sutou, S., Morita, T., Wakata, A., Asanami, S., Hosoya, S., Ozawa, S., Kondo, K., Nakajima, M., Shimada, H., *et al.* (2001). Evaluation of the rodent micronucleus assay by a 28-day treatment protocol: Summary of the 13th collaborative study by the collaborative study group for the micronucleus test (CSGMT)/Environmental Mutagen Society of Japan (JEMS)—Mammalian Mutagenicity Study Group (MMS). *Environ. Mol. Mutagen.* **37**, 93–110.
- Hayashi, M., Kodama, Y., Awogi, T., Suzuki, T., Asita, A. O., and Sufuni, T. (1992). The micronucleus assay using peripheral blood reticulocytes from mitomycin C- and cyclophosphamide-treated rats. *Mutat. Res.* **278**, 209–213.
- Hayashi, M., MacGregor, J. T., Gatehouse, D. G., Adler, I.-D., Blakey, D. H., Dertinger, S. D., Krishna, G., Morita, T., Russo, A., and Sutou, S. (2000). *In vivo* rodent erythrocyte micronucleus assay: Aspects of protocol design including repeated treatments, integration with toxicity testing, and automated scoring. A report from the International Workshop on Genotoxicity Test Procedures (IWGTP). *Environ. Mol. Mutagen.* **35**, 234–252.
- Hayashi, M., MacGregor, J. T., Gatehouse, D. G., Blakey, D. H., Dertinger, S. D., Abramson-Zetterberg, L., Krishna, G., Morita, T., Russo, A., Asano, N., *et al.* *In vivo* erythrocyte micronucleus assay: III. Validation and regulatory acceptance of automated scoring and the use of rat peripheral blood reticulocytes, with discussion of non-hematopoietic target cells and a single dose-level limit test. *Environ. Mol. Mutagen.* (in press).
- Hayashi, M., Morita, T., Kodama, Y., Sofuni, T., and Ishidate, M., Jr. (1990). The micronucleus assay with mouse peripheral blood reticulocytes using acridine orange-coated slides. *Mutat. Res.* **245**, 245–249.
- Hynes, G. M., Torous, D. K., Tometsko, C. R., Burlingson, B., and Gatehouse, D. G. (2002). The single laser flow cytometric micronucleus test: A time course study using colchicines and urethane in rat and mouse peripheral blood and acetaldehyde in rat peripheral blood. *Mutagenesis* **17**, 15–23.
- MacGregor, J. T., Tucker, J. D., Eastmond, D. A., and Wyrobek, A. J. (1995). Integration of cytogenetic assays with toxicology studies. *Environ. Mol. Mutagen.* **25**, 328–337.
- MacGregor, J. T., Wehr, C. M., Hiatt, R. A., Peters, B., Tucker, J. D., Langlois, R. G., Jacob, R. A., Jenson, R. H., Yager, J. W., Shigenaga, M. K., *et al.* (1997). "Spontaneous" genetic damage in man: Evaluation of interindividual variability, relationship among markers of damage, and influence of nutritional status. *Mutat. Res.* **377**, 125–135.
- Organisation for Economic Cooperation and Development (OECD) (1997). OECD Guideline 474. Guideline for the testing of chemicals. Mammalian erythrocyte micronucleus test.
- Romagna, F., and Staniforth, C. D. (1989). The automated bone marrow micronucleus test. *Mutat. Res.* **213**, 91–104.
- Schmid, W. (1976). The micronucleus test for cytogenetic analysis. In *Chemical Mutagens: Principles and Methods for their Detection*, A. Hollaender, (Ed.) Vol. 4, pp. 31–53. Plenum Press, New York.
- Tometsko, A. M., Torous, D. K., and Dertinger, S. D. (1993). Analysis of micronucleated cells by flow cytometry. 1. Achieving high resolution with a malaria model. *Mutat. Res.* **292**, 129–135.
- Torous, D., Asano, N., Tometsko, C., Sugunan, S., Dertinger, S., Morita, T., and Hayashi, M. (2006). Performance of flow cytometric analysis for the micronucleus assay—A reconstruction model using serial dilutions of malaria infected cells with normal mouse peripheral blood. *Mutagenesis* **21**, 11–13.
- Torous, D. K., Dertinger, S., Hall, N., and Tometsko, C. (2000). Enumeration of micronucleated reticulocytes in rat peripheral blood: A flow cytometric study. *Mutat. Res.* **465**, 91–99.
- Torous, D. K., Hall, N. E., Dertinger, S. D., Diehl, M., Illi-Love, A. H., Cederbrant, K., Sandelin, K., Bolcsfoldi, G., Ferguson, L. R., Pearson, A., *et al.* (2001). Flow cytometric enumeration of micronucleated reticulocytes: High transferability among 14 laboratories. *Environ. Mol. Mutagen.* **38**, 59–68.
- Torous, D. K., Hall, N. E., Murante, F. G., Gleason, S. E., Tometsko, C. R., and Dertinger, S. D. (2003). Comparative scoring of micronucleated reticulocytes in rat peripheral blood by flow cytometry and microscopy. *Toxicol. Sci.* **74**, 309–314.
- U.S. Food and Drug Administration (2000) Office of Food Additive Safety, Redbook (2000). Toxicological Principals for the Safety Assessment of Food Ingredients. Available at: <http://www.cfsan.fda.gov/~redbook/red-toca.html>. Updated on November 2003. U.S. Food and Drug Administration, Washington, D.C.
- Wakata, A., Miyamae, Y., Sato, S., Suzuki, T., Morita, T., Asano, N., Awogi, T., Kondo, K., and Hayashi, M. (1998). Evaluation of the rat micronucleus test with bone marrow and peripheral blood: Summary of the 9th collaborative study by CSGMT/JEMS.MMS. *Environ. Mol. Mutagen.* **32**, 84–100.

Commentary

Globally Harmonized System on Hazard Classification and Labeling of Chemicals and Other Existing Classification Systems for Germ Cell Mutagens

Takeshi Morita^{1,3}, Makoto Hayashi² and Kaoru Morikawa¹

¹Division of Safety Information on Drug, Food and Chemicals,

²Division of Genetics and Mutagenesis, National Institute of Health Sciences, Tokyo, Japan

(Received August 8, 2006; Revised August 26, 2006; Accepted August 28, 2006)

The Globally Harmonized System (GHS) on hazard classification and labeling of chemicals will be implemented globally by 2008. The GHS includes (a) harmonized criteria for classifying chemicals and chemical mixtures according to their health, environmental and physical hazards, and (b) harmonized hazard communication elements, including requirements for labeling and safety data sheets. Germ cell mutagenicity is included in the GHS health hazard classes in addition to carcinogenicity. This means increased significance for then results of genetic toxicology testing for the classification of chemicals. GHS requires the classification of chemicals if they are germ cell mutagens (categories 1A, 1B and 2) or not. Several classification systems for germ cell mutagens have been proposed in the EU, Germany, US, Canada, in advance of the adoption of the GHS. In this paper, these classification systems including GHS are introduced and summarized to provide the basis of the hazard classification of germ cell mutagens. Though the objectives, target audiences and criteria of these classification systems are different, the GHS will become standard for hazard classification. Hazard classification is a significant first step in risk communication. Further development of risk evaluation criteria and communication on germ cell mutagens is expected.

Key words: GHS, hazard classification, germ cell mutagenicity, germ cell mutagens

Introduction

The Globally Harmonized System (GHS) of classification and labeling of chemicals is a single, globally harmonized system to address classification of chemicals, labelling, and safety data sheets, which has been developed by the United Nations (UN). The GHS document has been prepared and published by the secretariat of the United Nations Economic Commission for Europe (1,2). The GHS covers all hazardous chemicals except for pharmaceuticals, food additives, cosmetics, and pesticide residues in food in terms of labeling at the point of intentional intake. The GHS is

based on currently available data and thus compliance with these criteria will not require retesting of chemicals for which acceptable test data already exists.

The goal of the GHS is to identify the intrinsic hazards found in chemicals and chemical mixtures and to convey the information about these hazards to the target audiences including consumers, workers, transport workers, and emergency responders. The World Summit on Sustainable Development in Johannesburg on 4th September, 2002 encouraged countries to implement the GHS as soon as possible with a view to having the system fully operational by 2008. In Japan, an inter-ministerial committee was organized in 2001 to share information about the GHS among ministries and to play a pivotal role in the UN-Subcommittee. Seven government offices, *i.e.*, Ministry of Health, Labour and Welfare (MHLW), Ministry of Economy, Trade and Industry (METI), Ministry of the Environment (MOE), Ministry of Internal Affairs and Communications (MIC), Ministry of Agriculture, Forestry and Fisheries (MAFF), Ministries of Land Infrastructure and Transport (MLIT) and Ministry of Foreign Affairs (MOFA), and experts from national laboratories and industries participated in the committee. The committee's activities include: (i) translation of the GHS into Japanese (3); (ii) information sharing among ministries with respect to the relevant domestic laws; (iii) classification of chemicals under each relevant domestic law (4); and (iv) deliberation on the agenda items and documents of the UN Sub-Committee meetings and decision-making about the Japanese position.

The GHS includes the following two elements: harmonized criteria for classifying substances and mixtures according to their health (10 hazard classes),

³Correspondence to: Takeshi Morita, Division of Safety Information on Drug, Food and Chemicals, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan. Tel: +81-3-3700-1141, Fax: +81-3-3700-1483, E-mail: morita-tk@nihs.go.jp

environmental (1 hazard class) and physical hazards (16 hazard classes); and harmonized hazard communication elements, including requirements for labeling and safety data sheets. The GHS requests classification of the following 10 hazard classes in health hazard, *i.e.*, (i) acute toxicity, (ii) skin corrosion/irritation, (iii) serious eye damage/eye irritation, (iv) respiratory or skin sensitization, (v) germ cell mutagenicity, (vi) carcinogenicity, (vii) reproductive toxicity, (viii) specific target organ systemic toxicity—single exposure, (ix) specific target organ systemic toxicity—repeated exposure, and (x) aspiration hazard.

It is striking that not mutagenicity per se but germ cell mutagenicity specifically is included in GHS health hazard class in addition to carcinogenicity. The GHS focuses on heritable effects by mutagens. Several classification systems of mutagens or germ cell mutagens have been proposed from European Union (EU), Germany, United States of America (US), Canada, etc. The criteria for germ cell mutagens of GHS and other systems are reviewed and summarized to aid understanding of the control of chemicals by these regulations.

Classification Systems for Germ Cell Mutagens

Classification systems for mutagens or germ cell mutagens in GHS, EU, Germany, USA, Canada, and Japan are described below.

GHS: In the GHS (2), the term "mutation" applies both to heritable genetic changes that may be manifested at the phenotypic level and to the underlying DNA modifications when known (including, for example, specific base pair changes and chromosomal translocations). The term "mutagenic" and "mutagen" will be used for chemicals giving rise to an increased occurrence of mutations in populations of cells and/or organisms. The more general terms "genotoxic" and "genotoxicity" apply to chemicals or processes which alter the structure, information content, or segregation of DNA, including those which cause DNA damage by interfering with normal replication processes, or which in a non-physiological manner (temporarily) alter its replication.

In order to achieve classification, GHS states that "Test results are considered from experiments determining mutagenic and/or genotoxic effects in germ and/or somatic cells of exposed animals. The system is hazard based, classifying chemicals on the basis of their intrinsic ability to induce mutations in germ cells. The scheme is, therefore, not meant for the (quantitative) risk assessment of chemical substances. Classification for heritable effects in human germ cells is made on the basis of well conducted, sufficiently validated tests, preferably as described in OECD Test Guidelines. Evaluation of the test results should be done using expert judgment and all the available evidence should be

weighed for classification. The classification of individual substances should be based on the total weight of evidence available, using expert judgment. In those instances where a single well-conducted test is used for classification, it should provide clear and unambiguously positive results. If new, well validated, tests arise these may also be used in the total weight of evidence to be considered. The relevance of the route of exposure used in the study of the chemical compared to the route of human exposure should also be taken into account."

The criteria for classification of germ cell mutagens places chemicals in one of three categories, category 1 being used for chemicals known to induce heritable mutations (category 1A) or known to be regarded as if they induce heritable mutations in germ cells of humans (category 1B); category 2 for chemicals which cause concern for humans owing to the possibility that they may induce heritable mutations in the germ cells of humans (2). The criterion for category 1A is positive evidence from human epidemiological studies. The criteria for category 1B are as follows: (i) positive result(s) from *in vivo* heritable germ cell mutagenicity tests in mammals; or (ii) positive result(s) from *in vivo* somatic cell mutagenicity tests in mammals, in combination with some evidence that the substance has the potential to cause mutations to germ cells. This supporting evidence may, for example, be derived from mutagenicity/genotoxic tests in germ cells *in vivo*, or by demonstrating the ability of the substance or its metabolite(s) to interact with the genetic material of germ cells; or (iii) positive results from tests showing mutagenic effects in the germ cells of humans, without demonstration of transmission to progeny; for example, an increase in the frequency of aneuploidy in the sperm cells of exposed people. The criteria for category 2 are positive evidence obtained from experiments in mammals and/or in some cases from *in vitro* experiments, obtained from: (i) somatic cell mutagenicity tests *in vivo*, in mammals; or (ii) other *in vivo* somatic cell genotoxicity tests which are supported by positive results from *in vitro* mutagenicity assays. In addition, following criteria are included as *Note*: Chemicals which are positive in *in vitro* mammalian mutagenicity assays, and which also show chemical structure activity relationship to known germ cell mutagens, should be considered for classification as Category 2 mutagens.

If there are not enough data for the evaluation of mutagenicity of the chemical, it regards as "classification not possible" (Fig. 1). If there is no concern of induction of heritable mutations in the germ cells of humans or insufficient evidence for inclusion in category 1 or 2, the chemicals are regarded as "not classified". Hazard categories and their criteria for germ cell mutagens in GHS are summarized in Table 1. For classification of chemical mixtures, the mixture will be




Fig. 1. Decision logic for the classification of germ cell mutagens in GHS (2)

classified as a mutagen when at least one ingredient has been classified as a category 1 or category 2 mutagen and is present at or above the cut-off value/concentration limits below for category 1 ($\geq 0.1\%$) or category 2 ($\geq 1.0\%$), respectively.

European Union (EU): The criteria for classification of mutagens in EU are described in Commission directive (5, 6). There are three categories: substances known to be mutagenic to human (category 1) for which there is sufficient evidence to establish a causal association between human exposure to the chemical and heritable genetic damage; substances which should be regarded as if they were mutagenic to human (category 2) for which there is sufficient evidence to provide a strong presumption that human exposure to the chemical may result in the development of heritable genetic damage, generally on the basis of appropriate animal studies and other relevant information; substances which cause concern for humans owing to possible mutagenic effect (category 3) for which there is evidence from appropriate mutagenicity studies, but it is insufficient to place the substance in category 2. EU criteria for classification of chemicals are summarized in Table 2.

This system is primarily based on intrinsic hazard, despite the statement in the Annex (7) that 'the object of classification is to identify all the physicochemical, toxicological and ecotoxicological properties of sub-

stances and preparations which may constitute a risk during normal handling or use' (4).

Germany: Maximale Arbeitsplatz-Konzentration (MAK) Commission in Germany proposed 5 categories for classification of germ cell mutagens at the workplace (8, 9). These are germ cell mutagens which have been shown to increase the mutant frequency in the progeny of exposed humans (category 1); germ cell mutagens which have been shown to increase the mutant frequency in the progeny of exposed mammals (category 2); chemicals which have been shown to induce genetic damage in germ cells of humans and/or animals, or which produce mutagenic effects in somatic cells of mammals *in vivo* and the chemicals have been shown to reach the germ cells in an active form (category 3A); chemicals which are suspected of being germ cell mutagens because of their genotoxic effects in mammalian somatic cells *in vivo*; in exceptional cases, chemicals for which there are no *in vivo* data but which are clearly mutagenic *in vitro* and structurally related to known *in vivo* mutagens (category 3B); and germ cell mutagens, the potency of which is considered to be so low that, provided the MAK value (Maximum Concentration at the Workplace) is observed, their contribution to genetic risk for man is expected not to be significant (category 5). Category 4 is not applicable in germ cell mutagenicity because this classification system has been

Table 1. GHS classification on germ cell mutagens (1,2)

Category	Classification	Criteria
Category 1A	Chemicals known to induce heritable mutations in germ cells of humans	Positive evidence from human epidemiological studies.
Category 1B	Chemicals which should be regarded as if they induce heritable mutations in the germ cells of humans	<p>—Positive result(s) from <i>in vivo</i> heritable germ cell mutagenicity tests in mammals; or</p> <p>—Positive result(s) from <i>in vivo</i> somatic cell mutagenicity tests in mammals, in combination with some evidence that the substance has potential to cause mutations to germ cells. This supporting evidence may, for example, be derived from mutagenicity/genotoxic tests in germ cells <i>in vivo</i>, or by demonstrating the ability of the substance or its metabolite(s) to interact with the genetic material of germ cells; or</p> <p>—Positive results from tests showing mutagenic effects in the germ cells of humans, without demonstration of transmission to progeny; for example, an increase in the frequency of aneuploidy in sperm cells of exposed people.</p> <p>Examples of <i>in vivo</i> heritable germ cell mutagenicity tests are:</p> <ul style="list-style-type: none"> — Rodent dominant lethal mutation test — Mouse heritable translocation assay — Mouse specific locus test <p>Examples of <i>in vivo</i> somatic cell mutagenicity test are:</p> <ul style="list-style-type: none"> — Mammalian bone marrow chromosome aberration test — Mouse spot test — Mammalian erythrocyte micronucleus test <p>Examples of mutagenicity/genotoxicity tests in germ cells are:</p> <p>(a) Mutagenicity tests:</p> <ul style="list-style-type: none"> — Mammalian spermatogonial chromosome aberration test — Spermatid micronucleus assay <p>(b) Genotoxicity tests:</p> <ul style="list-style-type: none"> — Sister chromatid exchange analysis in spermatogonia — Unscheduled DNA synthesis test (UDS) in testicular cells
Category 2	Chemicals which cause concern for humans owing to the possibility that they may induce heritable mutations in the germ cells of humans	<p>—Positive evidence obtained from experiments in mammals and/or in some cases from <i>in vitro</i> experiments, obtained from:</p> <p>—Somatic cell mutagenicity tests <i>in vivo</i>, in mammals; or</p> <p>—Other <i>in vivo</i> somatic cell genotoxicity tests which are supported by positive results from <i>in vitro</i> mutagenicity assays.</p> <p>Examples of genotoxicity tests in somatic cells are:</p> <ul style="list-style-type: none"> — Liver UDS <i>in vivo</i> — Mammalian bone marrow sister chromatid exchanges <p>Examples of <i>in vitro</i> mutagenicity tests are:</p> <ul style="list-style-type: none"> — <i>In vitro</i> mammalian chromosome aberration test — <i>In vitro</i> mammalian cell gene mutation test — Bacterial reverse mutation tests <p>Note: Chemicals which are positive in <i>in vitro</i> mammalian mutagenicity assays, and which also show chemical structure activity relationship to known germ cell mutagens, should be considered for classification as Category 2 mutagens.</p>

established in analogy to the categories for carcinogenic chemicals. Category 4 carcinogenic substances are those with non-genotoxic mechanisms of action. By definition, germ cell mutagens are genotoxic. MAK categories and criteria for classification of germ cell mutagens are summarized in Table 3.

The MAK Commission describes germ cell mutagenicity as follows (9): 'Germ cell mutagens produce heritable gene mutations, and heritable structural and numerical chromosome aberrations in germ cells. The

consequences of germ cell mutations in subsequent generations include genetically determined phenotypic alterations without signs of illness, reduction in fertility, embryonic or perinatal death, more or less severe congenital malformations, and genetic diseases with various degrees of health impairment. The term "germ cell mutagenicity" refers specifically to mutagenicity in male and female germ cells and is distinguished from mutagenicity in somatic cells, which can initiate cancer. Epidemiological studies, however, have been unable to

Table 2. EU criteria for classification of chemicals as mutagenic (6,7)

Category	Classification	Criteria
Category 1	Substances known to be mutagenic to human	Positive evidence from human mutation epidemiology studies will be needed. Examples of such substances are not known to date.
Category 2	Substances which should be regarded as if they are mutagenic to human	<p>Positive results from assays showing (a) mutagenic effects, or (b) other cellular interactions relevant to mutagenicity, in germ cells of mammals <i>in vivo</i>, or (c) mutagenic effects in somatic cells of mammals <i>in vivo</i> in combination with clear evidence that the substance or a relevant metabolite reaches the germ cells.</p> <p>With respect to placement in category 2, at present the following methods are appropriate: 2(a) <i>In vivo</i> germ cell mutagenicity assays:</p> <ul style="list-style-type: none"> — specific locus mutation test, — heritable translocation test, — dominant lethal mutation test. <p>These assays actually demonstrate the appearance of affected progeny or a defect in the developing embryo.</p> <p>2(b) <i>In vivo</i> assays showing relevant interaction with germ cells (usually DNA):</p> <ul style="list-style-type: none"> — assays for chromosomal abnormalities, as detected by cytogenetic analysis, including aneuploidy, caused by malsegregation of chromosomes, — test for sister chromatid exchanges (SCEs), — test for unscheduled DNA synthesis (UDS), — assay of (covalent) binding of mutagen to germ cell DNA, — assaying other kinds of DNA damage. <p>These assays provide evidence of a more or less indirect nature. Positive results in these assays would normally be supported by positive results from <i>in vivo</i> somatic cell mutagenicity assays, in mammals or in man.</p> <p>2(c) <i>In vivo</i> assays showing mutagenic effects in somatic cells of mammals, in combination with toxicokinetic methods, or other methodologies capable of demonstrating that the compound or a relevant metabolite reaches the germ cells.</p> <p>For 2(b) and 2(c), positive results from host-mediated assays or the demonstration of unequivocal effects in <i>in vitro</i> assays can be considered as supporting evidence.</p>
Category 3	Substances which cause concern for human owing to possible mutagenic effects.	<p>There is evidence from appropriate mutagenicity studies, but this is insufficient to place the substance in category 2.</p> <p>Assays showing (a) mutagenic effects or (b) other cellular interaction relevant to mutagenicity, in somatic cells in mammals <i>in vivo</i>. The latter would be supported by positive results from <i>in vitro</i> mutagenicity assays.</p> <p>For effects in somatic cells <i>in vivo</i> at present the following methods are appropriate:</p> <p>3(a) <i>In vivo</i> somatic cell mutagenicity assays:</p> <ul style="list-style-type: none"> — bone marrow micronucleus test or metaphase analysis, — metaphase analysis of peripheral lymphocytes, — mouse coat color spot test. <p>3(b) <i>In vivo</i> somatic cell DNA interaction assays:</p> <ul style="list-style-type: none"> — test for SCEs in somatic cells, — test for UDS in somatic cells, — assay for the (covalent) binding of mutagen to somatic cell DNA, — assay for DNA damage, e.g. by alkaline elution, in somatic cells. <p>Substances showing positive results only in one or more <i>in vitro</i> mutagenicity assays should normally not be classified. Their further investigation using <i>in vivo</i> assays, however, is strongly indicated. In exceptional cases, e.g., for a substance showing pronounced responses in several <i>in vitro</i> assays, for which no relevant <i>in vivo</i> data are available, and which shows resemblance to known mutagens/carcinogens, classification in category 3 could be considered.</p>

provide any evidence as yet that exposure to chemicals or to radiation results in hereditary diseases in man. Although structural changes have been demonstrated in the chromosomes of the germ cells of men exposed to radiation, even this finding can only provide indirect evidence that such exposures could lead to hereditary disorders in the offspring. The proof that an increased frequency of hereditary diseases is related to a particular

exposure would be associated with great methodological difficulties. In the human population there are a large number of hereditary diseases of unknown origin with frequencies that differ widely in different subpopulations. Since mutational events occur largely randomly in the genome, it is not to be expected that one particular substance would induce one characteristic genetic disease. Therefore, it is most unlikely that proof of a

Table 3. Categories for classification of germ cell mutagens by MAK commission (8,9)

Category	Classification	Criteria
Category 1	Substances shown to increase the mutant frequency in the progeny of exposed humans	In the section 'Epidemiological methods and their limitations' it is explained why epidemiological studies to date have not been able to prove that the exposure of a particular human population to a particular substance has resulted in an increase in the incidence of inherited mutations. This is true both for ionizing radiation and chemical mutagens. Even if epidemiological methods are improved further, it is unlikely that such proof will be available in the foreseeable future. Category 1 will therefore probably remain without any entries.
Category 2	Substances shown to increase the mutant frequency in the progeny of exposed mammals	Classified as category 2 are substances that increase the incidence of genetically modified live progeny in animal studies, for example in the specific locus test or in the test for heritable translocations. Likewise, substances that should be classified as category 2 are those that increase the incidence of embryos that die in utero, for example in the dominant lethal test.
Category 3A	Substances shown to induce genetic damage in germ cells of humans or animals, or which produce mutagenic effects in somatic cells of mammals <i>in vivo</i> and shown to reach the germ cells in an active form	The methods include tests for genotoxicity in germ cells of experimental animals, such as tests for induction of structural chromosomal changes in spermatogonia or spermatocytes, for sister chromatid exchange in spermatogonia, for micronuclei in round spermatids, for numerical chromosome changes in secondary spermatocytes or in spermatozoa, for DNA single strand breaks and for repair synthesis or for covalent binding to the DNA. Also relevant are the observations obtained from exposed human populations which provide evidence for structural or numerical chromosome changes in spermatozoa of exposed persons. The development of new methods, especially molecular genetic methods for the detection of gene mutations in germ cells is to be expected. Substances that yield positive results in tests with germ cells are classified as category 3A. Also taken into account are clearly positive results from <i>in vivo</i> tests for mutagenicity in somatic cells, for example chromosomal aberrations or micronuclei in bone marrow cells, somatic mutations in the mammalian spot test or transgenic animals, provided that it has been demonstrated that the active substance or an active metabolite reaches the germ cells after relevant exposure of the experimental animals. Such substances are also suspected of being mutagenic in germ cells. Therefore they are classified as category 3A.
Category 3B	Substances suspected of germ cell mutagens because of their genotoxic effects in mammalian somatic cells <i>in vivo</i> ; in exceptional cases, substances without <i>in vivo</i> data but with clearly mutagenic <i>in vitro</i> and structurally related to known <i>in vivo</i> mutagens	If the available data are not sufficient for classification in category 3A but the substance is clearly genotoxic in somatic cells of exposed animals or humans, the substance is also suspected of being mutagenic in germ cells. Substances that have yielded positive results in one or several <i>in vitro</i> mutagenicity tests generally not classified as category 3B. An exception is made for substances for which there are no relevant <i>in vivo</i> data but which are clearly genotoxic <i>in vitro</i> and also structurally related to substances known to be genotoxic <i>in vivo</i> . Such substances raise concern and are classified as category 3B.
Category 4	Not applicable	Category 4 carcinogenic substances are those with non-genotoxic mode of action. By definition, germ cell mutagens are genotoxic. Therefore, a category 4 for germ cell mutagens cannot exist. Depending on future research results, a category 4 could be defined at a later time for genotoxic substances with targets other than DNA (<i>i.e.</i> , pure aneugens).
Category 5	Substances considered the potency is considered so low, their contribution to genetic risk for man is expected not to be significant	Substances classified as category 5 are not expected to contribute significantly to the genetic risk for humans provided the MAK value is observed. For classification in this category, information on the spectrum of effects and their dose-dependence, and toxicokinetic data for species comparison are required. Biochemical and biological end-points can be used to characterize the contribution to genetic risk. The contribution to genetic risk is considered not to be significant after exposure at the workplace if the internal exposure level of the substance or its biomarkers is in the range of the background levels in a not specifically exposed reference population: ⊗ Under workplace conditions the levels of biochemical effect markers such as DNA and protein adducts are not significantly increased above the background levels. ⊗ Physiological-toxicokinetic model calculations based on animal data do not reveal a significant genetic risk for humans.

causal relationship between exposure to a chemical and occurrence of heritable diseases will become available in the foreseeable future. In this situation, for the identification of germ cell mutagens the results of animal experiments must be given particular attention. The mutagenic effect of chemicals on the germ cells of exposed parent animals can be demonstrated by observing an increased mutant frequency among the progeny. In addition, the demonstration of genotoxic effects of a substance in germ cells or somatic cells provides evidence of a potential hazard for subsequent generations.'

United States of America (US): US EPA: A classification using the following three categories of germ cell mutagens was proposed by the US Environmental Protection Agency (EPA) in 1984 for the evaluation of chemicals with respect to their ability to induce mutations in mammalian germ cells (10,11). Category I is based on sufficient evidence obtained from at least one *in vivo* mammalian germ cell mutation test or from at least two *in vivo* somatic cell mutation tests (point mutation and/or chromosomal aberrations), plus sufficient *in vivo* evidence that the chemical interacts with mammalian germ cells. Category II is based on suggestive evidence provided from positive results of *in vivo* somatic cell mutation tests plus evidence for interaction of the chemical with mammalian germ cells, but the evidence is insufficient to place the chemical in Category I. Category III is based on limited evidence of *in vivo* mutagenic activity or interaction of the chemical with mammalian germ cell DNA or other chromatin constituents. In 1986, guidelines for a "weight-of-evidence" approach to human germ cell mutagenicity were established, leading to eight categories with a decreasing order of strength of evidence (10). In addition, a "non-mutagen" category and a category for substances with inadequate evidence were described. The eight categories of evidence are as follows: (i) positive data derived from human germ cell mutagenicity studies; (ii) valid positive results from studies on heritable mutational events (of any kind) in mammalian germ cells; (iii) valid positive results from mammalian germ cell chromosome aberration studies that do not involve transmission from one generation to the next; (iv) sufficient evidence for a chemical's interaction with mammalian germ cells, together with valid positive mutagenicity test results from two assay systems, at least one of which is mammalian (*in vivo* or *in vitro*). The positive results may be both for gene mutation and chromosomal aberrations in mammalian systems; (v) suggestive evidence for a chemical's interaction with mammalian germ cells, together with valid positive mutagenicity evidence from two assay systems as described above under #iv. Alternatively, positive mutagenicity evidence of less strength than defined

under #iv, when combined with sufficient evidence for a chemical's interaction with mammalian germ cells; (vi) positive mutagenicity test results of less strength than defined under #iv, combined with suggestive evidence for a chemical's interaction with mammalian germ cells; (vii) although definitive proof of non-mutagenicity is not possible, a chemical could be operationally classified as a non-mutagen for human germ cells if it gives valid negative test results for all endpoints of concern; and (viii) inadequate evidence bearing on either mutagenicity or chemical interaction with mammalian germ cells.

This system is for the classification for transmissible germ cell genetic risk. The Guideline (10) describes 'Evidence that an agent induces heritable mutations in human beings could be derived from epidemiologic data indicating a strong association between chemical exposure and heritable effects. It is difficult to obtain such data because any specific mutation is a rare event, and only a small fraction of the estimated thousands of human genes and conditions are currently useful as markers in estimating mutation rates. Human genetic variability, small numbers of offspring per individual and long generation times further complicate such studies. In addition, only disorders caused by dominant mutations, some sex-linked recessive mutations, and certain chromosome aberrations can be detected in the first generation after their occurrence. Conditions caused by autosomal recessive disorders (which appear to occur more frequently than dominant disorders) or by polygenic traits may go unrecognized for many generations. Therefore, in the absence of human epidemiological data, it is appropriate to rely on data from experimental animal systems as long as the limitations of using surrogate and model systems are clearly stated. Despite species differences in metabolism, DNA repair, and other physiological processes affecting chemical mutagenesis, the virtual universality of DNA as the genetic material and of the genetic code provides a rationale for using various nonhuman test systems to predict the intrinsic mutagenicity of test chemicals. Additional support for the use of nonhuman systems is provided by the observation that chemicals causing genetic effects in one species or test system frequently cause similar effects in other species or systems. Evidence also exists that chemicals can induce genetic damage in somatic cells of exposed humans. Furthermore, a wide variety of different types of mutations have been observed in humans, including numerical chromosome aberrations, translocations, base-pair substitutions, and frameshift mutations. Although the cause of these mutations is uncertain, it is clear from these observations that human germ-cell DNA is subject to the same types of mutational events that are observed in other species and test systems.'

Recent US EPA proposals: Dearfield *et al.* from

Table 4. Proposed mutagenicity classification categories by US EPA researchers (12)

Classification	Criteria in Somatic cells	Criteria in Germ cells
Human mutagen	<p>Positive in human somatic cell mutagenicity studies as a result of human <i>in vivo</i> exposure. May be human carcinogens; unless the risk characterization suggest not as likely.</p> <p>This can include cytogenetic endpoints in tissues (such as lymphocytes) from exposed persons.</p>	<p>Positive in human <i>in vivo</i> germ cell mutagenicity studies. Human germ cell mutagens.</p> <p>This is based on positive <i>in vivo</i> findings from appropriate germ cell targets in exposed humans. It is recognized that a human germ cell mutagen is not currently identified.</p>
Probable human mutagen	<p>Clear evidence for genotoxic activity <i>in vivo</i> mammalian test(s), usually supported by <i>in vitro</i> test(s). Usually animal carcinogens and may be human carcinogens.</p> <p>Classification at this level usually means that some <i>in vivo</i> testing has been performed as follow-up to positive results from <i>in vitro</i> testing. Therefore, there is usually some supporting positive evidence from <i>in vitro</i> testing.</p>	<p>Sufficient evidence of interaction with mammalian germ cells with clear evidence for genotoxic activity. Includes valid positive results from studies on heritable mutational events in mammalian germ cells; or, valid positive results from mammalian germ cell chromosomal aberration studies that do not include an intergeneration test. Putative human germ cell mutagens if they reach target cells.</p> <p>Alternatively, it should produce positive results for DNA strand breaks (e.g. comet test), UDS, SCE and/or chromosome aberrations in germinal cells.</p>
Possible human mutagen	<p>Some evidence for genotoxic activity. May be carcinogenic through genotoxic mechanisms; possibly in humans.</p> <p>Confirmed positive results may be seen in the <i>in vitro</i> test systems without supporting evidence from the <i>in vivo</i> assays. Agents falling into this category are considered to have intrinsic mutagenic potential which is not detected <i>in vivo</i>.</p>	<p>Suggestive evidence of interaction with mammalian germ cells with some evidence for genotoxic activity. May be putative human germ cell mutagens if they reach target cells.</p> <p>For a test agent to be considered to present suggestive evidence of germ cell interaction, data are needed to demonstrate that: (i) the test agent shows some evidence of somatic cell mutagenicity and/or genotoxicity; (ii) the test agent reaches the gonads (e.g. data from pharmacokinetic/tissue distribution studies of the test agent and/or metabolites); (iii) the test agent interacts with germinal cells; these data come from subchronic or chronic toxicity tests showing gonadal pathology (e.g. sperm abnormalities); and (iv) the test agent causes adverse effects on reproductive parameters (e.g. decreased fertility, increased dead implants, reduced litter sizes).</p>
Equivocal evidence	Results from acceptable tests that cannot be convincingly called negative or positive.	
Negative evidence	Negative results in acceptable tests.	

the US EPA proposed 6 categories of mutagenicity classification that are divided to 3 categories each for somatic cells and germ cells (12). This includes (i) human somatic cell mutagens in which positive data in human somatic cells are derived from studies with exposed humans that gives positive data in human; (ii) human germ cell mutagen in which positive data from human *in vivo* germ cell studies are obtained; (iii) probable human somatic cell mutagen in which clear evidence for genotoxic activity from positive *in vivo* mammalian test(s); (iv) probable human germ cell mutagen in which sufficient evidence of interaction with mammalian germ cells with clear evidence for genotoxic activity; (v) possible human somatic cell mutagen in which confirmed positive results may be seen in the *in vitro* test systems without supporting evidence from the *in vivo* assays; and (vi) possible human germ cell mutagen in which suggestive evidence of interaction

with mammalian germ cells is seen. The mutagenicity classification categories proposed by Dearfield *et al.* (12) are summarized in Table 4.

The mutagenicity/genotoxicity data organized into "clear" and "some" evidence for mutagenicity and into "suggestive" and "sufficient" evidence germ cell interaction. The classification of mutagenicity results falls into the broad categories of inadequate, negative, equivocal, and positive data. With respect to the effects of mutagens, Dearfield *et al.* stated that 'In addition to cancer, adverse health effects from somatic cell mutations and/or germ cell mutations include sickle cell anemia, cardiovascular disease, reproductive/developmental effects, and neurobehavioral effects among many specific and general endpoints, as well as having impact on the aging process.' (12).

Canada: Health Canada: Health Canada proposed 6 categories of classification of chemicals with respect to

their mutagenic potential for germ cells in the Canadian Environmental Protection Act (CEPA), Human Health Risk Assessment for Priority Substances (13). Four categories have several subgroups based on the degree of evidence (Table 5). These classifications are as follows: human germ cell mutagen (group I), for which data from adequate epidemiological studies indicate that there is a causal relationship between exposure of humans to a chemical and an increased incidence of inherited mutations in live or dead offspring; probable human germ cell mutagen (group II), for which data from epidemiological studies to assess germ cell mutagenicity are inadequate; however, there is sufficient evidence of germ cell mutagenicity in animal species (*i.e.*, there is an increased incidence of gene mutations, structural or numerical chromosomal aberrations, or inherited congenital malformations in the live offspring of exposed animals; or an increase in dominant lethal mutations in the potential offspring of exposed animals); possible human germ cell mutagen (group III); unlikely to be a human germ cell mutagen (group IV); probably not a human germ cell mutagen (group V); unclassifiable with respect to germ cell mutagenicity in humans (group VI). Groups III, IV, V and VI have four, two, three and three subgroups, respectively (see Table 5).

The Canadian Environmental Protection Act (13) mentions 'These effects including mutagenic are manifested at the biochemical, cellular, histopathological and morphological levels.', and 'Chemical substances are classified, therefore, with respect to their potential carcinogenicity and mutagenicity to humans; this is accomplished on the basis of rigorous examination of the quantity, quality and nature of the results of available toxicological and epidemiological studies. The criteria by which Priority Substances are classified based on their weight of evidence of carcinogenicity and mutagenicity.'

Health Protection Branch: The Health Protection Branch defined toxicological findings that would be regarded as germ cell genotoxicity (14). The toxicological indication of germ cell genotoxicity are (i) *in vitro* test results and positive evidence for mutagenicity in somatic cells *in vivo*, and (ii) evidence from pharmacokinetic/tissue distribution studies that the test material and/or metabolites reaches the gonads; or (iii) evidence from subchronic or chronic treatment studies that gonadal pathology indicates germ cell damage; or (iv) evidence for reproductive/developmental effects showing reduced numbers of pregnancies, reduced litter sizes or increased time to mating following treatment in some cases. A significant proportion of agents that cause *in vivo* somatic cell mutation might also possess the ability to lead to mutation in germ cells that may be transmitted to offspring. When evidence for *in vivo*

somatic genotoxicity is demonstrated, along with tissue distribution, metabolic and/or pathologic evidence that the genotoxic chemical (or metabolites) reaches the germ lines (whether or not overt effects on fertility are found), the possibility of induced genetic damage to germ cells leading to heritable effects should be investigated (14).

Canadian Centre for Occupational Health and Safety: Controlled Products Regulations (CPR) has two mutagenicity criteria (11,15). According to CPR paragraph 57 in the classification of subdivision A (very toxic material), a chemical or chemical mixture is considered to present a hazard to man if (a) there is epidemiological evidence that shows a causal connection between exposure of persons to the substance or mixture and heritable genetic effects; or (b) there is evidence of mutagenicity in mammalian germ cells *in vivo* as shown by (i) positive results in a study that measures mutations transmitted to offspring, or (ii) positive results in an *in vivo* study showing chemical interaction with the genetic material of mammalian germ cells and positive results in an *in vivo* study assessing either gene mutation or chromosomal aberration in somatic cells. CPR paragraph 62 in the classification of subdivision B (toxic material) explains that a pure chemical or a chemical mixture is considered to present a hazard if evidence of mutagenicity in mammalian somatic cells is obtained in a test to assess either gene mutations or chromosomal aberrations.

Japan: The Japanese Industrial Safety and Health Law addresses the mutagenicity classification of substances in the workplace (11). Under this legislation, Japan does not classify mutagenic substances according to a weight of evidence approach or whether a compound may be a somatic or germ cell mutagen but rather recognizes one category: "mutagenic". A substance is classified as mutagenic when the results of a "bacterial reverse mutation assay" are positive. Workers' health impairment by the exposure to the 'strong' mutagenic chemicals or chemical mixtures (excluding those containing 1% or less by weight of mutagenic chemicals), which induce more than 1000 revertants/mg/plate in a bacterial reverse mutation assay, should be prevented in the work for manufacture or handling of these chemicals (16).

Future of Classification of Germ Cell Mutagens

Chemical evaluation of mutagenicity/genotoxicity has three major directions that are (i) screening of carcinogens, (ii) mechanistic investigation of carcinogenesis, and (iii) investigation of heritable adverse effects in germ cells including those in humans. Mutagenicity in GHS is focused on the last topic in terms of classification of germ cell mutagens. GHS and the other existing classification systems have different

Table 5. Criteria for classification of mutagenicity in germ cells in CEPA by Health Canada (13)

Category	Classification	Criteria
Group I	Human Germ Cell Mutagen	Data from adequate epidemiological studies indicate that there is a causal relationship between exposure of humans to a substance and an increased incidence of inherited mutations in live or dead offspring.
Group II	Probable Human Germ Cell Mutagen	Data from epidemiological studies to assess germ cell mutagenicity are inadequate; however, there is sufficient evidence of germ cell mutagenicity in animal species (i.e., there is an increased incidence of gene mutations, structural or numerical chromosomal aberrations, or inherited congenital malformations in the live offspring of exposed animals; or an increase in dominant lethal mutations in the potential offspring of exposed animals).
Group III. A	Possible Human Germ Cell Mutagen	Data from epidemiological studies indicate an association between exposure and human germ cell mutagenicity, but alternative explanations such as chance, bias, or confounding cannot be excluded.
Group III. B	ditto	Data from epidemiological studies to assess germ cell mutagenicity are inadequate; however, there is sufficient evidence of somatic cell mutagenicity (<i>in vivo</i> gene mutations or chromosomal aberrations) in humans or animal species, and sufficient evidence of exposure to germ cells in humans or animal species.
Group III. C	ditto	Data from epidemiological studies to assess germ cell mutagenicity in humans are inadequate or lacking. There is sufficient data in animals to indicate that the chemical is a germ cell mutagen, but available data indicate that the induction of mutations occurs through an epigenetic threshold-based mechanism.
Group III. D	ditto	Data from epidemiological studies to assess germ cell mutagenicity in humans are inadequate. There is sufficient evidence of mutagenicity of somatic cells in humans or animal species (<i>in vivo</i> gene mutations or chromosomal aberrations), but evidence of exposure to germ cells is inadequate or lacking.
Group IV. A	Unlikely to Be a Human Germ Cell Mutagen	There is no evidence of human germ cell mutagenicity in sufficiently powerful and well-designed epidemiological studies. There is evidence of mutagenicity of somatic cells in well-designed and well-conducted studies in humans or animals, but there is no evidence of exposure of human or animal germ cells in well-designed studies.
Group IV. B	ditto	Data on germ cell mutagenicity in epidemiological studies in humans are inadequate; there is no evidence of mutagenicity <i>in vivo</i> in germ or somatic cells in well-designed and properly conducted studies in animals.
Group V. A	Probably Not a Human Germ Cell Mutagen	There is no evidence of germ cell mutagenicity in sufficiently powerful and well-designed epidemiological studies; there is no evidence of germ cell mutagenicity in animal species.
Group V. B	ditto	There is no evidence of germ cell mutagenicity in sufficiently powerful and well-designed epidemiological studies; data in animal species are inadequate.
Group V. C	ditto	Data from epidemiological studies to assess germ cell mutagenicity in humans are inadequate, but evidence of the lack of germ cell mutagenicity in animal species is strongly supported by other data on mutagenicity <i>in vivo</i> .
Group VI. A	Unclassifiable with Respect to Germ Cell Mutagenicity in Humans	Data from epidemiological and/or animal studies are inadequate (i.e., because of major qualitative limitations, the studies cannot be interpreted as showing either the presence or absence of germ cell mutagenicity).
Group VI. B	ditto	There are no <i>in vivo</i> mutagenicity data available for evaluation.
Group VI. C	ditto	Results of epidemiological studies in human populations and experimental studies in animal species are conflicting, without an identifiable mechanistic basis.

objectives, target audiences and criteria. For example, the primary objective of the systems in GHS, EU and Germany MAK is for hazard classification, on the other hand, the systems of US EPA and Health Canada are for risk assessment. Target audiences are workers in the

systems of Germany MAK, Canadian CPR and Japan; consumers and workers in EU system; consumers, workers, transport workers, and emergency responders in GHS. As for criteria, the report from EPA researchers includes adverse effects on reproductive

parameters.

The fundamental purposes are different in these regulations; therefore, it is difficult to discuss the significance of the different approaches. Simple classification category and criteria will be useful for hazard classification. The GHS and EU systems on (germ cell) mutagenicity meet to this point. Application of the criteria needs expert judgment in a weight of evidence approach in GHS. However, the expert judgment leads sometimes different conclusion from expert to expert. This will be a critical issue in the classification of germ cell mutagens by GHS. Furthermore, hazard classification/evaluation is often confused as risk evaluation, especially in Japan. Further efforts including risk evaluation and communication on germ cell mutagenicity will be needed to make understanding of regulations global.

Recently, the European Commission proposed a new regulatory framework for chemicals called REACH for the Registration, Evaluation and Authorisation of Chemicals in October 2003 (17). The authorisation process pays particular attention to the risks that the substance poses due to any carcinogenic, mutagenic, and/or reproductive toxicity (CMR) properties. CMR's category 1 and 2 in the EU criteria, which correspond to the category 1A and 1B, respectively in GHS criteria, are subject to authorization (18,19). All substances imported in quantities over 1 tonne that contain more than 0.1 per cent CMR material must be authorised before gaining access to the EU market. Authorisation provides a permit for specific uses and can be requested by 'producer' or 'user'. The GHS itself is not legally binding, however, some national or regional laws including REACH may be legally binding. Now, classification of germ cell mutagens becomes an important issue.

As GHS criteria have been adopted in worldwide, it will become standard for hazard classification. After implementation of GHS in each country, the classification of chemicals on germ cell mutagenicity will be performed by chemical suppliers (manufacturers or importers). Understanding classification systems for germ cell mutagens will be helpful for scientifically sound classification of chemicals in the GHS.

Acknowledgement: We are grateful to Dr. D.J. Tweats for his review of this manuscript. This work was supported in part by the Health and Labor Sciences Research Grants (H18-Chemistry-General-009).

References

- 1 United Nations (UN). Globally Harmonized System of Classification and Labelling of Chemicals (GHS), New York and Geneva, 2003. (http://www.unece.org/trans/danger/publi/ghs/ghs_rev00/00files_e.html)
- 2 United Nations (UN). Globally Harmonized System of Classification and Labelling of Chemicals (GHS), First revised edition, New York and Geneva, 2005. (http://www.unece.org/trans/danger/publi/ghs/ghs_rev01/01files_e.html)
- 3 Japanese inter-ministerial committee for GHS. Japanese translation version of the first revised edition of GHS documents, 2005. (in Japanese). (http://www.meti.go.jp/policy/chemical_management/kokusai/GHS/GHStexts/kariyaku.htm)
- 4 Japanese inter-ministerial committee for GHS. Publication of classification of chemicals under each relevant domestic law (in progress), 2006. (in Japanese). (<http://www.safe.nite.go.jp/ghs/index.html>)
- 5 European Communities (EC). Commission directive 2001/59/EC of 6 August 2001, Official J. of Europe Commun. L225, 21.8.2001.
- 6 Pratt IS, Barron T. Regulatory recognition of indirect genotoxicity mechanisms in the European Union. *Toxicol Lett.* 2003; 140-141: 53-62.
- 7 European Communities (EC). Classification, packaging, and labeling of dangerous substances in the EU, Part 1, Office for Official Publications of the European Communities, Luxembourg, 1997.
- 8 Adler ID, Anrae U, Kreis P, Neumann HG, Their R, Wild D. Recommendation for the categorization of germ cell mutagens. *Int Arch Occup Environ Health.* 2000; 73: 428-32.
- 9 Deutsche Forschungsgemeinschaft (DFG). List of MAK and BAT values, Commission for the investigation of health hazards of chemical compounds in the work area, Report No. 39, Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim, Germany; 2003.
- 10 US Environmental Protection Agency (USEPA). Guidelines for Mutagenicity Risk Assessment, Federal Register 51(185):34006-34012, EPA/630/R-98/003; September 1986.
- 11 Organisation for Economic Co-operation and Development (OECD). OECD Series on testing and assessment, Number 12, Detailed Review Document on Classification Systems for Germ Cell Mutagenicity in OECD Member Countries, ENV/JM/MONO(99)2, Environment Directorate, OECD, Paris, 1999.
- 12 Dearfield KL, Cimino MC, McCarroll NE, Mauer I, Valcovic LR. Genotoxicity risk assessment: a proposed classification strategy. *Mutat. Res.* 2002; 26: 121-35.
- 13 Health Canada. Canadian Environmental Protection Act (CEPA). Human Health Risk Assessment for Priority Substances, Minister of Supply and Services Canada, Ottawa, Canada, Cat. No. En40-215/41E; 1994.
- 14 Health Protection Branch. The assessment of mutagenicity, Mutagenicity Guidelines, Ministry of Supply and Services Canada, Cat. H49-85/1993E; 1993.
- 15 Canadian Centre for Occupational Health and Safety (CCOHS). Controlled Products Regulations. *SOR/88-66*; 1988.
- 16 Guidelines for Preventing Health Impairment by Chemical Substances with Mutagenicity Recognizes, LSB Notification No. 312, Industrial Safety and Health Law, Japan, dated May 17, 1993.

- 17 European Commission. Proposal for a Regulation of the European parliament and of the council concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency and amending Directive 1999/45/EC and Regulation (EC), COM(2003) 644 final, Brussels; 2003.
- 18 European Commission Services. Comparison between EU and GHS criteria, Human health and environment (Draft), Version June 08, 2005.
- 19 Morita T, Ishimitsu S, Morikawa K. Genotoxicity in risk assessment, Global perspectives. *Environ. Mutagen Res.* 2005; 27, 47-56. (in Japanese)

Editorial

The International Workshops on Genotoxicity Testing (IWGT): History and achievements

Keywords: Genotoxicity testing; IWGT workshops; History

The objectives and structure of the IWGT

Three workshops have been organised previously under the auspices of the International Workshops on Genotoxicity Testing (IWGT). Recognising the success of these earlier workshops, the International Association of Environmental Mutagen Societies (IAEMS) formalised these workshops in 2002 under the IAEMS umbrella and agreed that they would be held on a continuing basis in conjunction with the International Conferences on Environmental Mutagens (ICEM) that are held every 4 years. In this way, an ongoing process of international discussion and harmonisation of testing methods and testing approaches has been established that can take advantage of the international experts who attend these meetings. These ongoing workshops will help to ensure that different recommendations for methodology in these new assays do not arise in different parts of the world, and thus avoid situations that could lead to:

- Unnecessary duplication of testing to satisfy local requirements.
- Variations in the test performance.
- Potential differences in test outcome.
- Unjustified differences in the use of test data for description, assessment and management of risk.

The IWGT process is implemented through working groups of recognised international experts from industry, academia and the regulatory sectors, with due attention to geographical, disciplinary and sector balance. For each working group, a chairperson, deputy chair, and rapporteur are appointed. Experts in the science of each topic are invited to bring experimental

data to bear on the discussions; the remit of each group is to derive recommendations based on data, and not on unsupported opinion or anecdotal information. Geographical as well as scientific balance has been attempted within these groups as can be seen from the composition of the groups. There are several objectives sought in bringing together representatives from around the world to share their experiences in generating and evaluating genotoxicity data from a variety of methodological and strategic approaches. We strive to:

- Attain a greater understanding of true test performance from a wide database.
- Provide recommendations that minimise misinterpretation.
- Recognise that no single assay can detect every genotoxin.
- Achieve compromise for the sake of harmonisation or acceptance that more than one approach is both reasonable and valid.

Because of the IWGT approach, in particular development of data-driven consensus by the key global experts from academia, government and industry, IWGT recommendations have been seen as state-of-the-art and have high credibility. These recommendations serve as important supplements to established regulatory guidelines and provide a sound basis for updating those guidelines as the state of science advances.

2. Achievements of previous workshops

The first IWGT Workshop was held in Melbourne, Australia as a satellite to the International Conference

on Environmental Mutagens in 1993. Several working groups addressed the methodology of test systems in widespread use for scientific research and regulatory practice. The consensus recommendations from those working groups were published in Mutation Research in 1994 (volume 312, pages 195–318), and had a significant impact on revisions to OECD guidelines for genotoxicity testing that were on-going at that time [1]. These IWGT discussions and conclusions were also recognised as valid contributions for the ICH guidances for pharmaceuticals for human use that were finalised in 1995 and 1997 [2,3].

At the second IWGT Workshop, held in Washington, DC in 1999, recommendations for the mouse lymphoma *tk* mutation assay and the *in vivo* micronucleus test were updated and some recommendations were made that were different from or extended beyond the published OECD guidelines for these test systems. In addition, several assays for which no OECD guidelines existed were discussed in detail and recommendations for their conduct made. These reports were published in Environmental and Molecular Mutagenesis in 2000 (volume 35, pages 159–263).

Recommendations for the conduct of the mouse lymphoma *tk* mutation assay, the *in vitro* micronucleus test and for *in vivo* transgenic mutation assays were updated at the third IWGT Workshop in Plymouth, UK in 2002 and published in Mutation Research in 2003 (volume 540, pages 119–181). Guidance was also introduced on the usefulness and techniques for molecular analysis of tumour and non-tumour tissues from carcinogenicity studies with the transgenic haploinsufficient p53 and *RasH2* mouse strains [4]. An important aspect of this meeting was the initiation of discussions and recommendations on strategic approaches to genotoxicity testing [5].

The published IWGT reports that have extended recommendations for testing beyond existing OECD guidelines are:

- Advances in conduct of the *in vivo* micronucleus test [6].
- Advances in conduct of the mouse lymphoma *tk* mutation assay [7–9].

For the *in vivo* micronucleus test (OECD Guideline 474) the key new recommendations are:

- Repeat dose studies (e.g. for 28 days) can be conducted in the rat, where the bone marrow is still the principal tissue but additional data can be obtained from scoring micronuclei (MN) in peripheral blood.

- Modern automated scoring methods such as flow cytometry and image analysis, allowing much greater numbers of cells to be scored, are acceptable, if the utility of the analytical system has been demonstrated in the performing laboratory.
- It is not necessary to treat a concurrent positive control group with every study, particularly in laboratories that run the test frequently, and when test chemical dosing solutions have been well characterised and systemic exposure has been demonstrated. Control of staining and scoring can be achieved by coding positive control slides from a previous GLP study into the current study. However, until more experience is gained, it is recommended to include positive control in repeat dose studies (e.g. for 28 days).
- CREST antibodies and FISH with pancromerid or chromosome-specific probes can be used under certain circumstances to distinguish clastogens from aneugens. Positive controls should be included with these types of investigations.
- Micronuclei can be measured in liver, colonic epithelium, skin, spleen, lung, spermatids and in foetal/neonatal tissue, although concurrent positive control treatments will be required with these tissues, at least until the methods become more widely established and published.

For the mouse lymphoma assay (OECD Guideline 476), which has become the preferred mammalian cell mutation test in many regulatory documents [3,10–12] the key new recommendations are:

- The IWGT Working Group confirmed as useful the recommendations made by ICH [3] that a 24 h continuous treatment in the absence of S9 should be conducted when short treatments give negative results.
- Relative total growth (RTG) is the recommended measure of toxicity.
- Alternative suggestions for positive control chemicals were made, and a recommendation made that positive control treatments should demonstrate adequate detection of small colony mutants.
- Acceptable negative (solvent) control mutant frequencies for the agar and microwell methods have been defined.
- No single statistical test was identified as appropriate for evaluation of responses as positive or negative, and a Global Evaluation Factor has been defined for both agar and microwell versions of the test to be used in conjunction with other statistical methods (e.g. dose-response) in evaluation of results.