

assigning the newly identified alleles. This work was supported by Health and Labour Sciences Research Grants (H18-kenki-012 to F. K. and H16-kenki-059 to T. E.).

REFERENCES

1. National Institute of Infectious Diseases and Tuberculosis and Infectious Diseases Control Division, Ministry of Health, Labour and Welfare. (2000) Legionellosis, April 1999–July 2000. IASR Infectious Agents Surveillance Report. **21**: 186–7 (<http://idsc.nih.gov/iasr/21/247/tpc247.html>).
2. Amemura-Maekawa J., Kura F., Chang B., Watanabe H. (2005) *Legionella pneumophila* serogroup 1 isolates from cooling towers in Japan form a distinct genetic cluster. *Microbiol Immunol* **49**: 1027–33.
3. Gaia V., Fry N.K., Afshar B., Lück P.C., Meugnier H., Etienne J. *et al.* (2005) Consensus sequence-based scheme for epidemiological typing of clinical and environmental isolates of *Legionella pneumophila*. *J Clin Microbiol* **43**: 2047–52.
4. Nagai T., Sobajima H., Iwasa M., Tsuzuki T., Kura F., Amemura-Maekawa J. *et al.* (2003) Neonatal sudden death due to *Legionella pneumonia* associated with water birth in a domestic spa bath. *J. Clin. Microbiol.* **41**: 2227–9.
5. Aurell H., Farge P., Meugnier H., Gouy M., Forey F., Lina G. *et al.* (2005) Clinical and environmental isolates of *Legionella pneumophila* serogroup 1 cannot be distinguished by sequence analysis of two surface protein genes and three housekeeping genes. *Appl Environ Microbiol* **71**: 282–9.
6. Huang B., Yuan Z., Heron B.A., Gray B.R., Eglezos S., Bates J.R. *et al.* (2006) Distribution of 19 major virulence genes in *Legionella pneumophila* serogroup 1 isolates from patients and water in Queensland, Australia. *J Med Microbiol* **55**: 993–7.
7. Ko K.S., Miyamoto H., Lee H.K., Park M.-Y., Fukuda K., Park B.J. *et al.* (2006) Genetic diversity of *Legionella pneumophila* inferred from *rpoB* and *dotA* sequences. *Clin Microbiol Infect* **12**: 254–61.