




ELSEVIER

ORIGINAL ARTICLE


<http://www.iof.org/oonet/aoaso.htm>

## Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase

Masayasu Yoneda<sup>a,1</sup>, Ying Guo<sup>a,1</sup>, Haruya Ono<sup>a</sup>, Yusuke Nakatsu<sup>a</sup>, Jun Zhang<sup>a</sup>, XingLong Cui<sup>a</sup>, Misaki Iwashita<sup>b</sup>, Sonoko Kumamoto<sup>b</sup>, Yoshihiro Tsuchiya<sup>a</sup>, Hideyuki Sakoda<sup>c</sup>, Midori Fujishiro<sup>c</sup>, Akifumi Kushiyama<sup>d</sup>, Yuko Koketsu<sup>c</sup>, Takako Kikuchi<sup>d</sup>, Hideaki Kamata<sup>a</sup>, Fusanori Nishimura<sup>b</sup>, Tomoichiro Asano<sup>a,\*</sup>

<sup>a</sup> Department of Medical Chemistry, Division of Molecular Medical Science, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan

<sup>b</sup> Department of Dental Science for Health Promotion, Division of Cervico-Gnathostomatology, Graduate School of Biomedical Sciences, Hiroshima University, Japan

<sup>c</sup> Department of Internal Medicine, Graduate School of Medicine, University of Tokyo, Japan

<sup>d</sup> Institute for Adult Diseases, Asahi Life Foundation, Japan

Received 13 October 2009; received in revised form 19 January 2010; accepted 1 February 2010

### KEYWORDS

AMP-activated protein kinase (AMPK);  
Sirtuin 1 (SIRT1);  
High-fat diet (HFD)

### Summary

**Aims:** Energy sensing systems including AMPK and SIRT1 play important roles in the regulation of hepatic gluconeogenesis and fatty acid oxidation. In this study, we investigated how hepatic LKB1-AMPK signaling and SIRT1 expression are altered after 2 or 8 weeks of HFD feeding.

**Methods:** The livers of male mice fed a HFD or a standard diet for 2 or 8 weeks were removed. The expression and phosphorylation levels of LKB1, AMPK, ACC and TORC2, and SIRT1 expression levels were examined by immunoblotting.

**Results:** In mice fed a HFD for 2 weeks, the phosphorylations of AMPK $\alpha$  and ACC were decreased without significant alterations in LKB1 phosphorylation or SIRT1 protein levels, while TORC2 protein levels were increased. In mice fed a HFD for 8 weeks, marked reductions in LKB1 phosphorylation and SIRT1 protein amount were observed in addition to the decreased phosphorylations of AMPK $\alpha$  and ACC.

\* Corresponding author. Tel.: +81 82 257 5135; fax: +81 82 257 5136.

E-mail address: [tasano@hiroshima-u.ac.jp](mailto:tasano@hiroshima-u.ac.jp) (T. Asano).

<sup>1</sup> These authors contributed equally to this work.

**Conclusions:** The mechanisms underlying impaired energy sensing signaling differ with the duration of HFD feeding. In the early phase of HFD feeding, LKB1 and SIRT1 were not impaired, while in the later phase of HFD feeding, decreased SIRT1 expression and LKB1 phosphorylation may be involved in the development of severe glucose and lipid intolerance.

© 2010 Published by Elsevier Ltd on behalf of Asian Oceanian Association for the Study of Obesity.

## Introduction

AMP-activated protein kinase (AMPK) is a key sensor of cellular energy status that regulates glucose and lipid metabolism. AMPK is activated through an increase in the intracellular AMP/ATP ratio induced by metabolic stresses such as starvation, hypoxia, or exercise [1], as well as stimulation with leptin [2], adiponectin [3] and the anti-diabetic agents metformin [4–6] and thiazolidinediones [6–8]. The process of AMPK activation requires phosphorylation at threonine residue (Thr) 172 within the catalytic  $\alpha$  subunit by LKB1 [9]. The phosphorylation of AMPK is reportedly suppressed in the livers [10,11] and skeletal muscles [12] of HFD-fed rats.

LKB1, an upstream kinase for AMPK and AMPK-related kinases including the salt-inducible kinase (SIK) cascade, is localized mainly in the nucleus under basal conditions. LKB1 is reportedly phosphorylated at serine residue (Ser) 431 in mice (Ser428 in humans) by protein kinase A (PKA) [13], p90 ribosomal S6 kinase [14] or PKC- $\zeta$  [15], and is translocated from the nucleus to the cytoplasm. In the cytoplasm, LKB1 forms a heterotrimeric complex with two adaptor proteins termed STE20-related adaptor (STRAD) and MO25 [16,17]. The binding and interaction of LKB1 with STRAD and MO25 increase the catalytic activity of LKB1, resulting in the phosphorylation and activation of AMPK [18]. Activated AMPK then induces the phosphorylation and inactivation of acetyl coenzyme A carboxylase (ACC), decreasing malonyl-CoA and thereby relieving inhibition of carnitine palmitoyl-transferase 1, and leading to an increase in fatty acid oxidation in the liver and skeletal muscle [19].

In addition, activated AMPK phosphorylates and inactivates the transducer of regulated cyclic AMP responsive element binding protein (CREB) 2 (TORC2, or CREB-regulated transcription coactivator, CRTC2), identified as a critical molecular switch in the regulation of hepatic gluconeogenesis [20]. TORC2 phosphorylated at Ser171 translocates from the nucleus to the cytoplasm, leading to its poly-ubiquitination and degradation [21,22], thereby reducing the expressions

of gluconeogenic genes, including peroxisome proliferator-activated receptor  $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ ), phosphoenolpyruvate carboxykinase and glucose-6-phosphatase [20].

Sirtuin 1 (SIRT1), a mammalian homologue of Sir2 (silent information regulator 2), is an NAD<sup>+</sup>-dependent histone deacetylase, and its expression and enzymatic activity are enhanced by an increase in the NAD<sup>+</sup>/NADH ratio [23]. In HFD-fed rodents, SIRT1 protein levels are reportedly decreased in the liver [24] and skeletal muscle [25]. Moreover, resveratrol, an activator of SIRT1, improves HFD-induced fat accumulation and insulin resistance in the liver by enhancing AMPK phosphorylation [10,26].

Thus, there exist multiple key molecules including AMPK and SIRT1 in the liver, which regulate glucose tolerance and insulin sensitivity in response to the nutrient conditions. In this study, we attempted to investigate how AMPK and SIRT1-mediated pathways are impaired after the different duration of high-fat diet feeding, and prepared the mice treated with short (2 weeks) or long-term (8 weeks) control or HFD feeding. Herein, we discuss the pathophysiological significance of our observations on HFD-induced insulin resistance.

## Materials and methods

### Animals and diets

Male 7-week-old ddY mice were purchased from Shimizu Laboratory Supplies (Kyoto, Japan) and fed a standard diet (SD,  $n=8$ ) or a HFD ( $n=8$ ) for 2 weeks and then the SD ( $n=4$ ) or HFD ( $n=4$ ) for 8 weeks. The nutrient composition of the SD was 54.4% carbohydrate, 23.6% protein, 5.3% fat, and a vitamin and mineral mixture, while that of the HFD purchased from Oriental Yeast (Tokyo, Japan) was 7.5% carbohydrate, 24.5% protein, 60% fat, and a vitamin and mineral mixture, as in our previous studies [27,28]. All mice were fed *ad libitum* and maintained on 12-h light and dark cycles under controlled environment settings. All mice were fasted for 6 h and then killed. The protocols for this animal

study were approved by the Animal Experimental Committee of Hiroshima University.

### Liver tissue samples

All mice were anesthetized with pentobarbital sodium (60mg/kg body weight) intraperitoneally. The livers were removed and homogenized in ice-cold lysis buffer [50 mM Tris–HCl (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 1 mM sodium orthovanadate, and 1 mM phenylmethylsulfonyl fluoride] and centrifuged at  $16,000 \times g$  for 30 min at 4 °C. The supernatants were used as liver lysates. The liver lysates were added to Sodium Dodecyl Sulfate (SDS) sample buffer [200 mM Tris–HCl (pH 6.8), 4% SDS, 10% glycerol, 0.1% bromophenol blue, 5% 2-mercaptoethanol], boiled for 5 min, and analyzed by Western blot assays.

### Western blot assays

For Western blot assays, the liver lysates were analyzed with antibodies against: LKB1 (Molecular Weight 51 kDa), SIRT1 (110 kDa) (Upstate, NY), phosphorylated LKB1-Ser431 (52 kDa), TORC2 (87 kDa) (Santa Cruz Biotechnology, CA), STRAD (20 kDa) (Imgenex, CA), MO25 $\alpha$  (39 kDa), phosphorylated AMPK $\alpha$ -Thr172 (62 kDa), phosphorylated ACC-Ser79 (280 kDa), phosphorylated TORC2-Ser171 (85 kDa) (Cell Signaling Technology, MA), or Actin (Sigma–Aldrich, MO), followed by the secondary antibodies: anti-rabbit or anti-mouse immunoglobulin-horseradish peroxidase-linked whole antibody, and also enhanced chemiluminescent reagents (GE Healthcare, UK).

### Quantitative real-time PCR

Total RNA from mouse liver tissue was extracted using Sepasol-RNA I (Nacalai Tesque, Kyoto, Japan), and cDNA was generated using a Transcriptor First Strand cDNA Synthesis Kit (Roche, Mannheim, Germany). Expression level of the *TORC2* gene were analyzed by quantitative real-time PCR using SYBR Premix Ex Taq<sup>TM</sup> (Takara Bio, Shiga, Japan) and an ABI PRISM 7700 Sequence Detector (Applied

Biosystems, CA). Primer sequences were as follows: *TORC2*, 5'-AGTGAGGGATCAATGAGTGG-3' (forward) and 5'-AGAGTCTCCTGTGAGTATGA-3' (reverse). All PCR data were normalized to expression levels of *GAPDH* in the corresponding sample.

### Statistical analysis

The data are expressed as means  $\pm$  standard deviation. The difference in the mean of any parameter between the two groups was tested using the unpaired *t* test. A *P* < 0.05 was considered statistically significant.

## Results

### Parameters of mice fed HFD for 2 or 8 weeks

As shown in the Table 1, after 2 weeks of HFD feeding, body weights and fasting blood glucose levels were higher than those of mice fed the SD, but the differences were not significant (*P* = 0.549 and 0.068, respectively). However, HFD feeding for 8 weeks significantly increased body weight and fasting blood glucose as compared with SD feeding.

### LKB1-AMPK-ACC/-TORC2 pathways in livers of mice fed HFD for 2 weeks

Regarding the LKB1-AMPK-ACC and -TORC2 pathways in the liver, we compared protein expressions and phosphorylation levels, by Western blotting, between mice fed the SD versus the HFD for 2 weeks. As shown in Fig. 1, AMPK $\alpha$  and TORC2 protein levels were significantly increased, but phosphorylated AMPK $\alpha$ -Thr172 and phosphorylated ACC-Ser79 protein levels were markedly decreased in HFD-fed mice as compared with SD-fed mice. Phosphorylated LKB1-Ser431 was slightly, but not significantly (*P* = 0.132), decreased in HFD-fed mice. Levels of two adaptor proteins of LKB1, STRAD and MO25, did not differ between SD and HFD-fed mice. Thus, these data show that HFD feeding for 2 weeks suppresses hepatic AMPK-ACC


**Table 1** Body weight and fasting blood glucose in ddY mice fed SD or HFD for 2 or 8 weeks.

|                       | 2 weeks        |                | 8 weeks        |                 |
|-----------------------|----------------|----------------|----------------|-----------------|
|                       | SD             | HFD            | SD             | HFD             |
| Body weight (g)       | 38.2 $\pm$ 3.0 | 39.3 $\pm$ 4.1 | 48.9 $\pm$ 1.7 | 66.5 $\pm$ 4.7* |
| Blood glucose (mg/dl) | 164 $\pm$ 31   | 205 $\pm$ 49   | 154 $\pm$ 18   | 363 $\pm$ 36*   |


Data are expressed as means  $\pm$  S.D. Male 7-week-old ddY mice were fed a standard diet (SD, *n* = 8) or a high-fat diet (HFD, *n* = 8) for 2 or 8 weeks (SD, *n* = 4; HFD, *n* = 4). Body weight and fasting blood glucose were measured at 9 and 15 weeks of age.

\* *P* < 0.05; SDvsHFD.

Please cite this article in press as: Yoneda M, et al. Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase. *Obes Res Clin Pract* (2010), doi:10.1016/j.orcp.2010.02.002




**Figure 1** LKB1-AMPK-ACC/-TORC2 pathways in livers of mice fed HFD for 2 weeks. Western blot analysis of LKB1, phosphorylated LKB1-Ser431 (P-LKB1), STRAD, MO25 $\alpha$ , AMPK $\alpha$ , phosphorylated AMPK $\alpha$ -Thr172 (P-AMPK $\alpha$ ), phosphorylated ACC-Ser79 (P-ACC), TORC2 and phosphorylated TORC2-Ser171 (P-TORC2) proteins as well as real-time PCR analysis of TORC2 mRNA level in mice fed the SD (white columns;  $n=4$ ) or HFD (black columns;  $n=4$ ) for 2 weeks ( $^{\dagger}P<0.05$ ; vs SD). The blots shown are representative of three blots from each group of mice.



**Figure 2** LKB1-AMPK-ACC/-TORC2 pathways in livers of mice fed HFD for 8 weeks. Western blot analysis of LKB1, phosphorylated LKB1-Ser431 (P-LKB1), STRAD, MO25 $\alpha$ , AMPK $\alpha$ , phosphorylated AMPK $\alpha$ -Thr172 (P-AMPK $\alpha$ ), phosphorylated ACC-Ser79 (P-ACC), TORC2 and phosphorylated TORC2-Ser171 (P-TORC2) proteins in mice fed the SD (white columns;  $n=4$ ) or HFD (black columns;  $n=4$ ) for 8 weeks ( $^{\dagger}P<0.05$ ; vs SD). The blots shown are representative of three blots from each group of mice.

Please cite this article in press as: Yoneda M, et al. Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase. *Obes Res Clin Pract* (2010), doi:10.1016/j.orcp.2010.02.002



**Figure 3** SIRT1 expressions in livers of mice fed HFD for 2 or 8 weeks. Western blot analysis of SIRT1 proteins in mice fed the SD (white columns;  $n=4$ ) or HFD (black columns;  $n=4$ ) for 2 weeks (A) or 8 weeks (B) ( $^{\dagger}P<0.05$ ; vs SD). The blots shown are representative of three blots from each group of mice.

182 signaling without changing LKB1 phosphorylation  
183 while upregulating TORC2 expression.

#### 184 LKB1-AMPK-ACC/-TORC2 pathways in livers 185 of mice fed HFD for 8 weeks

186 As shown in Fig. 2, phosphorylated LKB1-Ser431,  
187 STRAD, phosphorylated AMPK $\alpha$ -Thr172 and phospho-  
188 rylated ACC-Ser79 protein levels were signifi-  
189 cantly decreased in mice fed the HFD for 8 weeks as  
190 compared with SD-fed mice. TORC2 protein levels  
191 were significantly increased in HFD-fed mice, while  
192 TORC2 mRNA levels were not significantly altered,  
193 which suggests increased TORC2 protein not via  
194 the transcriptional regulation but rather via the  
195 increased translational regulation or the decreased  
196 degradation velocity. Thus, reduced hepatic LKB1  
197 phosphorylation was observed after 8 weeks of HFD  
198 feeding, while suppressed AMPK-ACC signaling was  
199 similar to that with 2 weeks of HFD feeding.

#### 200 SIRT1 expression in livers of mice fed HFD 201 for 2 or 8 weeks


202 It was recently reported that SIRT1 functions as an  
203 upstream regulator for LKB1 in cultured cells and  
204 in the livers of mice and rats [29,30]. As shown in  
205 Fig. 3A, in mice fed the HFD for 2 weeks, SIRT1  
206 protein levels were slightly decreased as compared  
207 with SD-fed mice, but the difference was not sig-  
208 nificant ( $P=0.117$ ). However, SIRT1 protein levels  
209 were drastically decreased in mice fed the HFD for 8  
210 weeks (Fig. 3B). These data indicate that long-term  
211 HFD feeding gradually reduces SIRT1 expression,  
212 which would then contribute to the declining LKB1  
phosphorylation.

#### Discussion

213  
214 In the energy deficient state, increased AMP binds  
215 to AMPK, and structurally modified AMPK is thereby  
216 rendered accessible to phosphorylation by LKB1.  
217 Thus, at the cellular level, increases in both the  
218 AMP/ATP ratio and LKB1 activity are considered  
219 to be responsible for AMPK phosphorylation. In  
220 this study, LKB1 phosphorylation was not signifi-  
221 cantly suppressed after 2 weeks of HFD feeding,  
222 although AMPK phosphorylation had already been  
223 attenuated. This suggests AMPK impairment in the  
224 relatively early phase of HFD feeding to be due  
225 to the decreased hepatic AMP/ATP ratio (Fig. 4A).  
226 However, after long-term HFD feeding, decreased  
227 LKB1 phosphorylation becomes overt. The phos-  
228 phorylation of LKB1 at Ser431 reportedly does  
229 not affect the kinase activity itself, but is neces-  
230 sary for the formation of a complex with STRAD  
231 and MO25 and its translocation to the cytoplasm  
232 [16,17]. In addition, SIRT1 expression was also  
233 markedly reduced after long-term HFD feeding.  
234 Thus, it is reasonable to speculate that not only a  
235 decreased AMP/ATP ratio but also decreased cyto-  
236 plasmic LKB1 activities are involved in the reduced  
237 AMPK phosphorylation seen after long-term HFD  
238 feeding (Fig. 4B).

239 According to recent studies, SIRT1 regulates  
240 LKB1 deacetylation and induces translocation of  
241 LKB1 to the cytosol from the nucleus [29,30].  
242 Translocated LKB1 associates with STRAD and MO25  
243 and exhibits kinase activity leading to the phos-  
244 phorylation of AMPK [18]. Moreover, SIRT1 also  
245 reportedly deacetylates TORC2 and promotes its  
246 ubiquitin-dependent degradation in the mouse liver  
247 after prolonged fasting and in primary hepato-

Please cite this article in press as: Yoneda M, et al. Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase. *Obes Res Clin Pract* (2010), doi:10.1016/j.orcp.2010.02.002



**Figure 4** Proposed hepatic LKB1-AMPK-ACC/-TORC2 pathways in mice fed HFD for 2 weeks (A) or 8 weeks (B). (A) In mice fed the HFD for 2 weeks, AMPK phosphorylation is decreased, without changing SIRT1 and LKB1, followed by reduced ACC phosphorylation, resulting in the suppression of fatty acid oxidation. TORC2 phosphorylation and degradation are reduced by the inactivation of AMPK, which in turn raises the de-phosphorylated form of TORC2 and enhances gluconeogenesis. (B) In mice fed the HFD for 8 weeks, a decrease in SIRT1 expression suppresses LKB1 deacetylation and the phosphorylations of LKB1, AMPK and ACC. TORC2 phosphorylation and degradation are reduced by the inactivation of AMPK, and in addition, TORC2 deacetylation may be impaired by the decrease in SIRT1 expression [32], leading to an extreme increase in de-phosphorylated TORC2 proteins and even greater enhancement of gluconeogenesis. P, phosphorylation; Ac, acetylation.

cytes exposed to glucagon [32]. Taken together, our results suggest that SIRT1 expression reduction observed after long-term HFD feeding would suppress hepatic LKB1-AMPK-ACC signaling and TORC2 degradation. As a result, increased gluconeogenesis and the resultant hyperglycemia may become increasingly evident, depending on the duration of overnutrition. In fact, with prolonged HFD feeding from 2 to 8 weeks, hyperglycemia was markedly exacerbated as compared with SD-fed controls.

AMPK and SIK are kinases located downstream from LKB1. These two kinases have conserved kinase domains, but SIK is phosphorylated by LKB1 without being affected by the AMP/ATP ratio [31]. In addition, SIK phosphorylates TORC2 very efficiently, but probably not ACC. Thus, while fatty acid oxidation through ACC phosphorylation is mediated by AMPK, increased hepatic gluconeogenesis mediated by the TORC2 pathway is induced by both SIK and AMPK. Taking these observations together, decreased LKB1 phosphorylation after long-term overnutrition would appear to impair both AMPK and SIK phosphorylation, possibly leading to more markedly reduced TORC2 phosphorylation.

In conclusion, we revealed that short-term HFD feeding suppressed hepatic AMPK-ACC signaling, while prolonged HFD feeding decreased SIRT1 expression and LKB1 phosphorylation. These clearly different impairments in energy sensing mechanisms may also develop gradually in the livers of human diabetic and obese subjects, possibly affecting the efficiency of AMPK activators including metformin and SIRT1 activators such as polyphenols like resveratrol.

**Conflict of interest**

The authors declare that they have no conflict of interest.

**Acknowledgements**

We are grateful to the Analysis Center of Life Science, Hiroshima University for the use of their facilities.

**References**

- [1] Towler MC, Hardie DG. AMP-activated protein kinase in metabolic control and insulin signaling. *Circ Res* 2007;100:328-41.
- [2] Minokoshi Y, Kim YB, Peroni OD, Fryer LG, Müller C, Carling D, et al. Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature* 2002;415:339-43.
- [3] Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, et al. Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med* 2002;8:1288-95.
- [4] Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, et al. Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* 2001;108:1167-74.
- [5] Musi N, Hirshman MF, Nygren J, Svanfeldt M, Bavenholm P, Rooyackers O, et al. Metformin increases AMP-activated protein kinase activity in skeletal muscle of subjects with type 2 diabetes. *Diabetes* 2002;51:2074-81.

Please cite this article in press as: Yoneda M, et al. Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase. *Obes Res Clin Pract* (2010), doi:10.1016/j.orcp.2010.02.002

- [6] Fryer LG, Parbu-Patel A, Carling D. The anti-diabetic drugs rosiglitazone and metformin stimulate AMP-activated protein kinase through distinct signaling pathways. *J Biol Chem* 2002;277:25226–32.
- [7] Ye JM, Dzamko N, Hoy AJ, Iglesias MA, Kemp B, Kraegen E. Rosiglitazone treatment enhances acute AMP-activated protein kinase-mediated muscle and adipose tissue glucose uptake in high-fat-fed rats. *Diabetes* 2006;55:2797–804.
- [8] Coletta DK, Sriwijitkamol A, Wajcberg E, Tantiwong P, Li M, Prentki M, et al. Pioglitazone stimulates AMP-activated protein kinase signalling and increases the expression of genes involved in adiponectin signalling, mitochondrial function and fat oxidation in human skeletal muscle in vivo: a randomised trial. *Diabetologia* 2009;52:723–32.
- [9] Lizcano JM, Göransson O, Toth R, Deak M, Morrice NA, Boudeau J, et al. LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. *EMBO J* 2004;23:833–43.
- [10] Shang J, Chen LL, Xiao FX, Sun H, Ding HC, Xiao H. Resveratrol improves non-alcoholic fatty liver disease by activating AMP-activated protein kinase. *Acta Pharmacol Sin* 2008;29:698–706.
- [11] Park KG, Min AK, Koh EH, Kim HS, Kim MO, Park HS, et al. Alpha-lipoic acid decreases hepatic lipogenesis through adenosine monophosphate-activated protein kinase (AMPK)-dependent and AMPK-independent pathways. *Hepatology* 2008;48:1477–86.
- [12] Liu Y, Wan Q, Guan Q, Gao L, Zhao J. High-fat diet feeding impairs both the expression and activity of AMPKa in rat skeletal muscle. *Biochem Biophys Res Commun* 2006;339:701–7.
- [13] Collins SP, Reoma JL, Gamm DM, Uhler MD. LKB1, a novel serine/threonine protein kinase and potential tumour suppressor, is phosphorylated by cAMP-dependent protein kinase (PKA) and prenylated in vivo. *Biochem J* 2000;345:673–80.
- [14] Sapkota GP, Kieloch A, Lizcano JM, Lain S, Arthur JS, Williams MR, et al. Phosphorylation of the protein kinase mutated in Peutz–Jeghers cancer syndrome, LKB1/STK11, at Ser431 by p90<sup>RSK</sup> and cAMP-dependent protein kinase, but not its farnesylation at Cys433, is essential for LKB1 to suppress cell growth. *J Biol Chem* 2001;276:19469–82.
- [15] Xie Z, Dong Y, Scholz R, Neumann D, Zou MH. Phosphorylation of LKB1 at serine 428 by protein kinase C- $\zeta$  is required for metformin-enhanced activation of the AMP-activated protein kinase in endothelial cells. *Circulation* 2008;117:952–62.
- [16] Baas AF, Boudeau J, Sapkota GP, Smit L, Medema R, Morrice NA, et al. Activation of the tumour suppressor kinase LKB1 by the STE20-like pseudokinase STRAD. *EMBO J* 2003;22:3062–72.
- [17] Boudeau J, Baas AF, Deak M, Morrice NA, Kieloch A, Schutkowski M, et al. MO25 $\alpha/\beta$  interact with STRAD $\alpha/\beta$  enhancing their ability to bind, activate and localize LKB1 in the cytoplasm. *EMBO J* 2003;22:5102–14.
- [18] Boudeau J, Scott JW, Resta N, Deak M, Kieloch A, Komander D, et al. Analysis of the LKB1-STRAD-MO25 complex. *J Cell Sci* 2004;117:6365–75.
- [19] Winder WW, Wilson HA, Hardie DG, Rasmussen BB, Hutber CA, Call GB, et al. Phosphorylation of rat muscle acetyl-CoA carboxylase by AMP-activated protein kinase and cAMP-dependent protein kinase. *J App Physiol* 1997;82:219–25.
- [20] Koo SH, Flechner L, Qi L, Zhang X, Srean RA, Jeffries S, et al. The CREB coactivator TORC2 is a key regulator of fasting glucose metabolism. *Nature* 2005;437:1109–11.
- [21] Dentin R, Liu Y, Koo SH, Hedrick S, Vargas T, Heredia J, et al. Insulin modulates gluconeogenesis by inhibition of the coactivator TORC2. *Nature* 2007;449:366–9.
- [22] Shaw RJ, Lamia KA, Vasquez D, Koo SH, Bardeesy N, Depinho RA, et al. The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* 2005;310:1642–6.
- [23] Rodgers JT, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1 $\alpha$  and SIRT1. *Nature* 2005;434:113–8.
- [24] Deng XQ, Chen LL, Li NX. The expression of SIRT1 in non-alcoholic fatty liver disease induced by high-fat diet in rats. *Liver Int* 2007;27:708–15.
- [25] Sun C, Zhang F, Ge X, Yan T, Chen X, Shi X, et al. SIRT1 improves insulin sensitivity under insulin-resistant conditions by repressing PTP1B. *Cell Metab* 2007;6:307–19.
- [26] Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, et al. Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 2006;444:337–42.
- [27] Fujio J, Kushiyaama A, Sakoda H, Fujishiro M, Ogihara T, Fukushima Y, et al. Regulation of gut-derived resistin-like molecule beta expression by nutrients. *Diabetes Res Clin Pract* 2008;79:2–10.
- [28] Anai M, Funaki M, Ogihara T, Terasaki J, Inukai K, Katagiri H, et al. Altered expression levels and impaired steps in the pathway to phosphatidylinositol 3-kinase activation via insulin receptor substrates 1 and 2 in Zucker fatty rats. *Diabetes* 1998;47:13–23.
- [29] Lan F, Caciccedo JM, Ruderman N, Ido Y. SIRT1 modulation of the acetylation status, cytosolic localization, and activity of LKB1. Possible role in AMP-activated protein kinase activation. *J Biol Chem* 2008;283:27628–35.
- [30] Hou X, Xu S, Maitland-Toolan KA, Sato K, Jiang B, Ido Y, et al. SIRT1 regulates hepatocyte lipid metabolism through activating AMP-activated protein kinase. *J Biol Chem* 2008;283:20015–26.
- [31] Takemori H, Kajimura J, Okamoto M. TORC-SIK cascade regulates CREB activity through the basic leucine zipper domain. *FEBS J* 2007;274:3202–9.
- [32] Liu Y, Dentin R, Chen D, Hedrick S, Ravnskaer K, Schenk S, et al. A fasting inducible switch modulates gluconeogenesis via activator/coactivator exchange. *Nature* 2008;456:269–73.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)



ScienceDirect

Please cite this article in press as: Yoneda M, et al. Decreased SIRT1 expression and LKB1 phosphorylation occurs with long-term high-fat diet feeding, in addition to AMPK phosphorylation impairment in the early phase. *Obes Res Clin Pract* (2010), doi:10.1016/j.orcp.2010.02.002

