

ers who infected their children should have a marked abundance of non-PPP forms. In fact, our phylogenetic analysis demonstrated no PPP virus clusters while there are some clusters of non-PPP viruses (Fig. S1), suggesting more frequent transmission of non-PPP viruses than PPP virus. We are currently examining these possibilities.

If the PPP p26 molecule is associated with a number of fitness-decreasing properties, what maintains this less fit gene in the population? One possibility is that there may be direct HLA selection for proline residues at these p26 positions. All three proline sites lie within or adjacent to known HLA epitopes. The presence of these key proline residues could either block the host immune recognition of these epitopes or interfere with processing to release the epitope. HIV-1 clearly adapts to its current host's HLA system by changing recognized epitopes [41–45] and it is likely that HIV-2 is subject to the same host HLA selection. Thus virus evolution may be driven by a shortsighted response to HLA selection resulting in PPP p26 that in the long term results in reduced viral replication. An abundance of HLA alleles in Caio that select for PPP p26 may be responsible for the high frequency of controlled HIV-2 infections in Caio. A cross-sectional study on HLA associations with p26 variation is underway. Adaptation to the current host's HLA haplotype has important consequences for the design of T cell based vaccines and could be exploited in vaccines to encourage the evolution of less aggressive variants.

Acknowledgements

Contributors: John Townend provided statistical expertise. We appreciate the help and support of Ramu Sarge-Njie for providing serological data and Abraham Alabi for providing viral load data for these studies. We thank Tim Vincent and the staff of the Caio field station for facilitating the field work and Carla van Tienen, Bouke DeJong and Neil Berry for critical reading of the manuscript.

Appendix A. Supplementary data


Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.vaccine.2009.08.060.

References

- Barin F, M'Boup S, Denis F, Kanki P, Allan JS, Lee TH, et al. Serological evidence for virus related to simian T-lymphotropic retrovirus III in residents of west Africa. *Lancet* 1985;2(8469–70):1387–9.
- Clavel F, Guetard D, Brun-Vezinet F, Chamaret S, Rey MA, Santos-Ferreira AO, et al. Isolation of a new human retrovirus from West African patients with AIDS. *Science* 1986;233(4761):343–6.
- Lemey P, Pybus OG, Wang B, Saksena NK, Salemi M, Vandamme AM. Tracing the origin and history of the HIV-2 epidemic. *Proc Natl Acad Sci USA* 2003;100(11):6588–92.
- Rowland-Jones SL, Whittle HC. Out of Africa: what can we learn from HIV-2 about protective immunity to HIV-1? *Nat Immunol* 2007;8(4):329–31.
- Poulsen AG, Kvinesdal B, Aaby P, Molbak K, Frederiksen K, Dias F, et al. Prevalence of and mortality from human immunodeficiency virus type 2 in Bissau, West Africa. *Lancet* 1989;1(8642):827–31.
- Berry N, Ariyoshi K, Balfe P, Tedder R, Whittle H. Sequence specificity of the human immunodeficiency virus type 2 (HIV-2) long terminal repeat U3 region in vivo allows subtyping of the principal HIV-2 viral subtypes A and B. *AIDS Res Hum Retroviruses* 2001;17(3):263–7.
- Berry N, Ariyoshi K, Jaffar S, Sabally S, Corrah T, Tedder R, et al. Low peripheral blood viral HIV-2 RNA in individuals with high CD4 percentage differentiates HIV-2 from HIV-1 infection. *J Hum Virol* 1998;1(7):457–68.
- Ariyoshi K, Jaffar S, Alabi AS, Berry N, Schim van der Loeff M, Sabally S, et al. Plasma RNA viral load predicts the rate of CD4 T cell decline and death in HIV-2-infected patients in West Africa. *AIDS* 2000;14(4):339–44.
- Gao F, Yue L, Robertson DL, Hill SC, Hui H, Biggar RJ, et al. Genetic diversity of human immunodeficiency virus type 2: evidence for distinct sequence subtypes with differences in virus biology. *J Virol* 1994;68(11):7433–47.
- Ricard D, Wilkins A, N'Gum PT, Hayes R, Morgan G, Da Silva AP, et al. The effects of HIV-2 infection in a rural area of Guinea-Bissau. *AIDS* 1994;8(7):977–82.
- Grassly NC, Xiang Z, Ariyoshi K, Aaby P, Jensen H, Schim van der Loeff M, et al. Mortality among human immunodeficiency virus type 2-positive villagers in rural Guinea-Bissau is correlated with viral genotype. *J Virol* 1998;72(10):7895–9.
- Norrgrén H, Marquina S, Leitner T, Aaby P, Melbye M, Poulsen AG, et al. HIV-2 genetic variation and DNA load in asymptomatic carriers and AIDS cases in Guinea-Bissau. *J Acquir Immune Defic Syndr Hum Retrovirol* 1997;16(1):31–8.
- Padua E, Jenkins A, Brown S, Bootman J, Paixao MT, Almond N, et al. Natural variation of the Nef gene in human immunodeficiency virus type 2 infections in Portugal. *J Gen Virol* 2003;84(Pt 5):1287–99.
- Leligdowicz A, Yindom LM, Onyango C, Sarge-Njie R, Alabi A, Cotten M, et al. Robust Gag-specific T cell responses characterize viremia control in HIV-2 infection. *J Clin Invest* 2007;117(10):3067–74.
- Mortuza GB, Haire LF, Stevens A, Smerdon SJ, Stoye JP, Taylor IA. High-resolution structure of a retroviral capsid hexameric amino-terminal domain. *Nature* 2004;431(7007):481–5.
- Briggs JA, Grunewald K, Glass B, Forster F, Krausslich HG, Fuller SD. The mechanism of HIV-1 core assembly: insights from three-dimensional reconstructions of authentic virions. *Structure* 2006;14(1):15–20.
- Ganser BK, Li S, Klishko VY, Finch JT, Sundquist WI. Assembly and analysis of conical models for the HIV-1 core. *Science* 1999;283(5398):80–3.
- Ganser-Pornillos BK, Yeager M, Sundquist WI. The structural biology of HIV assembly. *Curr Opin Struct Biol* 2008;18(2):203–17.
- Gamble TR, Yoo S, Vajdos FF, von Schwedler UK, Worthylyake DK, Wang H, et al. Structure of the carboxyl-terminal dimerization domain of the HIV-1 capsid protein. *Science* 1997;278(5339):849–53.
- Mammiano F, Ohagen A, Hoglund S, Gottlinger HG. Role of the major homology region of human immunodeficiency virus type 1 in virion morphogenesis. *J Virol* 1994;68(8):4927–36.
- Alabi AS, Jaffar S, Ariyoshi K, Blanchard T, Schim van der Loeff M, Awasana AA, et al. Plasma viral load, CD4 cell percentage, HLA and survival of HIV-1, HIV-2, and dually infected Gambian patients. *AIDS* 2003;17(10):1513–20.
- Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 1999;41:95–8.
- Song H, Nakayama EE, Yokoyama M, Sato H, Levy JA, Shioda T. A single amino acid of the human immunodeficiency virus type 2 capsid affects its replication in the presence of cynomolgus monkey and human TRIM5alphas. *J Virol* 2007;81(13):7280–5.
- Kinomoto M, Yokoyama M, Sato H, Kojima A, Kurata T, Ikuta K, et al. Amino acid 36 in the human immunodeficiency virus type 1 gp41 ectodomain controls fusogenic activity: implications for the molecular mechanism of viral escape from a fusion inhibitor. *J Virol* 2005;79(10):5996–6004.
- Kinomoto M, Appiah-Opong R, Brandful JA, Yokoyama M, Nii-Trebi N, Ugly-Kwame E, et al. HIV-1 proteases from drug-naïve West African patients are differentially less susceptible to protease inhibitors. *Clin Infect Dis* 2005;41(2):243–51.
- Shirakawa K, Takaori-Kondo A, Yokoyama M, Izumi T, Matsui M, Ito K, et al. Phosphorylation of APOBEC3G by protein kinase A regulates its interaction with HIV-1 Vif. *Nat Struct Mol Biol* 2008;15(11):1184–91.
- Monaco-Malbet S, Berthet-Colominas C, Novelli A, Bataï N, Cheynet V, et al. Mutual conformational adaptations in antigen and antibody upon complex formation between an Fab and HIV-1 capsid protein p24. *Structure* 2000;8(10):1069–77.
- Baker D, Sali A. Protein structure prediction and structural genomics. *Science* 2001;294(5540):93–6.
- Ponder JW, Case DA. Force fields for protein simulations. *Adv Protein Chem* 2003;66:27–85.
- Lee K, Chu CK. Molecular modelling approach to understanding the mode of action of L-nucleosides as antiviral agents. *Antimicrob Agents Chemother* 2001;45(1):138–44.
- Guyader M, Emerman M, Sonigo P, Clavel F, Montagnier L, Alizon M. Genome organization and transactivation of the human immunodeficiency virus type 2. *Nature* 1987;326(6114):662–9.
- Stremlau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J. The cytoplasmic body component TRIM5alpha restricts HIV-1 infection in old World monkeys. *Nature* 2004;427(6977):848–53.
- Goff SP. HIV: replication trimmed back. *Nature* 2004;427(6977):791–3.
- Towers G, Bock M, Martin S, Takeuchi Y, Stoye JP, Danos O. A conserved mechanism of retrovirus restriction in mammals. *Proc Natl Acad Sci USA* 2000;97(22):12295–9.
- Stremlau M. GE Prize-winning essay, Why old World monkeys are resistant to HIV-1. *Science* 2007;318(5856):1565–6.
- Cowan S, Hatzioannou T, Cunningham T, Muesing MA, Gottlinger HG, Bieniasz PD. Cellular inhibitors with Fv1-like activity restrict human and simian immunodeficiency virus tropism. *Proc Natl Acad Sci USA* 2002;99(18):11914–9.
- Stremlau M, Perron M, Lee M, Li Y, Song B, Javanbakht H, et al. Specific recognition and accelerated uncoating of retroviral capsids by the TRIM5alpha restriction factor. *Proc Natl Acad Sci USA* 2006;103(14):5514–9.
- Creighton TE. *Proteins: Structures and Molecular Properties*. 2nd ed. New York: W.H. Freeman; 1992.
- Howard BR, Vajdos FF, Li S, Sundquist WI, Hill CP. Structural insights into the catalytic mechanism of cyclophilin A. *Nat Struct Biol* 2003;10(6):475–81.
- Berry N, Jaffar S, Schim van der Loeff M, Ariyoshi K, Harding E, N'Gom PT, et al. Low level viremia and high CD4% predict normal survival in a cohort of HIV type-2-infected villagers. *AIDS Res Hum Retroviruses* 2002;18(16):1167–73.
- Moore CB, John M, James IR, Christiansen FT, Witt CS, Mallal SA. Evidence of HIV-1 adaptation to HLA-restricted immune responses at a population level. *Science* 2002;296(5572):1439–43.

- [42] Brumme ZL, Tao I, Szeto S, Brumme CJ, Carlson JM, Chan D, et al. Human leukocyte antigen-specific polymorphisms in HIV-1 Gag and their association with viral load in chronic untreated infection. *AIDS* 2008;22(11):1277–86.
- [43] Bhattacharya T, Daniels M, Heckerman D, Foley B, Frahm N, Kadie C, et al. Founder effects in the assessment of HIV polymorphisms and HLA allele associations. *Science* 2007;315(5818):1583–6.
- [44] Kawashima Y, Pfafferoth K, Frater J, Matthews P, Payne R, Addo M, et al. Adaptation of HIV-1 to human leukocyte antigen class I. *Nature* 2009;458(7238):641–5.
- [45] Klenerman P, McMichael A. AIDS/HIV. Finding footprints among the trees. *Science* 2007;315(5818):1505–7.
- [46] Kimura M. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 1980;16(2):111–20.
- [47] Tamura K, Dudley J, Nei M, Kumar S. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 2007;24(8):1596–9.
- [48] Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987;4(4):406–25.
- [49] Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbour-joining method. *Proc Natl Acad Sci USA* 2004;101(30):11030–5.
- [50] Wilkins A, Ricard D, Todd J, Whittle H, Dias F, Paulo Da Silva A. The epidemiology of HIV infection in a rural area of Guinea-Bissau. *AIDS* 1993;7(8):1119–22.

REVIEW



Anti-retroviral activity of TRIM5 α

Emi E. Nakayama* and Tatsuo Shioda

Department of Viral Infections, Research Institute for Microbial Disease, Osaka University, 3-1 Yamadaoka, Suita, Osaka 565-0871, Japan

SUMMARY

Human immunodeficiency virus type 1 (HIV-1) shows a very narrow host range limited to humans and chimpanzees. Experimentally, HIV-1 does not infect Old World monkeys, such as rhesus (Rh) and cynomolgus (CM) monkeys, and fails to replicate in activated CD4 positive T lymphocytes obtained from these monkeys. In contrast, simian immunodeficiency virus isolated from a macaque monkey (SIVmac) can replicate well in both Rh and CM. In 2004, tripartite motif 5 α (TRIM5 α) was identified as a host factor which plays an important role in the restricted host range of HIV-1. Rh and CM TRIM5 α restrict HIV-1 infection but not SIVmac, while in comparison, anti-viral activity of human TRIM5 α against those viruses is very weak. TRIM5 α consists of the RING, B-box 2, coiled-coil and SPRY (B30.2) domains. The RING domain is frequently found in E3 ubiquitin ligase and TRIM5 α is degraded via the ubiquitin-proteasome pathway during HIV-1 restriction. TRIM5 α recognises the multimerised capsid (viral core) of an incoming virus by its α -isoform specific SPRY domain and is believed to be involved in innate immunity to control retroviral infection. Differences in amino acid sequences in the SPRY domain of TRIM5 α of different monkey species were found to affect species-specific restriction of retrovirus infection, while differences in amino acid sequences in the viral capsid protein determine viral sensitivity to restriction. Accurate structural analysis of the binding surface between the viral capsid protein and TRIM5 α SPRY is thus required for the development of new antiretroviral drugs that enhance anti-HIV-1 activity of human TRIM5 α . Copyright © 2009 John Wiley & Sons, Ltd.

Received: 24 August 2009; Revised: 10 September 2009; Accepted: 10 September 2009

INTRODUCTION

Human immunodeficiency virus type 1 (HIV-1), a major causative agent of acquired immunodeficiency syndrome (AIDS), belongs to the genera lentivirus of the family *Retroviridae*. HIV-1 is thought to have been introduced into the human

population from chimpanzees [1] and has a very narrow host range limited to humans and chimpanzees. Experimentally, HIV-1 fails to replicate in activated CD4 positive T lymphocytes obtained from Old World monkey (OWM)s, such as rhesus (Rh) [2,3] and cynomolgus (CM) monkeys [4,5]. In contrast, another lentivirus simian immunodeficiency virus isolated from sooty mangabey (SIVsm) and simian immunodeficiency virus isolated from African green monkey (SIVagm) replicate in their natural hosts [6]. Simian immunodeficiency virus isolated from a macaque monkey (SIVmac), with a genome that has 55% nucleotide sequence homology to that of HIV-1, was evolved from SIVsm in macaques in captivity, and replicates efficiently in Rh [2,3] and CM [4,5]. The restricted host range of HIV-1 has greatly hampered its use in animal experiments, and, thus the development of prophylactic vaccines against HIV-1 infection. In 2004, tripartite motif 5 α (TRIM5 α) was identified as a host factor that plays an important role in the restricted host range of HIV-1. In this review

*Corresponding author: E. E. Nakayama, Department of Viral Infections, Research Institute for Microbial Disease, Osaka University, 3-1 Yamadaoka, Suita, Osaka 565-0871 Japan.
E-mail: emien@biken.osaka-u.ac.jp

Abbreviations used

A, alanine; AGM, African green monkey; AIDS, acquired immune deficiency syndrome; APOBEC, ApoB mRNA editing catalytic subunit; CA, capsid protein; CM, cynomolgus monkey; CsA, cyclosporine A; CypA, cyclophilin A; HIV-1, human immunodeficiency virus type 1; HIV-2, human immunodeficiency virus type 2; L4/5, a loop between α -helices 4 and 5 of CA; L6/7, a loop between α -helices 6 and 7 of CA; NC, nucleocapsid protein; N-MLV, N-tropic murine leukemia viruses; OWM, Old World monkey; P, proline; Q, glutamine; R, arginine. Rh, rhesus monkey; RING, really interesting new gene; SPRY, a sequence repeat in the dual-specificity kinase *splA* and ryanodine receptors; SHIV, Chimeric virus between SIVmac and HIV-1; SIVmac, simian immunodeficiency virus isolated from macaque; TRIM5 α , tripartite motif 5 α ; TRIMCyp, TRIM5 and CypA fusion protein; VL, viral load; VSV-G, vesicular stomatitis virus glycoprotein

article, recent findings in TRIM5 α research are summarised and details of the molecular mechanisms of HIV-1 restriction by TRIM5 α are discussed.

IDENTIFICATION OF TRIM α AS A RESTRICTION FACTOR AGAINST HIV-1 IN OWM CELLS

Several earlier studies have suggested that the block of HIV-1 replication in OWM cells occurs at a post-entry step [2,3,7] and appears to result from a failure to initiate reverse transcription [3]. Studies of HIV-1 and SIVmac chimera have suggested that restriction determinants lie within the HIV-1 p24 capsid protein (CA) [8–11]. The block was still observed in CD4-negative monkey cells infected with HIV-1 pseudotyped with vesicular stomatitis virus glycoprotein (VSV-G) (Figure 1) but was overridden by high-dose infection with

VSV-G-pseudotyped virus or virus-like particles lacking genomic RNA [12–15]. Importantly, resistance against HIV-1 infection was shown to be dominant in heterokaryons between human and OWM cells, suggesting the presence of inhibitory factor(s) against HIV-1 infection in OWM cells [14].

In 2004, the screening of a Rh cDNA library identified TRIM5 α , a component of cytoplasmic bodies, as a factor that confers resistance to HIV-1 infection [16]. Rh and CM TRIM5 α restrict HIV-1 infection but not SIVmac [16,17]. In contrast, human TRIM5 α is almost powerless to restrict the aforementioned viruses, but potently restricts N-tropic murine leukemia viruses (N-MLV), which belong to genera Gammaretrovirus. It was previously shown that human cells are resistant to infection with N-MLV and the presence in human




Figure 1. Old World monkey (OWM) cells are resistant to HIV-1 infection. Human C143 and 293T cell line were highly sensitive to vesicular stomatitis virus glycoprotein-pseudotyped HIV-1 infection, while rhesus monkey (Rh) LLC-MK2 and African green monkey (AGM) CV1 cell lines were resistant. When extremely high doses of virus were inoculated, cells became sensitive to infection, suggesting that the intrinsic restriction factor(s) were saturated with virions

Anti-retroviral activity of TRIM5 α

cells of a virtual restriction factor known as Ref1 was posited. It is now widely accepted that the presence of human TRIM5 α substantiates that of the restriction factor Ref1 [18–21]. On the other hand, African green monkey (AGM) cells have been shown to possess another factor, Lv1, which restricts both HIV-1 and SIVmac infection and we and others identified the factor as AGM TRIM5 α [17,18]. AGM TRIM5 α fails to restrict SIVagm. Unlike humans and other OWMs, pig-tailed monkeys lack expression of TRIM5 α , but instead express TRIM5 θ and TRIM5 η , which lack anti-HIV-1 activity [22]. It is now known that type I interferons up-regulate the transcription of TRIM5 α in human [23] and monkey cells [24] and this in turn enhances restriction activity against N-MLV [24,25].

TRIPARTITE MOTIF OF TRIM5 α

The human genome contains approximately 70 genes of the TRIM family, which characteristically encode a tripartite protein motif [26–29]. This tripartite motif consists of a really interesting new gene (RING) zinc-finger domain, one or two B-box zinc-finger domains and an α -helical coiled-coil domain. *TRIM* genes are scattered throughout the human genome, while the *TRIM5* locus lies in a small cluster of four related *TRIM* genes including *TRIM6*, *TRIM34* and *TRIM22* [30]. Although the functions of most TRIM family members are still unknown, several TRIM proteins including TRIM1, TRIM19 (PML), TRIM22 and TRIM32 reportedly have anti-viral effects (reviewed in Reference [26]). Especially, TRIM19 can suppress broad spectrum of viruses such as herpes simplex virus type 1 and lymphocytic choriomeningitis virus [26]. TRIM21 is a trimeric protein that binds IgG Fc via the C-terminal of the B30.2 domain [31]. Subcellular localisation of the TRIM proteins varies among members of the TRIM family [27]: TRIM19, 24 and 27 are associated with nuclear bodies and TRIM1 and 18 with microtubules. TRIM5 α was first identified as a cytoplasmic body protein [27], but diffuse expression in cytoplasm has proved important for its anti-viral activity [32].

As shown in Figure 2, TRIM5 α consists of RING, B-box 2, coiled-coil and SPRY (B30.2) domains [27]. The RING domain containing proteins possess E3 ubiquitin ligase activity [33] and the intact RING domain of TRIM5 α was thought to be essential for retrovirus restriction (see below). The intact




Figure 2. TRIM5 α protein. TRIM5 α protein contains RING, B-box 2, and coiled-coil domains, the three signature domains of the TRIM gene family. The α isoform possesses a SPRY domain sometimes referred to as a B30.2 domain. V1, V2 and V3 denote variable regions 1, 2 and 3 among monkey species, respectively. A histidine-to-tyrosine substitution at position 43 (H43Y) and a glycine-to-arginine substitution at position 110 (G110R) of human TRIM5 (shown in red) modulate anti-viral activity against retroviruses *in vitro*. Position 332 in human TRIM5 α is arginine (R) and no polymorphism was reported in human genome. In contrast, proline (P)-to-glutamine (Q) substitution in rhesus monkey (Rh) and R-to-P or -Q substitution in sooty mangabeys were found (R332P/Q). A 339th-TFP-341st to -Q polymorphism (TFP339Q) that reduces the anti-human immunodeficiency virus type 2 (HIV-2) activity was found in Rh TRIM5 α .

B-box 2 domain is also required for TRIM5 α mediated antiviral activity since the restrictive activity of TRIM5 α is diminished by several amino acid substitutions in the B-box 2 domain [34]. TRIM5 α has been shown to form a trimer [35,36] or a dimer [37,38], while the B-box 2 domain mediates higher-order self-association of Rh TRIM5 α oligomers. This self-association increases the efficiency of TRIM5 α binding to the retroviral CA, thus potentiating restriction of retroviral infection [39,40]. The coiled-coil domain of TRIM5 α has been identified as important for the formation of homo-oligomers [35], and homo-oligomerisation of TRIM5 α as essential for antiviral activity [36,41]. The SPRY domain is specific for an α isoform among at least three splicing variants transcribed from the *TRIM5* gene. *TRIM5 γ* and *TRIM5 δ* lack the SPRY domain because of alternative splicing and their functions remain unknown. Exogenously expressed TRIM5 γ is unstable [42] and over-expression of the TRIM5 protein lacking the SPRY domain dominant-negatively suppresses the anti-viral activity of the intact TRIM5 α through hetero-oligomerisation [41,42].

SPRY (B30.2) DOMAIN OF TRIM5 α

The C-terminal halves of TRIM family proteins are variable and half of them, including the TRIM5 α , encode B30.2 (SPRY or PRYSPRY) domain [27]. Soon after identification of TRIM5 α as a restriction

factor of Rh, many studies found that differences in the amino acid sequences in the TRIM5 α SPRY domain of different monkey species affect the species-specific restriction of retrovirus infection [17,43–50].

Studies on human and Rh recombinant TRIM5 α s have shown that the determinant of the species-specific restriction against HIV-1 infection resides in variable region 1 (V1) of the SPRY domain [43,44]. We found that 17-amino-acid residues and adjacent 20-amino-acid duplication in the V1 of AGM TRIM5 α determined species-specific restriction against SIVmac [17]. Interestingly, a study comparing human and Rh TRIM5 α showed that a single change from arginine (R) to proline (P) at the 332nd position in the V1 of human TRIM5 α (R332P) conferred potent restriction ability against not only HIV-1 but also SIVmac239 [49,50]. In the case of human immunodeficiency virus type 2 (HIV-2) infection, we found that three amino acid residues of TFP at the 339th to 341st positions of Rh TRIM5 α V1 are important for restricting particular HIV-2 strains which are still resistant to CM TRIM5 α [45].

A study comparing orangutan and gorilla TRIM5 α s showed that two amino acid residues at the 385th and 389th positions in the variable region 2 (V2) of SPRY domain of orangutan TRIM5 α are important for restriction against HIV-1 and SIVmac [46]. We found that one amino acid residue at the 385th (baboon) or 383rd (CM) position in V2 of the SPRY domain of TRIM5 α also affects its restriction ability against HIV-2 [47]. A computer-assisted 3-D model of the TRIM5 α SPRY domain showed that V1 and V2 are located in the loops at the surface of the SPRY domain and the structure composed of the V1 and V2 regions is thought to be important for TRIM5 α restriction (Figure 3).

Furthermore, a comparison of human and Rh TRIM5 α restriction of N-MLV showed that the amino acid residues of human TRIM5 α at the 409th and 410th positions in the variable region 3 (V3) of SPRY domain are important for restricting N-MLV [48].

Finally, biochemical studies have shown that TRIM5 α associates with CA in detergent-stripped N-MLV virions [51] or with an artificially constituted HIV-1 core structure composed of the capsid-nucleocapsid (CA-NC) fusion protein in a SPRY domain dependent manner [52]. The SPRY domain is thus thought to recognise viral cores.




Figure 3. Structure of SPRY domain of TRIM5 α . This 3-D model of the baboon SPRY domain was constructed with a homology-modeling technique based on mouse TRIM21 [131]. Position 332 in human TRIM5 α is the same as position 334 in baboon TRIM5 α . SFP motif in V1 and P residue at position 385 in V2 are shown

VIRAL DETERMINANT OF TRIM5 α SENSITIVITY

To determine the CA region that interacts with TRIM5 α , we focused on HIV-2, which strongly resembles SIVmac [53]. Previous studies have shown that HIV-2 strains vary widely in their ability to grow in OWM cells such as baboon, Rh and CM [54–58] and HIV-2 isolates with various growth capabilities in OWM cells were evaluated for their sensitivity to CM TRIM5 α [59]. We found that viral sensitivity to CM TRIM5 α inversely correlates with growth capability in OWM cells. Sequence analysis showed that the CM TRIM5 α -sensitive viruses had proline (P) at the 119th or 120th position of CA, while the CM TRIM5 α -resistant viruses had either alanine (A) or glutamine (Q) at the same position. Replacing the proline of a CM TRIM5 α -sensitive HIV-2 molecular clone with either alanine or glutamine changed the phenotype from sensitive to resistant (Figure 4) and the mutant viruses replicated well in the presence of CM TRIM5 α . The reverse was observed when the glutamine of a resistant SIVmac molecular clone was replaced with proline. Similar results, although to a lesser extent, were observed when human TRIM5 α was used [59]. These results indicate that a single amino acid at the 119th or 120th




Figure 4. A single amino acid of HIV-2 capsid (CA) affects its replication in the presence of cynomolgus monkey (CM) TRIM5 α . HIV-2 GH123 replication was restricted in CM-TRIM5 α expressing human CD4 T cell line (CM-TRIM5 α ; black circles). The mutant GH123-P120A and GH123-P120Q viruses were generated by changing a single amino acid proline (P) at position 120 of GH123 CA to alanine (A) or glutamine (Q). These mutant viruses replicated in CM-TRIM5 α expressing cells as efficiently as in cells lacking TRIM5 α expression (CM-SPRY(-); white circles)



position of HIV-2 CA drastically affects viral sensitivity to TRIM5 α .

A computer-assisted 3-D model of the HIV-2 CA showed that the 119th or 120th position is located in the loop between α -helices 6 and 7 (L6/7, Figure 5A). Previously, a single amino acid substitution at the 110th position of N-MLV CA had been shown to determine viral susceptibility to Fv1 [60], another restriction factor present in certain strains of mice [61] as well as to Ref1 (human TRIM5 α) [18,20,21]. The recently published 3-D structure of MLV CA [62,63] revealed that the 110th position of N-MLV CA is located at a position in the surface-exposed loop analogous to the 119th or 120th position of HIV-2 CA. As mentioned above, V3 in the SPRY domain of human TRIM5 α reportedly plays an important role in the restriction of N-MLV [48], whereas V1 in the SPRY domain of OWM TRIM5 α s determines restriction specificity against HIV-1 and SIVmac [17,43,44,49,50]. These results indicate that the surface-exposed loop of CA is important for recognition by cellular restriction factors, even though critical amino acid residues in human TRIM5 α for N-MLV restriction are different from those in CM TRIM5 α for HIV-2.

HIV-2 is assumed to have originated from SIVsm as the result of zoonotic events involving monkeys and humans [53]. Almost all the SIV isolates in the Los Alamos database contain glutamine at the position corresponding to the 119th or 120th position of HIV-2 CA. In contrast, HIV-2

strains possess a mixture of glutamine, alanine and proline at the corresponding position. It is thus likely that glutamine-to-alanine or glutamine-to-proline substitutions occurred after the hypothesised zoonotic transfer of virus from monkeys to humans. According to this hypothesis, a single nucleotide change in the second position of the glutamine codons (CAA or CAG) would generate proline codons (CCN), and an additional single nucleotide change in the first position in CCN would generate alanine codons (GCN). Most likely, the glutamine residue first changed into proline residue in humans as a result of certain pressures from the human immune system in the absence of strong pressure from OWM TRIM5 α . Since HIV-2 strains with proline residue show moderate sensitivity to human TRIM5 α , an additional change would have to have occurred to generate alanine residue for better replication in human populations.

Does amino acid residue at the 119th or 120th position in HIV-2 CA affect HIV disease in infected individuals? It is known that HIV-1 and HIV-2 have distinct natural histories, levels of viremia, transmission rates and disease associations despite strong sequence homology between the two viruses [64]. Although some HIV-2-infected patients progress to AIDS, the infection is controlled in the majority of patients [65,66] and those with low viral load (VL) have a much longer survival [67]. A detailed sequence analysis of HIV-2 CA variations within a large community cohort



in Guinea Bissau comprising both high- and low-VL patients demonstrated that CA from viruses in low-VL patients had proline residues at the 119th or 120th position, but in patients with higher VL, the 119th or 120th position was frequently occupied by non-proline residues. Stratification of the subjects according to the presence or absence of proline at the 119th or 120th position showed a 3-fold difference in the median VL of the two groups (Figure 5B). These results demonstrate that HIV-2 replication in infected individuals can be linked to CA variation and human TRIM5 α sensitivity [68].

In a case of Rh TRIM5 α , Ylinen *et al.* mapped one of the determinants of TRIM5 α sensitivity in a loop between α -helices 4 and 5 (L4/5) of HIV-2 [69]. They replaced the L4/5 of the SIVmac CA (CA 78 to CA97) with that of HIV-2 CA (CA78 to CA98) in the SIVmac backbone and found that the resultant mutant virus showed impaired growth ability in Rh cells compared with the parental SIVmac239. Although we could confirm this finding, the reciprocal virus with SIVmac239 CA L4/5 in the HIV-2 backbone did not gain resistance against Rh TRIM5 α . However, we recently found that multiple components including L4/5 and L6/7 in the N-terminal half of SIVmac CA contributed to the escape of SIVmac from Rh TRIM5 α (Kono *et al.* manuscript in preparation). Lin and Emerman also reported that SIVagmTAN with both HIV-1 L4/5 and L6/7 was susceptible to Rh TRIM5 α restriction [70].

How do these regions in CA interact with specific regions in SPRY domain of TRIM5 α ? For this, it is necessary to obtain detailed and accurate structural information of the complex between viral CA

Figure 5. A single amino acid of HIV-2 CA affects 3-D structure of CA and its replication in infected individuals. (A) The 3-D models of six HIV-2 CAs were constructed with the homology-modeling technique. N and C indicate the amino termini and carboxyl termini, respectively. The ribbons represent the backbones of CA, and the 12 colour-coded-helices are numbered. P and A indicate a loop between α -helices 6 and 7 (L6/7) with a proline residue (in red) and an alanine residue (in blue), respectively, at position 119. (B) HIV-2 viral load (VL) correlates with amino acids variation at position 119 or 120 of CA. The differences in numbers were caused by the 1 amino acid insertion at the N-terminal part of CA in some HIV-2 isolates. Log-transformed VL for each patient in a West African community cohort is plotted. The patients were stratified according to the presence or absence of proline at position 119 or 120 (119P or non-P). Median VL of each group is indicated by a horizontal bar

and TRIM5 α SPRY. Structural data on the binding surface between HIV-1 CA and TRIM5 α SPRY of various primate species would also help to develop small molecular compounds that enhance anti-HIV-1 activity of human TRIM5 α , and such compounds could be used as new therapeutic measures in HIV-1 infection.

ROLE OF CYCLOPHILIN A (CYPA)

The hydrophobic pocket of CypA makes direct contact with the proline residue at the 90th position and adjacent residues on the L4/5 of HIV-1 CA [71–73]. This interaction can be disrupted experimentally by mutations that alter CA proline 90 or adjacent residues [72], and by competitive inhibitors of the interaction including cyclosporine A (CsA) [74–76]. Disruption of CypA-CA interaction reduces HIV-1 susceptibility in human cells [72,74,76–80], with the block occurring early, at the time of reverse transcription [74]. These data have led to the hypothesis that, by binding to CA, CypA protects HIV-1 against antiviral restriction activity in human cells [81,82].

In OWM cell lines, however, CsA treatment has been reported to increase HIV-1 replication [15,83,84]. Concerning the relationship between CypA and TRIM5 α , two groups reported that the simultaneous knock-down of CypA and TRIM5 in OWM cells caused minimal additional increase of HIV-1 infection compared with knock-down of TRIM5 α alone, suggesting that CypA acts in trans to promote TRIM5 α -mediated restriction of HIV-1 [85,86]. However, we demonstrated that exogenous expression of CypA suppresses HIV-1 infection in OWM cells in the absence of functional TRIM5 α and that the inhibitory activity of CypA depends upon the interaction of CypA moiety with HIV-1 CA, while disruption of CypA-CA interaction by CsA treatment enhanced the HIV-1 susceptibility of OWM cells even in the absence of functional TRIM5 α [41]. It is noteworthy that one of the two groups mentioned earlier observed slightly higher HIV-1 infection after simultaneous knock-down of CypA and TRIM5 than after knock-down of TRIM5 alone at the low-dose virus infection [85], suggesting that there is a TRIM5 α -independent anti-HIV-1 activity mediated by CypA in OWM cells.

As already mentioned, L4/5 of SIVmac and HIV-2 CA is one of the determinants for Rh TRIM5 α restriction, but it should be noted that

SIVmac and HIV-2 CA L4/5s, unlike HIV-1 CA L4/5, do not bind to CypA [72].

CHIMERIC VIRUSES BETWEEN SIVMAC AND HIV-1 (SHIV)

In order to establish a monkey model of HIV-1/AIDS study, various SHIVs have been constructed and tested for their replicative capability in simian cells. The first SHIV was generated with a genetic backbone of SIVmac containing HIV-1 *tat*, *rev*, *vpu* and *env* genes [11]. Although such an SHIV is useful for the analysis of humoral immune responses against Env protein [87–89], SHIVs containing other HIV-1 structural proteins, especially the Gag-Pol protein, were urgently needed since it is well known that the cellular immune response against Gag is important for disease control [90–92].

Construction of an SHIV with a minimal segment of SIVmac was reported by Kamada *et al.* [93]. This virus (NL-ScaVR, Figure 6A) contains the L4/5 of CA and the entire *vif* segment of SIVmac and was designed to escape from restrictions mediated by ApoB mRNA editing catalytic subunit (APOBEC) 3G and CypA in OWM cells. APOBEC3G modifies the minus strand viral DNA during reverse transcription, resulting in impairment of viral replication [94–96], but this activity could be counteracted with the viral protein Vif [97–99]. Although HIV-1 Vif can potently suppress human APOBEC3G, it is not effective against Rh APOBEC3G, which explains at least partly the restriction of HIV-1 replication in monkey cells. As mentioned above, CypA directly binds to L4/5 of HIV-1 CA but not to SIVmac CA and augments HIV-1 infection in human cells but inhibits its replication in OWM cells [15,84,100].

NL-ScaVR could replicate in pig-tailed monkey primary CD4+ T cells as well as in the CM T cell line HSC-F. In both HSC-F and primary CD4+ T cells, NL-ScaVR grew to lower titers than did SIVmac [93], and inoculation into pig-tailed monkey with this SHIV did not cause CD4+ T cell depletion or any clinical symptoms [101]. In contrast, stHIV-1, another SHIV carrying 202 amino acid residues of SIVmac CA and *vif* generated by Hatzioannou *et al.*, could replicate more efficiently in Rh cells [102] and contains more of the SIVmac-derived CA segment than the virus reported by Kamada *et al.*, suggesting the presence outside of L4/5 of another determinant of intrinsic restriction factor(s) in CA.




Figure 6. Modification of L6/7 of CA improves replication in CM cells of HIV-1 derivative. (A) Structure of chimeric viruses between simian immunodeficiency virus isolated from macaque (SIVmac) and HIV-1. White bars denote HIV-1 (NL4-3) and gray bars SIVmac239 sequences. (B) Equal amounts of NL-ScaVR (white triangles: virus with SIVmac L4/5 and *vif*), or NL-ScaVR6/7S (black triangles: virus with SIVmac L4/5, L6/7, and *vif*) were inoculated into CM CD4-positive T cell line HSC-F, and culture supernatants were collected periodically. The levels of reverse transcriptase (RT) in culture supernatants were measured by ELISA

As mentioned earlier, a single amino acid in L6/7 of HIV-2 CA was identified as a determinant of the susceptibility of HIV-2 to CM TRIM5 α . Introduction of the entire SIVmac L6/7 of CA into the previously constructed version of HIV-1 derivatives with SIVmac L4/5 of CA and *vif* (NL-ScaVR6/7S, Figure 5A) caused only four amino acid changes in CA but showed improvement of replication capability of HIV-1 in CM cell line HSC-F (Figure 6B) and primary CD4+ T cells. While the high-dose inoculation of particles of wild type HIV-1 into monkey cells saturated endogenous TRIM5 α and enhanced the subsequent infection with HIV-1, the introduction of both L4/5 and L6/7 of SIVmac greatly impaired the ability of HIV-1 particles to saturate TRIM5 α [103]. These observations support the notion that

TRIM5 α recognises the overall structures composed of L4/5 and L6/7 of HIV-1 CA.

PROTEASOME-DEPENDENT AND PROTEASOME-INDEPENDENT MECHANISMS OF RESTRICTION

The RING domain containing proteins possess E3 ubiquitin ligase activity [33]. In fact, Rh TRIM5 α was found to be poly-ubiquitinated and degraded rapidly via the ubiquitin-proteasome pathway [104], while disruption of the RING domain eliminated its auto-ubiquitination [104]. Furthermore, it was demonstrated that TRIM5 α is degraded via the ubiquitin-proteasome pathway during HIV-1 restriction [105]. As for the antiviral activity of TRIM5 α , however, deletion of the RING domain only partially attenuated its anti-HIV-1 activity [34,106], and modulation of E1 ubiquitin-activating enzyme expression did not affect TRIM5 α -mediated restriction activity in a temperature dependent cell line [106]. Moreover, proteasome inhibitors did not affect TRIM5 α mediated HIV-1 restriction but caused HIV-1 to generate late reverse transcripts during TRIM5 α mediated restriction [107,108]. Proteasome-dependent and -independent pathways may therefore be involved in HIV-1 restriction by Rh TRIM5 α (Figure 7). A proteasome-dependent HIV-1 restriction would mean co-degradation of TRIM5 α with incoming viral capsids in proteasomes but the exact molecular mechanism of the proteasome-independent pathway is still unclear at present. It was previously shown that the incubation of *in vitro* assembled CA composed of recombinant HIV-1 CA-NC fusion protein with the purified TRIM5-21R protein containing the Rh TRIM5 α B-box, coiled-coil and SPRY domains and the TRIM21 RING domain caused apparent breaks in the CA structure without any other cellular components [38]. It is thus likely that direct binding of Rh TRIM5 α proteins to incoming HIV-1 CA proteins causes CA disassembly, which is observed as proteasome-independent restriction.

We recently found that the contribution of the RING domain to retrovirus restriction by TRIM5 α differed among viral species. SIVmac completely escaped from attacks by RING mutants of AGM TRIM5 α that could still moderately restrict HIV-1 infection. Addition of proteasome inhibitor MG132 had no effect at all on the anti-HIV-1 activity of AGM TRIM5 α , but disrupted at




Figure 7. Current model of TRIM5 α restriction pathways. (Left) A proteasome-dependent pathway. Oligomerized TRIM5 α recognizes the incoming viral core through its SPRY domain and is poly-ubiquitinated in a RING-dependent manner. Ubiquitinated TRIM5 α and the viral core complex are degraded in proteasome, while MG132 inhibits the activity of proteasome. (Right) A proteasome-independent pathway. Although RING mutant TRIM5 α cannot ubiquitinate, HIV-1 CA and Rh or CM TRIM5 α binding via the SPRY domain causes destruction of the viral core without any other cellular factors

least partly the anti-SIVmac activity of AGM TRIM5 α . These results indicate that SIVmac is restricted by AGM TRIM5 α mainly in a proteasome-dependent manner, whereas HIV-1 restriction by AGM, Rh and CM TRIM5 α is both proteasome-dependent and -independent (Maegawa *et al.* submitted).

POLYMORPHISMS IN HUMAN TRIM5 GENE

HIV-1 infection in humans is generally characterised by a long-term, chronic disease course gradually progressing to AIDS. Polymorphisms in human CCR5 and other genes reportedly affect the susceptibility to HIV-1 transmission and/or the rate of disease progression to AIDS [109,110]. Sawyer *et al.* reported a common histidine-to-tyrosine polymorphism at the 43rd amino acid residue (H43Y) of the human TRIM5 α gene [111]. This single nucleotide polymorphism (SNP) is located in the RING domain (Figure 2) and was shown to greatly reduce the ability of TRIM5 α to restrict N-MLV [111]. Several studies have reported that the anti-HIV-1 activity of TRIM5 α with 43Y was lower than that with 43H *in vitro* [111,112] which

was confirmed in a previous study of ours, although the difference in anti-HIV-1 activity was very small [113].

Associations of H43Y with the rate of progression to AIDS have been tested in several studies, but with inconsistent results [111–115]. Previously, we compared 43Y frequencies of HIV-1 infected long-term non-progressors and standard progressors in France and Japan and failed to find any differences in 43Y frequency between the two groups in either country, indicating that this SNP does not affect disease progression [113]. In spite of the lower anti-HIV-1 activity of TRIM5 α with 43Y, Javanbakht *et al.* have found a paradoxical protective effect of TRIM5 α with 43Y against HIV-1 transmission in African-American [111,112]. Interestingly, we also found that the 43Y-allele was found less frequently in Japanese and Indian HIV-1 infected subjects than in the ethnic-matched controls [116]. The reasons for the discrepancy between the epidemiological and functional effects of H43Y remain unclear at present and further studies are required to clarify the impact of H43Y on susceptibility to HIV-1 transmission and/or rate of progression to AIDS.

In contrast to HIV-1, HIV-2 replication was moderately inhibited by the wild type human TRIM5 α [59], but not at all by human TRIM5 α with 43Y (Maegawa *et al.* submitted). This suggests that H43Y polymorphism has a stronger impact on HIV-2 than on HIV-1 in infected individuals. It would thus be of interest to investigate whether this allele affects the rate of disease progression in HIV-2-infected individuals.

In the B-box domain, we recently found in Japan a novel and rare glycine-to-arginine substitution at the 110th position of TRIM5 α (G110R) (Figure 2) and this 110R allele was observed more frequently in HIV-1-infected subjects than in controls. As was observed epidemiologically, this substitution weakened the anti-HIV-1 and anti-HIV-2 activity *in vitro* [116]. These findings together indicate that anti-HIV-1 activity of human TRIM5 α can hardly protect human beings from an HIV-1 pandemic, but it may affect HIV-1 transmission.

EVOLUTION OF TRIM5 GENE

TRIM5 homologs have been found in the genomes of primates, mouse, rat, dog, cow and pig, but not in chicken [30,117]. TRIM5 homolog genes are found in large numbers in cow, rat and mouse, but the human genome contains only a single TRIM5 gene, and the dog's homolog was inactivated by a transposon [118]. No antiviral activity against eight retroviruses has been reported for the TRIM5 homologs of mouse (TRIM12 and TRIM30) [117] and mouse TRIM30 targets TAB2 (TAK1-binding protein 2) for degradation [119]. It is speculated that human TRIM5 α also interact with a molecule of certain signal transduction cascade.

The TRIM5 gene varies considerably among primate species. The distribution of positively selected amino acid site is located in the SPRY domain and coiled-coil domains [44,120,121]. It is thus not surprising that the very beginning of SPRY domain (V1) is highly variable since TRIM5 α interacts with the retroviral core through this region as discussed above, while the main pressure for positive selection may be endogenous retroviruses [122]. In contrast to that for SPRY polymorphisms, the hypothesis to explain positive selection operating on the coiled-coil domain is too speculative, since the well-established function of the coiled-coil domains is only to promote TRIM5 α oligomerisation [35]. Interestingly, in Rh, there is a 339th-

TFP-341st to Q polymorphism in TRIM5 α (Figure 2) [121], which reduces the anti-HIV-2 activity [45]. Position 332 in human TRIM5 α is arginine (R) and there is no polymorphism in the human genome, but P/Q was found in Rh and R/P/Q polymorphism in sooty mangabeys [121,123] (Figure 2). H43Y polymorphism was frequently found in human but not in monkey species [118]. We recently found that H43Y mutation in AGM TRIM5 α failed to affect its antiviral activity (Nakayama *et al.*). Since H43Y substitution seems to be neutral in monkey TRIM5 α , there is no advantage in its accumulation in monkey population.

Among New World monkeys, owl monkeys possess CypA as a fusion protein with TRIM5 (TRIMCyp) as a result of LINE-1-mediated retrotranspositional insertion [124,125]. CypA can bind to the CA of HIV-1, so that the TRIMCyp expressed in owl monkey cells recognises the HIV-1 core and shows an anti-HIV-1 effect. Retrotransposition of CypA into the TRIM5 gene also occurred independently in OWM Rh and pig-tailed monkeys [22,126–129]. Rh TRIMCyp restricts infection of HIV-2 and feline immunodeficiency virus but not HIV-1 [129], and TRIMCyp expressed in pig-tailed monkey also lacks anti-HIV-1 activity [22,126,127]. A lack of functional TRIM5 α expression in pig-tailed monkey enabled Hatzioannou *et al.* to construct an SHIV strain that differs from HIV-1 only in the *vif* gene and can efficiently replicate in pig-tailed monkeys [130].

CONCLUSION

Within 5 years, many studies have emphasised the impact of TRIM5 α on retroviral infection. The variable regions of the SPRY domain of TRIM5 α evolved under the pressure of viral infection, and surface exposed loops of viral CA played a critical role in its escape from restriction of TRIM5 α . Detailed and accurate structural analysis of the binding surface between viral CA and TRIM5 α SPRY is thus needed for the development of new antiretroviral drugs that enhance anti-HIV-1 activity of human TRIM5 α .

ACKNOWLEDGEMENTS

Authors sincerely thank to our collaborators C. Onyango, M. Cotton, S. Rowland-Jones (Medical Research Council Laboratories, Gambia), K. Bozek,

F.S. Domingues (Max Plank Institute for Informatics, Germany), M. Yokoyama, H. Sato (National Institute of Infectious Diseases, Japan), A. Saito, H. Akari (National Institute of Biomedical Innovation, Japan), A. Adachi (Tokushima University, Japan), T. Nakajima, A. Kimura (Tokyo Medical and Dental University, Japan), I. Theodorou, P. Debre (Hôpital Pitié Salpêtrière, France), K. Yoshimura, S. Matsushita (Kumamoto University, Japan), A. Iwamoto (The University of Tokyo, Japan), Jay A. Levy (University of California, USA) and all members of our laboratory, especially, A. Kuroishi for providing the data for Figure 6B and H. Maegawa for his illustration in Figure 7.

REFERENCES

- Gao F, Bailes E, Robertson DL, *et al.* Origin of HIV-1 in the chimpanzee *Pan troglodytes troglodytes*. *Nature* 1999; **397**: 436–441.
- Shibata R, Sakai H, Kawamura M, Tokunaga K, Adachi A. Early replication block of human immunodeficiency virus type 1 in monkey cells. *J Gen Virol* 1995; **76** (Pt 11): 2723–2730.
- Himathongkham S, Luciw PA. Restriction of HIV-1 (subtype B) replication at the entry step in rhesus macaque cells. *Virology* 1996; **219**: 485–488.
- Akari H, Mori K, Terao K, *et al.* In vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its susceptibility to infection with simian immunodeficiency viruses. *Virology* 1996; **218**: 382–388.
- Akari H, Nam KH, Mori K, *et al.* Effects of SIVmac infection on peripheral blood CD4+ CD8+ T lymphocytes in cynomolgus macaques. *Clin Immunol* 1999; **91**: 321–329.
- VandeWoude S, Apetrei C. Going wild: lessons from naturally occurring T-lymphotropic lentiviruses. *Clin Microbiol Rev* 2006; **19**: 728–762.
- Chackerian B, Long EM, Luciw PA, Overbaugh J. Human immunodeficiency virus type 1 coreceptors participate in postentry stages in the virus replication cycle and function in simian immunodeficiency virus infection. *J Virol* 1997; **71**: 3932–3939.
- Owens CM, Yang PC, Gottlinger H, Sodroski J. Human and simian immunodeficiency virus capsid proteins are major viral determinants of early, post-entry replication blocks in simian cells. *J Virol* 2003; **77**: 726–731.
- Owens CM, Song B, Perron MJ, Yang PC, Stremlau M, Sodroski J. Binding and susceptibility to postentry restriction factors in monkey cells are specified by distinct regions of the human immunodeficiency virus type 1 capsid. *J Virol*. 2004; **78**: 5423–5437.
- Hatzioannou T, Cowan S, Von Schwedler UK, Sundquist WI, Bieniasz PD. Species-specific tropism determinants in the human immunodeficiency virus type 1 capsid. *J Virol* 2004; **78**: 6005–6012.
- Shibata R, Kawamura M, Sakai H, Hayami M, Ishimoto A, Adachi A. Generation of a chimeric human and simian immunodeficiency virus infectious to monkey peripheral blood mononuclear cells. *J Virol* 1991; **65**: 3514–3520.
- Hatzioannou T, Cowan S, Goff SP, Bieniasz PD, Towers GJ. Restriction of multiple divergent retroviruses by Lv1 and Ref1. *Embo J* 2003; **22**: 385–394.
- Besnier C, Takeuchi Y, Towers G. Restriction of lentivirus in monkeys. *Proc Natl Acad Sci USA* 2002; **99**: 11920–11925.
- Munk C, Brandt SM, Lucero G, Landau NR. A dominant block to HIV-1 replication at reverse transcription in simian cells. *Proc Natl Acad Sci USA* 2002; **99**: 13843–13848.
- Kootstra NA, Munk C, Tonnu N, Landau NR, Verma IM. Abrogation of postentry restriction of HIV-1-based lentiviral vector transduction in simian cells. *Proc Natl Acad Sci USA* 2003; **100**: 1298–1303.
- Stremlau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J. The cytoplasmic body component TRIM5 α restricts HIV-1 infection in Old World monkeys. *Nature* 2004; **427**: 848–853.
- Nakayama EE, Miyoshi H, Nagai Y, Shioda T. A specific region of 37 amino acid residues in the SPRY (B30.2) domain of African green monkey TRIM5 α determines species-specific restriction of simian immunodeficiency virus SIVmac infection. *J Virol* 2005; **79**: 8870–8877.
- Hatzioannou T, Perez-Caballero D, Yang A, Cowan S, Bieniasz PD. Retrovirus resistance factors Ref1 and Lv1 are species-specific variants of TRIM5 α . *Proc Natl Acad Sci USA* 2004; **101**: 10774–10779.
- Keckesova Z, Ylinen LM, Towers GJ. The human and African green monkey TRIM5 α genes encode Ref1 and Lv1 retroviral restriction factor activities. *Proc Natl Acad Sci USA* 2004; **101**: 10780–10785.
- Yap MW, Nisole S, Lynch C, Stoye JP. Trim5 α protein restricts both HIV-1 and murine leukemia virus. *Proc Natl Acad Sci USA* 2004; **101**: 10786–10791.
- Perron MJ, Stremlau M, Song B, Ulm W, Mulligan RC, Sodroski J. TRIM5 α mediates the postentry block to N-tropic murine leukemia viruses in human cells. *Proc Natl Acad Sci USA* 2004; **101**: 11827–11832.


22. Brennan G, Kozyrev Y, Kodama T, Hu SL. Novel TRIM5 isoforms expressed by *Macaca nemestrina*. *J Virol* 2007; **81**: 12210–12217.
23. Asaoka K, Ikeda K, Hishinuma T, Horie-Inoue K, Takeda S, Inoue S. A retrovirus restriction factor TRIM5alpha is transcriptionally regulated by interferons. *Biochem Biophys Res Commun* 2005; **338**: 1950–1956.
24. Carthagena L, Parise MC, Ringiard M, Chelbi-Alix MK, Hazan U, Nisole S. Implication of TRIM alpha and TRIMCyp in interferon-induced anti-retroviral restriction activities. *Retrovirology* 2008; **5**: 59.
25. Sakuma R, Mael AA, Ikeda Y. Alpha interferon enhances TRIM5alpha-mediated antiviral activities in human and rhesus monkey cells. *J Virol* 2007; **81**: 10201–10206.
26. Nisole S, Stoye JP, Saib A. TRIM family proteins: retroviral restriction and antiviral defence. *Nat Rev Microbiol* 2005; **3**: 799–808.
27. Reymond A, Meroni G, Fantozzi A, *et al.* The tripartite motif family identifies cell compartments. *Embo J* 2001; **20**: 2140–2151.
28. Meroni G, Diez-Roux G. TRIM/RBCC, a novel class of 'single protein RING finger' E3 ubiquitin ligases. *Bioessays* 2005; **27**: 1147–1157.
29. James LC, Keeble AH, Khan Z, Rhodes DA, Trowsdale J. Structural basis for PRYSPRY-mediated tripartite motif (TRIM) protein function. *Proc Natl Acad Sci USA* 2007; **104**: 6200–6205.
30. Sawyer SL, Emerman M, Malik HS. Discordant evolution of the adjacent antiretroviral genes TRIM22 and TRIM5 in mammals. *PLoS Pathog* 2007; **3**: e197.
31. Rhodes DA, Trowsdale J. TRIM21 is a trimeric protein that binds IgG Fc via the B30.2 domain. *Mol Immunol* 2007; **44**: 2406–2414.
32. Song B, Diaz-Griffero F, Park do H, Rogers T, Stremlau M, Sodroski J. TRIM5alpha association with cytoplasmic bodies is not required for antiretroviral activity. *Virology* 2005; **343**: 201–211.
33. Jackson PK, Eldridge AG, Freed E, *et al.* The lore of the RINGs: substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol* 2000; **10**: 429–439.
34. Javanbakht H, Diaz-Griffero F, Stremlau M, Si Z, Sodroski J. The contribution of RING and B-box 2 domains to retroviral restriction mediated by monkey TRIM5alpha. *J Biol Chem* 2005; **280**: 26933–26940.
35. Mische CC, Javanbakht H, Song B, *et al.* Retroviral restriction factor TRIM5alpha is a trimer. *J Virol* 2005; **79**: 14446–14450.
36. Javanbakht H, Yuan W, Yeung DF, *et al.* Characterization of TRIM5alpha trimerization and its contribution to human immunodeficiency virus capsid binding. *Virology* 2006; **353**: 234–246.
37. Kar AK, Diaz-Griffero F, Li Y, Li X, Sodroski J. Biochemical and biophysical characterization of a chimeric TRIM21-TRIM5alpha protein. *J Virol* 2008; **82**: 11669–11681.
38. Langelier CR, Sandrin V, Eckert DM, *et al.* Biochemical characterization of a recombinant TRIM5alpha protein that restricts human immunodeficiency virus type 1 replication. *J Virol* 2008; **82**: 11682–11694.
39. Li X, Sodroski J. The TRIM5alpha B-box 2 domain promotes cooperative binding to the retroviral capsid by mediating higher-order self-association. *J Virol* 2008; **82**: 11495–11502.
40. Diaz-Griffero F, Qin XR, Hayashi F, *et al.* A B-box 2 surface patch important for TRIM5(alpha) self-association, capsid-binding avidity and retrovirus restriction. *J Virol* 2009; **83**: 10737–10751.
41. Nakayama EE, Maegawa H, Shioda T. A dominant-negative effect of cynomolgus monkey tripartite motif protein TRIM5alpha on anti-simian immunodeficiency virus SIVmac activity of an African green monkey orthologue. *Virology* 2006; **350**: 158–163.
42. Maegawa H, Nakayama EE, Kuroishi A, Shioda T. Silencing of tripartite motif protein (TRIM) 5alpha mediated anti-HIV-1 activity by truncated mutant of TRIM5alpha. *J Virol Methods* 2008; **151**: 249–256.
43. Perez-Caballero D, Hatzioannou T, Yang A, Cowan S, Bieniasz PD. Human tripartite motif 5alpha domains responsible for retrovirus restriction activity and specificity. *J Virol* 2005; **79**: 8969–8978.
44. Sawyer SL, Wu LL, Emerman M, Malik HS. Positive selection of primate TRIM5alpha identifies a critical species-specific retroviral restriction domain. *Proc Natl Acad Sci USA* 2005; **102**: 2832–2837.
45. Kono K, Song H, Shingai Y, Shioda T, Nakayama EE. Comparison of anti-viral activity of rhesus monkey and cynomolgus monkey TRIM5alphas against human immunodeficiency virus type 2 infection. *Virology* 2008; **373**: 447–456.
46. Ohkura S, Yap MW, Sheldon T, Stoye JP. All three variable regions of the TRIM5alpha B30.2 domain can contribute to the specificity of retrovirus restriction. *J Virol* 2006; **80**: 8554–8565.
47. Kono K, Bozek K, Domingues FS, Shioda T, Nakayama EE. Impact of a single amino acid in the variable region 2 of the Old World monkey TRIM5alpha SPRY (B30.2) domain on anti-human immunodeficiency virus type 2 activity. *Virology* 2009; **388**: 160–168.
48. Perron MJ, Stremlau M, Sodroski J. Two surface-exposed elements of the B30.2/SPRY domain as potency determinants of N-tropic murine leukemia virus restriction by human TRIM5alpha. *J Virol* 2006; **80**: 5631–5636.

49. Yap MW, Nisole S, Stoye JP. A single amino acid change in the SPRY domain of human Trim5alpha leads to HIV-1 restriction. *Curr Biol* 2005; **15**: 73–78.
50. Stremlau M, Perron M, Welikala S, Sodroski J. Species-specific variation in the B30.2(SPRY) domain of TRIM5alpha determines the potency of human immunodeficiency virus restriction. *J Virol* 2005; **79**: 3139–3145.
51. Sebastian S, Luban J. TRIM5alpha selectively binds a restriction-sensitive retroviral capsid. *Retrovirology* 2005; **2**: 40.
52. Stremlau M, Perron M, Lee M, *et al.* Specific recognition and accelerated uncoating of retroviral capsids by the TRIM5alpha restriction factor. *Proc Natl Acad Sci USA* 2006; **103**: 5514–5519.
53. Hahn BH, Shaw GM, De Cock KM, Sharp PM. AIDS as a zoonosis: scientific and public health implications. *Science* 2000; **287**: 607–614.
54. Castro BA, Nepomuceno M, Lerche NW, Eichberg JW, Levy JA. Persistent infection of baboons and rhesus monkeys with different strains of HIV-2. *Virology* 1991; **184**: 219–226.
55. Fujita M, Yoshida A, Sakurai A, *et al.* Susceptibility of HVS-immortalized lymphocytic HSC-F cells to various strains and mutants of HIV/SIV. *Int J Mol Med* 2003; **11**: 641–644.
56. Castro BA, Barnett SW, Evans LA, Moreau J, Odehouri K, Levy JA. Biologic heterogeneity of human immunodeficiency virus type 2 (HIV-2) strains. *Virology* 1990; **178**: 527–534.
57. Locher CP, Witt SA, Herndier BG, *et al.* Increased virus replication and virulence after serial passage of human immunodeficiency virus type 2 in baboons. *J Virol* 2003; **77**: 77–83.
58. Locher CP, Blackburn DJ, Herndier BG, *et al.* Transient virus infection and pathogenesis of a new HIV type 2 isolate, UC12, in baboons. *AIDS Res Hum Retroviruses* 1998; **14**: 79–82.
59. Song H, Nakayama EE, Yokoyama M, Sato H, Levy JA, Shioda T. A single amino acid of the human immunodeficiency virus type 2 capsid affects its replication in the presence of cynomolgus monkey and human TRIM5alphas. *J Virol* 2007; **81**: 7280–7285.
60. Kozak CA, Chakraborti A. Single amino acid changes in the murine leukemia virus capsid protein gene define the target of Fv1 resistance. *Virology* 1996; **225**: 300–305.
61. Best S, Le Tissier P, Towers G, Stoye JP. Positional cloning of the mouse retrovirus restriction gene Fv1. *Nature* 1996; **382**: 826–829.
62. Mortuza GB, Dodding MP, Goldstone DC, Haire LF, Stoye JP, Taylor IA. Structure of B-MLV capsid amino-terminal domain reveals key features of viral tropism, gag assembly and core formation. *J Mol Biol* 2008; **376**: 1493–1508.
63. Mortuza GB, Haire LF, Stevens A, Smerdon SJ, Stoye JP, Taylor IA. High-resolution structure of a retroviral capsid hexameric amino-terminal domain. *Nature* 2004; **431**: 481–485.
64. Rowland-Jones SL, Whittle HC. Out of Africa: what can we learn from HIV-2 about protective immunity to HIV-1? *Nat Immunol* 2007; **8**: 329–331.
65. Poulsen AG, Kvinesdal B, Aaby P, *et al.* Prevalence of and mortality from human immunodeficiency virus type 2 in Bissau, West Africa. *Lancet* 1989; **1**: 827–831.
66. Berry N, Ariyoshi K, Balfe P, Tedder R, Whittle H. Sequence specificity of the human immunodeficiency virus type 2 (hiv-2) long terminal repeat u3 region in vivo allows subtyping of the principal hiv-2 viral subtypes a and b. *AIDS Res Hum Retroviruses* 2001; **17**: 263–267.
67. Ariyoshi K, Jaffar S, Alabi AS, *et al.* Plasma RNA viral load predicts the rate of CD4 T cell decline and death in HIV-2-infected patients in West Africa. *Aids* 2000; **14**: 339–344.
68. Onyango C, Leligdowicz A, Yokoyama M, *et al.* HIV-2 capsids distinguish high and low virus load patients in a West African community cohort. *Vaccine* 2009 (in press).
69. Ylinen LM, Keckesova Z, Wilson SJ, Ranasinghe S, Towers GJ. Differential restriction of human immunodeficiency virus type 2 and simian immunodeficiency virus SIVmac by TRIM5alpha alleles. *J Virol* 2005; **79**: 11580–11587.
70. Lin TY, Emerman M. Determinants of cyclophilin A-dependent TRIM5 alpha restriction against HIV-1. *Virology* 2008; **379**: 335–341.
71. Luban J, Bossolt KL, Franke EK, Kalpana GV, Goff SP. Human immunodeficiency virus type 1 Gag protein binds to cyclophilins A and B. *Cell* 1993; **73**: 1067–1078.
72. Franke EK, Yuan HE, Luban J. Specific incorporation of cyclophilin A into HIV-1 virions. *Nature* 1994; **372**: 359–362.
73. Gamble TR, Vajdos FF, Yoo S, *et al.* Crystal structure of human cyclophilin A bound to the amino-terminal domain of HIV-1 capsid. *Cell* 1996; **87**: 1285–1294.
74. Braaten D, Aberham C, Franke EK, Yin L, Phares W, Luban J. Cyclosporine A-resistant human immunodeficiency virus type 1 mutants demonstrate that Gag encodes the functional target of cyclophilin A. *J Virol* 1996; **70**: 5170–5176.
75. Thali M, Bukovsky A, Kondo E, *et al.* Functional association of cyclophilin A with HIV-1 virions. *Nature* 1994; **372**: 363–365.

76. Franke EK, Luban J. Inhibition of HIV-1 replication by cyclosporine A or related compounds correlates with the ability to disrupt the Gag-cyclophilin A interaction. *Virology*. 1996; **222**: 279–282.
77. Rosenwirth B, Billich A, Datema R, *et al.* Inhibition of human immunodeficiency virus type 1 replication by SDZ NIM 811, a nonimmunosuppressive cyclosporine analog. *Antimicrob Agents Chemother* 1994; **38**: 1763–1772.
78. Braaten D, Luban J. Cyclophilin A regulates HIV-1 infectivity, as demonstrated by gene targeting in human T cells. *Embo J* 2001; **20**: 1300–1309.
79. Sokolskaja E, Berthoux L, Luban J. Cyclophilin A and TRIM5 α independently regulate human immunodeficiency virus type 1 infectivity in human cells. *J Virol* 2006; **80**: 2855–2862.
80. Sokolskaja E, Sayah DM, Luban J. Target cell cyclophilin A modulates human immunodeficiency virus type 1 infectivity. *J Virol* 2004; **78**: 12800–12808.
81. Luban J. Cyclophilin A, TRIM5, and resistance to human immunodeficiency virus type 1 infection. *J Virol* 2007; **81**: 1054–1061.
82. Towers GJ. The control of viral infection by tripartite motif proteins and cyclophilin A. *Retrovirology* 2007; **4**: 40.
83. Towers GJ, Hatzioannou T, Cowan S, Goff SP, Luban J, Bieniasz PD. Cyclophilin A modulates the sensitivity of HIV-1 to host restriction factors. *Nat Med* 2003; **9**: 1138–1143.
84. Berthoux L, Sebastian S, Sokolskaja E, Luban J. Lv1 inhibition of human immunodeficiency virus type 1 is counteracted by factors that stimulate synthesis or nuclear translocation of viral cDNA. *J Virol* 2004; **78**: 11739–11750.
85. Berthoux L, Sebastian S, Sokolskaja E, Luban J. Cyclophilin A is required for TRIM5 α -mediated resistance to HIV-1 in Old World monkey cells. *Proc Natl Acad Sci USA* 2005; **102**: 14849–14853.
86. Keckesova Z, Ylinen LM, Towers GJ. Cyclophilin A renders human immunodeficiency virus type 1 sensitive to old world monkey but not human TRIM5 α antiviral activity. *J Virol* 2006; **80**: 4683–4690.
87. Mascola JR, Stiegler G, VanCott TC, *et al.* Protection of macaques against vaginal transmission of a pathogenic HIV-1/SIV chimeric virus by passive infusion of neutralizing antibodies. *Nat Med* 2000; **6**: 207–210.
88. Nishimura Y, Igarashi T, Haigwood N, *et al.* Determination of a statistically valid neutralization titer in plasma that confers protection against simian-human immunodeficiency virus challenge following passive transfer of high-titered neutralizing antibodies. *J Virol* 2002; **76**: 2123–2130.
89. Shibata R, Igarashi T, Haigwood N, *et al.* Neutralizing antibody directed against the HIV-1 envelope glycoprotein can completely block HIV-1/SIV chimeric virus infections of macaque monkeys. *Nat Med* 1999; **5**: 204–210.
90. Matano T, Shibata R, Siemon C, Connors M, Lane HC, Martin MA. Administration of an anti-CD8 monoclonal antibody interferes with the clearance of chimeric simian/human immunodeficiency virus during primary infections of rhesus macaques. *J Virol* 1998; **72**: 164–169.
91. Migueles SA, Sabbaghian MS, Shupert WL, *et al.* HLA B*5701 is highly associated with restriction of virus replication in a subgroup of HIV-infected long term nonprogressors. *Proc Natl Acad Sci USA* 2000; **97**: 2709–2714.
92. Nixon DF, Townsend AR, Elvin JG, Rizza CR, Gallwey J, McMichael AJ. HIV-1 gag-specific cytotoxic T lymphocytes defined with recombinant vaccinia virus and synthetic peptides. *Nature* 1988; **336**: 484–487.
93. Kamada K, Igarashi T, Martin MA, *et al.* Generation of HIV-1 derivatives that productively infect macaque monkey lymphoid cells. *Proc Natl Acad Sci USA* 2006; **103**: 16959–16964.
94. Harris RS, Bishop KN, Sheehy AM, *et al.* DNA deamination mediates innate immunity to retroviral infection. *Cell* 2003; **113**: 803–809.
95. Mangeat B, Turelli P, Caron G, Friedli M, Perrin L, Trono D. Broad antiretroviral defence by human APOBEC3G through lethal editing of nascent reverse transcripts. *Nature* 2003; **424**: 99–103.
96. Sheehy AM, Gaddis NC, Choi JD, Malim MH. Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* 2002; **418**: 646–650.
97. Mariani R, Chen D, Schrofelbauer B, *et al.* Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif. *Cell* 2003; **114**: 21–31.
98. Marin M, Rose KM, Kozak SL, Kabat D. HIV-1 Vif protein binds the editing enzyme APOBEC3G and induces its degradation. *Nat Med* 2003; **9**: 1398–1403.
99. Sheehy AM, Gaddis NC, Malim MH. The antiretroviral enzyme APOBEC3G is degraded by the proteasome in response to HIV-1 Vif. *Nat Med* 2003; **9**: 1404–1407.
100. Nakayama EE, Shingai Y, Kono K, Shioda T. TRIM5 α -independent anti-human immunodeficiency virus type 1 activity mediated by cyclophilin A in Old World monkey cells. *Virology* 2008; **375**: 514–520.

101. Igarashi T, Iyengar R, Byrum RA, *et al.* Human immunodeficiency virus type 1 derivative with 7% simian immunodeficiency virus genetic content is able to establish infections in pig-tailed macaques. *J Virol* 2007; **81**: 11549–11552.
102. Hatzioannou T, Princiotta M, Piatak M, *et al.* Generation of simian-tropic HIV-1 by restriction factor evasion. *Science* 2006; **314**: 95.
103. Kuroishi A, Saito A, Shingai Y, *et al.* Modification of a loop sequence between alpha-helices 6 and 7 of virus capsid (CA) protein in a human immunodeficiency virus type 1 (HIV-1) derivative that has simian immunodeficiency virus (SIVmac239) vif and CA alpha-helices 4 and 5 loop improves replication in cynomolgus monkey cells. *Retrovirology* 2009; **6**: 70.
104. Diaz-Griffero F, Li X, Javanbakht H, *et al.* Rapid turnover and polyubiquitylation of the retroviral restriction factor TRIM5. *Virology* 2006; **349**: 300–315.
105. Rold CJ, Aiken C. Proteasomal degradation of TRIM5alpha during retrovirus restriction. *PLoS Pathog* 2008; **4**: e1000074.
106. Perez-Caballero D, Hatzioannou T, Zhang F, Cowan S, Bieniasz PD. Restriction of human immunodeficiency virus type 1 by TRIM-CypA occurs with rapid kinetics and independently of cytoplasmic bodies, ubiquitin, and proteasome activity. *J Virol* 2005; **79**: 15567–15572.
107. Anderson JL, Campbell EM, Wu X, *et al.* Proteasome inhibition reveals that a functional preintegration complex intermediate can be generated during restriction by diverse TRIM5 proteins. *J Virol* 2006; **80**: 9754–9760.
108. Wu X, Anderson JL, Campbell EM, Joseph AM, Hope TJ. Proteasome inhibitors uncouple rhesus TRIM5alpha restriction of HIV-1 reverse transcription and infection. *Proc Natl Acad Sci USA* 2006; **103**: 7465–7470.
109. O'Brien SJ, Nelson GW. Human genes that limit AIDS. *Nat Genet* 2004; **36**: 565–574.
110. Shioda T, Nakayama EE. Human genetic polymorphisms affecting HIV-1 diseases. *Int J Hematol* 2006; **84**: 12–17.
111. Sawyer SL, Wu LI, Akey JM, Emerman M, Malik HS. High-frequency persistence of an impaired allele of the retroviral defense gene TRIM5alpha in humans. *Curr Biol* 2006; **16**: 95–100.
112. Javanbakht H, An P, Gold B, *et al.* Effects of human TRIM5alpha polymorphisms on antiretroviral function and susceptibility to human immunodeficiency virus infection. *Virology* 2006; **354**: 15–27.
113. Nakayama EE, Carpentier W, Costagliola D, *et al.* Wild type and H43Y variant of human TRIM5alpha show similar anti-human immunodeficiency virus type 1 activity both in vivo and in vitro. *Immunogenetics* 2007; **59**: 511–515.
114. Speelmon EC, Livingston-Rosanoff D, Li SS, *et al.* Genetic association of the antiviral restriction factor TRIM5alpha with human immunodeficiency virus type 1 infection. *J Virol* 2006; **80**: 2463–2471.
115. van Manen D, Rits MA, Beugeling C, van Dort K, Schuitemaker H, Kootstra NA. The effect of Trim5 polymorphisms on the clinical course of HIV-1 infection. *PLoS Pathog* 2008; **4**: e18.
116. Nakajima T, Nakayama EE, Kaur G, *et al.* Impact of novel TRIM5alpha variants, Gly110Arg and G176del, on the anti-HIV-1 activity and the susceptibility to HIV-1 infection. *Aids* 2009; **23**: 2091–2100.
117. Tareen SU, Sawyer SL, Malik HS, Emerman M. An expanded clade of rodent Trim5 genes. *Virology* 2009; **385**: 473–483.
118. Johnson WE, Sawyer SL. Molecular evolution of the antiretroviral TRIM5 gene. *Immunogenetics* 2009; **61**: 163–176.
119. Shi M, Deng W, Bi E, *et al.* TRIM30 alpha negatively regulates TLR-mediated NF-kappa B activation by targeting TAB2 and TAB3 for degradation. *Nat Immunol* 2008; **9**: 369–377.
120. Song B, Gold B, O'Huigin C, *et al.* The B30.2(SPRY) domain of the retroviral restriction factor TRIM5alpha exhibits lineage-specific length and sequence variation in primates. *J Virol* 2005; **79**: 6111–6121.
121. Newman RM, Hall L, Connole M, *et al.* Balancing selection and the evolution of functional polymorphism in Old World monkey TRIM5alpha. *Proc Natl Acad Sci USA* 2006; **103**: 19134–19139.
122. Kaiser SM, Malik HS, Emerman M. Restriction of an extinct retrovirus by the human TRIM5alpha antiviral protein. *Science* 2007; **316**: 1756–1758.
123. Wilson SJ, Webb BL, Maplanka C, *et al.* Rhesus macaque TRIM5 alleles have divergent antiretroviral specificities. *J Virol* 2008; **82**: 7243–7247.
124. Sayah DM, Sokolskaja E, Berthoux L, Luban J. Cyclophilin A retrotransposition into TRIM5 explains owl monkey resistance to HIV-1. *Nature* 2004; **430**: 569–573.
125. Nisole S, Lynch C, Stoye JP, Yap MW. A Trim5-cyclophilin A fusion protein found in owl monkey kidney cells can restrict HIV-1. *Proc Natl Acad Sci USA* 2004; **101**: 13324–13328.
126. Kuang YQ, Tang X, Liu FL, *et al.* Genotyping of TRIM5 locus in northern pig-tailed macaques (*Macaca leonina*), a primate species susceptible to Human Immunodeficiency Virus type 1 infection. *Retrovirology* 2009; **6**: 58.
127. Virgen CA, Kratovac Z, Bieniasz PD, Hatzioannou T. Independent genesis of chimeric TRIM5-cyclophilin proteins in two primate species. *Proc Natl Acad Sci USA* 2008; **105**: 3563–3568.

128. Newman RM, Hall L, Kirmaier A, *et al.* Evolution of a TRIM5-CypA splice isoform in old world monkeys. *PLoS Pathog* 2008; **4**: e1000003.
129. Wilson SJ, Webb BL, Ylinen LM, Verschoor E, Heeney JL, Towers GJ. Independent evolution of an antiviral TRIMCyp in rhesus macaques. *Proc Natl Acad Sci USA* 2008; **105**: 3557–3562.
130. Hatziioannou T, Ambrose Z, Chung NP, *et al.* A macaque model of HIV-1 infection. *Proc Natl Acad Sci USA* 2009; **106**: 4425–4429.
131. Keeble AH, Khan Z, Forster A, James LC. TRIM21 is an IgG receptor that is structurally, thermodynamically, and kinetically conserved. *Proc Natl Acad Sci USA* 2008; **105**: 6045–6050.



Contribution of RING domain to retrovirus restriction by TRIM5 α depends on combination of host and virus

Hikoichiro Maegawa, Tadashi Miyamoto, Jun-ichi Sakuragi, Tatsuo Shioda, Emi E. Nakayama*

Department of Viral Infections, Research Institute for Microbial Diseases, Osaka University, 3-1, Yamadaoka, Suita-shi, Osaka 565-0871, Japan

ARTICLE INFO

Article history:

Received 19 October 2009
Returned to author for revision
8 November 2009
Accepted 5 January 2010
Available online xxx

ABSTRACT

The anti-retroviral restriction factor TRIM5 α contains the RING domain, which is frequently observed in E3 ubiquitin ligases. It was previously proposed that TRIM5 α restricts human immunodeficiency virus type 1 (HIV-1) via proteasome-dependent and -independent pathways. Here we examined the effects of RING domain mutations on retrovirus restriction by TRIM5 α in various combinations of virus and host species. Simian immunodeficiency virus isolated from macaque (SIVmac) successfully avoided attacks by RING mutants of African green monkey (AGM)-TRIM5 α that could still restrict HIV-1. Addition of proteasome inhibitor did not affect the anti-HIV-1 activity of AGM-TRIM5 α , whereas it disrupted at least partly its anti-SIVmac activity. In the case of mutant human TRIM5 α carrying proline at the position 332, however, both HIV-1 and SIVmac restrictions were eliminated as a result of RING domain mutations. These results suggested that the mechanisms of retrovirus restriction by TRIM5 α vary depending on the combination of host and virus.

© 2009 Published by Elsevier Inc.

Introduction

Replication of retroviruses is influenced by several factors in host cells. Tripartite motif protein (TRIM) 5 α has been identified as a restriction factor of human immunodeficiency virus type 1 (HIV-1) in rhesus monkey (Rh) cells (Stremlau et al., 2004). Rh TRIM5 α potently restricts HIV-1 but only weakly does so simian immunodeficiency virus isolated from macaque (SIVmac) (Stremlau et al., 2004; Song et al., 2005), whereas African green monkey (AGM) TRIM5 α can potently restrict both HIV-1 and SIVmac (Nakayama et al., 2005; Song et al., 2005). TRIM5 α consists of the RING, B-box 2, coiled-coil, and SPRY (B30.2) domains (Reymond et al., 2001). Differences in the amino acid sequences in the SPRY domain of TRIM5 α of different monkey species were shown to affect the species-specific restriction of retrovirus infection (Pérez-Caballero et al., 2005a; Nakayama et al., 2005; Sawyer et al., 2005; Stremlau et al., 2005; Yap et al., 2005). In addition, biochemical studies have shown that TRIM5 α associates with retroviral capsid (CA) protein in detergent-stripped virions or with an artificially constituted core structure composed of capsid-nucleocapsid (CA-NC) fusion protein in a SPRY domain-dependent manner (Sebastian and Luban, 2005; Stremlau et al., 2006a). The SPRY domain is thus thought to recognize viral core. The coiled-coil domain

of TRIM5 α is important for the formation of homo-oligomers (Mische et al., 2005) and is essential for antiviral activity (Javanbakht et al., 2006). The intact B-box 2 domain is also required for TRIM5 α mediated antiviral activity, since the restrictive activity of TRIM5 α is diminished by amino acid substitutions in the B-box 2 domain (Javanbakht et al., 2005). RING containing proteins were frequently found to possess E3 ubiquitin ligase activity (Jackson et al., 2000). Indeed, Rh TRIM5 α was poly-ubiquitinated and degraded rapidly via the ubiquitin-proteasome pathway, while disruption of the RING domain eliminated its auto-ubiquitination (Diaz-Griffero et al., 2006). Furthermore, it was demonstrated that TRIM5 α is degraded via the ubiquitin-proteasome pathway during HIV-1 restriction (Rold and Aiken, 2008). However, deletion of the RING domain in TRIM5 α only partially attenuates anti-HIV-1 activity (Javanbakht et al., 2005; Perez-Caballero et al., 2005b). Moreover, modulation of E1 ubiquitin-activating enzyme expression did not affect TRIM5 α -mediated restriction activity in a temperature-dependent cell line (Perez-Caballero et al., 2005b) and finally, proteasome inhibitors did not affect TRIM5 α mediated HIV-1 restriction (Anderson et al., 2006; Perez-Caballero et al., 2005b; Rold and Aiken, 2008; Stremlau et al., 2006a; Wu et al., 2006) even though they allowed HIV-1 to generate viral late reverse transcripts under TRIM5 α mediated HIV-1 restriction (Anderson et al., 2006; Wu et al., 2006). The exact role of the TRIM5 α RING domain in retrovirus restriction thus remains unclear.

In the study presented here, we investigated the effects of RING domain mutations on HIV-1 and SIVmac restrictions by TRIM5 α and report that TRIM5 α restricts HIV-1 and SIVmac differently.

* Corresponding author. Fax: +81 6 6879 8347.
E-mail address: emien@biken.osaka-u.ac.jp (E.E. Nakayama).

80 **Results**

81 *Auto poly-ubiquitination of TRIM5 α impaired by mutations in*
82 *RING domain*

83 The RING finger domain of TRIM5 α comprises eight potential
84 metal ligands and binds two atoms of zinc, with each zinc atom ligated
85 tetrahedrally by either four cysteines or three cysteines and a single
86 histidine. Based on the three-dimensional structure of the RING
87 domains of TRIM5 (Abe et al., 2007) and the promyelocytic leukemia

(PML) protein (Borden et al., 1995; Borden and Freemont, 1996), the 88
first pair of metal ligands of the AGM TRIM5 α RING domain (C15 and 89
C18) would share a zinc atom with the third pair (C35 and C38), and 90
the second (C30 and H32) and fourth pairs (C56 and C59) would share 91
another zinc atom (Fig. 1A). To determine whether anti-HIV-1 and 92
anti-SIVmac activities of AGM TRIM5 α are similarly affected by RING 93
domain mutations, several AGM TRIM5 α constructs with mutations in 94
the RING domain were generated (Fig. 1B). In the mutant TRIM5 α 95
constructs with C15AC18A, C30AH32A, or C15AC18AC30AH32A, two 96
key amino acid residues in the first or second, or in both the first and 97




Fig. 1. Auto poly-ubiquitination of TRIM5 α was impaired by RING domain mutations. (A) The RING finger zinc binding motif. The numbered AGM TRIM5 α zinc-binding ligands are shown in circles. Each zinc atom is coordinated tetrahedrally by four ligands. Zinc-binding site 1 (**bold**) and site 2 (*italic*) are indicated. The numbers of amino acid residues between the zinc-binding cysteine and histidine ligands in AGM TRIM5 α are also indicated. (B) Primary amino acid sequences of the RING domains of AGM TRIM5 α (AGM T5aRING) and human TRIM5 α (Hu T5aRING) are aligned. Zinc-binding site 1 (**bold**), site 2 (*italic*), and cysteine and histidine ligands (large numbers) are indicated. (C) Schematic representation of TRIM5 α constructs. Black and white bars denote AGM and Hu sequences, respectively. Abbreviations for domains: R, RING; B, B-box 2; CC, Coiled-coil; S, SPRY. A dotted box denotes deletion of corresponding amino acid. The positions of individual amino acid changes are also indicated. (D) 293 T cells were transfected with plasmids encoding HA-tagged AGM TRIM5 α (TRIM5 α -HA) or its RING mutants together with a plasmid expressing myc-tagged ubiquitin (myc-Ub). Forty-eight hours after transfection, the cells were lysed and TRIM5 α proteins in the lysates were precipitated with an anti-HA antibody. The immunoprecipitates were Western blotted and probed with anti-HA antibody for TRIM5 α detection or with anti-myc antibody for ubiquitin detection. The representative results of two independent experiments with similar results are shown.

Please cite this article as: Maegawa, H., et al., Contribution of RING domain to retrovirus restriction by TRIM5 α depends on combination of host and virus, Virology (2010), doi:10.1016/j.virol.2010.01.003