

Impact of novel TRIM5 α variants, Gly110Arg and G176del, on the anti-HIV-1 activity and the susceptibility to HIV-1 infection

Toshiaki Nakajima^{a,b,*}, Emi E. Nakayama^{c,*}, Gurbinder Kaur^d, Hiroshi Terunuma^e, Jun-ich Mimaya^f, Hitoshi Ohtani^{a,b}, Narinder Mehra^d, Tatsuo Shioda^c and Akinori Kimura^{a,b}

Objective: TRIM5 α is one of the factors contributing to intracellular defense mechanisms against HIV-1 infection. We investigated the association of TRIM5 α sequence variations with the susceptibility to HIV-1 infection in Japanese and Indian.

Design: Sequence variations in TRIM5 α were investigated in HIV-1-infected patients and ethnic-matched controls. Functional alterations caused by rare variants were analyzed.

Methods: We sequenced TRIM5 α -exon 2 in both Japanese (94 HIV-1-infected patients and 487 controls) and Indian (101 HIV-1-infected patients and 99 controls). Frequency of variants and haplotypes were compared between the HIV-1-infected patients and controls. Functional analyses were performed for two rare variants, Gly110Arg and G176del.

Results: The frequency of 43Tyr-allele in the Indian HIV-1-infected patients was significantly lower than that in the ethnic-matched controls (odds ratio = 0.52, 95% confidence interval = 0.31–0.89, $P = 0.015$). A similar tendency was observed in Japanese sample, although it was not statistically significant (odds ratio = 0.67, 95% confidence interval = 0.43–1.05, $P = 0.095$). On the other hand, haplotype analyses revealed that the haplotype carrying the 43Tyr-allele was significantly associated with the reduced susceptibility to HIV-1 infection in both ethnic groups. Functional analysis revealed that Gly110Arg variant weakened the anti-HIV-1 and anti-HIV-2 activities of human TRIM5 α , whereas the truncated G176del-TRIM5 enhanced the antiviral activity of coexpressed TRIM5 α . Epidemiological data were consistent in that Gly110Arg and G176del were associated with the susceptibility to and protection from HIV-1 infection, respectively.

Conclusion: Both common and rare variants of TRIM5 α are associated with the susceptibility to HIV-1 infection.

© 2009 Wolters Kluwer Health | Lippincott Williams & Wilkins

AIDS 2009, 23:2091–2100

Keywords: association, HIV-1, polymorphism, susceptibility, TRIM5

^aDepartment of Molecular Pathogenesis, Medical Research Institute, ^bLaboratory of Genome Diversity, Graduate School of Biomedical Science, Tokyo Medical and Dental University, Tokyo, ^cDepartment of Viral Infection, Research Institute for Microbial Disease, Osaka University, Osaka, Japan, ^dDepartment of Transplant Immunology and Immunogenetics, All India Institute of Medical Sciences, New Delhi, India, ^eBiotherapy Institute of Japan, Tokyo, and ^fDivision of Hematology and Oncology, Shizuoka Children's Hospital, Shizuoka, Japan.

Correspondence to Akinori Kimura, MD, PhD, Department of Molecular Pathogenesis, Medical Research Institute, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8510, Japan.

Tel: +81 3 5803 4905; fax: +81 3 5803 4907; e-mail: akitis@mri.tmd.ac.jp

* T.N. and E.E.N. contributed equally to this study.

Received: 25 March 2009; revised: 22 July 2009; accepted: 29 July 2009.

DOI:10.1097/QAD.0b013e328331567a

Introduction

Although the cellular and humoral immune systems are known to play crucial roles in the defense against retroviral infection, mammals have also evolved defense mechanisms within cells. Several lines of evidence indicated that restriction factors within host cells inhibit viral replication more effectively than the immune system, and TRIM5 α is one of the factors involved in the intracellular defense against retroviruses [1,2]. It was reported that TRIM5 α from rhesus monkeys restricted HIV-1 production at a postentry, preintegration stage in the viral life cycle through rapid degradation of HIV-1 Gag polyproteins [3,4], whereas human TRIM5 α restricted HIV-1 only weakly and potently restricted N-tropic murine leukemia virus [5,6].

TRIM5 is a member of the tripartite-motif containing superfamily and includes a Really Interesting New Gene (RING) domain, B-box 2 domain and coiled-coil domain [7]. Alternative splicing of TRIM5 gene generates several isoforms of TRIM5 proteins. One isoform, TRIM5 α , contains the carboxy-terminal B30.2 (SPRY) domain that is essential for anti-HIV-1 activity, and sequence differences in the SPRY domain contribute to the differences in the anti-HIV-1 activity among primate species [4,8–13].

It is well known that the infection by HIV-1 and progression to AIDS are variable among human individuals, which are considered to be controlled by diversity in the human genome [14,15]. As TRIM5 α has crucial roles in the restriction of viral replication within the host cells, it is a good candidate gene controlling the susceptibility to or protection from HIV-1 infection and/or progression to AIDS. Actually, several studies have demonstrated that common TRIM5 α functional polymorphisms, His43Tyr and Arg136Gln, were associated with the susceptibility to HIV-1 infection. However, the significance of association has not been established [16–20].

In this study, we investigated two ethnic populations, Japanese and Indian, for the polymorphism in TRIM5 α -exon 2 and its association with the HIV-1 infection. We found that a TRIM5 α haplotype carrying the 43Tyr-allele was associated with the reduced susceptibility to HIV-1 infection in both ethnic groups. In addition, we identified two rare variants, G176del and Gly110Arg, which affected the anti-HIV-1 activity and showed suggestive associations with the HIV-1 infection.

Material and methods

Participants

Protocol of the present study was approved by the Ethics Review Board of the Medical Research Institute, Tokyo

Medical and Dental University and that of All India Institute of Medical Science. At the set-up of the cohort in 1995, all the HIV-1-infected Japanese hemophiliac patients had been infected for longer than 10 years and they were asymptomatic without any antiviral measures. Blood samples were collected from 94 well characterized patients who were selected from the cohort after obtaining written informed consent [21,22]. Control DNA samples were prepared from Epstein–Barr virus-transformed human B cell lines established from randomly selected healthy donors with written informed consent ($n = 487$), which were purchased from the Japan Health Sciences Foundation. The DNA samples from HIV-1-infected individuals were prepared from the blood sample by using QuickGene DNA whole blood kit S (FUJIFILM, Tokyo, Japan). In addition, blood DNA samples were obtained with written informed consent from 101 HIV-1-infected Indian patients and 99 healthy Indian volunteers in the related hospitals of All India Institute of Medical Sciences, New Delhi. DNA samples from whites ($n = 96$) and African–Americans ($n = 96$) were obtained from the Coriell Institute for Medical Research (Camden, New Jersey, USA).

Identification and genotyping of nucleotide variations in TRIM5 α -exon 2

Primer sets were designed to amplify the genomic segments covering the entire TRIM5 α -exon 2 as follows: sense primer (5'-TTGGTCCCATTTTAAACC TTCC-3') and antisense primer (5'-AAGGCAGT TAA TGTCAAAGGC-3'). Genomic DNA was subjected to PCR amplification followed by sequencing on both strands using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, California, USA). Polymorphisms were identified using the Sequencher program (Gene Code Co., Ann Arbor, Michigan, USA).

Cloning and expression of TRIM5 α

The generation of recombinant Sendai viruses (SeVs) expressing human TRIM5 α , human/African green monkey (AGM) chimeric TRIM5 α [18], cynomolgus monkey TRIM5 α lacking SPRY domain [CM-TRIM5 α -SPRY(-)-HA] [23] and AGM TRIM5 α lacking coiled-coil domain [AGM-TRIM5 α -CC(-)-HA] has been described previously [24]. All TRIM5 α variants carried a hemagglutinin (HA)-tag (YPYDVP-DYAA) at the C-terminus. The Gly110Arg mutation was introduced into both human TRIM5 α and human/AGM chimeric TRIM5 α by PCR site-directed mutagenesis. To generate SeV expressing G176del carrying an HA-tag at the N-terminus (HA-G176del-TRIM5), the amplified PCR fragment from genomic DNA carrying G176del was cloned into a pSeV18+b(+) vector. Recombinant SeVs expressing human 110Arg-TRIM5 α , human/AGM chimeric 110Arg-TRIM5 α and HA-G176del-TRIM5 were recovered as described previously [12]. The second passages in embryonated

chicken eggs were used as the stock viruses in all experiments.

Western blot analysis

MT4 cells (1×10^6) infected with recombinant SeVs expressing HA-tagged TRIM5 α proteins were lysed in lysis buffer (50 mmol/l Tris-HCl, pH 7.5, 150 mmol/l NaCl, 1% Nonidet P40, 0.5% sodium deoxycholate). Western blot analyses with anti-HA high-affinity rat monoclonal antibody (Roche, Indianapolis, Indiana, USA) and anti-CypA affinity rabbit polyclonal antibody (Sigma, St Louis, Missouri, USA) were performed as described previously [19].

Viral infection

MT4 or CEM-SS cells (1×10^5) were infected with SeV expressing human TRIM5 α , human 110Arg-TRIM5 α , human/AGM chimeric TRIM5 α , human/AGM chimeric 110Arg-TRIM5 α , CM-TRIM5 α -SPRY(-) or AGM-TRIM5 α -CC(-) at a multiplicity of infection of 10 plaque-forming units (PFUs) per cell and incubated at 37°C for 9 h. To examine the effects of G176del-TRIM5 on the full length TRIM5 α , 5 PFU per cell of each SeV expressing either CM-TRIM5 α -SPRY(-), AGM-TRIM5 α -CC(-) or HA-G176del-TRIM5 was simultaneously inoculated to MT4 cells with SeV expressing human/AGM chimeric TRIM5 α . Cells were then superinfected with 30 ng of p24 of an X4 HIV-1 strain NL43 or 30 ng of p25 of HIV-2 GH123. The culture supernatants were collected periodically and the level of p24 or p25 was measured by RETROtek antigen ELISA kit (ZeptoMetrix, Buffalo, New York, USA).

Statistical analysis

All statistical analyses in this study were performed using GraphPad InStat version 3.06 for Windows (GraphPad Software, San Diego, California, USA). Correction for multiple testing was done by multiplying the *P* value by the number of tested markers to obtain the corrected *P* (*P*_c) value. Haplotype association analyses were performed by using SNPalyze version 6.0 standard (DYNACOM Co., Ltd., Tokyo, Japan). Meta-analysis was performed using a Mantel-Haenszel method. *P* values less than 0.05 were considered to be statistically significant.

Results

Associations of TRIM5 α -exon 2 polymorphism with susceptibility to HIV-1 infection

We identified 10 different nucleotide variations in the TRIM5 α -exon 2 in this study and eight of them were reported in previous studies. Most of the sequence variations, except for His43Tyr, Val112Phe and Arg136Gln were observed with low frequencies (allele

frequency less than 0.05) in the tested populations. Associations between the TRIM5 α polymorphisms and the susceptibility to HIV-1 infection were summarized in Table 1. The frequency of 43Tyr-allele in Indian HIV-1-infected patients was significantly lower than that in the ethnic-matched controls [odds ratio (OR) (95% confidence interval (CI)) = 0.52 (0.31–0.89), *P* = 0.015 by χ^2 test]. A similar tendency was observed in Japanese, although it did not reach statistical significance [OR (95% CI) = 0.67 (0.43–1.05), *P* = 0.095 by χ^2 test]. A meta-analysis of data from two populations demonstrated the significant association corrected for multiple testing [OR (95% CI) = 0.61 (0.43–0.85), *P* = 0.0004, *P*_c = 0.004]. When we analyzed the data for HIV-1 loads after 7–8 years during the observation period, which were available for 75 Japanese HIV-1-infected patients, no significant correlation between the His43Tyr genotype and HIV-1 loads was observed (data not shown).

Two novel polymorphisms, Gly110Arg and G176del, were identified only in the Japanese samples. The Gly110Arg variant was more frequent in the HIV-1-infected patients than in the controls [OR (95% CI) = 13.12 (2.53–68.21), *P* = 0.002 by Fisher's exact test]. On the other hand, the G176del variant, a deletion of a G at the coding nucleotide position 176 from the initiation site of translation, which may result in a truncated TRIM5 protein product, was found only in the Japanese controls.

TRIM5 α haplotype and susceptibility to HIV-1 infection

The associations between the susceptibility to HIV-1 infection and TRIM5 α haplotypes composed of five sequence variations with relatively high frequency, His43Tyr, Gly110Arg, Val112Phe, Thr128Thr and Arg136Gln, were investigated in Japanese and Indian populations (Table 2). In both populations, the frequency of a common haplotype 43Tyr-110Gly-112Val-128Thr-136Arg was significantly low in the HIV-1-infected patients. This result was consistent with the association of 43Tyr with the reduced susceptibility to HIV-1 infection, because this haplotype was in tight linkage disequilibrium with the 43Tyr-allele.

Anti-HIV-1 activity of TRIM5 α was attenuated by Gly110Arg substitution

To investigate the functional significance of Gly110Arg on the anti-HIV activity of TRIM5 α , we constructed a SeV containing a C-terminal HA-tagged human 110Arg-TRIM5 α (Fig. 1a). As shown in Fig. 1b, expression level of variant 110Arg-TRIM5 α was comparable to that of wild-type human TRIM5 α . A human/AGM chimeric TRIM5 α , which possessed the SPRY domain of AGM TRIM5 α , was also generated to enhance the weak anti-HIV-1 activity of human TRIM5 α (Fig. 1a). As shown in Fig. 1b, the expression level of human/AGM chimeric

Table 1. Allele frequencies of TRIM5 α -exon 2 sequence variations and associations of them with HIV-1/AIDS susceptibility.

Sequence variations ^a	Japanese			Indian			White		African-American	
	Control (n = 487)	HIV-1-infected patients (n = 94)	Odds ratio (95% confidence interval)	Control (n = 99)	HIV-1-infected patients (n = 101)	Odds ratio (95% confidence interval)	Control (n = 96)	Odds ratio (95% confidence interval)	Control (n = 96)	Odds ratio (95% confidence interval)
Gly31Ser	0.000	0.000	ND	0.000	0.000	ND	0.000	0.032	0.000	0.032
His43Tyr	0.184	0.133	0.67 (0.42 - 1.05)	0.227	0.134***	0.52 (0.31 - 0.89)	0.115	0.068	0.115	0.068
Cys58Tyr	0.000	0.000	ND	0.000	0.000	ND	0.000	0.011	0.000	0.011
G176del	0.005	0.000	ND	0.000	0.000	ND	0.000	0.000	0.000	0.000
Asp109Asp	0.000	0.000	ND	0.000	0.000	ND	0.000	0.005	0.000	0.005
Gly110Arg	0.002	0.021***	13.14 (2.53 - 68.21)	0.000	0.000	ND	0.000	0.000	0.000	0.000
Gly110Glu	0.000	0.000	ND	0.000	0.000	ND	0.000	0.005	0.000	0.005
Val112Phe	0.052	0.043	0.80 (0.37 - 1.70)	0.192	0.198	1.04 (0.63 - 1.71)	0.052	0.021	0.052	0.021
Thr128Thr	0.000	0.011**	ND	0.000	0.000	ND	0.000	0.000	0.000	0.000
Arg136Gln	0.105	0.144	1.48 (0.94 - 2.32)	0.177	0.173	0.98 (0.58 - 1.64)	0.349	0.177	0.349	0.177

ND, not defined.

^aThe numbers of sequence variations, except for G176del, are referenced by the amino acid coding position of TRIM5 α . G176del is a deletion of a G at the coding nucleotide position 176 from the initiation site of translation.

** $P < 0.05$ in Fisher's exact test, when compared with control.

*** $P < 0.01$ in Fisher's exact test, when compared with control.

**** $P < 0.05$ in χ^2 test with Yates correction, when compared with control.

Table 2. Haplotype frequencies of four common haplotypes for TRIM5 α -exon 2 and association of them with HIV-1/AIDS susceptibility.

Haplotype (His43Tyr-Gly110Arg-Val112Phe-Thr128Thr-Arg136Gln)	Japanese			Indian		
	Control (n = 487)	HIV-1-infected patients (n = 94)	Odds ratio (95% confidence interval)	Control (n = 99)	HIV-1-infected patients (n = 101)	Odds ratio (95% confidence interval)
43His-110Gly-112Val-128Thr-136Arg	0.659	0.721	1.35 (0.96 - 1.91)	0.404	0.495	1.45 (0.97 - 2.15)
43Tyr-110Gly-112Val-128Thr-136Arg	0.184	0.089	0.44 (0.26 - 0.75)	0.227	0.134	0.52 (0.31 - 0.89)
43His-110Gly-112Val-128Thr-136Gln	0.103	0.090	0.87 (0.51 - 1.49)	0.177	0.173	0.98 (0.58 - 1.64)
43His-110Gly-112Phe-128Thr-136Arg	0.051	0.031	0.61 (0.26 - 1.44)	0.192	0.198	1.04 (0.64 - 1.71)

* $P < 0.01$ in permutation test.




Fig. 1. Expression of TRIM5 α constructs used in this study. (a) Schematic representation of TRIM5 α fused with a hemagglutinin (HA)-tag. Domain structures of TRIM5 α are shown at the top. White and black bars denote human and African green monkey (AGM) sequences, respectively. Gray bar denotes the G176del-specific 16 amino acid residues generated by the frameshift. A BgIII site was used to exchange carboxy-terminal B30.2 (SPRY) domains between human and AGM TRIM5 α . 'G' or 'R' denotes the amino acid residue at the 110th position. WT denotes wild type. (b) Western blot analysis of TRIM5 protein expressed by recombinant Sendai virus (SeV). MT4 cells were infected with a SeV containing a HA-tagged variant (110Arg) human TRIM5 α (lane 1), wild-type human TRIM5 α (lane 2), human/AGM chimeric 110Arg-TRIM5 α (lane 3), human/AGM chimeric wild-type-TRIM5 α (lane 4) and G176del-TRIM5 (lane 5). Sixteen hours after the infection, cells were lysed and subjected to SDS-PAGE. HA-tagged proteins were detected by anti-HA antibody.

110Arg-TRIM5 α was similar to that of human/AGM chimeric wild-type-TRIM5 α .

These TRIM5 α constructs were tested for their ability to restrict the X4-tropic HIV-1 strain NL43 and HIV-2 strain GH123. MT4 cells infected with recombinant SeV expressing each of the TRIM5 α constructs were superinfected with HIV-1 NL43 or HIV-2 GH123. We used SeV expressing cynomolgus monkey TRIM5 α lacking the SPRY domain CM-TRIM5 α -SPRY(-) as a negative control for functional TRIM5 α , as overexpression of TRIM5 α lacking the SPRY domain exerted a dominant

negative effect on the endogenous human TRIM5 α [24]. We also used SeV expressing AGM-TRIM5 α lacking the coiled-coil domain AGM-TRIM5 α -CC(-) as a non-interfering control [24]. As shown in Fig. 2a, both wild-type (110Gly) and variant (110Arg) human/AGM chimeric TRIM5 α strongly restricted HIV-1 NL43. On the other hand, both wild-type and variant human TRIM5 α showed only weak anti-HIV-1 activity. There was, however, a small increase of HIV-1 in cells expressing the human/AGM chimeric 110Arg-TRIM5 α than the cells with the human/AGM chimeric TRIM5 α . In the case of HIV-2, virus grew to higher titers in cells




Fig. 2. Effect of TRIM5 α variants on the anti-HIV-1 and anti-HIV-2 activities. Human MT4 (a, b) or CEM-SS (c) cells were infected with recombinant Sendai virus (SeV) carrying human wild-type TRIM5 α (■; hu), human 110Arg-TRIM5 α (□; 110R-hu), human/African green monkey (AGM) chimeric TRIM5 α (▲; hu/AGM), human/AGM chimeric 110-ArgTRIM5 α (△; 110R-hu/AGM), CM-TRIM5 α -SPRY(-) (○; SPRY(-)) or AGM-TRIM5 α -CC(-) (●; CC(-)). Nine hours after infection, cells were inoculated with HIV-1 NL43 (a and c) or HIV-2 GH123 (b). Culture supernatants were periodically assayed for levels of p24 (a and c) or p25 (b). MT4 cells were simultaneously infected with two recombinant SeVs at 5 plaque-forming unit (PFU) per cell for

expressing dominant negative TRIM5 α -SPRY(-) than in cells with noninterfering TRIM5 α -CC(-), demonstrating the anti-HIV-2 activity of endogenous human TRIM5 α (Fig. 2b). Both wild-type and variant human TRIM5 α exhibited weak but apparent anti-HIV-2 activity, and HIV-2 grew to higher titers in cells expressing the human 110Arg-TRIM5 α than in cells with the human wild-type-TRIM5 α (Fig. 2b). In human/AGM chimeric version, wild-type TRIM5 α completely restricted HIV-2 (Fig. 2b). In contrast, HIV-2 grew to slightly higher titers in cells expressing the human/AGM chimeric 110Arg-TRIM5 α than in cells expressing the wild-type human/AGM chimeric TRIM5 α (Fig. 2b). These results indicated that the Gly110Arg variant weakened the anti-HIV-1 and anti-HIV-2 activities of human TRIM5 α in MT4 cells.

We recently found that the expression of TRIM5 α protein introduced by SeV varied depending on cell types, that is, it was much lower in CEM-SS than in MT4 cells [25]. To evaluate the anti-HIV-1 activity of variant TRIM5 α at more physiological levels of expression, we performed experiments using CEM-SS (Fig. 2c). Neither wild-type nor variant human TRIM5 α exhibited anti-HIV-1 activity, probably due to the low level expression of TRIM5 α in CEM-SS cells. However, HIV-1 grew to approximately 10 times higher levels in cells expressing the human/AGM chimeric 110Arg-TRIM5 α than in cells with the wild-type chimeric TRIM5 α , suggesting that the anti-HIV-1 activity of TRIM5 α in CEM-SS cells was also reduced by the Gly110Arg substitution. Therefore, we concluded that the Gly110Arg polymorphism affected both the anti-HIV-1 and anti-HIV-2 activities of human TRIM5 α .

Truncated G176del-TRIM5 enhanced antiviral activity of coexpressed TRIM5 α

To express the G176del-TRIM5, we added an HA-tag at its N-terminus, because the expression of G176del-TRIM5 protein tagged with HA at the C-terminus could not be detected. Although the expression of HA-fused protein was clearly visualized by anti-HA antibody, its expression was much lower than the full-length TRIM5 α (Fig. 1b). In cells infected with SeV expressing the

G176del-TRIM5, HIV-2 grew to the same titers as those in cells infected with SeV expressing a nonfunctional mutant TRIM5 α -CC(-), indicating that the G176del-TRIM5 lost the anti-HIV-2 activity (data not shown). We then investigated whether the G176del-TRIM5 showed any effects on the anti-HIV activity of coexpressed full-length TRIM5 α , because all individuals carrying the G176del variant were in the heterozygous state. As shown in Fig. 2d and 2e, both HIV-1 and HIV-2 were restricted in cells simultaneously expressing the human/AGM chimeric TRIM5 α and TRIM5 α -CC(-). As expected, both HIV-1 and HIV-2 grew to high titers in cells expressing the human/AGM chimeric TRIM5 α and the dominant negative mutant TRIM5 α -SPRY(-) [24]. In contrast, both HIV-1 and HIV-2 were severely restricted in cells expressing the human/AGM chimeric TRIM5 α and G176del-TRIM5 as compared within cells expressing the human/AGM chimeric TRIM5 α and TRIM5 α -CC(-). These results suggested that the G176del-TRIM5 enhanced the antiviral activity induced by the full-length TRIM5 α .

Next, we investigated whether the truncated G176del-TRIM5 could affect the expression of TRIM5 α . Expressions of the human/AGM chimeric TRIM5 α in cells expressing either TRIM5 α -SPRY(-), TRIM5 α -CC(-) or G176del-TRIM5 are shown in Fig. 2f. Amount of human/AGM chimeric TRIM5 α in cells coexpressing the G176del-TRIM5 was 1.7 times higher than that in cells coexpressing the TRIM5 α -SPRY(-). When we infected a constant amount of SeV expressing the human/AGM TRIM5 α in combination with the increasing amounts of SeV expressing the G176del TRIM5 variant, we found that the expression level of human/AGM TRIM5 α was increased by the G176del TRIM5 (Fig. 2g).

Discussion

It is widely accepted that within host cells, there are restriction factors that oppose retroviral replication more effectively than the conventional arms of the immune

Fig. 2. (continued)

each SeV. CM-TRIM5 α -SPRY(-) and human/AGM chimeric TRIM5 α (○; S(-) + hu/AGM), AGM-TRIM5 α -CC(-) and human/AGM chimeric TRIM5 α (■; CC(-) + hu/AGM), or hemagglutinin (HA)-G176del-TRIM5 and human/AGM chimeric TRIM5 α (▲; G176del + hu/AGM) were simultaneously inoculated. Nine hours after the infection, cells were superinfected with HIV-1 NL43 (d) or HIV-2 GH123 (e) and culture supernatants were periodically assayed for levels of p24 (d) or p25 (e). The means with standard deviations of triplicate samples are shown. (f) Western blottings for TRIM5 protein and cyclophilin A from MT4 cells infected with SeV expressing the HA-tagged human/AGM chimeric TRIM5 α (human/AGM-TRIM5 α -HA) coexpressed with the AGM-TRIM5 α -CC(-)-HA (lane 1; TRIM5 α -CC(-)-HA), coexpressed with the CM-TRIM5 α -SPRY(-)-HA (lane 2; TRIM5 α -SPRY(-)-HA), or with the HA-G176del-TRIM5 (lane 3: HA-G176del-TRIM5). The relative amounts of human/AGM chimeric TRIM5 α are shown on the top with the standard deviation of six independent samples. (g) MT4 cells were infected with SeV expressing the HA-tagged human/AGM chimeric TRIM5 α coinfecting with SeV expressing the HA-G176del-TRIM5 or an empty vector parental Z strain. The multiplicity of infection in each SeV is shown on the bottom. The relative amounts of human/AGM chimeric TRIM5 α are shown on the top with standard deviation of triplicate samples.

system [1,2]. Because TRIM5 α has crucial roles in the intracellular defense mechanisms against HIV-1 [2–4], sequence variations in TRIM5 α might be associated with the susceptibility to HIV-1 infection and/or progression to AIDS. In this study, we demonstrated the association of 43Tyr-allele with the reduced susceptibility to HIV-1 infection in two ethnically distinct populations. In addition, we identified two novel rare variants, Gly110Arg and G176del, both of which had an impact on the anti-HIV-1 activity and susceptibility to HIV-1 infection.

The association of His43Tyr with the HIV-1 infection or AIDS progression has been tested in several studies, but the results were not consistent [16–20]. We found that the 43Tyr-allele was less frequent in the HIV-1-infected patients than in the ethnic-matched controls in both Japanese and Indian populations. The study sizes were not very large, but two independent ethnic populations did exhibit the same trends for the association with His43Tyr. We previously analyzed HIV-1-infected long-term nonprogressors and standard progressors in France and Japan for the TRIM5 α polymorphisms and failed to find any differences in the frequency of 43Tyr-allele between these two HIV-infected groups both in France and Japan [19]. However, the allele frequency of 43Tyr in the Japanese HIV-1-infected patients we analyzed in the present study was 0.143, which was similar to that in the present study (0.133, Table 1). Interestingly, several studies have reported that the anti-HIV-1 activity of TRIM5 α with 43Tyr was lower than that with 43His [16,18]. In our previous study, we also showed that the anti-HIV-1 activity of TRIM5 α with 43Tyr was lower than that with 43His, although the difference in anti-HIV-1 activity between the 43His-TRIM5 α and 43Tyr-TRIM5 α was very small [19]. In spite of the lower anti-HIV-1 activity of the 43Tyr-TRIM5 α , several epidemiological studies have shown that the 43Tyr-allele was associated with the reduced susceptibility to HIV-1 infection [16,18], as demonstrated in this study. The reasons for the discrepancy between the epidemiological and functional effects of His43Tyr remain unclear at the moment. On the other hand, van Manen *et al.* [20] recently reported that homozygous status for 43Tyr was associated with the accelerated disease progression in white populations, which was consistent with the effect of His43Tyr variation on the anti-HIV-1 activity. Further epidemiological studies will be required to clarify the impact of His43Tyr on the susceptibility to HIV-1 infection and AIDS progression.

We also showed that the impact of His43Tyr on the susceptibility to HIV-1 infection was slightly different between Japanese and Indian. The frequency of 43Tyr-allele in the Indian HIV-1-infected patients was significantly lower than that in the Indian controls, but the significant difference was not found in Japanese.

Different distribution of HIV-1 subtypes might be one of the reasons for the different contribution of 43Tyr-allele in the susceptibility, because all of the Indian patients examined in this study were infected with HIV-1 subtype C, whereas only subtype B was observed in our Japanese patients, as was found in the previous reports [26,27]. Kaumanns *et al.* [28] have reported that the antiretroviral activities of TRIM5 α differed among the HIV-1 subtypes, although the differences in the in-vitro antiretroviral effect of TRIM5 α between the subtypes C and B were not evident.

In this study, one focus was the functional impact of two rare TRIM5 α variants found in our epidemiological studies. First, our findings indicated that the 110Arg variant weakened the anti-HIV-1 and anti-HIV-2 activities of human TRIM5 α in human T-cell lines. This variant was observed more frequently in the Japanese HIV-1-infected patients than in the controls. This variation substitutes the smallest amino acid glycine with a positively charged amino acid arginine at the 110 amino acid position of TRIM5 α and is located next to the amino acid residue 109Gly, which is suspected to be a zinc-coordinating residue in the B-box 2 domain [29]. This drastic change in amino acid character might change the structure of TRIM5 α , in which an intact B-box 2 domain was essential for the antiretroviral activity of TRIM5 α and disruption of the TRIM5 α B-box domain by specific amino acid substitution resulted in loss of retroviral restriction [8,30–32]. The 3D structure of the amino acid residues 11–133 of TRIM5 α was modeled by SWISS-MODEL, an Automated Comparative Protein Modeling Server (<http://swissmodel.expasy.org/SWISS-MODEL.html>) [33]. As shown in Fig. 3, residue 110 constituted one of the β sheets in the N-terminal half of TRIM5 α . Interestingly, the location of residue 110 was close to residue 43 in the modeled 3D structure of TRIM5 α (Fig. 3b and 3c). As described previously, His43Tyr was reported to affect antiretroviral activity. These data suggested that residue 110 might be one of the key amino acid residues in the TRIM5 α structure like residue 43.

Second, we found that the truncated G176del-TRIM5 enhanced the antiviral activity of coexpressed full-length TRIM5 α . Coinfection of SeVs expressing the G176del-TRIM5 and human/AGM-TRIM5 α was accompanied by the increased protein level of full-length human/AGM-TRIM5 α . The amount of human/AGM chimeric TRIM5 α in cells coinfecting with SeVs expressing the G176del-TRIM5 was 1.7 times higher than in cells coinfecting with SeVs expressing the TRIM5 α -SPRY(–). These data suggested that the truncated TRIM5 α was degraded rapidly, resulting in a delay of the degradation process of full-length TRIM5 α and leading to the augmentation of protein levels. Recently, we observed that coexpression of a splice variant of TRIM5, TRIM5 γ , increased the amount of TRIM5 α .




Fig. 3. Structural model of variant TRIM5 α . (a) The primary structure is illustrated by the amino acid sequence of residues 11–133. One-letter amino acid code is used. The secondary structure is diagrammatically represented below the sequence showing the regions of the polypeptide chain, which are folded into the α helices (red boxes), the β sheets (green arrows) and random coils (black lines). (b) The 3D structure of amino acids 11–133 was modeled by SWISS-MODEL. (c) A magnified view, which is enclosed with a white square in (b), is shown. The arrows indicate the 43His and 110Gly residues.

TRIM5 γ itself was unstable and its expression was lower than TRIM5 α (Maegawa, unpublished data).

In this study, we identified three individuals harboring the G176del variant in the heterozygous state only in the Japanese controls, not in the HIV-1-infected patients. It appeared that the homozygous state for the G176del-allele would increase the susceptibility to HIV-1 infection, because it should result in null TRIM5 α activity. It follows from the enhanced anti-HIV-1 activity of full-length TRIM5 α by the truncated G176del-TRIM5 that the heterozygous state for the 176del-allele might mask the reduction in TRIM5 α gene number and thus might not have a serious effect on the susceptibility to HIV-1 infection.

We demonstrated the association of common variant 43Tyr with the reduced susceptibility to HIV-1 infection in Japanese and Indian. We also identified two rare variants, 110Arg and G176del, which decreased and increased, respectively, the anti-HIV-1 activity in human cells expressing TRIM5 α . We suggested that the sequence variations of TRIM5 α were tightly linked to the susceptibility to or protection against the HIV-1 infection. However, further epidemiological studies using larger population samples will be required to clarify the impact of these rare variants on the HIV-1/AIDS susceptibility. In an effort to understand the genetic factors controlling the HIV-1 infection and AIDS

progression, considerable attention should be paid to rare variants in addition to common variants in the candidate genes.

Acknowledgements

We are grateful for sample collection in Japan to Drs H. Hanabusa (Ogikubo Hospital), J. Matsuda (Teikyo University School of Medicine), M. Sakai (University of Occupational and Environmental Health), S. Ikeda (Sasebo Municipal Hospital) and T. Fujii (Hiroshima University School of Medicine) and for sample collection in India to Dr S.K. Sharma (All India Institute of Medical Sciences). We would like to thank Gaurav Sharma for DNA preparation. This work was supported in part by research grants from the Ministry of Health, Labor and Welfare, Japan, the Japan Health Science Foundation, the program of Founding Research Centers for Emerging and Reemerging Infection Disease, the Japan Health Science Foundation, the program of Research on Publicly Essential Drugs and Medical Devices, Grant-in-Aids for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan, a grant from Heiwa Nakajima Foundation, a grant for Japan-India collaboration research program from Japan Society for the Promotion of Science, and grants from the Department of Biotechnology

(DBT), Ministry of Science and Technology, Government of India and the Indian Council of Medical Research (ICMR).

References

1. Sheehy AM, Gaddis NC, Choi JD, Malim MH. Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* 2002; **418**:646–650.
2. Strelau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J. The cytoplasmic body component TRIM5 α restricts HIV-1 infection in Old World monkeys. *Nature* 2004; **427**:848–853.
3. Wu X, Anderson JL, Campbell EM, Joseph AM, Hope TJ. Proteasome inhibitors uncouple rhesus TRIM5 α restriction of HIV-1 reverse transcription and infection. *Proc Natl Acad Sci U S A* 2006; **103**:7465–7470.
4. Strelau M, Perron M, Lee M, Li Y, Song B, Javadakht H, et al. Specific recognition and accelerated uncoating of retroviral capsids by the TRIM5 α restriction factor. *Proc Natl Acad Sci U S A* 2006; **103**:5514–5519.
5. Yap MW, Nisole S, Lynch C, Stoye JP. Trim5 α protein restricts both HIV-1 and murine leukemia virus. *Proc Natl Acad Sci U S A* 2004; **101**:10786–10791.
6. Perron MJ, Strelau M, Lee M, Javanbakht H, Song B, Sodroski J. The human TRIM5 α restriction factor mediates accelerated uncoating of the N-tropic murine leukemia virus capsid. *J Virol* 2007; **81**:2138–2148.
7. Ozato K, Shin DM, Chang TH, Morse HC 3rd. TRIM family proteins and their emerging roles in innate immunity. *Nat Rev Immunol* 2008; **8**:849–860.
8. Perez-Caballero D, Hatzioannou T, Yang A, Cowan S, Bieniasz PD. Human tripartite motif 5 α domains responsible for retrovirus restriction activity and specificity. *J Virol* 2005; **79**:8969–8978.
9. Strelau M, Perron M, Welikala S, Sodroski J. Species-specific variation in the B30.2 (SPRY) domain of TRIM5 α determines the potency of human immunodeficiency virus restriction. *J Virol* 2005; **79**:3139–3145.
10. Yap MW, Nisole S, Stoye JP. A single amino acid change in the SPRY domain of human Trim5 α leads to HIV-1 restriction. *Curr Biol* 2005; **15**:73–78.
11. Sawyer SL, Wu LI, Emerman M, Malik HS. Positive selection of primate TRIM5 α identifies a critical species-specific retroviral restriction domain. *Proc Natl Acad Sci U S A* 2005; **102**:2832–2837.
12. Nakayama EE, Miyoshi H, Nagai Y, Shioda T. A specific region of 37 amino acid residues in the SPRY (B30.2) domain of African green monkey TRIM5 α determines species-specific restriction of simian immunodeficiency virus SIVmac infection. *J Virol* 2005; **79**:8870–8877.
13. Kono K, Song H, Shingai Y, Shioda T, Nakayama EE. Comparison of antiviral activity of rhesus and cynomolgus monkey TRIM5 α s against HIV-2 infection. *Virology* 2008; **373**:447–456.
14. O'Brien SJ, Nelson GW. Human genes that limit AIDS. *Nat Genet* 2004; **36**:565–574.
15. Kaslow RA, Dorak T, Tang J. Influence of host genetic variation on susceptibility to HIV type 1 infection. *J Infect Dis* 2005; **191**:S68–S77.
16. Sawyer SL, Wu LI, Akey JM, Emerman M, Malik HS. High-frequency persistence of an impaired allele of the retroviral defense gene TRIM5 α in humans. *Curr Biol* 2006; **16**:95–100.
17. Speelman EC, Livingston-Rosanoff D, Li SS, Vu Q, Bui J, Geraghty DE, et al. Genetic association of the antiviral restriction factor TRIM5 α with human immunodeficiency virus type 1 infection. *J Virol* 2006; **80**:2463–2471.
18. Javanbakht H, An P, Gold B, Petersen DC, O'Huigin C, Nelson GW, et al. Effects of human TRIM5 α polymorphisms on antiretroviral function and susceptibility to human immunodeficiency virus infection. *Virology* 2006; **354**:15–27.
19. Nakayama EE, Carpentier W, Costagliola D, Shioda T, Iwamoto A, Debre P, et al. Wild type and H43Y variant of human TRIM5 α show similar antihuman immunodeficiency virus type 1 activity both in vivo and in vitro. *Immunogenetics* 2007; **59**:511–515.
20. van Manen D, Rits MA, Beugeling C, van Dort K, Schuitemaker H, Kootstra NA. The effect of Trim5 polymorphisms on the clinical course of HIV-1 infection. *PLoS Pathog* 2008; **4**:e18.
21. Munkanta M, Terunuma H, Takahashi M, Hanabusa H, Miura T, Ikeda S, et al. HLA-B polymorphism in Japanese HIV-1 infected long-term surviving hemophiliacs. *Viral Immunol* 2005; **18**:500–505.
22. Nakajima T, Ohtani H, Naruse T, Shibata H, Mimaya J, Terunuma H, et al. Copy number variations of CCL3L1 and long-term prognosis of HIV-1 infection in asymptomatic HIV-infected Japanese with hemophilia. *Immunogenetics* 2007; **59**:793–798.
23. Song H, Nakayama EE, Yokoyama M, Sato H, Levy JA, Shioda T. A single amino acid of the human immunodeficiency virus type 2 capsid affects its replication in the presence of cynomolgus monkey and human TRIM5 α s. *J Virol* 2007; **81**:7280–7285.
24. Maegawa H, Nakayama EE, Kuroishi A, Shioda T. Silencing of tripartite motif protein (TRIM) 5 α mediated anti-HIV-1 activity by truncated mutant of TRIM5 α . *J Virol Methods* 2008; **151**:249–256.
25. Kono K, Bozek K, Domingues FS, Shioda T, Nakayama EE. Impact of a single amino acid in the variable region 2 of the Old World monkey TRIM5 α SPRY(B30.2) domain on antihuman immunodeficiency virus type 2 activity. *Virology* 2009; **388**:160–168.
26. Lakhshas S, Thakar M, Godbole S, Tripathy S, Paranjape R. HIV infection in India: epidemiology, molecular epidemiology and pathogenesis. *J Biosci* 2008; **33**:515–525.
27. Weniger BG, Takebe Y, Ou CY, Yamazaki S. The molecular epidemiology of HIV in Asia. *AIDS* 1994; **8**:S13–S28.
28. Kaumanns P, Hagmann I, Dittmar MT. Human TRIM5 α mediated restriction of different HIV-1 subtypes and Lv2 sensitive and insensitive HIV-2 variants. *Retrovirology* 2006; **3**:79.
29. Massiah MA, Matts JA, Short KM, Simmons BN, Singireddy S, Yi Z, Cox TC. Solution structure of the MID1 B-box2 CHC(D/C)C(2)H(2) zinc-binding domain: insights into an evolutionarily conserved RING fold. *J Mol Biol* 2007; **369**:1–10.
30. Diaz-Griffero F, Kar A, Perron M, Xiang SH, Javanbakht H, Li X, et al. Modulation of retroviral restriction and proteasome inhibitor-resistant turnover by changes in the TRIM5 α B-box 2 domain. *J Virol* 2007; **81**:10362–10378.
31. Javanbakht H, Diaz-Griffero F, Strelau M, Si Z, Sodroski J. The contribution of RING and B-box 2 domains to retroviral restriction mediated by monkey TRIM5 α . *J Biol Chem* 2005; **280**:26933–26940.
32. Li X, Sodroski J. The TRIM5 α B-box 2 domain promotes cooperative binding to the retroviral capsid by mediating higher-order self-association. *J Virol* 2008; **82**:11495–11502.
33. Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODEL workspace: a web-based environment for protein structure homology modeling. *Bioinformatics* 2006; **22**:195–201.

Research

Open Access

HLA-Cw*04 allele associated with nevirapine-induced rash in HIV-infected Thai patients

Sirirat Likanonsakul*¹, Tippawan Rattanatham¹, Siriluk Feangvad¹,
Sumonmal Uttayamakul¹, Wisit Prasithsirikul¹, Preecha Tunthanathip¹,
Emi E Nakayama² and Tatsuo Shioda²

Address: ¹Bamrasnaradura Infectious Diseases Institute, Department of Disease Control, Ministry of Public Health, Nonthaburi, Thailand and ²Research Institute for Microbial Diseases, Osaka University, Osaka, Japan

Email: Sirirat Likanonsakul* - imvbi@health2.moph.go.th; Tippawan Rattanatham - rattanatham@gmail.com;
Siriluk Feangvad - siriluktucky@gmail.com; Sumonmal Uttayamakul - sumonmal@health.moph.go.th;
Wisit Prasithsirikul - drwisit_p@yahoo.com; Preecha Tunthanathip - Tsainui@hotmail.com; Emi E Nakayama - emien@biken.osaka-u.ac.jp;
Tatsuo Shioda - shioda@biken.osaka-u.ac.jp

* Corresponding author

Published: 21 October 2009

Received: 19 August 2009

AIDS Research and Therapy 2009, 6:22 doi:10.1186/1742-6405-6-22

Accepted: 21 October 2009

This article is available from: <http://www.aidsrestherapy.com/content/6/1/22>

© 2009 Likanonsakul et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A high incidence of rash has been reported in HIV-1 patients who received the anti-retroviral drug nevirapine. In addition, several studies have suggested that polymorphisms of human leukocyte antigen (HLA) genes may play important roles in nevirapine-induced rash. The aim of the present study was to evaluate the effects of different HLA-C alleles on rash associated with nevirapine in patients who started highly active anti-retroviral therapy (HAART) containing nevirapine in Thailand.

Results: A case-control study was carried out involving HIV-1 patients under treatment at Bamrasnaradura Infectious Diseases Institute, Nonthaburi, Thailand between March 2007 and March 2008. The study included all HIV/AIDS patients being treated with nevirapine-containing regimens. The study population comprised 287 HIV/AIDS patients of whom 248 were nevirapine-tolerant and 39 developed rash after nevirapine treatment. From the nevirapine-tolerant patients, 60 were selected as the control group on the basis of age, sex, and therapy history matched for nevirapine-induced rash cases. We observed significantly more HLA-Cw*04 alleles in nevirapine-induced rash cases than in nevirapine-tolerant group, with frequencies of 20.51% and 7.50%, respectively ($P = 0.009$). There were no significant differences between the rash and tolerant groups for other HLA-C alleles except for HLA-Cw*03 ($P = 0.015$).

Conclusion: This study suggests that HLA-Cw*04 is associated with rash in nevirapine treated Thais. Future screening of patients' HLA may reduce the number of nevirapine-induced rash cases, and patients with alleles associated with nevirapine-induced rash should be started on anti-retroviral therapy without nevirapine.

Background

Highly active antiretroviral therapy (HAART) has significantly improved the prognosis of HIV-1-infected patients and prolonged AIDS-free survival [1]. HAART has also resulted in immune restoration and reduction of morbidity and mortality even for patients with advanced HIV-1 infection [1,2]. Nevirapine (NVP) is a non-nucleoside reverse transcriptase inhibitor (NNRTI) that has been shown to have high antiretroviral efficacy [3]. NVP-based HAART regimens have therefore been widely used in resource-limited countries because of their efficacy, availability and relatively low cost. In Thailand, the Government Pharmaceutical Organization (GPO) has produced GPOvir, a low cost (US\$ 30 per month) and fixed-dose combination of NVP, stavudine (D4T), and lamivudine (3TC), which has been commercially available since March 2002.

However, NVP-associated rash has been reported to be as high as 48% after the start of treatment with this inhibitor [4]. Nearly 90% of the side effects of GPOvir are thought to be due to NVP hypersensitivity [5]. Skin rash is the most common adverse drug reaction associated with NVP, and hypersensitivity reaction to NVP is rapid and severe when drug administration is suspended and re-challenged [6]. Most patients develop rash between the first and third week of treatment [7], including the more severe forms of rash such as extensive maculopapular rash, serum sickness-like reaction, hypersensitivity syndrome, Steven-Johnson syndrome and toxic epidermal necrosis [7,8]. NVP-induced rash has been reported in 4.3-36% of adults [9,10] with the incidence for Thai patients ranging from 6 to 21% [5], reflecting the comparatively high incidence of rash in Asians [11]. Several features of NVP hypersensitivity suggest that genetic factors may play an important predisposing role in NVP hypersensitivity, in which NVP

itself or NVP-induced antigens may trigger an immunological response that is dependent on CD4 T lymphocytes in susceptible hosts. This supports the hypothesis that the hypersensitivity reaction to NVP may be HLA-associated [9,12,13], while HLA-alleles have also been identified as clinically relevant susceptibility markers for hypersensitivity reaction to another antiretroviral drug [14]. Recent studies have shown that in Japan the *HLA-Cw*08* allele is associated with NVP hypersensitivity [15]. The objective of the study presented here was to compare allele frequency of *HLA-C* in Thai patients with rash who had to change from GPOvir to a regime containing efavirenz and those who were NVP-tolerant.

Results

A case-control study was carried out. The study population comprised 287 HIV/AIDS patients of whom 248 were nevirapine-tolerant and 39 developed rash. From the nevirapine-tolerant patients, 60 were selected as the control group on the basis of age, sex, and therapy history matched for nevirapine-induced rash cases. As shown in Table 1, the demographic and clinical characteristics of patients with NVP-induced rash were very similar to those of NVP-tolerant patients. The medians of CD4 cell counts of NVP-induced rash and NVP-tolerant group were comparable at both time points of immediately before NVP treatment and 6 month after treatment. It is known that HIV-infected patients frequently suffered from allergic drug reactions [16]. Nearly all of rash cases (37 out of 39) used steroids and more than half (35 out of 60 cases) of NVP-tolerant patients also used steroids. Four out of 35 NVP-tolerant steroid users had developed mild rash that could be controlled by steroid. Remaining 31 NVP-tolerant patients used steroid to suppress allergic reactions including chronic allergic skin, rhinitis, asthma, and drug reactions upon *Pneumocystis carinii* pneumonia treat-

Table 1: Demographics and immunological variables of the NVP-induced rash and NVP-tolerant groups

Variables	NVP-induced rash (n = 39)	NVP-tolerant (n = 60)	P value
Age (median IQR)	39.0 (34.0-44.0)	38.0 (35.0-41.75)	*0.71
Sex [n(%)]			**0.68
Male	22 (56.41%)	31 (51.67%)	
Female	17 (43.59%)	29 (48.33%)	
Duration of Treatment, year (median IQR)	1 (0-3)	1 (0-2.75)	*0.47
Pre-NVP-treatment CD4 T-cell count × 10 ⁶ /l (median IQR)	43.5 (19.50-135.50)	55 (28.25-137.50)	*0.71
Post-NVP-treatment CD4 T-cell count × 10 ⁶ /l (median IQR)	243 (155.00-328.00)	241 (185.00-312.25)	*0.98

*Mann-Whitney U-test

**Chi square test

ment, and immune reconstitution inflammatory syndrome. The NVP-induced rash cases manifested severe rash, which could not be suppressed by steroid, and had to change the regimen.

The frequencies of the *HLA-C* alleles identified in the 39 samples in the NVP-induced rash group and 60 samples in the NVP tolerant group are presented in Table 2. Frequency of *HLA-Cw*04* was approximately 21% for the patients with NVP-induced rash and 7.5% for the NVP-tolerant group, showing a statistically significant difference in *HLA-Cw*04* allele frequency ($P = 0.0088$, Fisher's exact test). The reported *HLA-Cw*04* allele frequency for the normal Thai population (0.102) [17] is higher than that of the NVP-tolerant group (0.075) and lower than that of the NVP-induced rash group (0.205). Although statistical significance of this difference was lost after stringent Bonferroni correction ($P_c = 0.088$), these results suggested that *HLA-Cw*04* was associated with NVP-induced rash in HIV-1 infected Thai patients. One-third of the patients with NVP-induced rash (13 out of 39) carried *HLA-Cw*04* in comparison with 15% of the NVP-tolerant patients (9 out of 60) ($P = 0.047$, Fisher's exact test; $P_c = 0.47$). When we compared 39 NVP-induced rash patients with 25 NVP-tolerant patients who did not receive steroid, the concentration of *HLA-Cw*04* allele in NVP rash cases was still apparent (0.205 vs 0.060, $P = 0.04$, Fisher's exact test).

In contrast to *HLA-Cw*04*, fewer *HLA-Cw*03* alleles were found in patients with NVP-induced rash than in NVP-tolerant ones (0.051 and 0.167) ($P = 0.015$, Fisher's exact test; $P_c = 0.15$). Approximately 7.7% of patients with NVP-induced rash (3 out of 39) carried *HLA-Cw*03* in comparison with 30% of the NVP-tolerant patients (18 out of 60) ($P = 0.011$ Fisher's exact test, $P_c = 0.11$). There were no significant differences between the NVP-induced rash and NVP-tolerant groups in allele frequencies of

*HLA-Cw*01*, *HLA-Cw*05*, *HLA-Cw*06*, *HLA-Cw*07*, *HLA-Cw*08*, *HLA-Cw*12*, *HLA-Cw*14*, and *HLA-Cw*15*. Other *HLA-Cw** alleles were not detected in the tested samples. The NVP-tolerant group allele frequencies of *HLA-Cw*03* (0.167), *HLA-Cw*08* (0.125) and *HLA-Cw*12* (0.067) were very similar to those of the normal Thai population (0.174, 0.144 and 0.06, respectively) [17], which suggests that our genotyping was accurate.

*HLA-DRB1*01* was reported to be associated with NVP hypersensitivity in Australian [13] and French [18] cohorts. We therefore performed PCR-sequence specific oligonucleotide probe (SSOP) method to detect *HLA-DRB1*01* alleles. Contrary to our expectation, we failed to detect *HLA-DRB1*01* allele in 39 NVP-induced rash cases, while we detected five of this allele in 60 NVP-tolerant patients. This result suggested that the *HLA-DRB1*01* allele may not be involved in NVP rash in Thai population.

The C allele of the SNP rs9264942, located in the 5'upstream region of the *HLA-C* gene, was reported to associate with higher levels of *HLA-C* expression and *HLA-Cw*04* allele [19]. To know whether or not higher levels of *HLA-C* gene expression associated with NVP-induced rash, we genotyped rs9264942 SNP by TaqMan SNP genotyping system. The C allele frequency in 39 NVP rash cases was 0.45, while it was 0.38 in 60 NVP-tolerant patients ($P = 0.36$, Chi square test). This result suggested that the *HLA-Cw*04* allele itself rather than the relative high levels of *HLA-C* expression was involved in NVP-induced rash.

Discussion

In the study reported here, we genotyped *HLA-C* alleles of 39 patients with NVP-induced rash and 60 NVP-tolerant Thai patients, and found that frequency of *HLA-Cw*04* was higher in NVP-induced rash Thai patients than in NVP-tolerant patients. While the number of samples in

Table 2: Occurrence of HLA-C alleles in the nevirapine (NVP)-rash cases and NVP-tolerant controls in Thailand

Allele	NVP-induced rash number (%)	NVP-tolerant number (%)	*P value
<i>HLA-Cw*01</i>	12 (15.38)	12 (10.00)	NS
<i>HLA-Cw*03</i>	4 (5.13)	20 (16.67)	0.01
<i>HLA-Cw*04</i>	16 (20.51)	9 (7.50)	0.009
<i>HLA-Cw*05</i>	2 (2.56)	3 (2.50)	NS
<i>HLA-Cw*06</i>	8 (10.26)	9 (7.50)	NS
<i>HLA-Cw*07</i>	19 (24.36)	39 (35.50)	NS
<i>HLA-Cw*08</i>	9 (11.54)	15 (12.50)	NS
<i>HLA-Cw*12</i>	6 (7.69)	8 (6.67)	NS
<i>HLA-Cw*14</i>	0	3 (2.50)	NS
<i>HLA-Cw*15</i>	2 (2.56)	2 (1.67)	NS
Total	78	120	

*Fisher's exact test
NS: $P > 0.05$

our study is small, the increased frequency of *HLA-Cw*04* in patients with NVP-induced rash suggested that this allele plays an important role in the development of rash after GPOvir treatment. *HLA-Cw*04* was found to be associated with rapid development of AIDS-defining conditions in Caucasians [20,21] but to have a protective effect in African Americans [22]. In hepatitis C virus infection cases, *HLA-Cw*04* was associated with viral persistence [23].

Previous studies [15,24] have suggested that *HLA-Cw*08* is associated with NVP hypersensitivity. Littera *et al.* studied 49 Sardinian HIV-1-positive patients treated with NVP and reported that *HLA-Cw*08* and/or *HLA-B*14(65)* is associated with NVP hypersensitivity [24]. Subsequently, Gatanaga *et al.* studied HIV-1 infected individuals in Japan [15]. In this study, 41 patients had a history of NVP treatment, 12 of whom showed NVP hypersensitivity. The frequency of *HLA-B*14* is nearly 0% in Japan. On the other hand, the frequency of *HLA-Cw*08*-positive patients in the NVP hypersensitive group was 42%, which was significantly higher than that the NVP tolerant group (10%) [15]. In our study, however, the frequencies of *HLA-Cw*08* in the NVP rash and tolerant groups were 0.115 and 0.125, respectively, without any significant difference between the two groups ($P = 1.000$, Fisher's test). Although the precise reason for the difference between the findings of these previous studies and ours is not clear at present, there were several differences between them. First, we genotyped approximately three times more patients with NVP-induced rash than was done in the previous studies. Second, our study focused on rash after NVP treatment but the other studies dealt with patients with rash and/or hepatotoxicity. Furthermore, it is possible that the levels of linkage disequilibrium between *HLA-C* alleles and those of other HLA locus and/or other genes differ among different ethnic groups. As described above, *HLA-Cw*04* was associated with rapid AIDS progression in Caucasians [20,21] but to have a protective effect in African Americans [22]. Thus, it is possible that other SNP(s) or HLA allele(s) responsible for NVP-induced rash is in a stronger linkage disequilibrium with *HLA-Cw*04* in Thais than the other ethnic groups tested previously.

Previous studies have also suggested that higher CD4 counts at baseline increased risk of NVP-induced rash [25-30]. In our study, however, the median baseline CD4 counts of NVP-induced rash cases (43.5 cells/ μ l) was very similar to that of 248 NVP-tolerant patients (43 cells/ μ l). The effects of high baseline CD4 counts on risk of NVP-induced rash were reportedly observed mainly in patients whose CD4 counts were over 250 cells/ μ l [29]. Accordingly, nearly all of patients in our study started NVP-containing regimen after their CD4 counts dropped below 250 cells/ μ l. Therefore, it is also possible that low baseline

CD4 counts in our study limit the ability to detect an HLA association reported previously [15,24]. Nevertheless, our results are practically meaningful since most of HIV-1-infected individuals in Thailand start antiretroviral treatment after their CD4 counts drop below 250 cells/ μ l.

One report from Thailand demonstrated that the effects of high baseline CD4 counts on risk of NVP-induced rash were still observed even in patients whose baseline CD4 counts were below 250 cells/ μ l, although the levels of such effects was very small in patients whose baseline CD4 counts were below 200 cells/ μ l [30]. Therefore, we divided patients according to their baseline CD4 counts. When we picked up patients whose baseline CD4 counts were over 100 cells/ μ l, *HLA-Cw*04* allele frequency was 0.25 in 14 NVP-induced rash cases and 0.025 in 20 NVP-tolerant patients. Statistical significance of this difference rather increased in these groups ($P = 0.0068$). When we picked up patients whose baseline CD4 were less than 100 cells/ μ l, *HLA-Cw*04* allele frequency was 0.2045 in 22 NVP-induced rash cases and 0.0897 in 39 NVP-tolerant patients. Although there was an apparent trend towards high *HLA-Cw*04* frequency in NVP-induced rash cases, statistical significance of the difference greatly reduced in these groups ($P = 0.094$). Therefore, the difference in the allele frequency was more prominent in patients with higher CD4 counts even in our patient group.

During the preparation of this manuscript, Chantarangsu *et al.* reported a strong association between *HLA-B*3505* and NVP-induced skin rash in HIV-infected Thai patients [31]. Our results at least partly confirm theirs since *HLA-Cw*0401* showed the second highest levels of difference in allele frequency between NVP-rash and NVP-tolerant patients in their study [31]. It is known that *HLA-Cw*04* is in a linkage disequilibrium with *HLA-B*35* in Thais [32]. It is, thus, necessary to investigate whether or not *HLA-B*3505* is also overrepresented in NVP-rash cases in our study.

Conclusion

We observed higher frequency of *HLA-Cw*04* in NVP-induced skin rash than in NVP-tolerant patients in Thailand. In addition to *HLA-Cw*04* and/or *HLA-B*3505*, future screening of patients' HLA and genes involved in hypersensitive reactions may identify other alleles responsible for the incidence of NVP-induced rash. Patients possessing alleles responsible for NVP-rash should be started on anti-retroviral therapy without NVP.

Methods

Clinical specimens

A case-control study was carried out with HIV-1 infected patients who were under treatment at Bamrasnaradura Infectious Diseases Institute, Ministry of Public Health,

Nonthaburi, Thailand. The targeted study population comprised 672 HIV-1/AIDS patients and the study period ran from March 2007 to March 2008. Patients who developed apparent skin rash anywhere on the body after NVP containing HAART and had to change their NVP-containing regime to efavirenz-containing ones were diagnosed as rash. There were 39 patients who matched these criteria. Most of these 39 patients developed rash within two months after NVP treatment (NVP-rash), and none of them showed liver toxicity. On the other hand, 248 patients showed reasonably good adherence to NVP and did not develop rash at all or developed only mild rash that could be controlled by steroid within the observation period. The remaining 385 patients were excluded because of treatment without NVP (184 cases), incomplete clinical records (101 cases), treatment interruptions (62 cases), adverse drug effects other than rash (18 cases), and incomplete HIV-1 suppression (20 cases). From the 248 NVP tolerant patients, we first tried to have two control patients for each rash case matching age, sex, and duration of therapy before NVP containing HAART. However, some rash cases have only one control mainly due to the limitation of available reagents. Total of 60 samples were thus selected for the control group (NVP-tolerant). Age, sex, treatment history and CD4 cell counts were not different between test and control groups as shown in Table 1. Two hundred μ l of whole blood was collected from each of those patients and kept at -20°C until DNA extraction with the QIAamp DNA Blood Mini Kit (QIAGEN, Hilden, Germany). All participants signed informed consent forms. The present study was approved by the institutional ethics committees of the Bamrasnaradura Infectious Diseases Institute and the Department of Diseases Control, Ministry of Public Health, Thailand.

HLA-C Typing

Medium-high resolution HLA-C typing was performed with an HLA-C typing kit (MPH-2 HLA-C typing kit, Wakunaga, Japan) according to the manufacturer's instructions. Any ambiguous results were checked by nucleotide sequence determination of PCR-amplified DNA fragments of HLA-C exons 2, and 3 [33]. GeneAmp[®] PCR system 9600 (Applied Biosystems, Foster City, CA) was used for all the PCR reactions and DNA sequencer 373 (Applied Biosystems) was used for determination of the nucleotide sequence of an amplified fragment.

HLA-DRB1*01 detection

We performed HLA class II DNA-based typing of DR1 as described in the 12th International Histocompatibility Working Group version 1.5. Amplified DNA with primer pair 2DRBAMP1 (5'-TTCTTGTGGCAGCTTAAGTT-3') and 2DRBAMP-B (5'-CCGCTGCACITGTGAAGCTCT-3') in exon 2 was treated with NaOH and the denatured DNA was loaded onto a nylon membrane manually using a

milliblot system with a vacuum manifold. UV light was used to crosslink the DNA to the membrane. For hybridization with DR1 specific probe, we used DRB 1001w (5'-TGGCAGCTTAAGTTGAA-3') digoxigenin-labeled SSOP for detection of HLA-DRB1*01 positive sample. After stringent wash procedure, the membrane was incubated with an antibody to digoxigenin coupled with alkaline phosphatase. Addition of a substrate for alkaline phosphatase caused light to be emitted by the Lumiphos. This light was detected by exposure of X-ray film.

HLA-C 5' SNP genotyping

The rs9264942 SNP genotyping was performed by TaqMan SNP genotyping system with ABI real time PCR 7300. A validated primer and probe mix (C_29901957_10) were purchased from Applied Biosystems.

Statistical analysis

Differences in age, duration of therapy before start of NVP containing HAART, and pre- and post-therapy CD4 cell counts between case and control groups were evaluated by Mann-Whitney U test. A difference in proportion of sex was evaluated by Chi square test. Differences in the allele frequencies between the two groups were evaluated by Fisher's exact test. P values less than 0.05 were considered to be statistically significant. The corrected P (Pc) values were calculated by using Bonferroni's correction.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SL conceived of the study, participated in the design and coordination of the study, and drafted the manuscript, TR carried out genotyping and analysis of clinical data, SF carried out genotyping and analysis of clinical data, SU participated in coordination of the study and helped to draft the manuscript. WP participated in collection of clinical data and helped to draft the manuscript, PT participated in coordination of the study and helped to draft the manuscript, EEN supervised genotyping, participated in study design and helped to draft the manuscript, TS participated in the design of the study, performed the statistical analysis, and helped to draft the manuscript. All authors read and approved the final manuscript.

Author's information

SL is a chief of Immunology and Virology Laboratory, Bamrasnaradura Infectious Diseases Institute, which is a governmental institute with the largest infectious disease hospital in Thailand. TR and SF are research assistants of the study. SU is a sub-chief of Immunology and Virology Laboratory and working on HIV-1 diagnosis. WP is a clinician who is taking care of HIV-1 infected patients. PT is a director of Bamrasnaradura Infectious Diseases Institute.

EEN is an assistant professor of Osaka University, Japan. TS is a professor of Osaka University working on HIV-1 infection and host genome.

Acknowledgements

This work was supported by grants from The Health Science Foundation, the Ministry of Health, Labour, and Welfare, and the Ministry of Education, Culture, Sports, Science, and Technology, Japan and Bamrasnaradura Infectious Diseases Institute, Department of Disease Control, Ministry of Public Health, Nonthaburi, Thailand. We thank all the HIV-infected individuals who participated in this study. We thank Dr. Achara Chaovanich, the former Director, and Dr. Boosun Chua-intra from Bamrasnaradura Infectious Diseases Institute, Nonthaburi for her support; Mr. Nopphanath Chumpathad for his part advice regarding the statistical analysis; and Dr. Komon Luangtrakool, Department of Transfusion Medicine, Faculty of Medicine Siriraj Hospital, Mahidol University, for his technical assistance of sequence based typing.

References

- Palella FJ Jr, Delaney KM, Moorman AC, Loveless MO, Fuhrer J, Sat-ten GA, Aschman DJ, Holmberg SD: **Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators.** *N Engl J Med* 1998, **338**:853-860.
- Manosuthi W, Chottanapand S, Thongyen S, Chaovanich A, Sungkanuparph S: **Survival rate and risk factors of mortality among HIV/tuberculosis-coinfected patients with and without antiretroviral therapy.** *J Acquir Immune Defic Syndr* 2006, **43**:42-46.
- Sabbatani S, Manfredi R, Biagetti C, Chiodo F: **Antiretroviral Therapy in the Real World: Population-Based Pharmacoeconomic Analysis of Administration of Anti-HIV Regimens to 990 Patients.** *Clin Drug Investig* 2005, **25**:527-535.
- Havir D, Cheeseman SH, McLaughlin M, Murphy R, Erice A, Spector SA, Greenough C, Sullivan JL, Hall D, Myers M, et al: **High-dose nevirapine: safety, pharmacokinetics, and antiviral effect in patients with human immunodeficiency virus infection.** *J Infect Dis* 1995, **171**:537-545.
- Anekthananon T, Ratanasuwon W, Techasathit W, Sonjai A, Suwanagool S: **Safety and efficacy of a simplified fixed-dose combination of stavudine, lamivudine and nevirapine (GPO-VIR) for the treatment of advanced HIV-infected patients: a 24-week study.** *J Med Assoc Thai* 2004, **87**:760-767.
- Stern JO, Robinson PA, Love J, Lanes S, Imperiale MS, Mayers DL: **A comprehensive hepatic safety analysis of nevirapine in different populations of HIV infected patients.** *J Acquir Immune Defic Syndr* 2003, **34**(Suppl 1):S21-33.
- Carr A, Cooper DA: **Adverse effects of antiretroviral therapy.** *Lancet* 2000, **356**:1423-1430.
- Kappelhoff BS, van Leth F, MacGregor TR, Lange J, Beijnen JH, Huitema AD: **Nevirapine and efavirenz pharmacokinetics and covariate analysis in the 2NN study.** *Antivir Ther* 2005, **10**:145-155.
- Gangar M, Arias G, O'Brien JG, Kemper CA: **Frequency of cutaneous reactions on challenge with nevirapine and delavirdine.** *Ann Pharmacother* 2000, **34**:839-842.
- Pollard RBRP, Dransfield K: **Safety profile of nevirapine, a non-nucleoside reverse transcriptase inhibitor for the treatment of human immunodeficiency virus infection.** *Clin Ther* 1998, **20**:1071-1092.
- Ho TT, Wong KH, Chan KC, Lee SS: **High incidence of nevirapine-associated rash in HIV-infected Chinese.** *Aids* 1998, **12**:2082-2083.
- Pirmohamed M, Park BK: **HIV and drug allergy.** *Curr Opin Allergy Clin Immunol* 2001, **1**:311-316.
- Martin AM, Nolan D, James I, Cameron P, Keller J, Moore C, Phillips E, Christiansen FT, Mallal S: **Predisposition to nevirapine hypersensitivity associated with HLA-DRB1*0101 and abrogated by low CD4 T-cell counts.** *Aids* 2005, **19**:97-99.
- Mallal S, Nolan D, Witt C, Masel G, Martin AM, Moore C, Sayer D, Castley A, Mamotte C, Maxwell D, James I, Christiansen FT: **Association between presence of HLA-B*5701, HLA-DR7, and HLA-DQ3 and hypersensitivity to HIV-1 reverse-transcriptase inhibitor abacavir.** *Lancet* 2002, **359**:727-732.
- Gatanaga H, Yazaki H, Tanuma J, Honda M, Genka I, Teruya K, Tachikawa N, Kikuchi Y, Oka S: **HLA-Cw8 primarily associated with hypersensitivity to nevirapine.** *Aids* 2007, **21**:264-265.
- Coopman SA, Johnson RA, Platt R, Stern RS: **Cutaneous disease and drug reactions in HIV infection.** *N Engl J Med* 1993, **328**:1670-1674.
- Leetrakool N, Kunachiwa W, Dettrairat S, Kohreanudom S: **Distribution of class I molecular HLA-A*, -B* and -Cw* in people living with HIV-1/AIDS in Chiang Mai province of Northern Thailand [Abstract].** Paper presented at: *Int Conf AIDS, 11-16 July, 2004; Bangkok.*
- Vitezica ZG, Milpied B, Lonjou C, Borot N, Ledger TN, Lefebvre A, Hovnanian A: **HLA-DRB1*01 associated with cutaneous hypersensitivity induced by nevirapine and efavirenz.** *Aids* 2008, **22**:540-541.
- Fellay J, Shianna KV, Ge D, Colombo S, Ledergerber B, Weale M, Zhang K, Gumbs C, Castagna A, Cossarizza A, Cozzi-Lepri A, De Luca A, Easterbrook P, Francioli P, Mallal S, Martinez-Picado J, Miro JM, Obel N, Smith JP, Wyniger J, Descombes P, Antonarakis SE, Letvin NL, McMichael AJ, Haynes BF, Telenti A, Goldstein DB: **A whole-genome association study of major determinants for host control of HIV-1.** *Science* 2007, **317**:944-947.
- Carrington M, Nelson GW, Martin MP, Kissner T, Vlahov D, Goedert JJ, Kaslow R, Buchbinder S, Hoots K, O'Brien SJ: **HLA and HIV-1: heterozygote advantage and B*35-Cw*04 disadvantage.** *Science* 1999, **283**:1748-1752.
- Kaslow RA, Carrington M, Apple R, Park L, Munoz A, Saah AJ, Goedert JJ, Winkler C, O'Brien SJ, Rinaldo C, Detels R, Blattner W, Phair J, Erlich H, Mann DL: **Influence of combinations of human major histocompatibility complex genes on the course of HIV-1 infection.** *Nat Med* 1996, **2**:405-411.
- Cruse JM, Brackin MN, Lewis RE, Meeks W, Nolan R, Brackin B: **HLA disease association and protection in HIV infection among African Americans and Caucasians.** *Pathobiology* 1991, **59**:324-328.
- Thio CL, Gao X, Goedert JJ, Vlahov D, Nelson KE, Hilgartner MW, O'Brien SJ, Karacki P, Astemborski J, Carrington M, Thomas DL: **HLA-Cw*04 and hepatitis C virus persistence.** *J Virol* 2002, **76**:4792-4797.
- Littera R, Carcassi C, Masala A, Piano P, Serra P, Ortu F, Corso N, Casula B, La Nasa G, Contu L, Manconi PE: **HLA-dependent hypersensitivity to nevirapine in Sardinian HIV patients.** *Aids* 2006, **20**:1621-1626.
- de Maat MM, ter Heine R, Mulder JW, Meenhorst PL, Mairuhu AT, van Gorp EC, Huitema AD, Beijnen JH: **Incidence and risk factors for nevirapine-associated rash.** *Eur J Clin Pharmacol* 2003, **59**:457-462.
- Ananworanich J, Moor Z, Siangphoe U, Chan J, Cardiello P, Duncombe C, Phanuphak P, Ruxrungtham K, Lange J, Cooper DA: **Incidence and risk factors for rash in Thai patients randomized to regimens with nevirapine, efavirenz or both drugs.** *Aids* 2005, **19**:185-192.
- Kiertiburanakul S, Sungkanuparph S, Charoenyingwattana A, Mahasirimongkol S, Sura T, Chantratita W: **Risk factors for nevirapine-associated rash among HIV-infected patients with low CD4 cell counts in resource-limited settings.** *Curr HIV Res* 2008, **6**:65-69.
- Antinori A, Baldini F, Girardi E, Cingolani A, Zaccarelli M, Di Giambenedetto S, Barracchini A, De Longis P, Murri R, Tozzi V, Ammassari A, Rizzo MG, Ippolito G, De Luca A: **Female sex and the use of anti-allergic agents increase the risk of developing cutaneous rash associated with nevirapine therapy.** *Aids* 2001, **15**:1579-1581.
- van Leth F, Andrews S, Grinsztejn B, Wilkins E, Lazanas MK, Lange JM, Montaner J: **The effect of baseline CD4 cell count and HIV-1 viral load on the efficacy and safety of nevirapine or efavirenz-based first-line HAART.** *Aids* 2005, **19**:463-471.
- Manosuthi W, Sungkanuparph S, Tansuphaswadikul S, Inthong Y, Prathitsirikul W, Chottanapund S, Mankatitham W, Chimsuntorn S, Sitibusaya C, Moolasart V, Chumpathat N, Termvise P, Chaovanich A: **Incidence and risk factors of nevirapine-associated skin rashes among HIV-infected patients with CD4 cell counts <250 cells/microL.** *Int J STD AIDS* 2007, **18**:782-786.

31. Chantarangsu S, Mushiroda T, Mahasirimongkol S, Kiertiburanakul S, Sungkanuparph S, Manosuthi W, Tantisiriwat W, Charoenyingwattana A, Sura T, Chantratita W, Nakamura Y: **HLA-B*3505 allele is a strong predictor for nevirapine-induced skin adverse drug reactions in HIV-infected Thai patients.** *Pharmacogenet Genomics* 2009, **19**:139-146.
32. Imanishi T, Akaza T, Kimura A, Tokunaga K, Gojobori T: **Allele and haplotype frequencies for HLA and complement loci in various ethnic groups.** Paper presented at: Eleventh International Histocompatibility Workshop and Conference; 6-13 November, 1991, Yokohama, Japan 1991.
33. Delfino L, Morabito A, Longo A, Ferrara GB: **HLA-C high resolution typing: analysis of exons 2 and 3 by sequence based typing and detection of polymorphisms in exons 1-5 by sequence specific primers.** *Tissue Antigens* 1998, **52**:251-259.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp






Contents lists available at ScienceDirect

Vaccine

journal homepage: www.elsevier.com/locate/vaccine



HIV-2 capsids distinguish high and low virus load patients in a West African community cohort

Clayton O. Onyango^a, Aleksandra Leligdowicz^{a,b}, Masaru Yokoyama^c, Hironori Sato^c, Haihan Song^d, Emi E. Nakayama^d, Tatsuo Shioda^{d,*}, Thushan de Silva^a, John Townsend^a, Assan Jaye^a, Hilton Whittle^a, Sarah Rowland-Jones^{a,b}, Matthew Cotten^{a,*}

^a Medical Research Council Laboratories, Fajara, Atlantic Road, P.O. Box 273, The Gambia

^b Weatherall Institute of Molecular Medicine, Medical Research Council Human Immunology Unit, John Radcliffe Hospital, Oxford, OX3 9DS, United Kingdom

^c Laboratory of Viral Genomics, Center for Pathogen Genomics, National Institute of Infectious Diseases, Tokyo 208-0011, Japan

^d Department of Viral Infections, Research Institute for Microbial Diseases, Osaka University, Osaka 565-0871, Japan

ARTICLE INFO

Article history:

Received 15 May 2009

Received in revised form 13 August 2009

Accepted 17 August 2009

Available online xxx

Keywords:

HIV-2

Capsid

TRIM5 α

ABSTRACT

HIV-2 causes AIDS similar to HIV-1, however a considerable proportion of HIV-2 infected patients show no disease and have low plasma virus load (VL). An analysis of HIV-2 capsid (p26) variation demonstrated that proline at p26 positions 119, 159 and 178 are more frequent in lower VL subjects while non-proline residues at all three sites are more frequent in subjects with high VL. *In vitro* replication levels of viruses bearing changes at the three sites suggested that these three residues influence virus replication by altering susceptibility to TRIM5 α . These results provide new insights into HIV-2 pathogenesis.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

HIV-2 was discovered in West Africa patients [1,2] shortly after the discovery of HIV-1, with HIV-2 entering humans via zoonoses distinct from the entry of HIV-1 some time in the early 20th century [3]. The two viruses have 60–80% sequence homology and have similar genomic organization, yet the viruses have distinct transmission rates and disease associations (reviewed in Ref. [4]). Although some HIV-2-infected patients progress to AIDS, the majority control infection [5–7] and patients with low VL survive longer [8]. Early descriptions of HIV-2 observed differences in the virus genetics [9] and noted that the HIV-2 epidemic behaved like a mixture of pathological and non-pathological viruses [5,10]. Studying and understanding HIV-2 represents a possibility to discover how a potentially lethal lentivirus infection can be controlled by humans.

There are limited data on HIV-2 sequence variation associated with either long-term control or progression to disease [9,11,12]. Although some correlation of Nef variation with disease has been observed [13], most HIV-2 sequences in the public databases are derived from viruses isolated from AIDS patients where high VLs facilitate virus isolation and sequencing which could introduce bias in the data. Sequences derived from patients with low VL are essential for understanding disease non-progression in HIV-2 infection. A community cohort to study HIV was established in Caio, Guinea Bissau in 1988 [50], reviewed in Ref. [4]. HIV screening of the entire adult population in the Caio community cohort have identified HIV-2-positive subjects with a wide spectrum of clinical symptoms and survival [4,10,14]; a sequence analysis of the virus among these patients could identify virus variations that influence outcome.

The capsid (CA) protein (p24 in HIV-1; p26 in HIV-2) has a structure that is conserved among retroviruses [15,16]. p26 accumulates during replication as a Gag polyprotein, assembles into spherical structures that package viral RNA and is subsequently processed by the viral protease and reassembled into the mature virus cores [16–18]. The CA has a distinct amino-terminal domain (residues 1–145), that is exposed on the surface of cores, and a C-terminal domain (residues 146–230) which is required for oligomerization [19]. The 20 amino acid major homology region (MHR) in the C-terminal domain is highly conserved; changes in this motif can interfere with CA assembly, maturation and early stages of infec-

* Corresponding authors.

E-mail addresses: conyango@mrc.gm (C.O. Onyango),

aleksandra.leligdowicz@mail.mcgill.ca (A. Leligdowicz), yokoyama@nih.go.jp

(M. Yokoyama), hirosato@nih.go.jp (H. Sato), haihansong@hotmail.com

(H. Song), emien@biken.osaka-u.ac.jp (E.E. Nakayama), shioda@biken.osaka-u.ac.jp

(T. Shioda), tdesilva@mrc.gm (T. de Silva), jtownend@mrc.gm

(J. Townsend), ajaye@mrc.gm (A. Jaye), hwhittle@mrc.gm (H. Whittle),

sarah.rowland-jones@ndm.ox.ac.uk (S. Rowland-Jones), mcotten@mrc.gm

(M. Cotten).

0264-410X/\$ – see front matter © 2009 Elsevier Ltd. All rights reserved.

doi:10.1016/j.vaccine.2009.08.060

Please cite this article in press as: Onyango CO, et al. HIV-2 capsids distinguish high and low virus load patients in a West African community cohort. Vaccine (2009), doi:10.1016/j.vaccine.2009.08.060

Table 1
Comparison of the cohort p26 residues 119, 159 and 178 to HIV-2_{ROD} p26; variation with viral load.

ID	Source	VL	119	159	178	ID	Source	VL	119	159	178
ROD			P	P	P	ROD			P	P	P
CO310	RNA	<100	.	.	.	CO309	RNA	283	Q	S	.
CO315	DNA	<100	.	.	.	CO316	RNA	372	.	.	.
CO318	PRO	<100	.	.	.	CO364	RNA	387	A	.	.
CO319	RNA	<100	.	.	.	CO314	DNA	393	.	S	.
CO324	RNA	<100	.	.	.	CO354	DNA	413	.	.	.
CO326	RNA	<100	.	.	.	CO323	RNA	497	A	.	.
CO327	DNA	<100	.	.	.	CO365	RNA	523	A	S	A
CO339	RNA	<100	.	.	.	CO350	RNA	540	A	S	A
CO349	RNA	<100	.	.	.	CO322	RNA	610	.	S	A
CO359	RNA	<100	.	.	.	CO355	RNA	651	A	.	A
CO360	RNA	<100	.	.	.	CO357	RNA	813	A	S	Q
CO344	DNA	<100	A	.	.	CO332	RNA	1085	A	.	V
CO346	DNA	<100	A	.	.	CO305	RNA	1343	A	.	.
CO351	RNA	<100	A	.	.	CO340	RNA	1587	.	.	.
CO308	RNA	<100	Q	.	.	CO321	RNA	1608	.	.	.
CO337	DNA	<100	.	T	.	CO342	RNA	1907	A	S	.
CO361	RNA	<100	A	S	.	CO325	RNA	1999	.	.	.
CO366	RNA	<100	A	S	.	CO330	RNA	2653	A	S	.
CO367	DNA	<100	A	S	.	CO331	RNA	2949	A	S	A
CO363	DNA	<100	Q	S	.	CO320	RNA	3241	A	.	.
CO335	RNA	<100	A	.	A	CO307	RNA	3764	G	S	.
CO336	RNA	<100	A	.	A	CO368	RNA	6431	A	T	S
CO341	RNA	<100	A	.	A	CO345	RNA	9659	A	.	.
CO362	DNA	<100	A	.	A	CO312	RNA	9979	A	S	A
CO301	DNA	<100	A	S	.	CO334	RNA	10752	.	S	.
CO306	DNA	<100	A	S	A	CO311	RNA	14104	A	S	A
CO333	RNA	<100	A	S	A	CO313	RNA	17067	A	S	A
CO302	RNA	109	.	.	.	CO369	RNA	22446	Q	S	.
CO304	DNA	114	.	.	.	CO343	RNA	25836	A	S	.
CO303	RNA	154	A	S	A	CO338	RNA	28581	Q	S	A
CO356	DNA	182	.	.	.	CO317	RNA	37503	A	S	A
CO328	RNA	198	A	.	.	CO347	RNA	146284	A	.	.
CO353	DNA	234	.	.	.	CO348	RNA	148593	A	.	A
CO329	RNA	267	A	S	S	CO358	RNA	283542	A	S	.
CO352	RNA	275	.	S	.						

Samples from 69 HIV-2 singly-infected patients collected in 2006 were used for sequencing (see Section 2). Patient plasma VL was assayed by RT-PCR with a lower limit of detection of 100 virus copies/ml. For statistical analysis, undetectable viremia was given a value of 50 copies/ml. Viral loads (copies/ml) are indicated in second column. The source of template for the PCR reaction (circulation RNA (RNA) or proviral DNA (DNA)) is indicated. The reference sequence derived from HIV-2_{ROD} p26 is listed in the first row, in the subsequent rows identity with the HIV-2_{ROD} p26 sequence is indicated with a period (.).

tion [19,20]. The multiple functions impose strict constraints on the amino acid changes allowed in CA. Studying the limited variations that do occur in p26 can provide important information on both protein function and the selective forces acting on HIV-2. In this study, we have tested the hypothesis that HIV-2 p26 capsid variants modulate HIV-2 viral load.

2. Methods

2.1. Patient cohort.

Study subjects were recruited from the community-based cohort in Caio, Guinea Bissau (study subjects described in Ref. [14]). Briefly, HIV screening was performed with a Murex ICE HIV-1.2.0 immunoassay (Murex Diagnostics), confirmed and differentially diagnosed (HIV-1/HIV-2) using HEXAGON HIV (Human GmbH). Dually HIV-1+2 positive samples were subjected to peptide-based assays (Pepti-Lav 1-2, Sanofi Diagnostics Pasteur) and HIV-1 and HIV-2-specific polymerase chain reaction (PCR) with primers targeted to the long terminal repeats [7,8]. HIV-1+2 infected patients were excluded from the study. HIV-2 plasma viral load was quantified by PCR based method [7,21] with a lower limit of detection of 100 copies/ml, for analytic purposes, undetectable samples were assigned a value of 50 copies/ml. All participants were antiretroviral naïve at the time of the study and provided informed consent. The study was approved by the Gambian Government/MRC Ethics Committee, The Republic of Guinea Bissau Ministry of Health,

and the Oxford Tropical Research Ethics Committee (OXTREC), UK.

2.2. p26 gene amplification strategy, optimization of primers.

p26 gene amplification primers were designed by selecting conserved sequences flanking the p26 coding region from all HIV-2 isolates in the Los Alamos HIV Database to be used as targets for PCR primers. Coupled reverse transcriptase and nested PCR reactions were performed with plasma-isolated virus RNA as template. The following primer sequences in various combinations were used for amplification:

MO017(OF, outer forward) GTCTGCGTCATTTGGTGCAT
 MO018(IF, inner forward) CTGAGAGAAAATGCCAAGCA
 MO019(OR outer reverse) GGGCAGTTTGTATGATGTGTCC
 MO020(IR, inner reverse) GCCCTTCCTTCCACAGTTCCA
 MO021(IR) GCCCTTCCTTCCACAAATCCA
 MO030(OF)CACGCAGAAGAGAAAGTGAAAG
 MO031(OR) CGGGGAAGTTGCGRGGCTT
 MO032(IF) AGTAGACCAACAGCACCACC
 MO036(OF) GTGGGCAGCGAATGAATTGG
 MO037(OF) GTGGGCAGCGAACGAATTGG
 MO038(OR) AAAGAGAGAATTGAGGTGCAGCA.

In case none of the primer combinations amplified a product from RNA, amplification of provirus was performed with the same

series of primers using genomic DNA from the patients' peripheral blood mononuclear cells (PBMCs) as template. The origin of the sequence (viral RNA or proviral DNA) is indicated in Table 1. PCR products were excised from preparative gels, purified by QiaQuick columns, sequenced using the inner PCR primers from both directions and aligned using ClustalW; ambiguities were resolved by direct analysis of sequencing scans using a sequence alignment editor BioEdit [22].

The p26 sequences were monitored to rule out cross-contamination during sample handling or PCR. Contamination among different samples would generate multiple sequences with 100% identity. A phylogenetic analysis showed no identical sequences, with the closest pairs (CO336 and CO335) showing 0.58% difference (4 differences in 690 nucleotides). BLAST analyses revealed no homology greater than 94% with any sequence in the GenBank database (results not shown). All the p26 sequences were consistent with HIV-2 subtype A infections.

2.3. HIV-2 envelope gene amplification

Envelope sequences were obtained from 34 HIV-2 isolates selected randomly from the set of 69 Caio viruses (see below). The following PCR primers targeted to conserved envelope flanking sequences were used:

MO080 (outer forward) CAGTCATCACAGAGTCATGTG
 MO076 (outer reverse) TCCTTGTGGATAYGAYCTGT
 MO072 (inner forward) TCATGTGAYAAGCACTATTGGGA
 MO077 (inner reverse) GGAAGAGAAAACAGGCCTATAGCC.

An approximately 1500 bp fragment (HIV-2_{ROD} C2 to gp41, positions 6783–8285) was amplified in a nested RT-PCR from plasma virus. PCR products were purified as described above, sequenced using inner PCR primers from both directions. If the initial four primers failed, these additional primers were used:

MO122a GTGGACTAACTGCAGAGGAGAATT
 MO125 AGTTCTGCCACCTCTGCACT
 MO125a AGAAAACCAAGAACCCTAGCAC.

A fragment of approximately 1300 bp corresponding to HIV-2_{ROD} positions 6849–8210 was used for the phylogenetic analysis. The p26 and Env sequences described in this manuscript were submitted to GenBank (accession numbers GQ485448–GQ485550).

2.4. Molecular modelling

Three-dimensional (3-D) models of HIV-2 CAs were constructed by the homology-modelling technique using the Molecular Operating Environment (MOE) (Chemical Computing Group Inc., Quebec, Canada) as described in Refs. [23–26]. The two crystal structures of the HIV-1 CA proteins were used as templates for the modelling; a CA monomer at a resolution of 3.00 Å (Protein Data Bank (PDB) code: 1E6J [27]) and the dimer of CA C-terminal domain at a resolution of 1.70 Å (PDB code: 1A80 [19]). The amino acid sequence identity of HIV-1 (1E6J) and HIV-2 CA (CO310 in this study) is about 70.5%. The sequence similarity is sufficient to construct a structural model with an r.m.s. deviation of approximately 1.5 Å for the main chain between the predicted and actual structures [28]. The 3-D structures were optimized thermodynamically by energy minimization using MOE and an AMBER99 force field [29] and further refined the physically unacceptable local structures on the basis of evaluation of unusual dihedral angles, ψ and ϕ , by the Ramachandran plot using MOE.

The binding energies of the p26 dimer models, E_{bind} , were calculated as described elsewhere [25] [30], using the formula

$E_{\text{bind}} = E_{\text{dimer}} - 2E_{\text{monomer}}$, where E_{dimer} is the energy of the p26 dimer; E_{monomer} is the energy of the p26 monomer. Spearman's rank correlation coefficient for viral load and capsid binding energy and its statistical evaluation were calculated by SPSS ver 14 (SPSS Inc, Chicago).

2.5. Expression of TRIM5 α

Construction of recombinant Sendai viruses (SeVs) carrying human MT4-TRIM5 α -tag (Hu-TRIM5 α -Sev) and cynomolgus monkey TRIM5 α lacking the SPRY domain (CM-SPRY(-)-Sev) was described previously [23].

2.6. Construction of HIV-2 GH123 capsid variants

Mutant DNA constructs of infectious molecular clones of HIV-2 GH123 carrying alanine 119 (GH123/119A) was described previously [23]. Note that because of an insertion in the GH123 p26 protein, positions 120, 160 and 179 from Ref. [23] correspond to positions 119, 159 and 178 in this work. PCR-based mutagenesis with primer pairs P1, P2 and P3, P4 was used to generate mutant GH123 carrying alanine 178 (GH123/178A). Mutant GH123 carrying serine 159 and 178A (GH123/159S-178A) was then constructed using primer pairs P1, P5 and P4, P6 with GH123/178A as a template. Mutant GH123 carrying 119A, 159S and 178A (GH123/119A-159S-178A) was constructed with primer pairs P1, P5 and P4, P6; and both GH123/119A and GH123/178A as templates. Infectious viruses were prepared by transfection of 293T cells with mutant proviral DNA clones.

P1:CTTCCCTGTACAACAGACA, P2:TTTACTGCTGCATCTGTTGTTCTG,
 P3:CAGAACAACAGATGCAGCAGTAAA,
 P4:GTGCAGCAAGTCCTCTGTC
 P5:TAGCTCTGGAATGATTCCTTTGGTCC,
 P6:GGACCAAAGGAATCATTCCAGAGCTA

3. Results

3.1. p26 CA variation in the Caio cohort.

HIV-2 positive subjects were selected from the Caio community cohort to assess HIV-2 sequence variation among a wide range of VL (≤ 100 to $>280,000$ copies/ml). PCR was attempted from a total of 92 subjects and p26 sequences were obtained from 69 subjects; 53 sequences were derived from viral RNA and 16 were from proviral DNA. Additional clinical features of these patients were previously described [14].

Caio p26 sequences were compared to either the Caio 69 p26 consensus sequence (Fig. 1 upper) or to the reference strain HIV-2_{ROD} p26 [31] (Fig. 1 lower) to identify amino acid polymorphisms in p26. HIV-2_{ROD} was chosen as a reference because the virus was isolated early in the epidemic in Cape Verde, islands historically linked to Guinea Bissau. Both comparisons show a similar pattern with variant amino acid sites clustering outside of the alpha helices (marked in black) required for CA folding (Fig. 1). Variations that associate with high or low VL (Mann–Whitney test, $p < 0.05$, except for position 178) are highlighted with a dotted line. The polymorphisms at position 119, 159 and 178 were of special interest because they showed 35–70% variation from HIV-2_{ROD} and involved changes from the reference sequence proline, a change expected to strongly alter protein structure. The non-proline residues were either alanine (38 cases), glutamine (5), or glycine (1) at position 119, serine (29) or threonine (2) at 159, and alanine (18), serine (2), glutamine (1), or valine (1) at 178. Viruses in low VL patients often had proline at these three positions while in higher VL samples these sites were frequently occupied by