

LLC-MKII, naïve CRFK and TRIM5 $\alpha$ -expressing CRFK cells, plated at a density of  $4 \times 10^4$  on the day before infection. On day 3 post-infection, cells were collected by trypsin treatment and used for FACS analysis.

### 2.7. Intervention of interaction between CypA and HIV-1 CA

To determine the effect of intervention of CypA–CA interaction on viral single-cycle infectivity, CsA (SIGMA–ALDRICH) was added to cell cultures 4 h before virus inoculation at a final concentration of 5  $\mu$ M, and the concentration was maintained throughout the experiment. For gene-knockdown of CypA, cells were transfected with siRNA against CypA (siCypA, 5'-GUGA-CUUCACACHCCAUAUG-3'). As a negative control for the siRNA, siPerfect<sup>®</sup> Negative control (SIGMA–ALDRICH), well-characterized non-target control siRNA, was used. For siRNA transfection, we used the Lipofectamine<sup>™</sup> RNAi MAX (Invitrogen). Expression of CypA in cells was monitored by immunoblotting with an anti-CypA antibody (BIOMOL).

## 3. Results

### 3.1. Construction of CypA-binding loop mutants from NL-DT5R

Cellular CypA binds to CA in cells infected with HIV-1, and are incorporated into progeny virions. CypA-binding loop in HIV-1 CA is located at aa residues 85–93 (Fig. 1A), and quite variable. Moreover, CypA does not bind to CA proteins of SIVmac and SIVagm. To determine aa residues in CypA-binding loop of HIV-1 important for species-tropism, we constructed a series of site-specific mutants in this region of NL-DT5R (Fig. 1A). We firstly evaluated CypA content of mutant virions. Various proviral clones were transfected into 293T cells, and on day 2 post-transfection, virions produced were partially purified and concentrated by ultra-centrifugation to remove free CypA as previously described [25]. Upon transfection, all the clones produced progeny virions similarly, and no abnormality was observed in the Gag processing within virions (data not shown). Virions thus obtained were examined for CypA by immunoblotting analysis. As shown in Fig. 1B, while virions of DT5R/4-3 carrying the CypA-binding sequence of HIV-1 NL4-3 [27] contained a comparable level of CypA with those of NL4-3, NL-DT5R carrying the corresponding region of SIVmac did not at all. In addition, no incorporation of CypA into virions of DT5R/agm was observed as described above. These results indicated that sequence within the proline-rich region (9 aa) is necessary and sufficient for encapsidation of CypA, and that NL-DT5R is indistinguishable from SIVmac in this aspect. Based on mutant phenotypes in Fig. 1B, overall structure of the region, rather than particular aa, appeared to be important for virion-incorporation of CypA. In this regard, the results for SV1, SV2 and SV3 were particularly striking. They clearly showed that the insertion of two amino acids into the SIV sequence converts the binding phenotype. It can be concluded that




Fig. 1. Mutants of HIV-1 CypA-binding region in this study. (A) Aa sequence alignment of Gag-CA CypA-binding region (aa nos. 85–93 for HIV-1 NL4-3 clone [27]) of various clones. DT5R/4-3, DT5R/BaL, DT5R/agm, and NL-DT5R have sequences of HIV-1 NL4-3, HIV-1 BaL, SIVagm and SIVmac MA239, respectively, as indicated in parentheses. (B) Immunoblot analysis of CypA in virions. Virions produced from transfected 293T cells were prepared as described in Section 2, and analyzed for CypA and Gag-CA p24 as indicated. As controls, lysates of cell and ultra-centrifuged precipitates (pUC) prepared from 293T cells transfected with pUC19 were used. The experiments were repeated three times with similar results, and the representative data are shown here.

amino acids neighboring proline 90 are also important for CypA–CA binding and that the entire structure of the loop is critical. In summary, our mutants contained various levels of CypA in their virions (from negative to an NL4-3 level). While virions of DT5R/4-3, /H87Q, /H87Q/A88P and /SV3 had the NL4-3 level of CypA, those of NL-DT5R, DT5R/agm, /G89A, /SV1, /SV2, /SV4, /SV5, /SV6, /SV7 and /SV8 contained no detectable CypA. Mutants DT5R/BaL, /A88P and /P90A carried a low level of CypA in their virions.

### 3.2. Infectivity of CypA-binding site mutants

To know whether the CypA-binding inability of the mutants is correlated to or reflected in their infectivity in simian cells, viral growth kinetics was determined for all the clones in Fig. 1. Cell-free viruses were prepared from 293T cells transfected with various proviral mutant clones, and inoculated into CyM HSC-F cells using SIVmac239, NL-DT5R and NL-DT5R/4-3 as controls for viral growth rate. As shown in Fig. 2, SIVmac239 and NL-DT5R readily established a spreading infection, which peaked on days 6 and 18, respectively. In contrast, the infection by NL-DT5R/4-3 having the CypA-binding




Fig. 2. Growth kinetics in CyM HSC-F cells of CypA-binding loop mutants. Cell-free viruses were prepared from transfected 293T cells, and equal RT units were inoculated into HSC-F cells. RT activity in the culture supernatants was monitored at intervals as indicated. Results were shown separately (A–D) for clarity.

sequence of NL4-3 did not reach a plateau until day 33 post-infection. All test viruses examined productively infected HSC-F cells but with different kinetics. Whereas viral clones DT5R/agm, /SV5, and /SV8 grew similarly well with NL-DT5R, viruses DT5R/A88P, /G89A, /P90A, and /SV3 propagated similarly or even more poorly than DT5R/4-3. The other viral clones grew better than DT5R/4-3 but not NL-DT5R. Considering the strong binding activity to CypA (Fig. 1B), the results for DT5R/H87Q and DT5R/H87Q/A88P were noteworthy. The His87 could be an important determinant on viral infectivity in simian cells. In total, the CypA level in virions and the viral infectivity in simian cells did not necessarily show a negative correlation. We also determined a single-cycle infectivity of the mutants in rhesus monkey LLC-MKII cells to compare their growth rate using human 293T cells as control. GFP-viruses pseudotyped with VSV-G were prepared from 293T cells co-transfected with *env*-minus proviral reporter constructs and pCMV-G, and inoculated into 293T and LLC-MKII cells. On day 3 post-infection, cells were analyzed for GFP expression by FACS. As shown in Fig. 3, in control 293T cells, DT5R/4-3 showed the highest infectivity, and DT5R/agm and /H87Q were the second best. In contrast, in LLC-MKII cells, DT5R/4-3 showed the lowest infectivity except for DT5R/SV3 in good agreement with the results in Fig. 1. Also as expected, there were no viral clones that are significantly more infectious for LLC-MKII cells than NL-DT5R. Interestingly, DT5R/H87Q was highly infectious for both cell lines.

### 3.3. Susceptibility of CypA-binding site mutants to species-specific TRIM5 $\alpha$

Previous studies have shown a functional link between CypA and anti-retroviral factor TRIM5 $\alpha$  to inhibit HIV-1 replication in simian cells [3–5]. We therefore determined single-cycle infectivity of the CypA-binding loop mutants in

CRFK cells with/without human or CyM TRIM5 $\alpha$  (Fig. 4). Cell-free GFP-virus samples were prepared from transfected 293T cells as above, and inoculated into naïve and stably transfected CRFK cells. In human TRIM5 $\alpha$ -expressing CRFK cells, DT5R/4-3 and /H87Q showed an infectivity similar to that in naïve cells. In contrast, in CyM TRIM5 $\alpha$ -expressing CRFK cells, the infectivity of all test viruses was severely suppressed relative to that in naïve CRFK cells. These results strongly suggested that, although some mutants did overcome anti-viral action of human TRIM5 $\alpha$ , none of the variants examined here effectively countered the CyM TRIM5 $\alpha$ . Sakuma et al. reported that rhesus monkey TRIM5 $\alpha$  suppresses HIV-1 production through degradation of HIV-1 Gag [28]. We monitored a virus production level in 293T cells co-transfected with HIV-1 NL4-3 [27] or NL-DT5R and increasing amounts of expression vectors for human or CyM TRIM5 $\alpha$ . No clear difference was observed (data not shown).

### 3.4. Effect of CypA on viral replication

To further analyze the cell-dependent activity of CypA, we determined viral infectivity in various cell lines in the presence or absence of CsA. For this experiment, NL-DT5R, DT5R/4-3, /H87Q, /G89A and /SV3 with distinct characteristics (Figs. 1–4) were selected. As shown in Fig. 5, the effect of CypA on viral infectivity was very different by a combination of virus clones and target cells. In human 293T cells and feline CRFK cells expressing human TRIM5 $\alpha$ , by treating with CsA, infectivity of DT5R/4-3 decreased and that of NL-DT5R increased. In simian LLC-MKII cells and CRFK cells expressing CyM TRIM5 $\alpha$ , infectivity of DT5R/4-3 increased by the treatment. Almost no effects of CsA was noted for DT5R/87Q in 293T and LLC-MKII cells, but it behaved similarly with NL-DT5R and DT5R/4-3 in CyM TRIM5 $\alpha$ -CRFK cells. Both DT5R/G89A and /SV3 exhibited a low




Fig. 3. Single-cycle infectivity in human and simian cells of CypA-binding loop mutants. VSV-pseudotyped viruses were prepared from transfected 293T cells, and an equal amount of viruses as determined by RT assay was inoculated into human 293T and rhesus monkey LLC-MKII cells as described in Section 2. On day 3 post-infection, cells were assayed for GFP by a flow cytometer. The assays were repeated at least three times, and the mean values are presented.

infectivity relative to the other clones and were poor responders to CsA, suggesting the binding of CA to CypA is not an only important determinant for CypA-mediated control of HIV-1 replication. To verify the results obtained by the CsA treatment, we carried out CypA-specific siRNA experiments in 293T and LLC-MKII cells. Cell-free GFP-virus samples were prepared from transfected 293T cells as above, and inoculated into 293T and LLC-MKII cells which had been transfected with CypA-specific or control siRNA. CypA-siRNA

effectively reduced CypA expression in both cell lines as monitored by immunoblotting (approximately 80% reduction). As shown in Fig. 6, results were generally in good agreement with those in the CsA experiment. Of note, the CypA-knockdown significantly enhanced infectivity of all virus clones in LLC-MKII cells. In particular, DT5R/4-3 restored its infectivity up to the NL-DT5R level. We also noticed that the data for DT5R/4-3 in Figs. 5 and 6 were different. This difference may be related to the experimental methods used.




Fig. 4. Single-cycle infectivity in TRIM5 $\alpha$ -expressing feline cells of CypA-binding loop mutants. VSV-pseudotyped viruses were prepared from transfected 293T cells, and an equal amount of viruses as determined by RT assay was inoculated into naïve and human (Hu) or CyM TRIM5 $\alpha$ -expressing CRFK cells as described in Section 2. On day 3 post-infection, cells were assayed for GFP by a flow cytometer. The assays were repeated at least three times, and the mean values are presented.




Fig. 5. Effect of CsA treatment on viral infectivity in human, simian and TRIM5 $\alpha$ -expressing feline cells. Target cells were cultured in the presence of CsA (diluted in EtOH) from 4 h before infection to the end of the experiment as indicated. VSV-pseudotyped viruses were prepared from transfected 293T cells as described in Section 2, and an equal amount of viruses determined by RT assay was inoculated into various cell lines as indicated. On day 3 post-infection, cells were assayed for GFP by a flow cytometer. The assays were repeated at least three times, and the mean and standard error are presented. Statistical significance relative to non-treated samples as calculated by *t*-test is shown (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ).

CsA competitively inhibits the CypA–CA binding, while siRNA suppresses the expression level of CypA.

#### 4. Discussion

In this report, we have generated a series of CypA-binding loop mutants of HIV-1 CA from a prototype mt HIV-1 designated NL-DT5R [8] to improve viral replication in simian cells. Although they displayed various phenotypes with respect to the binding ability to CypA (Fig. 1) and viral infectivity at both single- and multi-cycle infection levels (Figs. 2 and 3), none of them were found to grow significantly better than NL-DT5R in simian cells. While the alterations introduced did affect viral

infectivity oppositely in human and simian cells as expected from published results [3,4,8,15–18], they were demonstrated not to antagonize the restriction in simian cells conferred by some factor(s) other than CypA (Figs. 4–6). Coupled with previous results by us and others [8,9], these data have strongly suggested that evasion from CypA- and APOBEC-mediated restrictions is still insufficient for HIV-1 to completely overcome the species barrier. In this regard, we noticed that NL-DT5R did not escape from the restriction by CyM TRIM5 $\alpha$  (Fig. 4) and that DT5R/4-3 restored its infectivity up to the NL-DT5R level in simian cells by CypA-knockdown (Fig. 6). Therefore, it is quite reasonable to assume that there is a region in CA independent of CypA-binding site responsible for the




Fig. 6. Effect of CypA-knockdown on viral infectivity in human and simian cells. Target cells were transfected with CypA-specific or control siRNA 30 h before infection. VSV-pseudotyped viruses were prepared from transfected 293T cells as described in Section 2, and an equal amount of viruses determined by RT assay was inoculated into target human 293T and rhesus monkey LLC-MKII cells as indicated. On day 3 post-infection, cells were assayed for GFP by a flow cytometer. The assays were repeated at least three times, and the mean and standard error are presented. Statistical significance relative to siControl samples as calculated by *t*-test is shown (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ).

TRIM5 $\alpha$ -mediated restriction of HIV-1 in simian cells. Experiments to identify the region are in progress in our laboratory to improve the growth ability of NL-DT5R in simian cells.

The underlying basis for CypA-mediated control of HIV/SIV replication is presently unknown, although it is clear that CypA promotes the replication of viruses with HIV-1 type CA in human cells but suppresses the viruses in simian cells as described above. The species-dependent regulation needs to be clarified at a molecular level. This issue is also intriguing from an evolutionary point of view. Almost all primate immunodeficiency viruses such as SIVmac and SIVagm do not require CypA for their replication whereas HIV-1 and SIVcpz do [29,30]. Our results described here showed that subtle changes in the CypA region can drastically alter viral phenotypes. It is thus interesting to analyze various primate immunodeficiency viruses based on sequence and/or structure of the CypA region.

#### Acknowledgements

We thank Ms. Kazuko Yoshida for editorial assistance. We also thank the members of our department for their helpful discussion. This work was supported in part by a Grant-in-Aid for Scientific Research on Priority Areas (19041051) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to A. A.), and by a Health Sciences Research Grant (Research on HIV/AIDS 2007–2009) from the Ministry of Health, Labour and Welfare of Japan (to A. A.).

#### References

- [1] M. Nomaguchi, N. Doi, K. Kamada, A. Adachi, Species barrier of HIV-1 and its jumping by virus engineering, *Rev. Med. Virol.* 18 (2008) 261–275.
- [2] R. Mariani, D. Chen, B. Schrofelbauer, F. Navarro, R. Koenig, B. Bollman, C. Munk, H. Nymark-McMahon, N.R. Landau, Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif, *Cell* 114 (2003) 21–31.
- [3] L. Berthou, S. Sebastian, E. Sokolskaja, J. Luban, Cyclophilin A is required for TRIM5 $\alpha$ -mediated resistance to HIV-1 in Old World monkey cells, *Proc. Natl. Acad. Sci. USA* 102 (2005) 14849–14853.
- [4] Z. Keckesova, L.M. Ylinen, G.J. Towers, Cyclophilin A renders human immunodeficiency virus type 1 sensitive to Old World monkey but not human TRIM5 $\alpha$  antiviral activity, *J. Virol.* 80 (2006) 4683–4690.
- [5] M. Stremlau, B. Song, H. Javanbakht, M. Perron, J. Sodroski, Cyclophilin A: an auxiliary but not necessary cofactor for TRIM5 $\alpha$  restriction of HIV-1, *Virology* 351 (2006) 112–120.
- [6] G.J. Towers, The control of viral infection by tripartite motif proteins and cyclophilin A, *Retrovirology* 4 (2007) 40.
- [7] M. Stremlau, C.M. Owens, M.J. Perron, M. Kiessling, P. Autissier, J. Sodroski, The cytoplasmic body component TRIM5 $\alpha$  restricts HIV-1 infection in Old World monkeys, *Nature* 427 (2004) 848–853.
- [8] K. Kamada, T. Igarashi, M.A. Martin, B. Khamsri, K. Hachio, T. Yamashita, M. Fujita, T. Uchiyama, A. Adachi, Generation of HIV-1 derivatives that productively infect macaque monkey lymphoid cells, *Proc. Natl. Acad. Sci. USA* 103 (2006) 16959–16964.
- [9] T. Hatzioannou, M. Princiotto, M. Piatak Jr., F. Yuan, F. Zhang, J.D. Lifson, P.D. Bieniasz, Generation of simian-tropic HIV-1 by restriction factor evasion, *Science* 314 (2006) 95.
- [10] D.A. Bosco, E.Z. Eisenmesser, S. Pochapsky, W.I. Sundquist, D. Kern, Catalysis of cis/trans isomerization in native HIV-1 capsid by human cyclophilin A, *Proc. Natl. Acad. Sci. USA* 99 (2002) 5247–5252.
- [11] B.R. Howard, F.F. Vajdos, S. Li, W.I. Sundquist, C.P. Hill, Structural insights into the catalytic mechanism of cyclophilin A, *Nat. Struct. Biol.* 10 (2003) 475–481.
- [12] J. Luban, K.L. Bossolt, E.K. Franke, G.V. Kalpana, S.P. Goff, Human immunodeficiency virus type 1 Gag protein binds to cyclophilins A and B, *Cell* 73 (1993) 1067–1078.
- [13] E.K. Franke, H.E. Yuan, J. Luban, Specific incorporation of cyclophilin A into HIV-1 virions, *Nature* 372 (1994) 359–362.
- [14] M. Thali, A. Bukovsky, E. Kondo, B. Rosenwirth, C.T. Walsh, J. Sodroski, H.G. Gottlinger, Functional association of cyclophilin A with HIV-1 virions, *Nature* 372 (1994) 363–365.
- [15] D. Braaten, E.K. Franke, J. Luban, Cyclophilin A is required for an early step in the life cycle of human immunodeficiency virus type 1 before the initiation of reverse transcription, *J. Virol.* 70 (1996) 3551–3560.
- [16] A.A. Bukovsky, A. Weimann, M.A. Accola, H.G. Gottlinger, Transfer of the HIV-1 cyclophilin-binding site to simian immunodeficiency virus from *Macaca mulatta* can confer both cyclosporin sensitivity and cyclosporin dependence, *Proc. Natl. Acad. Sci. USA* 94 (1997) 10943–10948.
- [17] B. Ackerson, O. Rey, J. Canon, P. Krogstad, Cells with high cyclophilin A content support replication of human immunodeficiency virus type 1

- Gag mutants with decreased ability to incorporate cyclophilin A, *J. Virol.* 72 (1998) 303–308.
- [18] D. Braaten, J. Luban, Cyclophilin A regulates HIV-1 infectivity, as demonstrated by gene targeting in human T cells, *EMBO J.* 20 (2001) 1300–1309.
- [19] M. Stremlau, M. Perron, M. Lee, Y. Li, B. Song, H. Javanbakht, F. Diaz-Griffero, D.J. Anderson, W.I. Sundquist, J. Sodroski, Specific recognition and accelerated uncoating of retroviral capsids by the TRIM5alpha restriction factor, *Proc. Natl. Acad. Sci. USA* 103 (2006) 5514–5519.
- [20] J.L. Anderson, E.M. Campbell, X. Wu, N. Vandegraaff, A. Engelman, T.J. Hope, Proteasome inhibition reveals that a functional preintegration complex intermediate can be generated during restriction by diverse TRIM5 proteins, *J. Virol.* 80 (2006) 9754–9760.
- [21] F. Diaz-Griffero, X. Li, H. Javanbakht, B. Song, S. Welikala, M. Stremlau, J. Sodroski, Rapid turnover and polyubiquitylation of the retroviral restriction factor TRIM5, *Virology* 349 (2006) 300–315.
- [22] X. Wu, J.L. Anderson, E.M. Campbell, A.M. Joseph, T.J. Hope, Proteasome inhibitors uncouple rhesus TRIM5alpha restriction of HIV-1 reverse transcription and infection, *Proc. Natl. Acad. Sci. USA* 103 (2006) 7465–7470.
- [23] M.J. Pineda, B.R. Orton, J. Overbaugh, A TRIM5alpha-independent post-entry restriction to HIV-1 infection of macaque cells that is dependent on the path of entry, *Virology* 363 (2007) 310–318.
- [24] R.L. Willey, D.H. Smith, L.A. Lasky, T.S. Theodore, P.L. Earl, B. Moss, D.J. Capon, M.A. Martin, In vitro mutagenesis identifies a region within the envelope gene of the human immunodeficiency virus that is critical for infectivity, *J. Virol.* 62 (1988) 139–147.
- [25] H. Akari, T. Fukumori, A. Adachi, Cell-dependent requirement of human immunodeficiency virus type 1 gp41 cytoplasmic tail for Env incorporation into virions, *J. Virol.* 74 (2000) 4891–4893.
- [26] J.K. Yee, A. Miyanohara, P. LaPorte, K. Bouic, J.C. Burns, T. Friedmann, A general method for the generation of high-titer, pantropic retroviral vectors: highly efficient infection of primary hepatocytes, *Proc. Natl. Acad. Sci. USA* 91 (1994) 9564–9568.
- [27] A. Adachi, H.E. Gendelman, S. Koenig, T. Folks, R. Willey, A. Rabson, M.A. Martin, Production of acquired immunodeficiency syndrome-associated retrovirus in human and nonhuman cells transfected with an infectious molecular clone, *J. Virol.* 59 (1986) 284–291.
- [28] R. Sakuma, J.A. Noser, S. Ohmine, Y. Ikeda, Rhesus monkey TRIM5alpha restricts HIV-1 production through rapid degradation of viral Gag polyproteins, *Nat. Med.* 13 (2007) 631–635.
- [29] J.L. Heeney, A.G. Dalgleish, R.A. Weiss, Origins of HIV and the evolution of resistance to AIDS, *Science* 313 (2006) 462–466.
- [30] T.Y. Lin, M. Emerman, Cyclophilin A interacts with diverse lentiviral capsids, *Retrovirology* 3 (2006) 70.

Research

Open Access

## Modification of a loop sequence between $\alpha$ -helices 6 and 7 of virus capsid (CA) protein in a human immunodeficiency virus type 1 (HIV-1) derivative that has simian immunodeficiency virus (SIVmac239) *vif* and CA $\alpha$ -helices 4 and 5 loop improves replication in cynomolgus monkey cells

Ayumu Kuroishi<sup>1</sup>, Akatsuki Saito<sup>2</sup>, Yasuhiro Shingai<sup>1</sup>, Tatsuo Shioda<sup>1</sup>, Masako Nomaguchi<sup>3</sup>, Akio Adachi<sup>3</sup>, Hirofumi Akari<sup>2</sup> and Emi E Nakayama\*<sup>1</sup>

Address: <sup>1</sup>Department of Viral Infections, Research Institute for Microbial Diseases, Osaka University, Osaka 565-0871, Japan, <sup>2</sup>Tsukuba Primate Research Center, National Institute of Biomedical Innovation, Ibaraki 305-0843, Japan and <sup>3</sup>Department of Virology, Institute of Health Biosciences, University of Tokushima Graduate School, Tokushima 770-8503, Japan

Email: Ayumu Kuroishi - kuroishi@biken.osaka-u.ac.jp; Akatsuki Saito - a-saito@nibio.go.jp; Yasuhiro Shingai - chokobo918@tcct.zaq.ne.jp; Tatsuo Shioda - shioda@biken.osaka-u.ac.jp; Masako Nomaguchi - nomaguchi@basic.med.tokushima-u.ac.jp; Akio Adachi - adachi@basic.med.tokushima-u.ac.jp; Hirofumi Akari - akari@nibio.go.jp; Emi E Nakayama\* - emien@biken.osaka-u.ac.jp

\* Corresponding author

Published: 3 August 2009

Received: 12 March 2009

Retrovirology 2009, 6:70 doi:10.1186/1742-4690-6-70

Accepted: 3 August 2009

This article is available from: <http://www.retrovirology.com/content/6/1/70>

© 2009 Kuroishi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Human immunodeficiency virus type 1 (HIV-1) productively infects only humans and chimpanzees but not cynomolgus or rhesus monkeys while simian immunodeficiency virus isolated from macaque (SIVmac) readily establishes infection in those monkeys. Several HIV-1 and SIVmac chimeric viruses have been constructed in order to develop an animal model for HIV-1 infection. Construction of an HIV-1 derivative which contains sequences of a SIVmac239 loop between  $\alpha$ -helices 4 and 5 (L4/5) of capsid protein (CA) and the entire SIVmac239 *vif* gene was previously reported. Although this chimeric virus could grow in cynomolgus monkey cells, it did so much more slowly than did SIVmac. It was also reported that intrinsic TRIM5 $\alpha$  restricts the post-entry step of HIV-1 replication in rhesus and cynomolgus monkey cells, and we previously demonstrated that a single amino acid in a loop between  $\alpha$ -helices 6 and 7 (L6/7) of HIV type 2 (HIV-2) CA determines the susceptibility of HIV-2 to cynomolgus monkey TRIM5 $\alpha$ .

**Results:** In the study presented here, we replaced L6/7 of HIV-1 CA in addition to L4/5 and *vif* with the corresponding segments of SIVmac. The resultant HIV-1 derivatives showed enhanced replication capability in established T cell lines as well as in CD8<sup>+</sup> cell-depleted primary peripheral blood mononuclear cells from cynomolgus monkey. Compared with the wild type HIV-1 particles, the viral particles produced from a chimeric HIV-1 genome with those two SIVmac loops were less able to saturate the intrinsic restriction in rhesus monkey cells.

**Conclusion:** We have succeeded in making the replication of simian-tropic HIV-1 in cynomolgus monkey cells more efficient by introducing into HIV-1 the L6/7 CA loop from SIVmac. It would be of interest to determine whether HIV-1 derivatives with SIVmac CA L4/5 and L6/7 can establish infection of cynomolgus monkeys *in vivo*.

## Background

Human immunodeficiency virus type 1 (HIV-1) productively infects only humans and chimpanzees but not Old World monkeys (OWM) such as cynomolgus (CM) and rhesus (Rh) monkeys [1]. Unlike the simian immunodeficiency virus isolated from macaques (SIVmac), HIV-1 replication is blocked early after viral entry, before the establishment of a provirus in OWM cells [1-3]. This restricted host range of HIV-1 has greatly hampered its use in animal experiments and has caused difficulties for developing prophylactic vaccines and understanding HIV-1 pathogenesis. In order to establish a monkey model of HIV-1/AIDS, various chimeric viral genomes between SIVmac and HIV-1 (SHIV) have been constructed and tested for their replicative capabilities in simian cells. The first SHIV was generated on a genetic background of SIVmac with HIV-1 *tat*, *rev*, *vif*, and *env* genes [4]. Although such a SHIV is useful for the analysis of humoral immune responses against the Env protein [5-7], SHIVs containing other HIV-1 structural proteins, especially the Gag-Pol protein, have become highly desirable, since cellular immune response against Gag is generally believed to be important for disease control [8-10].

In recent years, several host factors involved in HIV-1 restriction in OWM cells have been identified. ApoB mRNA editing catalytic subunit (APOBEC) 3 G modifies the minus strand viral DNA during reverse transcription, resulting in an impairment of viral replication [11-13]. This activity could be counteracted with the viral protein Vif [14-17]. Although HIV-1 Vif can potentially suppress human APOBEC3G, it is not effective against Rh APOBEC3G, which explains at least partly why HIV-1 replication is restricted in monkey cells. It is well known that Cyclophilin A (CypA) binds directly to the exposed loop between  $\alpha$ -helices 4 and 5 (L4/5) of HIV-1 capsid protein (CA), but not to the SIVmac CA. Several studies have found that CypA augments HIV-1 infection in human cells but inhibits its replication in OWM cells [18-20]. A construction of a SHIV with a minimal segment of SIVmac was reported recently by Kamada et al. [21]. This SHIV was designed to evade the restrictions mediated by APOBEC3G and CypA in OWM cells and contains the 7-aa segment corresponding to the L4/5 of CA and the entire *vif* of SIVmac. The SHIV was found to be able to replicate in primary CD4+ T cells from pig-tailed monkey as well as in the CM HSC-FT cell line. Both in HSC-F and in primary CD4+ T cells, this chimeric virus grew to lower titers than did SIVmac [21]; and when inoculated into pig-tailed monkeys, this SHIV did not cause CD4+ T cell depletion or any clinical symptoms in the inoculated animals [22]. Another SHIV, stHIV-1 (a virus carrying 202 amino acid residues of SIVmac CA and *vif* generated by Hatzioannou et al.) could replicate efficiently in Rh cells [23]. However, long-term passaging in Rh cells was necessary to generate

an efficiently replicating stHIV-1, and this adapted virus has not yet been fully characterized; so it may be that further modifications of the viral genome are necessary for optimal replication of HIV-1 genomes in OWM cells.

TRIM5 $\alpha$ , a member of the tripartite motif (TRIM) family proteins, was identified in 2004 as another intrinsic restriction factor of HIV-1 in OWM cells [24]. Rh and CM TRIM5 $\alpha$  were found to restrict HIV-1 but not SIVmac [25,26]. TRIM5 $\alpha$  recognizes the multimerized CA of an incoming virus by its  $\alpha$ -isoform specific SPRY domain [27-29] and is believed to be involved in innate immunity to control retroviral infection [30]. Previously, Ylinen et al. mapped one of the determinants of TRIM5 $\alpha$  sensitivity in L4/5 of HIV type 2 (HIV-2) CA [31]. In addition, we identified a single amino acid of the surface-exposed loop between  $\alpha$ -helices 6 and 7 (L6/7) of HIV-2 CA as a determinant of the susceptibility of HIV-2 to CM TRIM5 $\alpha$  [32]. We hypothesized that the L6/7 of HIV-1 CA also determines susceptibility to CM TRIM5 $\alpha$ . Here, we investigated whether an additional replacement of L6/7 of HIV-1 CA with that of SIVmac would enhance the replication capability of a SHIV genome in established T cell line HSC-F and in CD8+ cell depleted peripheral blood mononuclear cells (PBMCs) from CMs.

## Materials and methods

### DNA constructions

The HIV-1 derivatives were constructed on a background of infectious molecular clone NL4-3 [33]. NL-ScaVR, a virus containing SIVmac239 L4/5 and the entire *vif* gene, was constructed according to the procedure described by Kamada et al. [21]. A single amino acid His (H) at the 120th position of NL-ScaVR CA was replaced with Gln (Q) by means of site-directed mutagenesis with the PCR-mediated overlap primer extension method [34], and the resultant construct was designated NL-ScaVRA1. The L6/7 of CA (HNPPIP) of NL-SVR, NL-ScaVR, or NL-DT5R was also replaced with the corresponding segments of SIVmac239 CA (RQQNPIP) by means of site-directed mutagenesis, and the resultant constructs were designated NL-SVR6/7S, NL-ScaVR6/7S, or NL-DT5R6/7S, respectively. The BssHII-ApaI fragment of NL-ScaVR, NL-SVR6/7S, or NL-ScaVR6/7S, which corresponds to matrix (MA) and CA, was transferred to *env* deleted NL4-3 (NL-Nhe) to generate the *env* (-) version of each of the constructs.

### Cells and Virus propagation

The 293 T (human kidney), LLC-MK2 (Rh kidney), and TK-ts13 (hamster kidney) adherent cell lines were cultured in Dulbecco's modified Eagle medium supplemented with 10% heat-inactivated FBS. The CD4+ CXCR4+ CM T cell line HSC-F [35] was maintained in RPMI 1640 medium containing 10% FBS. Virus stocks were prepared by transfection of 293 T cells with HIV-1



NL4-3 derivatives using the calcium phosphate co-precipitation method. Viral titers were measured with the p24 or p27 RetroTek antigen ELISA kit (ZeptoMetrix, Buffalo, NY), and viral reverse transcriptase (RT) was quantified with the Reverse Transcriptase Assay kit (Roche Applied Science, Mannheim Germany).

#### **Green fluorescence protein (GFP) vector**

The HIV-1 vector expressing GFP was prepared as described previously [36,37]. To construct the HIV-1-WT-GFP and HIV-1-L4/5S-GFP vector, we replaced the Eco RI-Apa I fragment corresponding to MA and CA of the pMDLg/p.RRE packaging vector with those fragments from NL4-3 and NL-ScaVR, respectively. The GFP viruses were prepared from 293 T cells in a 15-cm dish by co-transfection with a combination of 24 µg of pMDLg/p.RRE derivatives, 36 µg of CS-CDF-CG-PRE (GFP encoding viral genomic plasmid), 10 µg of pMD.G (vesicular stomatitis virus glycoprotein (VSV-G) expressing plasmid), and 10 µg of pRSV-Rev (Rev expressing plasmid). Forty-eight hours after transfection, the culture supernatants were collected and used for infection.

#### **Viral infections**

$3 \times 10^5$  MT4 or HSC-F cells were infected with 20 ng of p24 of NL4-3, NL-ScaV, NL-ScaVR, NL-ScaVR6/7S, NL-DT5R, or NL-DT5R6/7S. The culture supernatants were collected periodically, and p24 levels were measured with an ELISA kit.

#### **Particle purification and Western blotting**

The culture supernatant of 293 T cells transfected with plasmids encoding HIV-1 NL4-3 derivatives was clarified by means of low speed centrifugation. Nine ml of the resultant supernatants were layered onto a 2 ml cushion of 20% sucrose (made in PBS) and centrifuged at 35,000 rpm for 2 h in a Beckman SW41 rotor. After centrifugation, the virion pellets were resuspended in PBS, and p24 antigen concentrations were measured with ELISA. SDS-polyacrylamide gel electrophoresis was applied to 120 ng of p24 of HIV-1 derivatives, and virion-associated proteins were transferred to a PVDF membrane. CA and CypA proteins were visualized with the anti-p24 antibody (Biosdesign International, Saco, ME) and the anti-CypA antibody (Affinity BioReagents, Golden, CO), respectively.

#### **Saturation assay**

HIV-1 derivatives or SIVmac particles were prepared by transfecting each of the env-deleted HIV-1 NL4-3 derivatives or SIVmac plasmids with a plasmid encoding VSV-G into 293 T cells, and culture supernatants were collected two days after transfection. One day before infection, Rh LLC-MK2 and hamster TK-ts13 were plated at a density of  $5 \times 10^4$  cells per well in a 24-well plate. Prior to GFP virus

infection, the cells were pretreated for 2 hours with 200 ng of p24 of each of the HIV-1 or SIVmac particles pseudotyped with VSV-G. Immediately after the pre-treatment, the cells were washed and infected with the HIV-1-WT-GFP or HIV-1-L4/5S-GFP virus. Two hours after infection, the inoculated GFP viruses were washed, and the cells were cultivated in fresh media. Two days after infection, the cells were fixed by formaldehyde, and GFP expressing cells were counted with a flowcytometer. To suppress endogenous TRIM5 $\alpha$  activity, the cells were first infected with Sendai (SeV) expressing TRIM5 lacking the SPRY domain at a multiplicity of infection of 10 plaque forming units per cell. Sixteen hours after SeV infection, the cells were treated with 200 ng of p24 of the particles and then infected with the HIV-1-L4/5S-GFP vector as described above.


#### **Preparation of CD8-depleted CM PBMCs and viral infection**

CM PBMCs were suspended in RPMI medium 1640 supplemented with 10% (vol/vol) FBS, and the CD8+ cells were removed with a magnetic bead system (Miltenyi Biotec, Auburn, CA) and stimulated for 1 day with 1 µg/ml of PHA-L (Sigma, St. Louis, MO). For prolonged stimulation, CD8-depleted CM PBMCs were first stimulated with 1 µg/ml of PHA-L for 2 days and then with human IL2 100 U/ml for 2 more days.  $3 \times 10^5$  cells were then inoculated with 200 ng of p24 of NL-DT5R, NL-DT5R6/7S or with 200 ng of p27 of SIVmac239 and incubated at 37°C in a medium containing 100 U/ml of human IL2. The culture supernatants were collected periodically, and the levels of p24 or p27 were measured with an antigen capture assay (Advanced BioScience Laboratories, Kensington, MD)

## **Results**

#### **Construction and characterization of HIV-1 molecular clones containing CA and Vif sequences from SIVmac239**

Several proviral DNA constructs have been generated to counteract the restriction of HIV-1 replication in CMT cell line HSC-F [38] (Fig. 1). We first generated NL-SVR and NL-ScaVR according to the procedure described by Kamada et al. [21]. NL-ScaVR, a virus with SIVmac239 L4/5 CA and *vif*, could replicate slowly in HSC-F and replicated well in MT4 as previously reported (Fig. 2A). We recently discovered that the 120th amino acid of CA affected the sensitivity of HIV-2 to CM TRIM5 $\alpha$  [32]. We, therefore, introduced an additional amino acid substitution, His to Gln, at this position in NL-ScaVR. The resultant virus was designated NL-ScaVRA1; but this virus unexpectedly showed less efficient replication than did the parental NL-ScaVR in both MT4 and HSC-F cells (Fig. 2A), probably due to a reduced viral fitness created by this mutation. We, therefore, replaced the entire L6/7 CA of NL-ScaVR (HNPPPIP) with the corresponding loop from SIVmac239 (RQQNPIP), and the resultant virus was des-




**Figure 1**

**Structure of the chimeric HIV-1/SIVmac clones and a summary of their replication capabilities.** White bars denote HIV-1 (NL4-3) and gray bars SIVmac239 sequences. +++, ++, +, and - denote the peak titer of virus growth in human (Hu) and cynomolgus monkey (CM) cells, respectively, to more than 1000 ng/ml, 100–1000 ng/ml, 10–100 ng/ml, 1–10 ng/ml, and less than 1 ng/ml concentration of capsid (CA) protein in the culture supernatants. \* denotes that NL-DT5R6/7S replicated faster in HSC-F than did the parental NL-DT5R (see Fig. 2C).

ignated NL-ScaVR6/7S. The amount of RT per 1 ng of CA of NL-ScaVR (0.083 ng) was comparable to that of NL-ScaVR6/7S (0.081 ng), indicating that the replacement of L6/7 in HIV-1 with the corresponding loop of SIVmac did not affect the reactivity of CA antigen. Although NL-ScaVR6/7S grew slightly slower in MT4 cells, it could replicate more efficiently in HSC-F cells than the parental NL-ScaVR could (Fig. 2A). Similar results were obtained when we inoculated 20 ng of RT equivalent of NL-ScaVR or NL-ScaVR6/7S into HSC-F cells and measured the periodic RT production in culture supernatants (data not shown).

These findings demonstrated that L6/7 CA of SIVmac improved the replication in CM cells of an HIV-1 derivative that already contained a SIVmac L4/5 and *vif*. We then generated NL-SVR6/7S, in which the L4/5 sequence was from HIV-1, but the L6/7 and *vif* came from SIVmac. NL-SVR6/7S showed better replication than NL-ScaVR6/7S in MT4 cells, but lost its replicative capability in HSC-F cells (Fig. 2B). NL-SVR, a virus with SIVmac *vif*, could replicate in MT4, but failed to do so in HSC-F (Fig. 2B). These results indicated that both L4/5 and L6/7 of SIVmac are required for efficient replication in HSC-F.



**Figure 2**

**Replication properties of HIV-1 derivatives.** Equal amounts of (A) NL-ScaVR (white diamonds: virus with SIVmac L4/5 and *vif*), and NL-ScaVRA1 (gray diamonds: virus with additional replacement of the 120th amino acid His with Gln in NL-ScaVR), and NL-ScaVR6/7S (black diamonds: virus with SIVmac L4/5, L6/7, and *vif*) (B) NL-SVR, NL-ScaVR6/7S, and NL-SVR6/7S (gray diamonds: virus with SIVmac L6/7 and *vif*), and (C) NL-DT5R (white squares) and NL-DT5R6/7S (black squares), were inoculated into human MT4 or CM HSC-F cells, and culture supernatants were collected periodically. p24 antigen levels were measured by ELISA.


We then introduced SIVmac L6/7 into NL-DT5R, a molecularly cloned virus with two nonsynonymous changes in the *env* gene gained during long-term passages of NL-ScaVR in HSC-F cells [21]. The resultant virus was designated NL-DT5R6/7S. Although the peak titer of NL-DT5R6/7S was almost the same as that of NL-DT5R, NL-DT5R6/7S could replicate faster in HSC-F than the parental NL-DT5R (Fig. 2C). This finding confirmed that SIVmac L6/7 CA sequence improved the replication in CM cells of HIV-1 derivatives that contained SIVmac L4/5 and *vif*. The finding suggested that HIV-1 L6/7 and L4/5 CA sequences are important for intrinsic restriction in CM cells.

#### **CypA incorporation into virus particles was not affected by replacement of HIV-1 L6/7 with that of SIVmac**

Several studies have demonstrated that CypA augments HIV-1 infection in human cells [39], but inhibits its replication in OWM cells [18-20]. CypA was packaged in HIV-1 but not in SIVmac virus particles. To determine whether the replacement of HIV-1 L6/7 with that of SIVmac affects CypA binding of HIV-1 CA, we performed Western blot analysis of viral particles from HIV-1 derivatives. As shown in Fig. 3 (upper panel), CypA proteins were clearly detected in the NL-SVR particles (lane 1) but not in those of NL-ScaVR (lane 3), thus confirming that the L4/5 sequence of HIV-1 but not of SIVmac is required for CypA incorporation into viral particles. CypA proteins were detected in NL-SVR6/7S (lane 2) but not in NL-ScaVR6/7S (lane 4), indicating that the additional replacement of HIV-1 L6/7 with that of SIVmac had little effect on CypA incorporation. This finding suggests that the effect of L6/7 replacement on viral growth was independent from CypA binding of HIV-1 CA. When we used anti-p24 antibody (Fig. 3, lower panel), p55 Gag precursors and p24 proteins were clearly detected. There were no differences in the amount of p24 or the ratio of p24 to p55 among the four HIV-1 derivatives, indicating that the HIV-1 Gag precursor proteins with SIVmac L4/5 and L6/7 were processed normally by the viral protease.


#### **Replacement of both L4/5 and L6/7 of HIV-1 CA with the corresponding loops from SIVmac impaired the CA binding activity of TRIM5 $\alpha$ in Rh cells**

It is known that the intrinsic restriction factors working against HIV-1 in CM and Rh cells can be saturated by inoculation of a high dose of HIV-1 particles [19,40-42]. To determine whether alteration in the CA of HIV-1 would affect its ability to saturate restriction factors, Rh LLC-MK2 cells were pre-treated with equal amounts of VSV-G pseudotyped HIV-1 particles that were with or without SIVmac L4/5 and/or L6/7 CA to saturate intrinsic restriction factor(s). The pre-treated cells were then infected with GFP-expressing HIV-1 carrying SIVmac L4/5 CA (HIV-1-L4/5S-GFP), since we wanted to exclude any effects of CypA on



**Figure 3**  
**Western blot analysis of CA and CypA in particles of HIV-1 derivatives.** The viral particles of NL-SVR (lane 1), NL-SVR6/7S (lane 2), NL-ScaVR (lane 3) and NL-ScaVR6/7S (lane 4) were purified by ultracentrifugation through a 20% sucrose cushion. CypA (upper panel) and p24 and p55 proteins (lower panel) were visualized by Western blotting (WB) using anti-CypA and anti-p24 antibody, respectively. "H" and "S" denote the amino acid sequences derived from HIV-1 and SIVmac, respectively.

the GFP expressing virus in LLC-MK2 cells. The susceptibility of particle-treated cells to virus infection was determined by the percentage of GFP-positive cells. The cells treated with the wild type (WT) particles showed greatly enhanced susceptibility to HIV-1 infection compared with non-treated cells (Fig. 4A, left), demonstrating that the intrinsic restriction factor(s) in LLC-MK2 cells were saturated by a high dose of particles. The cells treated with the particles carrying SIVmac L4/5 and those treated with particles carrying SIVmac L6/7 also showed enhanced susceptibility to HIV-1 infection (Fig. 4A, left). The cells treated with particles carrying both SIVmac L4/5 and L6/7 showed only slight enhancement of HIV-1 susceptibility (Fig. 4A, left;  $p = 0.007$  compared by means of paired t test using all data points with the WT particle treated cells). Similarly, the cells treated with SIVmac particles showed only minor enhancement in HIV-1 susceptibility (Fig. 4A, left). Hamster TK-ts13 cells which lack TRIM5 $\alpha$  expres-



**Figure 4**  
**Saturation of intrinsic antiviral factors resulting from inoculation of high dose of virus particles.** (A) Rhesus LLC-MK2 cells or hamster TK-ts13 cells were pre-treated with equal amounts of VSV-G pseudotyped particles with WT HIV-1 (white squares: Wt), with SIVmac L4/5 (white triangles: 4/5S), with SIVmac L6/7 (white circles: 6/7S), with SIVmac L4/5 and L6/7 (white diamonds: 4/5S6/7S), with SIVmac239 (pluses: SIVmac) or none (crosses) for 2 hours. The cells were then infected with the GFP expressing HIV-1 vector carrying SIVmac L4/5 (A: HIV-1-L4/5S-GFP) or GFP expressing HIV-1 vector with WT capsid (B: HIV-1-WT-GFP). Representative data of four independent experiments are shown. (C) Saturation activities were assessed in the presence or absence of functional TRIM5α. Before particle treatment, cells were infected with Sendai virus (SeV) expressing TRIM5 without the SPRY domain (black symbols), or an empty vector, parental Z strain of SeV (white symbols). Sixteen hours after SeV infection, cells were treated with particles for 2 hours and then infected with HIV-1-L4/5S-GFP. Representative data from six independent experiments are shown.


sion, on the other hand, showed no difference in HIV-1 susceptibility among cells treated with various HIV-1 derivatives or SIVmac particles (Fig. 4A, right). As shown in Fig. 4B, similar results were obtained when we used a GFP-expressing virus with WT HIV-1 capsid (HIV-1-WT-GFP). These results indicate that both HIV-1 L4/5 and L6/7 are important for CA binding to antiviral factor(s) in Rh cells. As described previously [20], HIV-1-WT-GFP could induce infection in only small numbers of LLC-MK2 cells. In contrast, more TK-ts13 cells were infected with HIV-1-WT-GFP than with HIV-1-L4/5-GFP. It is thus possible that CypA is a supporting factor for HIV-1 replication in hamster cells as well as in human cells.

Endogenous TRIM5 $\alpha$  seems to be a likely candidate for the antiviral factor saturated by a high dose of HIV-1 particles (Fig. 4A and 4B). To confirm this, we assessed the ability of WT and mutant HIV-1 particles to saturate the intrinsic restriction factor in the presence or absence of functional TRIM5 $\alpha$ . The dominant negative effect of an over-expressed TRIM5 mutant lacking SPRY domain [43] was used to suppress the function of cell endogenous TRIM5 $\alpha$ . As shown in Fig. 4C, the infection of a recombinant SeV expressing TRIM5 without the SPRY domain caused marked enhancement of HIV-1-L4/5S-GFP virus infection without prior particle treatment (crosses vs. asterisks). This indicates that this dominant negative

TRIM5 mutant successfully suppressed the restriction activity of endogenous TRIM5 $\alpha$ . Treatment with the WT HIV-1 particles also saturated the restriction factors in the cells infected with the empty vector virus (parental Z strain of SeV), while the additional effect of the dominant negative mutant TRIM5 $\alpha$  remained unclear (Fig. 4C left, white vs. black squares). These results suggest that the intrinsic factors saturated by the WT particles were mainly endogenous TRIM5 $\alpha$ . In contrast to the effect of the WT particle treatment, the effect of the dominant negative TRIM5 mutant on HIV-1 infection was evident when we used particles with SIVmac L4/5 and L6/7 (Fig. 4C, right, white vs. black diamonds,  $p = 0.007$ , paired t test). These findings suggest that the diminished capability of particles with SIVmac L4/5 and L6/7 to saturate restriction factors was mainly due to their loss of interaction with TRIM5 $\alpha$ . We, therefore, concluded that the ability of HIV-1 with SIVmac L4/5 and L6/7 to bind to TRIM5 $\alpha$  is diminished in LLC-MK2 cells.

#### **HIV-1 derivative with SIVmac L4/5, L6/7, and *vif* sequences can replicate efficiently in monkey primary cells**

To verify the effect of additional replacement of HIV-1 L6/7 with that of SIVmac in primary CM cells, we prepared PBMCs from CM and removed CD8 $^{+}$  cells by means of magnetic beads. The cells were then stimulated for 1 day with 1  $\mu$ g/ml of PHA-L. NL-DT5R6/7S showed more efficient replication than did the parental NL-DT5R in these cells and reached its peak titer 8 days after infection (Fig. 5A). For prolonged stimulation, CD8-depleted CM PBMCs were first stimulated with 1  $\mu$ g/ml of PHA-L for 2 days and then with human IL2 100 U/ml for 2 more days. In these cells, NL-DT5R with HIV-1 L6/7 did not grow at all. On the other hand, NL-DT5R with SIVmac L6/7 (NL-DT5R6/7S) grew in CM primary cells in response to prolonged stimulation by PHA and IL-2 to reach titers, similar to those attained in cells with short stimulation, up to 8 days after infection (Fig. 5A and 5B). Furthermore, NL-DT5R6/7S continued to grow to much higher titers and reached its peak titer 16 days after infection; this higher peak may be due to better proliferation of these cells than those cells receiving short term stimulation (Fig. 5B). These results confirmed that the replicative capability of HIV-1 in CM cells was augmented by the additional replacement of L6/7 of CA with the corresponding sequence from SIVmac.



**Figure 5**  
**Replication capabilities of HIV-1 derivatives in peripheral blood mononuclear cells (PBMC) from CM.** (A) PBMCs were obtained from CM, after which the CD8 $^{+}$  cells were removed, and the cells were stimulated with PHA-L for 1 day. (B) CD8-depleted CM PBMC were first stimulated with 1  $\mu$ g/ml of PHA-L for 2 days and then with human IL2 100 U/ml for 2 more days. Equal amounts of p24 of NL-DT5R (white squares) or NL-DT5R6/7S (black squares) were inoculated, and the culture supernatants were collected periodically. p24 antigen levels were measured by ELISA. Values represent means with actual fluctuations of duplicate samples added. The values for mock infected cell culture supernatants were zero in the ELISA assay.

#### **Discussion**

We created simian-tropic HIV-1 with more efficient replication capability in CM cells using the knowledge obtained from our previous study of TRIM5 $\alpha$  and HIV-2 capsid sequence variations [32]. Introduction of the entire SIVmac L6/7 CA into the previously constructed version of HIV-1 derivatives containing SIVmac L4/5 CA and *vif* [21] caused only a four amino acid change in CA but

showed improved replication capability of HIV-1 in the CM cell line HSC-F. Introduction of the entire SIVmac L6/7 CA into NL-DT5R, which has two additional amino acid mutations in the *env* gene, enhanced replication in CD8+ cells-depleted CM PBMCs. After prolonged stimulation of CM PBMCs, replication of the original version of NL-DT5R was suppressed while that of NL-DT5R with SIVmac L6/7 was not. It would thus be of interest to test whether those HIV-1 derivatives with both L4/5 and L6/7 from SIVmac can induce infection of CM *in vivo*.

While the high-dose inoculation of WT HIV-1 particles into Rh cells saturated endogenous TRIM5 $\alpha$  and enhanced subsequent infection with HIV-1, the introduction of HIV-1 particles that contained both L4/5 and L6/7 from SIVmac greatly impaired the ability of the particles to saturate TRIM5 $\alpha$ . When we replaced either HIV-1 L4/5 or L6/7 with the corresponding sequence from SIVmac, these particles still saturated TRIM5 $\alpha$ . These findings suggest that TRIM5 $\alpha$  recognized the overall structure composed of both L4/5 and L6/7 of HIV-1 CA. Our previous results from computational 3D-structure modeling analysis of HIV-2 CA support this hypothesis [32]. The 120th amino acid of HIV-2 CA, which affects viral susceptibility to TRIM5 $\alpha$  restriction, was located in L6/7. It is especially worth noting that the amino acid substitution at the 120th position was previously predicted to induce marked changes in the configuration of L6/7 and the L6/7 with the CM TRIM5 $\alpha$ -sensitive Pro positioned most closely to L4/5 of HIV-2 [32]. It would, therefore, be interesting to investigate whether monkey TRIM5 $\alpha$  proteins recognize CypA bound-L4/5 of HIV-1 CA.

During the preparation of our manuscript, Lin and Emerman reported that SIVagmTAN with both HIV-1 L4/5 and L6/7 was susceptible to Rh-TRIM5 $\alpha$  restriction [44]. Our result is consistent with their finding, since the HIV-1 particles with both SIVmac L6/7 and SIVmac L4/5 showed reduced saturation activity for TRIM5 $\alpha$  in Rh cells compared with HIV-1 particles with SIVmac L4/5 alone. Hatzioannou et al. very recently reported that stHIV-1 strains, which differ from HIV-1 only in the *vif* gene, could efficiently replicate in pig-tailed monkey and proposed a pig-tail monkey model of HIV-1 infection [45]. This is not surprising, since pig-tailed monkeys lack a TRIM5 $\alpha$  protein, and the dominant form of TRIM5 expressed in this monkey species is a TRIMCyp fusion protein lacking anti-HIV-1 activity [46-48].

When we subjected CD8-depleted CM PBMC to prolonged stimulation, NL-DT5R6/7S grew efficiently but NL-DT5R did not. Since the expression levels of TRIM5 $\alpha$  mRNA in human PBMC increased after stimulation with PHA and IL2 for 3 days (data not shown), we speculated that the higher expression levels of CM-TRIM5 $\alpha$  in fully

stimulated CM cells resulted in efficient restriction of NL-DT5R. However, no clear enhancement of CM TRIM5 $\alpha$  mRNA expression could be detected in the CM cells subjected to prolonged stimulation (data not shown). The reason why NL-DT5R failed to grow in CM cells with prolonged stimulation is not yet clear, but it is possible that fully stimulated CM cells exerted stronger intrinsic inhibitory activity against HIV-1 infection than those with short-term stimulation.

NL-DT5R6/7S and NL-ScaVR6/7S replicated less efficiently in human MT4 cells than did the parental NL-DT5R and NL-ScaVR. One possible explanation is that the virus with SIVmac L6/7 became resistant to CM TRIM5 $\alpha$  but became more sensitive to human TRIM5 $\alpha$ , since the latter can restrict SIVmac more efficiently than HIV-1. Another possibility is that replacement of CA allowed the virus to evade the intrinsic inhibitory factors in CM cells but impaired viral replication *per se*.

We used the CM T cell line HSC-F and CD8+ cell-depleted PBMC from CM but not from Rh for our replication experiments. Although we observed an improvement of viral replication in CM cells, we cannot assume that the replacement of L4/5 and L6/7 is enough for HIV-1 to replicate to high titers in Rh cells since the CM TRIM5 $\alpha$  resistant HIV-2 mutant virus GH123 (Q) was found to be restricted by Rh TRIM5 $\alpha$  [34]. NL-DT5R6/7S and NL-ScaVR6/7S also showed less efficient replication capability than did SIVmac (Fig. 1). We are currently trying to adapt these viruses to CM and Rh cells by means of long-term passaging in the hope of introducing compensating mutations that can overcome these disadvantages and further augment their replicative capabilities in human and simian cells to reach a similar level as seen with SIVmac.

## Conclusion

We have succeeded in improving simian-tropic HIV-1 for more efficient replication in CM cells by introduction of the SIVmac L6/7 CA sequence. It will be of interest to determine whether the HIV-1 derivatives with SIVmac L4/5 and L6/7 can induce infection in cynomolgus monkeys *in vivo*. Even if they fail to do so, further modification and/or adaptation of the current version of simian-tropic HIV-1 in monkey cells might be expected to lead to the development of an HIV-1 infection model in OWMs. This model has been long-awaited as a tool for vaccine development and as a model for better understanding of AIDS pathogenesis.

## Abbreviations

OWM: old world monkey; CM: cynomolgus monkey; Rh: rhesus monkey; SHIV: HIV-1/SIV chimeric virus; CypA: cyclophilin A; TRIM: tripartite motif; CA: capsid; PBMC: peripheral blood mononuclear cell; GFP: green fluores-

cence protein; VSV-G: vesicular stomatitis virus glycoprotein; SeV: Sendai virus; L4/5: a loop between  $\alpha$ -helices 4 and 5; L6/7: a loop between  $\alpha$ -helices 6 and 7.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

TS and EEN designed the research, AK, AS, YS, and EEN performed the research, TS, MN, AA, and EEN analyzed the data, and AA, HA, TS, and EEN wrote the paper.

### Acknowledgements

The authors wish to thank Mss. Setsuko Bandou and Noriko Teramoto for their helpful assistance.

This work was supported by grants from the Health Science Foundation, the Ministry of Education, Culture, Sports, Science, and Technology, and the Ministry of Health, Labour and Welfare, Japan.

### References

- Shibata R, Sakai H, Kawamura M, Tokunaga K, Adachi A: **Early replication block of human immunodeficiency virus type 1 in monkey cells.** *J Gen Virol* 1995, **76**(Pt 11):2723-2730.
- Himathongkham S, Luciw PA: **Restriction of HIV-1 (subtype B) replication at the entry step in rhesus macaque cells.** *Virology* 1996, **219**:485-488.
- Hofmann W, Schubert D, LaBonte J, Munson L, Gibson S, Scammell J, Ferrigno P, Sodroski J: **Species-specific, postentry barriers to primate immunodeficiency virus infection.** *J Virol* 1999, **73**:10020-10028.
- Shibata R, Kawamura M, Sakai H, Hayami M, Ishimoto A, Adachi A: **Generation of a chimeric human and simian immunodeficiency virus infectious to monkey peripheral blood mononuclear cells.** *J Virol* 1991, **65**:3514-3520.
- Mascola JR, Stiegler G, VanCott TC, Katinger H, Carpenter CB, Hanson CE, Beary H, Hayes D, Frankel SS, Birx DL, Lewis MG: **Protection of macaques against vaginal transmission of a pathogenic HIV-1/SIV chimeric virus by passive infusion of neutralizing antibodies.** *Nat Med* 2000, **6**:207-210.
- Nishimura Y, Igarashi T, Haigwood N, Sadjadpour R, Plishka RJ, Buckler-White A, Shibata R, Martin MA: **Determination of a statistically valid neutralization titer in plasma that confers protection against simian-human immunodeficiency virus challenge following passive transfer of high-titered neutralizing antibodies.** *J Virol* 2002, **76**:2123-2130.
- Shibata R, Igarashi T, Haigwood N, Buckler-White A, Ogert R, Ross W, Willey R, Cho MV, Martin MA: **Neutralizing antibody directed against the HIV-1 envelope glycoprotein can completely block HIV-1/SIV chimeric virus infections of macaque monkeys.** *Nat Med* 1999, **5**:204-210.
- Matano T, Shibata R, Siemon C, Connors M, Lane HC, Martin MA: **Administration of an anti-CD8 monoclonal antibody interferes with the clearance of chimeric simian/human immunodeficiency virus during primary infections of rhesus macaques.** *J Virol* 1998, **72**:164-169.
- Migueles SA, Sabbaghian MS, Shupert WL, Bettinotti MP, Marincola FM, Martino L, Hallahan CV, Selig SM, Schwartz D, Sullivan J, Connors M: **HLA B\*5701 is highly associated with restriction of virus replication in a subgroup of HIV-infected long term nonprogressors.** *Proc Natl Acad Sci USA* 2000, **97**:2709-2714.
- Nixon DF, Townsend AR, Elvin JG, Rizza CR, Gallwey J, McMichael AJ: **HIV-1 gag-specific cytotoxic T lymphocytes defined with recombinant vaccinia virus and synthetic peptides.** *Nature* 1988, **336**:484-487.
- Harris RS, Bishop KN, Sheehy AM, Craig HM, Petersen-Mahrt SK, Watt IN, Neuberger MS, Malim MH: **DNA deamination mediates innate immunity to retroviral infection.** *Cell* 2003, **113**:803-809.
- Mangeat B, Turelli P, Caron G, Friedli M, Perrin L, Trono D: **Broad antiretroviral defence by human APOBEC3G through lethal editing of nascent reverse transcripts.** *Nature* 2003, **424**:99-103.
- Sheehy AM, Gaddis NC, Choi JD, Malim MH: **Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein.** *Nature* 2002, **418**:646-650.
- Mariani R, Chen D, Schrofelbauer B, Navarro F, Konig R, Bollman B, Munk C, Nymark-McMahon H, Landau NR: **Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif.** *Cell* 2003, **114**:21-31.
- Marin M, Rose KM, Kozak SL, Kabat D: **HIV-1 Vif protein binds the editing enzyme APOBEC3G and induces its degradation.** *Nat Med* 2003, **9**:1398-1403.
- Sheehy AM, Gaddis NC, Malim MH: **The antiretroviral enzyme APOBEC3G is degraded by the proteasome in response to HIV-1 Vif.** *Nat Med* 2003, **9**:1404-1407.
- Goila-Gaur R, Strelbel K: **HIV-1 Vif, APOBEC, and intrinsic immunity.** *Retrovirology* 2008, **5**:51.
- Berthoux L, Sebastian S, Sokolskaja E, Luban J: **Lv1 inhibition of human immunodeficiency virus type 1 is counteracted by factors that stimulate synthesis or nuclear translocation of viral cDNA.** *J Virol* 2004, **78**:11739-11750.
- Kootstra NA, Munk C, Tonnu N, Landau NR, Verma JM: **Abrogation of postentry restriction of HIV-1-based lentiviral vector transduction in simian cells.** *Proc Natl Acad Sci USA* 2003, **100**:1298-1303.
- Nakayama EE, Shingai Y, Kono K, Shioda T: **TRIM5alpha-independent anti-human immunodeficiency virus type 1 activity mediated by cyclophilin A in Old World monkey cells.** *Virology* 2008, **375**:514-520.
- Kamada K, Igarashi T, Martin MA, Khamsri B, Hatcho K, Yamashita T, Fujita M, Uchiyama T, Adachi A: **Generation of HIV-1 derivatives that productively infect macaque monkey lymphoid cells.** *Proc Natl Acad Sci USA* 2006, **103**:16959-16964.
- Igarashi T, Iyengar R, Byrum RA, Buckler-White A, Dewar RL, Buckler CE, Lane HC, Kamada K, Adachi A, Martin MA: **Human immunodeficiency virus type 1 derivative with 7% simian immunodeficiency virus genetic content is able to establish infections in pig-tailed macaques.** *J Virol* 2007, **81**:11549-11552.
- Hatzioannou T, Princiotta M, Piatak M Jr, Yuan F, Zhang F, Lifson JD, Bieniasz PD: **Generation of simian-tropic HIV-1 by restriction factor evasion.** *Science* 2006, **314**:95.
- Stremelau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J: **The cytoplasmic body component TRIM5alpha restricts HIV-1 infection in Old World monkeys.** *Nature* 2004, **427**:848-853.
- Luban J: **Cyclophilin A, TRIM5, and resistance to human immunodeficiency virus type 1 infection.** *J Virol* 2007, **81**:1054-1061.
- Towers GJ: **The control of viral infection by tripartite motif proteins and cyclophilin A.** *Retrovirology* 2007, **4**:40.
- Li Y, Li X, Stremelau M, Lee M, Sodroski J: **Removal of arginine 332 allows human TRIM5alpha to bind human immunodeficiency virus capsids and to restrict infection.** *J Virol* 2006, **80**:6738-6744.
- Nakayama EE, Miyoshi H, Nagai Y, Shioda T: **A specific region of 37 amino acid residues in the SPRY (B30.2) domain of African green monkey TRIM5alpha determines species-specific restriction of simian immunodeficiency virus SIVmac infection.** *J Virol* 2005, **79**:8870-8877.
- Ohkura S, Yap MW, Sheldon T, Stoye JP: **All three variable regions of the TRIM5alpha B30.2 domain can contribute to the specificity of retrovirus restriction.** *J Virol* 2006, **80**:8554-8565.
- Ozato K, Shin DM, Chang TH, Morse HC 3rd: **TRIM family proteins and their emerging roles in innate immunity.** *Nat Rev Immunol* 2008, **8**:849-860.
- Ylisen LM, Keckesova Z, Wilson SJ, Ranasinghe S, Towers GJ: **Differential restriction of human immunodeficiency virus type 2 and simian immunodeficiency virus SIVmac by TRIM5alpha alleles.** *J Virol* 2005, **79**:11580-11587.
- Song H, Nakayama EE, Yokoyama M, Sato H, Levy JA, Shioda T: **A single amino acid of the human immunodeficiency virus type 2 capsid affects its replication in the presence of cynomolgus monkey and human TRIM5alphas.** *J Virol* 2007, **81**:7280-7285.



33. Adachi A, Gendelman HE, Koeing S, Folks T, Willey R, Rabson A, Martin MA: **Production of acquired immunodeficiency syndrome-associated retrovirus in human and nonhuman cells transfected with an infectious molecular clone.** *J Virol* 1986, **59**:284-291.
34. Kono K, Song H, Shingai Y, Shioda T, Nakayama EE: **Comparison of anti-viral activity of rhesus monkey and cynomolgus monkey TRIM5alphas against human immunodeficiency virus type 2 infection.** *Virology* 2008, **373**:447-456.
35. Akari H, Mori K, Terao K, Otani I, Fukasawa M, Mukai R, Yoshikawa Y: **In vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its susceptibility to infection with simian immunodeficiency viruses.** *Virology* 1996, **218**:382-388.
36. Miyoshi H, Takahashi M, Gage FH, Verma IM: **Stable and efficient gene transfer into the retina using an HIV-based lentiviral vector.** *Proc Natl Acad Sci USA* 1997, **94**:10319-10323.
37. Miyoshi H, Blomer U, Takahashi M, Gage FH, Verma IM: **Development of a self-inactivating lentivirus vector.** *J Virol* 1998, **72**:8150-8157.
38. Akari H, Fukumori T, Iida S, Adachi A: **Induction of apoptosis in Herpesvirus saimiri-immortalized T lymphocytes by blocking interaction of CD28 with CD80/CD86.** *Biochem Biophys Res Commun* 1999, **263**:352-356.
39. Lin TY, Emerman M: **Cyclophilin A interacts with diverse lentiviral capsids.** *Retrovirology* 2006, **3**:70.
40. Besnier C, Takeuchi Y, Towers G: **Restriction of lentivirus in monkeys.** *Proc Natl Acad Sci USA* 2002, **99**:11920-11925.
41. Cowan S, Hatzioannou T, Cunningham T, Muesing MA, Gottlinger HG, Bieniasz PD: **Cellular inhibitors with Fv1-like activity restrict human and simian immunodeficiency virus tropism.** *Proc Natl Acad Sci USA* 2002, **99**:11914-11919.
42. Hatzioannou T, Cowan S, Goff SP, Bieniasz PD, Towers GJ: **Restriction of multiple divergent retroviruses by Lvl and Ref1.** *Embo J* 2003, **22**:385-394.
43. Maegawa H, Nakayama EE, Kuroishi A, Shioda T: **Silencing of tripartite motif protein (TRIM) 5alpha mediated anti-HIV-1 activity by truncated mutant of TRIM5alpha.** *J Virol Methods* 2008, **151**:249-256.
44. Lin TY, Emerman M: **Determinants of cyclophilin A-dependent TRIM5 alpha restriction against HIV-1.** *Virology* 2008, **379**:335-341.
45. Hatzioannou T, Ambrose Z, Chung NP, Piatak M Jr, Yuan F, Trubey CM, Coalter V, Kiser R, Schneider D, Smedley J, Pung R, Gathuka M, Estes JD, Veazey RS, KewalRamani VN, Lifson JD, Bieniasz PD: **A macaque model of HIV-1 infection.** *Proc Natl Acad Sci USA* 2009, **106**:4425-4429.
46. Kuang YQ, Tang X, Liu FL, Jiang XL, Zhang YP, Gao G, Zheng YT: **Genotyping of TRIM5 locus in northern pig-tailed macaques (Macaca leonina), a primate species susceptible to Human Immunodeficiency Virus type 1 infection.** *Retrovirology* 2009, **6**:58.
47. Virgen CA, Kratovac Z, Bieniasz PD, Hatzioannou T: **Independent genesis of chimeric TRIM5-cyclophilin proteins in two primate species.** *Proc Natl Acad Sci USA* 2008, **105**:3563-3568.
48. Brennan G, Kozyrev Y, Kodama T, Hu SL: **Novel TRIM5 isoforms expressed by Macaca nemestrina.** *J Virol* 2007, **81**:12210-12217.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)



BioMed Central

## Polyfunctional CD4<sup>+</sup> T-Cell Induction in Neutralizing Antibody-Triggered Control of Simian Immunodeficiency Virus Infection<sup>∇</sup>

Takuya Yamamoto,<sup>1,2,†</sup> Nami Iwamoto,<sup>3,†</sup> Hiroyuki Yamamoto,<sup>3,†</sup> Tetsuo Tsukamoto,<sup>3</sup>  
Tetsuya Kuwano,<sup>3</sup> Akiko Takeda,<sup>3</sup> Miki Kawada,<sup>3</sup> Yasuko Tsunetsugu-Yokota,<sup>1</sup>  
and Tetsuro Matano<sup>3,4,5\*</sup>

*Department of Immunology<sup>1</sup> and AIDS Research Center,<sup>4</sup> National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-Ku, Tokyo 162-8640, Japan; Division of Cellular and Molecular Biology<sup>2</sup> and International Research Center for Infectious Diseases,<sup>3</sup> Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-Ku, Tokyo 108-8639, Japan; and Tsukuba Primate Research Center, National Institute of Biomedical Innovation, 1 Hachimandai, Tsukuba 305-0843, Japan<sup>5</sup>*

Received 22 January 2009/Accepted 6 March 2009

Rapid depletion of memory CD4<sup>+</sup> T cells and delayed induction of neutralizing antibody (NAb) responses are characteristics of human immunodeficiency virus (HIV) and simian immunodeficiency virus (SIV) infections. Although it was speculated that postinfection NAb induction could have only a limited suppressive effect on primary HIV replication, a recent study has shown that a single passive NAb immunization of rhesus macaques 1 week after SIV challenge can result in reduction of viral loads at the set point, indicating a possible contribution of postinfection NAb responses to virus control. However, the mechanism accounting for this NAb-triggered SIV control has remained unclear. Here, we report rapid induction of virus-specific polyfunctional T-cell responses after the passive NAb immunization postinfection. Analysis of SIV Gag-specific responses of gamma interferon, tumor necrosis factor alpha, interleukin-2, macrophage inflammatory protein 1β, and CD107a revealed that the polyfunctionality of Gag-specific CD4<sup>+</sup> T cells, as defined by the multiplicity of these responses, was markedly elevated in the acute phase in NAb-immunized animals. In the chronic phase, despite the absence of detectable NABs, virus control was maintained, accompanied by polyfunctional Gag-specific T-cell responses. These results implicate virus-specific polyfunctional CD4<sup>+</sup> T-cell responses in this NAb-triggered virus control, suggesting possible synergism between NABs and T cells for control of HIV/SIV replication.

Virus-specific CD4<sup>+</sup> and CD8<sup>+</sup> T-cell responses are crucial for the control of pathogenic human immunodeficiency virus type 1 (HIV-1) and simian immunodeficiency virus (SIV) infections (5, 6, 20, 23, 30, 39, 40). However, CD4<sup>+</sup> T cells, especially CCR5<sup>+</sup> memory CD4<sup>+</sup> T cells, are themselves targets for these viruses, which may be an obstacle to potent virus-specific CD4<sup>+</sup> T-cell induction (10, 47, 52). Indeed, HIV-1/SIV infection causes rapid, massive depletion of memory CD4<sup>+</sup> T cells (26, 31), and host immune responses fail to contain viral replication and allow persistent chronic infection, although virus-specific CD8<sup>+</sup> T-cell responses exert suppressive pressure on viral replication (15).

Recently, the importance of T-cell quality in virus containment has been highlighted, and T-cell polyfunctionality, which is defined by their multiplicity of antigen-specific cytokine production, has been analyzed as an indicator of T-cell quality (4, 8, 11, 41). However, there has been no evidence indicating an association of polyfunctional T-cell responses in the acute phase with HIV-1/SIV control. Even in the chronic phase, whether polyfunctional CD4<sup>+</sup> T-cell responses may be associ-

ated with virus control has been unclear, although an inverse correlation between polyfunctional CD8<sup>+</sup> T-cell responses and viral loads has been shown in HIV-1-infected individuals (4).

Another characteristic of HIV-1/SIV infections is the absence of potent neutralizing antibody (NAb) induction during the acute phase (7). This is mainly due to the unusually neutralization-resistant nature of the virus, such as masking of target epitopes in viral envelope proteins (24). Whether this lack of effective NAb response contributes to the failure to control the virus, and whether NAb induction in the acute phase can contribute to virus control, remains unclear. Previous studies documenting virus escape from NAb recognition suggested that NABs can also exert selective pressure on viral replication to a certain extent (38, 45, 49), but it was speculated that postinfection NAb induction could have only a limited suppressive effect on primary HIV-1/SIV replication (34, 37).

By passive NAb immunization of rhesus macaques after SIV challenge, we recently provided evidence indicating that the presence of NABs during the acute phase can result in SIV control (50). In that study, passive NAb immunization 1 week after SIVmac239 challenge resulted in transient detectable NAb responses followed by reduction in set point viral loads compared to unimmunized macaques. However, the mechanism of this virus control has remained unclear. In the present study, we found rapid appearance of polyfunctional Gag-specific CD4<sup>+</sup> T-cell responses after such passive NAb immunization postinfection. These animals maintained virus control

\* Corresponding author. Mailing address: International Research Center for Infectious Diseases, Institute of Medical Science, University of Tokyo, 4-6-1 Shirokanedai, Minato-Ku, Tokyo 108-8639, Japan. Phone: 81-3-6409-2078. Fax: 81-3-6409-2076. E-mail: matano@ims.u-tokyo.ac.jp.

† T.Y., N.I., and H.Y. contributed equally to this work.

∇ Published ahead of print on 18 March 2009.




FIG. 1. Follow up of NAb-immunized macaques. (A) Plasma viral loads (SIV *gag* RNA copies/ml plasma) in six unimmunized macaques (black lines) and five NAb-immunized animals (red lines) after SIVmac239 challenge. The plasma viral loads were measured as described previously (29). The lower limit of detection was approximately  $4 \times 10^2$  copies/ml. The MHC-I haplotypes are shown in parentheses following the macaque numbers as follows: E, haplotype 90-010-Ie; D, 90-010-Id; G, 90-030-Ig; J, 90-088-Ij; and H, 90-030-Ih. Below are comparisons of plasma viral loads in unimmunized (non-Tx) and NAb-immunized (NAb-Tx) macaques at weeks (wk) 1, 2, 8, 30, and 60. The bars indicate the geometric mean of each group. The comparisons at weeks 8, 30, and 60 (indicated by asterisks) showed significant differences between two groups ( $P = 0.841$  at week 1,  $P = 0.238$  at week 2,  $P = 0.002$  at week 8,  $P = 0.005$  around week 30, and  $P < 0.001$  around week 60 by *t* test). (B) Peripheral CD4<sup>+</sup> T-cell counts (per  $\mu$ l) in unimmunized controls (black lines) and NAb-immunized macaques (red lines) after SIVmac239 challenge. The ratios of the counts around week 60 to those at week 0 in NAb-immunized macaques were significantly higher than in unimmunized controls ( $P = 0.028$  by *t* test).

for more than 1 year in the absence of detectable plasma NABs, which was accompanied by potent Gag-specific T-cell responses. These results implicate virus-specific polyfunctional CD4<sup>+</sup> T-cell responses in this NAb-triggered primary and long-term SIV control.

#### MATERIALS AND METHODS

**Animal experiments.** We previously showed a reduction in set point viral loads by passive NAb immunization of rhesus macaques (*Macaca mulatta*) 1 week after SIVmac239 challenge (50). In the present study, we monitored these animals, including one additional NAb-immunized macaque (R06-023), for more than 1

year. Thus, five NAb-immunized rhesus macaques and six unimmunized controls were used in this study. Unimmunized macaque R02-021 was euthanized at week 32 because the animal showed loss of body weight, diarrhea, and general weakness. NAb-immunized macaque R02-020 was euthanized at week 42 because of a limitation on available cage numbers. Major histocompatibility complex class I (MHC-I) haplotypes were determined by reference strand-mediated conformation analysis as described previously (2, 29). A group of rhesus macaques possessing the MHC-I haplotype 90-120-Ia with the potential to efficiently elicit potent Gag-specific CD8<sup>+</sup> T-cell responses (21, 29) were not included in this study. All animals were maintained in accordance with the guidelines for animal experiments performed at the National Institute of Infectious Diseases (33).

For passive NAb immunization, immunoglobulin G (IgG) was purified from plasma samples from SIVmac239-infected macaques with SIV-specific NAb re-




FIG. 2. Gag-specific CD4<sup>+</sup> T-cell responses in the acute phase. PBMCs at week 2 postchallenge were stimulated with a recombinant SIV Gag p55, and the specific induction of five marker cytokines (IFN- $\gamma$ , TNF- $\alpha$ , IL-2, MIP-1 $\beta$ , and CD107a) in CD4<sup>+</sup> T cells was examined. PBMCs from macaque R02-004 were unavailable and therefore could not be included in this analysis. (A) Representative dot plots showing responses of IFN- $\gamma$ , TNF- $\alpha$ , IL-2, MIP-1 $\beta$ , and CD107a in CD4<sup>+</sup> T cells in macaque R03-013 after Gag-specific stimulation. Gag-specific IFN- $\gamma$  responses were undetectable at week 2 in this animal. (B) Frequencies of total Gag-specific CD4<sup>+</sup> T cells that showed Gag-specific induction of IFN- $\gamma$ , TNF- $\alpha$ , IL-2, MIP-1 $\beta$ , or CD107a in total CD4<sup>+</sup> T cells. (C) Percentages of cells exhibiting Gag-specific induction of single or multiple marker cytokines in total Gag-specific CD4<sup>+</sup> T cells. (Top) Mean percentages of Gag-specific CD4<sup>+</sup> T cells producing one (dark blue), two (light blue), three (green), four (yellow), or five (red) marker cytokines in unimmunized (non-Tx) and NAb-immunized (NAb-Tx) macaques. (Bottom) Mean percentages of individual Gag-specific CD4<sup>+</sup> T-cell subsets divided by the patterns of marker cytokine induction in unimmunized (blue bars) and NAb-immunized (red bars) macaques. (D) Frequencies of polyfunctional Gag-specific CD4<sup>+</sup> T cells that showed Gag-specific induction of  $\geq 3$  marker cytokines in total CD4<sup>+</sup> T cells. On the left, the bars indicate the geometric mean of each group. The frequencies in NAb-immunized macaques ( $n = 5$ ) were significantly higher than in unimmunized controls ( $n = 5$ ) ( $P = 0.008$ ).

sponses in the chronic phase, as described previously (50). The SIVmac239-specific NAb titer of this IgG preparation (30 mg/ml) was 1:16 on MT-4 cells. Animals were intravenously infused with 300 mg of IgG 1 week after challenge with 1,000 50% tissue culture infective doses (TCID<sub>50</sub>) of SIVmac239 (22). Two unimmunized controls, R02-021 and R06-038, received 300 mg of control IgG

prepared from noninfected rhesus macaques at week 1. Neutralizing titers were measured as described previously (50). In brief, serial twofold dilutions of heat-inactivated plasma in duplicate were incubated with 10 TCID<sub>50</sub> of SIVmac239 for 45 min (5  $\mu$ l of diluted sample was incubated with 5  $\mu$ l of virus in each mixture) and added to 5  $\times 10^4$  MT-4 cells/well in 96-well plates. Progeny virus