

- Simons, K., Toomre, D., 2000. Lipid rafts and signal transduction. *Nat. Rev. Mol. Cell Biol.* 1, 31–39.
- Szente, L., Szejtli, J., 1999. Highly soluble cyclodextrin derivatives: chemistry, properties, and trends in development. *Adv. Drug Deliv. Rev.* 36, 17–28.
- Taatjes, D.J., Sobel, B.E., Budd, R.C., 2008. Morphological and cytochemical determination of cell death by apoptosis. *Histochem. Cell Biol.* 129, 33–43.
- Tu, H.C., Ren, D., Wang, G.X., Chen, D.Y., Westergard, T.D., Kim, H., Sasagawa, S., Hsieh, J.J., Cheng, E.H., 2009. The p53-cathepsin axis cooperates with ROS to activate programmed necrotic death upon DNA damage. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1093–1098.
- Uekama, K., Hirayama, F., Irie, T., 1998. Cyclodextrin drug carrier systems. *Chem. Rev.* 98, 2045–2076.
- Wang, X., 2001. The expanding role of mitochondria in apoptosis. *Genes Dev.* 15, 2922–2933.
- Yang, E., Zha, J., Jockel, J., Boise, L.H., Thompson, C.B., Korsmeyer, S.J., 1995. Bad, a heterodimeric partner for Bcl-X_L and Bcl-2, displaces Bax and promotes cell death. *Cell* 80, 285–291.
- Yang, J., Liu, X., Bhalla, K., Kim, C.N., Ibrado, A.M., Cai, J., Peng, T.I., Jones, D.P., Wang, X., 1997. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science* 275, 1129–1132.
- Yunomae, K., Arima, H., Hirayama, F., Uekama, K., 2003. Involvement of cholesterol in the inhibitory effect of dimethyl- β -cyclodextrin on P-glycoprotein and MRP2 function in Caco-2 cells. *FEBS Lett.* 536, 225–231.
- Zha, J., Harada, H., Yang, E., Jockel, J., Korsmeyer, S.J., 1996. Serine phosphorylation of death agonist BAD in response to survival factor results in binding to 14-3-3 not BCL-X(L). *Cell* 87, 619–628.
- Zong, W.X., Ditsworth, D., Bauer, D.E., Wang, Z.Q., Thompson, C.B., 2004. Alkylating DNA damage stimulates a regulated form of necrotic cell death. *Genes Dev.* 18, 1272–1282.

