

Results

Spontaneous $[Ca^{2+}]_c$ Spiking Activities in Fluo-4 AM Loaded SCN Slices

We prepared and stained the acute slices of SCN with a bath application of 20 μ M fluo-4 AM (Fig. 1A and B) as described in Materials and Methods, and used a laser-scanning confocal microscope to monitor their short-term $[Ca^{2+}]_c$ activities over a duration of 800 seconds. Time traces exhibiting Ca^{2+} spikes are shown in Fig. 1C and D. Typically, the traces showed neither a discernable temporal pattern nor cell-specific characteristics: the width of each spike and the inter-spike interval both varied significantly in time and from one cell to the other (Fig. 1C and D). The observation time was limited due to dye bleaching. Therefore, we monitored a total of 2639 stained cells in 82 slices during various circadian phases and found that $18 \pm 1\%$ (mean \pm SEM) of them exhibited spontaneous Ca^{2+} spikes. Collectively, our extensive set of data showed no indication of circadian variation. In other words, unlike the AP firing frequency of SCN neurons, the Ca^{2+} spiking frequency showed no circadian variation (Fig. 1E).

Right after the $[Ca^{2+}]_c$ imaging experiments were done, some slices were double-stained with an astrocyte maker, sulforhodamine 101 to determine if the active

cells producing Ca^{2+} spikes were a neuron or glia [23, 26]. The majority of the active cells turned out to be neurons ($76 \pm 5\%$, 81 out of 106 cells in 9 slices).

Absence of Spontaneous Ca^{2+} Spikes in Cameleon-Transfected SCN Slices

We also looked for the occurrence of spontaneous Ca^{2+} spikes in rat and mouse slice cultures transfected with pCMV/YC or pNSE/YC. In the rat slice cultures, Ca^{2+} spikes were observed only in $2.8 \pm 1.2\%$ of the cells (number of slices = 5) transfected with pCMV/YC and in $0.6 \pm 0.6\%$ of the cells (number of slices = 5) transfected with pNSE/YC. In the mouse slice cultures, Ca^{2+} spikes were observed only in $1.2 \pm 0.7\%$ of the cells (number of slices = 5) transfected with pCMV/YC and were not observed at all in cells transfected with pNSE/YC (number of slices = 5). These percentages were much smaller than those for the acute rat SCN slices that were stained with fluo-4 AM ($F_{4,97} = 8.5$, $P < .01$ by one way ANOVA; Fig. 2A).

Using mouse slice cultures transfected with pNSE/YC, we confirmed the presence of circadian rhythms in $[\text{Ca}^{2+}]_c$ in $62.8 \pm 3.8\%$ of the SCN neuronal population (27 out of 44 neurons; number of slices = 6; Fig. 2B). After monitoring the circadian $[\text{Ca}^{2+}]_c$ rhythms for several cycles, we imaged the SCN neurons at a much higher sampling rate (1 frame per every 3 seconds) for 5 minutes during different circadian

phases (Fig. 2C and D). There was no Ca^{2+} spiking activity even at the plateau phase.

Inducible Ca^{2+} Spikes in Cameleon-Expressed SCN Cells

To understand the lack of Ca^{2+} spikes in cameleon-transfected SCN cells, we examined the effect of additional fluo-4 AM loading on the cameleon-transfected SCN cells (Fig. 3A and B). At a concentration of 2.5 μM , fluo-4 AM evoked Ca^{2+} spikes (Fig. 3C) in 14% of the SCN cells examined (17 out of 118 cameleon-expressed cells in 4 slices). Similarly, the treatment with BAPTA-AM (2.5 μM) could also induce Ca^{2+} spikes in 13% of the SCN cells (20 out of 155 cells in 5 slices) (Fig. 3D). Since both were dissolved in DMSO, we also performed several DMSO control experiments to find no Ca^{2+} spikes (9 slices).

The Modulations in the Level of $[\text{Ca}^{2+}]_c$ and the AP Firing Pattern of SCN Neurons by Fluo-4 Loading

We made fluo-4 imaging and current clamp recording simultaneously using fluo-4 dye loaded micro-pipettes in order to see the direct effect of fluo-4 calcium dye on the level of $[\text{Ca}^{2+}]_c$ and the AP firing pattern in individual SCN neurons. Figure 4A shows a typical pattern of spontaneous AP firing in a SCN neuron in the absence of

fluo-4 dye. It is quite regular and shows a stable resting membrane potential (V_m). In many cases, this regular pattern changed significantly with an addition of fluo-4 (pipette concentration in the range of 80–120 μM). As the dye diffused from the patch pipette into the target neuron, during the first 100 seconds or so, the AP firing rate gradually decreased from 7 Hz to 3Hz, and at around 150 seconds the AP activity disappeared rather quickly (Fig. 4B). This change was accompanied by a decrease in V_m from -45.0 ± 1.6 mV to -54.0 ± 4.0 mV (Fig. 4C). Subsequently, the AP firing rate showed a large change, producing a random sequence of sharp peaks with each peak matched with a Ca^{2+} spike in one-to-one manner (Fig. 4B). During each peak, the value of V_m recovered more or less to the initial level (Fig. 4C).

Regarding the Ca^{2+} spikes in the SCN cells induced by fluo-4 dye, the following facts were further confirmed. First, the results were concentration-dependent. When the pipette dye concentration was below 60 μM , no Ca^{2+} spikes were apparent, and there was no significant change in either the AP firing rate or V_m (10 neurons in 6 slices; Table 1). Conversely, when the dye concentration was more than 150 μM , V_m and $[\text{Ca}^{2+}]_c$ increased steadily until the SCN neurons died (8 neurons in 4 slices). When the dye concentration was in the intermediate range of 80–120 μM , about half (9 out of 18 cells in 8 different slices) of the tested SCN neurons exhibited Ca^{2+} spikes, and they

showed a decrease in V_m of 6.7% ($P < .05$ by Student's t -test).

Fluo-4 AM Concentration Dependence of Ca^{2+} Spikes

The frequency of induced Ca^{2+} spikes was not a simple function of fluo-4 AM concentration. We observed quite heterogeneous responses following a stepwise increase of fluo-4 AM concentration (0.5-10 μM ; Fig. 5). Based on these responses, SCN cell populations could be divided broadly into three different groups. For the majority of the cells (48.7%, 37 out of 76 cells in 3 slices), the amplitude as well as the frequency of Ca^{2+} spikes were an increasing function of fluo-4 AM concentration (Fig. 5A and B; Frequency: $F_{5,105} = 14.6$, Amplitude: $F_{5,105} = 3.8$, $P < .05$). For the next majority of the cells (35.5%, 27 out of 76 cells in 3 slices), there was no systematic trend (Fig. 5C and D) both in the frequency and the amplitude. Finally, for the rest of the cell population (15.8%, 12 out of 76 cells in 3 slices), the frequency was a decreasing function of fluo-4 AM concentration (Fig. 5E and F; $F_{5,33} = 7.74$, $P < .05$), while the amplitude could be decreasing, unchanging, or increasing, depending on the individual cells. In other words, the systematic trend in the frequency did not guarantee the same trend in the amplitude.

Discussion

The Rate of “Spontaneous Ca²⁺ Spikes” Showed No Circadian Variation

The spontaneous AP firing rates and the cytosolic concentration of free calcium ions $[Ca^{2+}]_c$ of SCN neurons are known to have circadian rhythms. Also reported was the existence of $[Ca^{2+}]_c$ transients in SCN having a typical duration of tens of seconds. Here, we termed these Ca²⁺ transients as “Ca²⁺ spikes.” One immediate question following these observations would be if the Ca²⁺ spiking activity of SCN neurons can exhibit any circadian variation similar to the circadian AP firing activity, and here we studied that issue. The level of $[Ca^{2+}]_c$ was visualized by Ca²⁺ sensitive fluorescent protein cameleon as well as fluo-4 AM dye. Many fluo-4-loaded SCN cells exhibited Ca²⁺ spikes spontaneously, but our extensive set of fluo-4 imaging experiments based on 82 acute SCN slices found no day/night variation in the Ca²⁺ spiking rates (Fig. 1E): the Ca²⁺ spiking activities of SCN cells during the night time were as strong (or weak) as those of day time, or vice versa. In our earlier studies that investigated the progressive changes in $[Ca^{2+}]_c$ for the duration longer than a circadian cycle, we used organotypic cultures of SCN slices that were transfected by pNSE/YC genes [14,16,17]. The level of $[Ca^{2+}]_c$ was estimated with YC2.1 at every 10 minute in order to minimize photo-

bleaching: with this slow image acquisition rate, the presence or the absence of Ca^{2+} spikes could not be determined [16]. In the present study, we used YC3.6 (either pNSE/YC or pCMV/YC), and $[\text{Ca}^{2+}]_c$ images were acquired at every 3 seconds to monitor Ca^{2+} spiking events. The cameleon YC3.6 is a newer version of YC2.1 giving a better spatiotemporal resolution [25]. We observed that YC3.6-transfected organotypic cultures of SCN could exhibit well-defined circadian rhythms in the level of $[\text{Ca}^{2+}]_c$, just as in the YC2.1-transfected cultures. However, in the same preparations almost no Ca^{2+} spike was observed even at the peak of $[\text{Ca}^{2+}]_c$ circadian oscillation (Fig. 2). This was definitely not due to the lack of sensitivity of YC3.6 because spontaneous Ca^{2+} spikes were observed in other hypothalamic neurons neighboring SCN (Figure S1).

“Spontaneous Ca^{2+} Spikes” Were Induced by Fluo-4 Ca^{2+} Dye Loading

Our study found Ca^{2+} spikes to be present in about 18% of fluo-4-loaded SCN cells, while Ca^{2+} spikes were observed only in 1-3% of cultured SCN cells, which expressed pCMV/YC genes, and in 0-0.6% of cultured SCN neurons, which expressed pNSE/YC genes. In addition, the population of cultured SCN cells that displayed Ca^{2+} spikes was significantly increased to about 15% by an additional loading of fluo-4 AM or BAPTA-AM (Fig. 3). Taken all together, we could conclude that the observed

spontaneous SCN Ca^{2+} spikes were not intrinsic to SCN but induced by fluo-4 dye loading.

In order to verify the effect of calcium dye loading on the level of $[\text{Ca}^{2+}]_c$ and the spontaneous AP firing activity in SCN neuron, more directly, we performed a patch-clamp recording along with fluo-4 imaging. Most notably, the loading of fluo-4 through the patching micro-pipette to the cell initially decreased both the AP firing rate and the level of resting membrane potential (Fig. 4) of SCN neurons. The decreases were not due to the whole-cell patching itself, as no such decrease was observed in our control experiments. Soon after the spontaneous AP firing activity of SCN neurons disappeared completely, an intermittent sequence of AP firings in groups, each of which well matched to a Ca^{2+} spike in one-to-one manner, emerged (Fig. 4B). The bursting mode of AP firings is quite unusual for SCN neurons, since their spontaneous AP firing pattern is typically regular as shown in Fig. 4A.

It has been proposed that Ca^{2+} influx via L-type Ca^{2+} channels [9] or BK type calcium-activated potassium channel activities [27] act to regulate spontaneous AP firing frequencies in SCN neurons. On the other hand, Aguilar-Roblero et al. [28] have demonstrated that AP firing frequencies in SCN neurons can be increased by ryanodine receptor activators (e.g., caffeine) and decreased by ryanodine receptor blockers (e.g.,

dantrolene): in other words, the mobilization of Ca^{2+} store by ryanodine-sensitive intracellular Ca^{2+} channels can modulate AP firing rate. In any case, all of these studies indicate a close connection between the level of $[\text{Ca}^{2+}]_c$ and the AP firing activity of SCN neurons, and thus it is not too surprising that in our whole-cell patching experiment each Ca^{2+} spike matches with a group of bursting AP spikes in one-to-one manner.

Although our current investigation did not answer how exactly the chelation of Ca^{2+} by dye loading interfered with the cell's intrinsic calcium kinetics to produce Ca^{2+} spikes, there is no doubt that they were induced by fluo-4 dye loading. The observed Ca^{2+} spikes could be a natural response of SCN neurons trying to recover their original cytosolic $[\text{Ca}^{2+}]_c$ levels. Some mechanisms were discussed earlier for the cause of SCN Ca^{2+} spikes. For example, van den Pol et al. [18] and Obrietan and van den Pol [19] found that they could be induced by the modulation of neurotransmitter, such as gamma-aminobutyric acid (GABA) and glutamate. Also, more recently, Irwin and Allen [20] demonstrated that the Ca^{2+} influx through L-type Ca^{2+} channels following the electrical stimulation of retinohypothalamic tracts could induce Ca^{2+} spikes in SCN neurons. The Ca^{2+} spikes observed in our experiments were neither induced by neurotransmitter modulations nor electrical stimulations. Yet, they were visually very

similar to the ones reported earlier [18-20]. In any case, the SCN Ca^{2+} spikes all were not an intrinsic spontaneous activity but induced.

Differences between Cameleon and Fluo-4 Calcium Dye

It has been known that BAPTA-based high affinity calcium dyes, such as fura-2, fluo-3, and fluo-4 can have a significant intracellular Ca^{2+} buffering and induce some non-intrinsic phenomena. For example, Ca^{2+} waves supported by the network of astrocytes can be completely blocked by the chelating action of fura-2 [29]. They can also delay the calcium signals initiated by extracellular calcium influx or release of calcium from intracellular stores by 5- to 20-fold [30]. Therefore, it is not too surprising that Ca^{2+} spikes in SCN neurons have appeared as an artifact of the Ca^{2+} chelation of fluo-4 dye.

When the chelation by indicator dye becomes an issue, its affinity (the inverse of dissociation constant K_d) and concentration are two important factors to be considered. High-affinity chelators trap calcium efficiently, and the presence of them at high concentration can sequester intracellular free Ca^{2+} significantly. We believe that the affinity ($K_d = 345 \text{ nM}$) and the concentration ($5 \sim 20 \text{ }\mu\text{M}$) of fluo-4 were high enough to interfere with the intrinsic calcium kinetics of SCN neurons and resulted in non-intrinsic

Ca²⁺ spikes. The affinity of cameleon YC3.6 (K_d= 250 nM) is 1.73 times higher than that of fluo-4 (K_d = 345 nM). Therefore, we suspect that the cytosolic concentrations of cameleon protein expressed within the SCN neurons were quite lower than that of fluo-4 dye that we used. As for the concentration, another important factor to be considered is the stability of the dye. It is well known that cameleon is far more stable than any BAPTA-based calcium indicators with respect to dye leakage, exhaustion, or cumulative photodestruction [31]. Therefore, the effective concentration of cameleon inside the cells could have been much lower than that of the fluo-4 dye.

The concerns with BAPTA based dyes are well known and our extensive set of data provides a clear case for concern, especially, regarding SCN calcium dynamics. However, our result does not invalidate the use of these dyes.

SCN Cells Are Heterogeneous with Regard to Their Cytosolic Ca²⁺ Buffering Systems

Previous research has shown that Ca²⁺ buffering molecules, such as calbindin D-28k (K_d is about 300 nM), may play a critical role for the function of SCN neurons [32,33]. The type of Ca²⁺ buffering proteins and their localization within the SCN may differ depending on the particular animal species [34,35], the developmental stages [36],

and the circadian timings [37]. For example, Ikeda et al. (2003a) has shown that the nucleus and the cytoplasm of SCN neurons can support quite different Ca^{2+} kinetics at a given time.

Several earlier studies found BAPTA-based fluorescent Ca^{2+} dyes often compete with the Ca^{2+} buffering system of involved cells. In fact, the Ca^{2+} chelating effects of fura-2 could be used to estimate the capacity of the Ca^{2+} buffering systems in various types of cells [38-40]. Accordingly, we suspect that the heterogeneous effect of fluo-4 AM on the generation of Ca^{2+} spikes (Fig. 5 and Table 1) may reflect the different Ca^{2+} buffering capacities of SCN cells. The SCN neurons have been categorized into several different subgroups [41,42], and it is quite possible that the heterogeneity of the Ca^{2+} buffering systems may underlie the existence of these subgroups.

Concluding Remarks

Our experimental studies demonstrated that Ca^{2+} spikes, having a typical bandwidth of tens of seconds, could be induced in SCN neurons by the Ca^{2+} chelating effect of BAPTA-based dyes. The firing rate of these induced SCN Ca^{2+} spikes did not show a circadian variation. Furthermore, Ca^{2+} spikes were very rarely observed when a reporter protein was used instead. Based on these facts, we conclude that spontaneous

Ca^{2+} spikes in the presence of BAPTA-based dyes are most likely not related to the intrinsic circadian rhythm of $[\text{Ca}^{2+}]_c$ in SCN neurons. At this point, however, we need to indicate several other possibilities as well. First of all, our study does not exclude the potential importance of Ca^{2+} spikes, for example, which are related to the modulation of neurotransmitters, for various functions of SCN. Second, it is also possible that there could have been a small subset of cells whose Ca^{2+} spiking rate indeed showed a circadian variation but masked by other non-circadian cells. We also realize that some cortical neurons can support very small and fast Ca^{2+} transients having a bandwidth of about a second or less. Such very fast Ca^{2+} transients could have existed in our experiments but unnoticed due to the limitation of our image acquisition rate (~ 1 second). On the other hand, the possible existence of such neurons in SCN has never been addressed nor been documented in the past.

References

1. Reppert SM, Weaver DR (2002) Coordination of circadian timing in mammals. *Nature* 418: 935-941.
2. Welsh DK, Yoo SH, Liu AC, Takahashi JS, Kay, SA (2004) Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Curr Biol* 14: 2289-2295.
3. Kornmann B, Schaad O, Bujard H, Takahashi JS, Schibler U (2007) System-driven and oscillator-dependent circadian transcription in mice with a conditionally active liver clock. *PloS Biol* 5: e34.
4. van den Pol AN (1980) The hypothalamic suprachiasmatic nucleus of the rat: intrinsic anatomy. *J Comp Neurol* 191: 661-702.
5. Klein D, Moore R, Reppert S (1991) *Suprachiasmatic nucleus. The Mind's Clock*. Oxford: Oxford University Press.
6. Green DJ, Gillette R (1982) Circadian rhythm of firing rate recorded from single cells in the rat suprachiasmatic brain slice. *Brain Res* 245: 198-200.
7. Herzog ED, Geusz ME, Khalsa SBS, Straume M, Block GD (1997) Circadian rhythms in mouse suprachiasmatic nucleus explants on multimicroelectrode plates.

Brain Res 757: 285-290.

8. Schaap J, Bos NP, de Jeu MT, Geurtsen AM, Meijer JH, et al. (1999) Neurons of the rat suprachiasmatic nucleus show a circadian rhythm in membrane properties that is lost during prolonged whole-cell recording. Brain Res 815: 154-166.
9. Pennartz CM, De Jeu MS, Bos NP, Schaap J, de Jeu MT, et al. (2002) Diurnal modulation of pacemaker potentials and calcium current in the mammalian circadian clock. Nature 416: 286-290.
10. Kohsaka A, Bass J (2006) A sense of time: how molecular clocks organize metabolism. Trends Endocrinol Metab 18: 4-11.
11. Haak LL (1999) Metabotropic glutamate receptor modulation of glutamate responses in the suprachiasmatic nucleus. J Neurophysiol 81: 1308-1317.
12. Honma S, Honma K (2003) The biological clock: Ca²⁺ links the pendulum to the hands. Trend Neurosci 26: 650-653.
13. Park D, Lee S, Jun K, Hong YM, Kim DY, et al. (2003) Translation of clock rhythmicity into neural firing in suprachiasmatic nucleus requires mGluR-PLCb4 signaling. Nat Neurosci 6: 337-338.
14. Ikeda M (2004) Calcium dynamics and circadian rhythms in suprachiasmatic nucleus neurons. Neuroscientist 10: 315-324.

15. Colwell CS (2000) Circadian modulation of calcium levels in cells in the suprachiasmatic nucleus. *Eur J Neurosci* 12: 571-576.
16. Ikeda M, Sugiyama T, Wallace CS, Gompf HS, Yoshioka T, et al. (2003) Circadian dynamics of cytosolic and nuclear Ca^{2+} in single suprachiasmatic nucleus neurons. *Neuron* 38: 253-263.
17. Sugiyama T, Yoshioka T, Ikeda M (2004) mPer2 antisense oligonucleotides inhibit mPER2 expression but not circadian rhythms of physiological activity in cultured suprachiasmatic nucleus neurons. *Biochem Biophys Res Commun* 323: 479-483.
18. van den Pol AN, Finkbeiner SM, Cornell-Bell AH (1992) Calcium excitability and oscillations in suprachiasmatic nucleus neurons and glia in vitro. *J Neurosci* 12: 2648-2664.
19. Obrietan K, van den Pol AN (1996) Neuropeptide Y depresses GABA-mediated calcium transients in developing suprachiasmatic nucleus neurons: a novel form of calcium long-term depression. *J Neurosci* 16: 3521-3533.
20. Irwin RB, Allen CN (2007) Calcium response to retinohypothalamic tract synaptic transmission in suprachiasmatic nucleus neurons. *J Neurosci* 27: 11748-11757.

21. Ikeda M, Yoshioka T, Allen CN (2003) Developmental and circadian changes in Ca^{2+} mobilization mediated by GABAA and NMDA receptors in the suprachiasmatic nucleus. *Eur J Neurosci* 17: 57-70.
22. Kim DY, Choi HJ, Kim JS, Kim YS, Jeong DU, et al. (2005) Voltage-gated calcium channels play crucial roles in the glutamate-induced phase shifts of the rat suprachiasmatic circadian clock. *Eur J Neurosci* 21: 1215-1222.
23. Ikegaya Y, Le Bon-Jego M, Yuste R (2005) Large-scale imaging of cortical network activity with calcium indicators. *Neurosci Res* 52: 132-138.
24. Eilers J, Konnerth A (2005) A practical guide: Dye loading with patch pipettes. In: Yuste R, Konnerth A, editors. *Imaging in Neuroscience and Development: a laboratory manual*. New York: Cold Spring Harbor Lab. pp. 277-281.
25. Nagai T, Yamada S, Tominaga T, Ichikawa M, Miyawaki A (2004) Expanded dynamic range of fluorescent indicators for Ca^{2+} by circularly permuted yellow fluorescent proteins. *Proc Natl Acad Sci USA* 101: 10554-10559.
26. Nimmerjahn A, Kirchhoff F, Kerr JN, Helmchen F (2004) Sulforhodamine 101 as a specific marker of astroglia in the neocortex in vivo. *Nat Methods* 1: 31-37.
27. Meredith AL, Wiler SW, Miller BH, Takahashi JS, Fodor AA, et al. (2006) BK calcium-activated potassium channels regulate circadian behavioral rhythms and

- pacemaker output. *Nat Neurosci* 9: 1041-1049.
28. Aguilar-Roblero R, Mercado C, Alamilla J, Laville A, Díaz-Muñoz M (2007) Ryanodine receptor Ca^{2+} -release channels are an output pathway for the circadian clock in the rat suprachiasmatic nuclei. *Eur J Neurosci* 26: 575-582.
 29. Wang Z, Tymianski M, Jones OT, Nedergaard M (1997) Impact of cytoplasmic calcium buffering on the spatial and temporal characteristics of intercellular calcium signals in astrocytes. *J Neurosci* 17: 7359-7371.
 30. Grant SK, Bansal A, Mitra A, Feighner SD, Dai G, et al. (2001) Delay of intracellular calcium transients using a calcium chelator: application to high-throughput screening of the capsaicin receptor ion channel and G-protein-coupled receptors. *Anal. Biochem* 294: 27-35.
 31. Miyawaki A, Griesbeck O, Hiem R, Tsien RY (1999) Dynamic and quantitative Ca^{2+} measurements using improved cameleons. *Proc Natl Acad Sci USA* 96: 2135-2140.
 32. Silver R, Romero MT, Besmer HR, Leak R, Nunez JM, et al. (1996) calbindin-D28k cells in the hamster SCN express light-induced Fos. *Neuroreport* 7: 1224-1228.
 33. LeSauter J, Silver R (1999) Localization of a suprachiasmatic nucleus subregion regulating locomotor rhythmicity. *J Neurosci* 19: 5574-5585.

34. Silver R, Sookhoo AI, LeSauter J, Stevens P, Jansen HT, et al. (1999) Multiple regulatory elements result in regional specificity in circadian rhythms of neuropeptide expression in mouse SCN. *Neuroreport* 10: 3165-3174.
35. Mahoney MM, Nunez AA, Smale L (2000) Calbindin and Fos within the suprachiasmatic nucleus and the adjacent hypothalamus of *Arvicanthis Niloticus* and *Rattus Norvegicus*. *Neuroscience* 99: 565-575.
36. Ikeda M, Allen CN (2003) Developmental changes in calbindin-D28k and calretinin expression in the mouse suprachiasmatic nucleus. *Eur J Neurosci* 17: 1111-1118.
37. Hamada T, LeSauter J, Lokshin M, Romero M-T, Yan L, et al. (2003) Calbindin influences response to photic input in suprachiasmatic nucleus. *J Neurosci* 23: 8820-8826.
38. Neher E, Augustine GJ (1992) Calcium gradients and buffers on bovine chromaffin cells. *J Physiol (Lond)* 450: 273-301.
39. Regehr WD, Tank DW (1992) Dendritic calcium dynamics. *Curr Opin Neurobiol* 4: 373-382.
40. Neher E (1995) The use of fura-2 for estimating Ca^{2+} buffers and Ca^{2+} fluxes. *Neuropharmacology* 34: 1423-1442.

41. Pennartz CM, Geurtsen AMS, De Jeu MTG, Sluiter AA, Hermes MHLJ (1998)
Electrophysiological and morphological heterogeneity of neurons in slices of rat
suprachiasmatic nucleus. *J Physiol (London)* 506: 775-793.
42. Laemle LK, Hori N, Strominger NL, Tan Y, Carpenter DO (2002) Physiological
and anatomical properties of the suprachiasmatic nucleus of an anophthalmic mouse.
Brain Res 953: 73-81.