

respectively, of CaM-M13 (Figure 1A). Of these constructs, Venus-CaM-M13-RLuc8 had a 2-fold higher dynamic range than RLuc8-CaM-M13-Venus and RLuc8-CaM-M13-cp157Venus while other constructs showed a lower dynamic range (Figure 2)e. This result was consistent with previous results showing that orientation in which the donor luciferase was located at the C-terminus of the fusion generated a higher BRET signal compared to the reverse orientation [1,13]. Overall, Venus-CaM-M13-RLuc8, which we designated as BRAC (BRET-based Autoluminescent Ca²⁺ indicator), had the largest dynamic range (60%) among the constructs that we tested (Table 1).

Physicochemical properties of BRAC

To evaluate the chemical properties of BRAC, Ca²⁺ titration was carried out. As shown in Figure 3A, emission from acceptor Venus increased in accordance with [Ca²⁺] while emission from donor RLuc8 was almost constant. This result closely corresponds to the result from the BRET-induced luminescence enhancement in BAF-Y as reported previously¹. Ca²⁺ titration revealed that BRAC gave a monophasic Ca²⁺ response curve with a Hill coefficient of 1.3, and apparent dissociation constant of 1.9 μM (Figure 3B). We also perform pH titration of BRAC, which indicated that the emission ratio of 530 to 480 nm in BRAC was stable in physiological pH between 6.5 and 8.0 (Figure 3C). We then measured the Ca²⁺-association kinetics of BRAC by stopped-flow photometry system. However, the time course data we obtained was composed of at least two exponential decay components ($\tau < 0.1$ sec) which were thought to be derived from both Ca²⁺ and coelenterazine-h binding to BRAC. Because both kinetics were quite similar, we could not

distinguish which component was derived from Ca^{2+} binding to BRAC. Thus, we then measured Ca^{2+} -dissociation kinetics of BRAC (Figure 3D). The dissociation time constant of BRAC ($\tau=0.21$ sec) was independent on Ca^{2+} concentration and much smaller than that of YC3.60 ($\tau=2.9$ sec, $k_{\text{off}}=0.34$ s⁻¹) (Figure 3E, F) and comparable to that of G-CaMP ($\tau=0.19$ sec) [14]. This result was in good agreement with the fact that Ca^{2+} affinity of BRAC ($K_d = 1.9$ μM) is about 10 times lower than that of YC3.60 ($K_d = 0.25$ μM), and indicated that association/dissociation kinetics of BRAC are fast enough to detect Ca^{2+} dynamics in living cells and organelles.

Ca^{2+} imaging by BRAC in living HeLa cells

In order to determine the validity of BRAC in living cells, we transfected HeLa cells with the cDNA encoding BRAC. Direct excitation of the Venus moiety with blue light revealed that BRAC was localized in the cytoplasmic compartment, and slightly in the nucleus (Figure 4A). Figure 4B, 4C and 4D show the time course of RLuc8 and Venus, and pseudo-colored Venus/RLuc8 ratio images, respectively (see Supporting Video 1). Upon stimulation with 10 μM histamine, RLuc8 signals did not show a spike-like shape and sometimes increased or decreased slightly while Venus signals showed an oscillated spike after histamine stimulation as well as drifting of basal signal intensity, as seen in RLuc8 signals (Figure. 4E). The drift observed in both RLuc8 and Venus signals might be caused by the uptake and consumption of coelenterazine-h or the change in cell shape. However, due to the ratiometric imaging of BRAC, this signal drift was successfully canceled out by dividing the Venus signal with that of RLuc8 (Figure 4F). We also examined the dynamic

range of BRAC in HeLa cells by measuring minimum and maximum BRET ratio values which were obtained by treatment of HeLa cells with ionomycin/1mM EGTA and ionomycin/5mM CaCl₂, respectively. The dynamic range obtained was 44% which was comparable to that measured *in vitro*.

For further validation of BRAC as an auto-luminescent Ca²⁺ indicator, we compared performance of BRAC with other BRET-based Ca²⁺ indicator, G5A [5] in HeLa cells. The signal to noise ratio of G5A was much better than that of BRAC (Figure 4G) because G5A did not emit any luminescence before Ca²⁺ binding. However, G5A failed to report histamine-induced Ca²⁺ oscillations that were reproducibly observed by using BRAC (Figure 4F, 4G), probably due to much slow regeneration rate of aequorin from apoaequorin [6,7]. This result indicates the superiority of BRAC in terms of long term imaging of Ca²⁺ dynamics.

Ca²⁺ imaging by BRAC in plant tissue

To further examine BRAC performance, we tried Ca²⁺ imaging in plant leaves, in which fluorescence-based Ca²⁺ imaging is difficult owing to the chloroplasts-derived strong auto-fluorescence, which disturbs reliable detection of fluorescence signal from Ca²⁺ indicator. Ca²⁺ is a ubiquitous intracellular second messenger involved in numerous signaling pathways in plants. Elevation of cytosolic Ca²⁺ concentration ([Ca²⁺]_i) are common early event in plant defense signaling, and Ca²⁺ plays a essential role in activating the plant's surveillance system against attempted microbial invasion [15]. *Arabidopsis thaliana* has resistance to *Pseudomonas syringae* pv. *Tomato* (*Pst*) DC3000 having the

avirulence gene *avrRpm1* (*Pst* DC3000/*avrRpm1*) but susceptible to *Pst* DC3000 [16]. To visualize the $[Ca^{2+}]_i$ increase during the plant defense signaling, we inoculated bacterial pathogens into the leaves of BRAC-expressing *Arabidopsis* lines. We monitored the changes in $[Ca^{2+}]_i$ during 30-45 and 80-95 minutes after the bacterial inoculation (Figure 5). Increase in $[Ca^{2+}]_i$ was observed at approximate 90 minutes after the inoculation of *Pst* DC3000/*avrRpm1* but not *Pst* DC3000. This result is consistent with the finding performed using aequorin-expressing *Arabidopsis* plants [17]. In that paper, aequorin could monitor $[Ca^{2+}]_i$ change in whole leaves every 5 sec or visualize $[Ca^{2+}]_i$ change in single leaf every 50 minutes. On the other hand, BRAC successfully visualized the $[Ca^{2+}]_i$ change in a region of single leaf every 10 sec.

Discussion

We successfully designed and constructed the BRET-based ratiometric Ca^{2+} indicator, BRAC, which could visualize intracellular Ca^{2+} dynamics with 1-second temporal resolution at the single-cell level. Taking advantage of the bioluminescence imaging property that does not require external excitation light, BRAC is a potentially powerful tool for elucidating the cellular implementation of systems-level neural processes such as action, thought and emotion by combining its use with an optogenetic technology for genetically targeted, millisecond-timescale optical excitation of neurons expressing a light-driven cation channel and pump such as Channelrhodopsin-2, and chloride pump, NpHR, respectively [18].

To date, many FRET-based indicators have been developed [19]. Generally, these

indicators use CFP and YFP as the FRET donor and acceptor, respectively. Given the similar emission spectrum between CFP and RLuc8, we could construct BRET-based indicators simply by substituting RLuc8 for the CFP moiety in the FRET-based indicators. However, this substitution of donor molecule may not yield high-performance indicators with a large dynamic range. To enhance the dynamic range, it may be necessary to use circularly permuted Venus variants to enable optimization of relative orientation between RLuc8 and Venus. Also, the development and application of circularly permuted RLuc8 variants might greatly improve BRET-based indicators in future.

Materials and Methods

Reagents

Coelenterazine-h was purchased from Promega (Madison, WI), Ni-NTA agarose from Qiagen GmbH (Hilden, Germany), Lipofectamine 2000 from Invitrogen (Carlsbad, CA), Dulbecco's Modified Eagle's Medium (DMEM) from Sigma (St. Louis, MO), ionomycin from Calbiochem (San Diego, CA) and fetal bovine serum from BioWest (Nuaille, France).

Gene construction

pGL4.70 [hRLuc] (Promega) was used as the template for introducing eight mutations, A55T/C124A/S130A/K136R/A143M/M185V/M253L/S287L, by means of a site-directed mutagenesis method to yield RLuc8 [2]. A series of BRET-based Ca²⁺ indicators was constructed by replacing ECFP and cp173Venus in YC3.60 with Venus or circularly permuted Venus variants, and RLuc8, respectively. G5A gene was constructed by fusion of

GFP gene with aequorin coding sequences through 5 times repeating SGGSGGGQ peptide coding sequence as shown previously [5].

Protein expression, purification and Ca²⁺ titration in vitro

Recombinant BRAC protein with N-terminal polyhistidine tags was expressed in *Escherichia coli* [JM109(DE3)] at 23°C, purified using an Ni-NTA column (Qiagen) and then diluted to 20 nM in 50 mM HEPES, pH 8.0. Emission spectra of BRAC were measured using a spectrophotometer (F-2500; Hitachi) and a microplate reader (SH-9000; Corona Electric). Final concentration of 1–10 μM coelenterazine-h (Promega) was used as the luminescent substrate for RLuc8. Ca²⁺ titration was performed by reciprocal dilution of Ca²⁺-free and Ca²⁺-saturated buffers prepared using *O,O'*-bis(2-aminoethyl)ethyleneglycol-*N,N,N',N'*-tetraacetic acid (EGTA), *N*-(2-Hydroxyethyl)ethylenediamine-*N,N',N'*-triacetic acid (EDTA-OH), or Nitrilotriacetic acid (NTA) in 100 mM KCl, 10 mM MOPS (pH 7.2). Free Ca²⁺ concentrations were calculated by using 0.15, 4.3 and 170 μM as the K_d value of EGTA, EDTA-OH and NTA for Ca²⁺, respectively [20]. A Ca²⁺ titration curve was used to calculate apparent K_d value by non-linear regression analysis. The averaged data from eight independent measurements were fitted to the Hill equation using Origin7 software (OriginLab).

Measurement of Ca²⁺ binding kinetics

Measurements of Ca²⁺ binding kinetics of BRAC were performed by using stopped-flow photometry system consisting of RX.2000 rapid mixing stopped-flow unit (Leatherhead,

UK) and FP-750 spectrophotometer (JASCO, Japan). Emission intensity of Venus (530 nm) from BRAC were monitored at 1 kHz just after rapid mixing of 5 nM BRAC protein with 20 μM coelenterazine-h in various concentration of Ca^{2+} buffer. In this experiment, we did not mix coelenterazine-h with BRAC prior to measurement to avoid undesirable consumption of coelenterazine-h by Rluc8 in BRAC during sample preparation. Thus, time course of emission intensity in the stopped-flow experiments consists of three components of kinetics derived from Ca^{2+} binding to BRAC, coelenterazine-h binding to BRAC, and catalytic oxidation of coelenterazine-h by BRAC. To estimate the catalytic oxidation of coelenterazine-h by Rluc8 in BRAC, we measured time course of emission intensity change after mixing BRAC with 20 μM coelenterazine-h in Ca^{2+} -free solution, and used the obtained data as a “base line”. Then, we measured time course of both association and dissociation of Ca^{2+} to and from BRAC by mixing 1 volume of BRAC in Ca^{2+} -free buffer with 25 volume of solution containing 1.69 μM Ca^{2+} , and 1 volume of BRAC in 1.69 μM Ca^{2+} solution with 25 volume of Ca^{2+} -free buffer, respectively. The averaged data from at least 5 independent measurements were used for following analysis. The averaged time course data for association and dissociation kinetics were subtracted by the base line to remove the fraction derived from autonomous catalytic oxidation of coelenterazine-h by BRAC. Then, the time constants (τ) were calculated by means of curve fitting in single exponential equation using the data from 0.2 sec to 2.0 sec to minimize contribution of signal derived from association of coelenterazine-h with BRAC just after mixing. Measurements of Ca^{2+} binding kinetics of YC3.60 were performed as shown previously [11]. In the stopped-flow experiment, final Ca^{2+} concentration was controlled by reciprocal

dilution of Ca^{2+} -free and Ca^{2+} -saturated buffers prepared using *O,O'*-bis(2-aminoethyl)ethyleneglycol-*N,N,N',N'*-tetraacetic acid (EGTA) in 100 mM KCl, 10 mM MOPS (pH 7.2). Free Ca^{2+} concentration in every solution was confirmed with Ca^{2+} -sensitive electrode (Metrohm, Herisau, Switzerland) which is calibrated with a CaCl_2 standard solution (Orion Research, Cambridge, MA).

pH titration

pH titration was performed using a series of 20 mM buffers with 100 mM KCl in pH 5.7 and 6.2 (MES), 6.6 and 7.0 (MOPS), 7.4 (Tris), 7.8 and 8.0 (HEPES), and 8.6 (Glycin).

Cell culture and transfection

Hela cells were cultured in a homemade 35-mm glass-bottom dish in DMEM (Sigma) containing 10% fetal bovine serum (BioWest). Cells were transfected with plasmids by means of Lipofectamine 2000 (Invitrogen). At 1 to 2 days after transfection, cells expressing BRAC or G5A were subjected to imaging. 10 μM coelenterazine-h were added to the culture medium just before observation of BRAC and 1-4 hours before observation of G5A.

Plant growth and pathology test conditions

We used homozygous *Arabidopsis thaliana* (ecotype Columbia, Col-0) expressing BRAC under control of the 35S promoter. Plants were grown at 22 °C under the 8-h-light in

a day for 5-6 weeks. For pathology tests, we used *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) and an avirulent strain (*Pst* DC3000/*avrRpm1*) that was kindly provided by Dr. Jeffery L. Dangl. We cultured the bacteria in KB medium with kanamycin (30 µg/ml) and rifampicin (100 µg/ml) [21]. After washing the cells twice in 10 mM MgCl₂, we suspended them at 5x10⁷ cfu/ml for pathology tests. We inoculated the suspensions into the abaxial surfaces of leaves with needleless syringes.

Live cell Imaging

An inverted microscope (Ti-E, Nikon) was used to observe BRET and G5A signals in living cells maintained at 37°C with a continuous supply of 95% air and 5% carbon dioxide by using a stage-top incubator (Tokai Hit, Fujinomiya, Japan). Light emission from the samples was focused by a 40× oil-immersion objective lens (Nikon Plan Fluor, numerical aperture 1.3) on a cooled electron-multiplying charge-coupled device (EM-CCD) camera (ImagEM; Hamamatsu Photonics). A 2×2 binning setting was used for every 1-second imaging. To simultaneously acquire both RLuc8 and Venus images, we used a W-View system (Hamamatsu Photonics) composed of 510-nm short pass and 510-nm long pass dichroic mirrors, and F01-479/40 and F01-536/40 bandpass filters (both from Semrock) to

split the RLuc8 and Venus images. Image analysis was performed using Aquacosmos (Hamamatsu Photonics). Maximum and minimum BRET ratio values in HeLa cells were measured after treatment with 5 μ M ionomycin/1 mM EGTA, and 5 μ M ionomycin/5 mM Ca^{2+} , respectively.

Live plant imaging

We used macro zoom microscopy (MVX10, Olympus) with 1 \times objective lens (Olympus MVPLAPO1X, numerical aperture 0.25) on a cooled electron-multiplying charge-coupled device (EM-CCD) camera (iXon DU-897E-BV, Andor Technology). A 1 \times 1 binning setting was used for every 10-second imaging. To simultaneously acquire both RLuc8 and Venus images, we used W-View system and mirrors as aforementioned in section of living cell imaging.

Supporting Information

Supporting Video 1

Video showing cytosolic Ca^{2+} oscillation in HeLa cells expressing BRAC in response to the application of 10 μ M histamine. Video corresponds to the data in Figure 4D. Resolution is 128x256, Quick-Time, 2.93 MB.

Acknowledgments

This work was partly supported by grants from Scientific Research on Advanced Medical Technology of the Ministry of Labor, Health and Welfare of Japan, and Precursory Research for Embryonic Science and Technology of the Japan Science and Technology Agency. We are grateful to Drs. Jeff L. Dangl at University of North Carolina and Junji Yamaguchi at Hokkaido University for donating *Pst* DC3000 strains.

Author Contributions

Conceived and designed the experiments: TN. Acquired and analyzed the data: KS KH. Wrote the paper: KS TN. Prepared plant and performed plant experiments: NH. Measure the sensor kinetics: TM and KM.

References

1. Hoshino H, Nakajima Y, Ohmiya Y (2007) Luciferase-YFP fusion tag with enhanced emission for single-cell luminescence imaging. *Nat Methods* 4: 637–639.
2. Loening AM, Fenn TD, Wu AM, Gambhir SS. (2006) Consensus guided mutagenesis of Renilla luciferase yields enhanced stability and light output. *Protein Eng Des Sel* 19: 391–400.
3. De A, Ray P, Loening AM, Gambhir SS (2009) BRET3: a red-shifted bioluminescence resonance energy transfer (BRET)-based integrated platform for imaging protein-protein interactions from single live cells and living animals. *Faseb J* 23: 2702–2709.
4. Brini M, Marsault R, Bastianutto C, Alvarez J, Pozzan T, et al. (1995) Transfected aequorin in the measurement of cytosolic CA^{2+} concentration ($[CA^{2+}]_C$). *J Biol Chem* 270: 9896–9903.

5. Baubet V, Le Mouellic H, Campbell AK, Lucas-Meunier E, Fossier P, et al. (2000) Chimeric green fluorescent protein-aequorin as bioluminescent Ca²⁺ reporters at the single-cell level. *Proc Natl Acad Sci* 97: 7260–7265.
6. Shimomura O, Kishi Y, Inouye S (1993) The relative rate of aequorin regeneration from apoaequorin and coelenterazine analogs. *Biochem J* 296: 549–551.
7. Shimomura O (2006) The jellyfish *Aequoria* and other luminous coelenterates. In: *Bioluminescence*. Singapore: World Scientific. pp. 115-116.
8. Kaihara A, Umezawa Y, Furukawa T (2008) Bioluminescent indicators for Ca²⁺ based on split Renilla luciferase complementation in living cells. *Anal Sci* 24: 1405–1408.
9. Kim SB, Sato M, Tao H (2009) Split Gaussia luciferase-based bioluminescence template for tracing protein dynamics in living cells. *Anal Chem* 81: 67–74.
10. Nagai T, Ibata K, Park ES, Kubota M, Mikoshiba K, et al. (2002) A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nat Biotechnol* 20: 87–90.
11. Miyawaki A, Llopis J, Heim R, McCaffery JM, Adams JA, et al. (1997) Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin. *Nature* 388: 882–887.
12. Nagai T, Yamada S, Tominaga T, Ichikawa M, Miyawaki A (2004) Expanded dynamic range of fluorescent indicators for Ca²⁺ by circularly permuted yellow fluorescent proteins. *Proc Natl Acad Sci* 101: 10554–10559.
13. Dacres H, Dumancic MM, Horne I, Trowell SC (2009) Direct comparison of bioluminescence-based resonance energy transfer methods for monitoring of proteolytic cleavage. *Anal Biochem* 385: 194–202.
14. Nakai J, Ohkura M, Imoto K (2001) A high signal-to-noise Ca²⁺ probe composed of a single green fluorescent protein. *Nat Biotechnol* 19: 137-141.
15. Lecourieux D, Ranjeva R, Pugin A (2006) Calcium in plant defence-singalling pathways. *New Phytologist* 171: 249-269.
16. Mackey D, Holt III BF, Wiig A, Dangl JL (2002) RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* 108: 743-754.

17. Grant M, Brown I, Adams S, Knight M, Ainslie A, et al. (2000) The *PRMI* plant disease resistance gene facilitates a rapid and sustained increase in cytosolic calcium that is necessary for the oxidative burst and hypersensitive cell death. *The Plant J* 23: 441-450.
18. Zhang F, Wang LP, Brauner M, Liewald JF, Kay K, et al. (2007) Multimodal fast optical interrogation of neural circuitry. *Nature* 446: 633–636.
19. Frommer WB, Davidson MW, Campbell RE (2009) Genetically encoded biosensors based on engineered fluorescent proteins. *Chem Soc Rev* 38: 2833–2841
20. Tsien RY, Pozzan T (1989) Measurement of cytosolic free Ca^{2+} with Quin2. *Methods Enzymol* 172: 230-262
21. Katagiri F, Thilmony R, He SY (2002) The Arabidopsis thaliana-Pseudomonas syringae interaction. In: Somerville C, Meyerowitz E, editors. *The Arabidopsis Book*. Boston: American Society for Plant Biology. pp. 1-35.

Figure Legends

Figure 1. Schematic structures of yellow cameleon 3.60 (YC3.60) and BRAC derivatives. (A) Domain structure of YC3.60 and BRAC derivatives. Circularly permuted Venus variants are indicated as cp50, 157, 173, 195, and 229Venus, which have different translation start sites from the original Venus. (B) Schematic three-dimensional structure of BRAC, which is composed of the Ca^{2+} -sensing domain, calmodulin (CaM) and M13, sandwiched between Venus and RLuc8. In the Ca^{2+} free state (left panel), CaM-M13 has an extended conformation so Venus is not located close to RLuc8 and thus only weak emission can be seen from Venus due to low BRET efficiency. Upon Ca^{2+} binding to CaM (right panel), Ca^{2+} -CaM makes a compact complex with M13, which induces efficient

BRET from RLuc8 to Venus resulting in fluorescence emission peaking at 530 nm.

Figure 2. Spectral properties of BRAC derivatives. Emission spectra of BRAC derivatives, (A-F) RLuc8-CaM-M13-Venus or cpVenus variants, (E-J) Venus or cpVenus variants-CaM-M13-RLuc8, under zero Ca^{2+} (dashed line) and saturated Ca^{2+} (solid line) conditions.

Figure 3. Chemical properties of BRAC. (A) Normalized emission spectra of BRAC at indicated free Ca^{2+} concentrations. All spectra were normalized at 480 nm. (B) Ca^{2+} titration curve of emission ratio (530 nm/480 nm) in BRAC. (C) pH titration curves of BRAC. pH titration curves of BRAC at zero (dashed line) and saturated (solid line) Ca^{2+} . (D) Time traces of luminescence intensity of BRAC after the rapid mixing of Ca^{2+} -saturated BRAC with Ca^{2+} -free buffer. A fitted curve of single-exponential function is shown as red lines. (E) The dissociation time constant at various Ca^{2+} concentration. (F) Relaxation rate constant (k_{obs}) for reaction of YC3.60 with Ca^{2+} . Association rate (k_{on}) and dissociation rate (k_{off}) can be calculated from fitting equation $k_{\text{obs}} = k_{\text{on}} [\text{Ca}^{2+}] + k_{\text{off}}$ and dissociation constant (K_{d}) can be calculated from the following equation $K_{\text{d}} = k_{\text{off}} / k_{\text{on}}$.

Figure 4. Ca^{2+} imaging of HeLa cells by using BRAC. (A) Fluorescence image of HeLa cells expressing BRAC. Emission signal from Venus (B) and RLuc8 (C). (D) A series of

pseudo-colored BRET images showing Ca^{2+} dynamics taken every 1 second. (E) Time course of averaged intensity of RLuc8 (gray line) and Venus (black line) at a cell shown in (B). (F) Time course of $\Delta R/R_0$ at the same cell. (G) Time course of averaged intensity of Hela cell expressing G5A. 10 μM histamine treatment were done at ~ 100 second. Scale bar, 20 μm .

Figure 5. Ca^{2+} imaging of plant leaves by using BRAC. A series of pseudo-colored BRET images showing Ca^{2+} dynamics in plant leaves 30 minutes (A) and 80 minutes (B) after inoculation of *Pst* DC3000/*avrRpm1* and *Pst* DC3000 (control). (C, D) Time course of $\Delta R/R_0$ at the plant leaves inoculated *Pst* DC3000/*avrRpm1* (black line) and *Pst* DC3000 (gray line). 0 min represents the time of bacteria inoculation. Scale bars, 1 mm.

Table 1. Emission ratio (530 nm/480 nm) without and with Ca²⁺ and dynamic range of newly developed constructs.

Construct	- Ca ²⁺	+ Ca ²⁺	Dynamic range (%)
Rluc8-CaM-M13-cp173Venus	0.8	0.9	10
Rluc8-CaM-M13-Venus	0.8	1.0	30
Rluc8-CaM-M13-cp50Venus	0.5	0.5	0
Rluc8-CaM-M13-cp157Venus	0.9	1.2	30
Rluc8-CaM-M13-cp195Venus	0.7	0.8	10
Rluc8-CaM-M13-cp229Venus	0.7	0.7	0
Venus-CaM-M13-Rluc8	1.1	1.8	60
cp50Venus-CaM-M13-Rluc8	1.2	1.0	20
cp157Venus-CaM-M13-Rluc8	1.1	1.4	20
cp173Venus-CaM-M13-Rluc8	1.1	1.3	10
cp195Venus-CaM-M13-Rluc8	1.1	1.2	10
cp229Venus-CaM-M13-Rluc8	1.1	1.5	30

Figure 1 Saito et al.




Figure 2 Saito et al.




Figure 3 Saito et al.




Figure 4 Saito et al.

