

Fig. 3. A: efferent sympathetic nerve stimulation-induced release of myocardial interstitial NE before and during hypothermia. †Hypothermia significantly attenuated the stimulation-induced NE release. B: efferent vagal nerve stimulation-induced release of myocardial interstitial ACh before and during hypothermia. †Hypothermia significantly attenuated the stimulation-induced ACh release.

did decrease HR from 202 beats/min (SD 24) to 179 beats/min (SD 15) ( $P < 0.05$ ) during the prestimulation, unpaced condition. MAP during the stimulation was 105 mmHg (SD 19) under normothermia and 93 mmHg (SD 33) under hypothermia.

#### DISCUSSION

A cardiac microdialysis is a powerful tool to estimate neurotransmitter levels in the myocardial interstitium in vivo (2, 3, 14, 19, 20, 31). The present study demonstrated that hypothermia significantly attenuated the myocardial interstitial releases of NE and ACh in the ischemic region during the LAD occlusion. In contrast, the increasing response in the ACh level from its baseline level and the decreasing response in the NE level from its baseline level observed in the nonischemic region were maintained under hypothermia. To our knowledge, this is the first report showing the effects of hypothermia on the myocardial interstitial releases of NE and ACh during acute myocardial ischemia in vivo. In addition, the present study showed that hypothermia significantly attenuated nerve stimulation-induced myocardial interstitial NE and ACh releases in vivo.

#### *Effects of Hypothermia on Ischemia-induced NE and ACh Releases in the Ischemic Region*

Acute myocardial ischemia causes energy depletion, which leads to myocardial interstitial NE release in the ischemic

region (Fig. 1A). The NE release can be classified as exocytotic or nonexocytotic (18, 24). Exocytotic release indicates NE release from synaptic vesicles, which normally occurs in response to nerve discharge and subsequent  $\text{Ca}^{2+}$  influx through voltage-dependent  $\text{Ca}^{2+}$  channels. On the other hand, nonexocytotic release indicates NE release from the axoplasm, such as that mediated by a reverse transport through the NE transporter. A neuronal uptake blocker, desipramine, can suppress the ischemia-induced NE release (19, 24). Whereas exocytotic release contributes to the ischemia-induced NE release in the initial phase of ischemia (within  $\sim 20$  min), carrier-mediated nonexocytotic release becomes predominant as the ischemic period is prolonged (1). Hypothermia significantly attenuated the ischemia-induced NE release (Fig. 1A). The NE level during the 45- to 60-min period of ischemia under hypothermia was  $\sim 20\%$  of that obtained under normothermia. The NE uptake transporter is driven by the  $\text{Na}^+$  gradient across the cell membrane (23). The loss of the  $\text{Na}^+$  gradient due to ischemia causes NE to be transported out of the cell by reversing the action of the NE transporter. Hypothermia inhibits the action of the NE transporter and also suppresses the intracellular  $\text{Na}^+$  accumulation (8), thereby reducing nonexocytotic NE release during ischemia. The present results are in line with an in vitro study that showed hypothermia suppressed nonexocytotic NE release induced by deprivation of oxygen and glucose (30). The present results are also consistent with a previous study from our laboratory that showed hypothermia attenuated the nonexocytotic NE release induced by ouabain, tyramine, or cyanide (16).

Acute myocardial ischemia increases myocardial interstitial ACh level in the ischemic region, as reported previously (Fig. 1B) (13). The level of ischemia-induced ACh release during 0- to 15-, 15- to 30-, 30- to 45-, or 45- to 60-min period of ischemia is comparable to that evoked by 4-min electrical stimulation of the bilateral vagi (Fig. 3B). Compared with the normothermic condition, hypothermia significantly attenuated the ischemia-induced myocardial interstitial release of ACh in the ischemic region. Our laboratory's previous study indicated that intracellular  $\text{Ca}^{2+}$  mobilization is essential for the ischemia-induced release of ACh (13). Hypothermia may have prevented the  $\text{Ca}^{2+}$  overload, thereby reducing the ischemia-induced ACh release. Alternatively, hypothermia may reduce the extent of the ischemic injury, which in turn suppressed the ischemia-induced ACh release. Because ACh has protective effects on the cardiomyocytes against ischemia (11), the suppression of ischemia-induced ACh release during hypothermia itself may be unfavorable for cardioprotection.

There is considerable controversy regarding the cardioprotective effects of  $\beta$ -adrenergic blockade during severe ischemia, with studies demonstrating a reduction of infarct size (10, 17) or no effects (7, 27). The  $\beta$ -adrenergic blockade seems effective to protect the heart only when the heart is reperfused within a certain period after the coronary occlusion. The  $\beta$ -adrenergic blockade would reduce the myocardial oxygen consumption through the reduction of HR and ventricular contractility and delay the progression of ischemic injury. Hence the infarct size might be reduced when the heart is reperfused before the ischemic damage becomes irreversible. The ischemia-induced NE release reached nearly 100 times the baseline NE level under normothermia (Fig. 1A), which by far exceeded the NE level attained by electrical stimulation of the

bilateral stellate ganglia (Fig. 3A). Because high NE levels have cardiotoxic effects (22), ischemia-induced NE release might aggravate the ischemic injury. However, catecholamine depletion by a reserpine treatment fails to reduce the infarct size (26, 29), throwing a doubt on the involvement of catecholamine toxicity in the progression of myocardial damage during ischemia. It is, therefore, most likely that the hypothermia-induced reductions in NE and ACh are the result of reduced myocardial damage or a direct effect on nerve endings.

Van den Doel et al. (28) showed that hypothermia does not abolish necrosis, but rather delays necrosis during sustained ischemia, so that hypothermia protected against infarction produced by a 30-min occlusion but not against infarction produced by a 60-min occlusion in the rat heart. At the same time, they mentioned that hypothermia was able to reduce the infarct size after a 60-min coronary occlusion in the dog, possibly because of the significant collateral flow in the canine hearts. Because the feline hearts are similar to the canine hearts in that they have considerable collateral flow compared with the rat hearts (21), hypothermia should have protected the feline heart against the 60-min coronary occlusion in the present study.

#### *Effects of Hypothermia on the NE and ACh Releases in the Nonischemic Region and on the Electrical Stimulation-induced NE and ACh Releases*

The NE and ACh levels in the nonischemic region may reflect the sympathetic and parasympathetic drives to this region. As an example, myocardial interstitial ACh levels increase during activations of the arterial baroreflex and the Bezold-Jarisch reflex (14). In the present study, acute myocardial ischemia decreased the NE level from its baseline level, whereas it increased the ACh level from its baseline level (Fig. 2). Ischemia also decreased MAP and HR (Tables 1 and 2), suggesting that the Bezold-Jarisch reflex was induced by the LAD occlusion under both normothermia and hypothermia. Taking into account the fact that electrical stimulation-induced ACh release was attenuated to ~70% (Fig. 3), similar ACh levels during ischemia imply the enhancement of the parasympathetic outflow via the Bezold-Jarisch reflex under hypothermia. These results are in line with the study by Zheng et al. (32), where pulmonary chemoreflex-induced bradycardia was maintained under hypothermia. Hypothermia increased the NE level in the nonischemic region, suggesting that sympathetic drive to this region also increased. Hypothermic stress is known to cause sympathetic activation, accompanying increases in MAP, HR, plasma NE, and epinephrine levels (4). In the present study, because the effect of hypothermia on MAP was insignificant (Table 1) and HR decreased under hypothermia (Table 2), the sympathetic activation observed in the nonischemic region might have been regional and not systemic.

Hypothermia attenuated the releases of NE and ACh in response to respective nerve stimulation to ~70% of that observed under normothermia (Fig. 3). The suppression of the exocytotic NE release by hypothermia is consistent with a previous study from our laboratory, where hypothermia attenuated the myocardial interstitial NE release in response to vena cava occlusion or to a local high  $K^+$  administration (15). The suppression of NE release by hypothermia is consistent with an

in vitro study by Kao and Westhead (12) in which catecholamine secretion from adrenal chromaffin cells induced by elevated  $K^+$  levels increased as the temperature increased from 4 to 37°C. On the other hand, because hypothermia inhibits the neuronal NE uptake, the NE concentration at the synaptic cleft is expected to be increased if the level of NE release remains unchanged. Actually, Vizi (30) demonstrated that hypothermia increased NE release in response to field stimulation in vitro. In the present study, however, the suppression of NE release might have canceled the potential accumulation of NE due to NE uptake inhibition. The present study also demonstrated that the ACh release was suppressed by hypothermia. In the rat striatum, hypothermia decreases the extracellular ACh concentration and increases the choline concentration (5). Hypothermia may inhibit a choline uptake transporter in the same manner as it inhibits a NE uptake transporter. The inhibition of the choline transporter by hypothermia may have hampered the replenishment of the available pool of ACh and thereby contributed to the suppression of the stimulation-induced ACh release.

#### *Limitations*

In *protocol 1*, because we did not measure the infarct size in the present study, the degree of myocardial protection by hypothermia was undetermined. Whether the reduction of ischemia-induced neurotransmitter release correlates with the reduction of infarct size requires further investigations. In *protocols 2* and *3*, baseline NE and ACh levels were not measured. The reduction of stimulation-induced NE and ACh release by hypothermia might be partly due to the reduction of baseline NE and ACh levels. However, because transection of the stellate ganglia (31) or vagi (3) reduces the baseline NE and ACh levels, changes in the baseline NE and ACh levels by hypothermia in *protocols 2* and *3* could not be as large as those observed under innervated conditions in *protocol 1* (Figs. 1 and 2).

In conclusion, hypothermia attenuated the ischemia-induced releases of NE and ACh in the ischemic region to ~30 and 20% of those observed under normothermia, respectively. Hypothermia also attenuated the nerve stimulation-induced releases of NE and ACh to ~70% of those observed during normothermia. In contrast, hypothermia did not affect the decreasing response in the NE level and the increasing response in the ACh level in the nonischemic region, suggesting that the Bezold-Jarisch reflex evoked by the LAD occlusion was maintained.

#### **GRANTS**

This study was supported by Health and Labour Sciences Research Grant for Research on Advanced Medical Technology, Health and Labour Sciences Research Grant for Research on Medical Devices for Analyzing, Supporting and Substituting the Function of Human Body, and Health and Labour Sciences Research Grant H18-Iryo-Ippan-023 from the Ministry of Health, Labour and Welfare of Japan; Program for Promotion of Fundamental Studies in Health Science from the National Institute of Biomedical Innovation; a grant provided by the Ichiro Kanehara Foundation; Ground-based Research Announcement for Space Utilization promoted by the Japan Space Forum; and Industrial Technology Research Grant Program 03A47075 from the New Energy and Industrial Technology Development Organization of Japan.

#### **REFERENCES**

1. Akiyama T, Yamazaki T. Norepinephrine release from cardiac sympathetic nerve endings in the in vivo ischemic region. *J Cardiovasc Pharmacol* 34: S11-S14, 1999.

2. Akiyama T, Yamazaki T, Ninomiya I. In vivo monitoring of myocardial interstitial norepinephrine by dialysis technique. *Am J Physiol Heart Circ Physiol* 261: H1643–H1647, 1991.
3. Akiyama T, Yamazaki T, Ninomiya I. In vivo detection of endogenous acetylcholine release in cat ventricles. *Am J Physiol Heart Circ Physiol* 266: H854–H860, 1994.
4. Chernow B, Lake CR, Zaritsky A, Finton CK, Casey L, Rainey TG, Fletcher JR. Sympathetic nervous system “switch off” with severe hypothermia. *Crit Care Med* 11: 677–680, 1983.
5. Damsma G, Fibiger HC. The effects of anaesthesia and hypothermia on interstitial concentrations of acetylcholine and choline in rat striatum. *Life Sci* 48: 2469–2474, 1991.
6. Duncker DJ, Klassen CL, Ishibashi Y, Herrlinger SH, Pavek T, Bache R. Effect of temperature on myocardial infarction in swine. *Am J Physiol Heart Circ Physiol* 270: H1189–H1199, 1996.
7. Genth K, Hofmann M, Hofmann M, Schaper W. The effect of  $\beta$ -adrenergic blockade on infarct size following experimental coronary occlusion. *Basic Res Cardiol* 76: 144–151, 1981.
8. Gerevich Z, Tretter L, Adam-Vizi V, Baranyi M, Kiss JP, Zelles T, Vizi ES. Analysis of high intracellular  $[Na^+]_i$ -induced release of  $[^3H]$ noradrenaline in rat hippocampal slices. *Neuroscience* 104: 761–768, 2001.
9. Hale SL, Kloner RA. Myocardial temperature in acute myocardial infarction: protection with mild regional hypothermia. *Am J Physiol Heart Circ Physiol* 273: H220–H227, 1997.
10. Jang IK, Van de Werf F, Vanhaecke J, De Geest H. Coronary reperfusion by thrombolysis and early beta-adrenergic blockade in acute experimental myocardial infarction. *J Am Coll Cardiol* 14: 1816–1823, 1989.
11. Kakinuma Y, Ando M, Kuwabara M, Katare RG, Okudela K, Kobayashi M, Sato T. Acetylcholine from vagal stimulation protects cardiomyocytes against ischemia and hypoxia involving additive nonhypoxic induction of HIF-1 $\alpha$ . *FEBS Lett* 579: 2111–2118, 2005.
12. Kao LS, Westhead EW. Temperature dependence of catecholamine secretion from cultured bovine chromaffin cells. *J Neurochem* 43: 590–592, 1984.
13. Kawada T, Yamazaki T, Akiyama T, Sato T, Shishido T, Inagaki M, Takaki H, Sugimachi M, Sunagawa K. Differential acetylcholine release mechanisms in the ischemic and non-ischemic myocardium. *J Mol Cell Cardiol* 32: 405–414, 2000.
14. Kawada T, Yamazaki T, Akiyama T, Shishido T, Inagaki M, Uemura K, Miyamoto T, Sugimachi M, Takaki H, Sunagawa K. In vivo assessment of acetylcholine-releasing function at cardiac vagal nerve terminals. *Am J Physiol Heart Circ Physiol* 281: H139–H145, 2001.
15. Kitagawa H, Akiyama T, Yamazaki T. Effects of moderate hypothermia on in situ cardiac sympathetic nerve endings. *Neurochem Int* 40: 235–242, 2002.
16. Kitagawa H, Yamazaki T, Akiyama T, Mori H, Sunagawa K. Effects of moderate hypothermia on norepinephrine release evoked by ouabain, tyramine and cyanide. *J Cardiovasc Pharmacol* 41: S111–S114, 2003.
17. Ku DD, Lucchesi BR. Effects of dimethyl propranolol (UM-272: SC-27761) on myocardial ischemic injury in the canine heart after temporary coronary artery occlusion. *Circulation* 57: 541–548, 1978.
18. Kurz T, Richardt G, Hagl S, Seyfarth M, Schömig A. Two different mechanisms of noradrenaline release during normoxia and simulated ischemia in human cardiac tissue. *J Mol Cell Cardiol* 27: 1161–1172, 1995.
19. Lameris TW, de Zeeuw S, Alberts G, Boomsma F, Duncker DJ, Verdouw PD, Veld AJ, van den Meiracker AH. Time course and mechanism of myocardial catecholamine release during transient ischemia in vivo. *Circulation* 101: 2645–2650, 2000.
20. Lameris TW, de Zeeuw S, Duncker DJ, Alberts G, Boomsma F, Verdouw PD, van den Meiracker AH. Exogenous angiotensin II does not facilitate norepinephrine release in the heart. *Hypertension* 40: 491–497, 2002.
21. Maxwell MP, Hearse DJ, Yellon DM. Species variation in the coronary collateral circulation during regional myocardial ischaemia: a critical determinant of the rate of evolution and extent of myocardial infarction. *Cardiovasc Res* 21: 737–746, 1987.
22. Rona G. Catecholamine cardiotoxicity. *J Mol Cell Cardiol* 17: 291–306, 1985.
23. Schwartz JH. Neurotransmitters. In: *Principles of Neural Science* (4th Ed.), edited by Kandel ER, Schwartz JH, Jessell TM. New York: McGraw-Hill, 2000, p. 280–297.
24. Schömig A, Kurz T, Richardt G, Schömig E. Neuronal sodium homeostasis and axoplasmic amine concentration determine calcium-independent noradrenaline release in normoxic and ischemic rat heart. *Circ Res* 63: 214–226, 1988.
25. Simkhovich BZ, Hale SL, Kloner RA. Metabolic mechanism by which mild regional hypothermia preserves ischemic tissue. *J Cardiovasc Pharmacol Ther* 9: 83–90, 2004.
26. Toombs CF, Wiltse AL, Shebuski RJ. Ischemic preconditioning fails to limit infarct size in reserpinized rabbit myocardium. Implication of norepinephrine release in the preconditioning effect. *Circulation* 88: 2351–2358, 1993.
27. Torr S, Drake-Holland AJ, Main M, Hynd J, Isted K, Noble MIM. Effects on infarct size of reperfusion and pretreatment with  $\beta$ -blockade and calcium antagonists. *Basic Res Cardiol* 84: 564–582, 1989.
28. Van den Doel MA, Gho BC, Duval SY, Schoemaker RG, Duncker DJ, Verdouw PD. Hypothermia extends the cardioprotection by ischaemic preconditioning to coronary artery occlusions of longer duration. *Cardiovasc Res* 37: 76–81, 1998.
29. Vander Heide RS, Schwartz LM, Jennings RB, Reimer KA. Effect of catecholamine depletion on myocardial infarct size in dogs: role of catecholamines in ischemic preconditioning. *Cardiovasc Res* 30: 656–662, 1995.
30. Vizi ES. Different temperature dependence of carrier-mediated (cytoplasmic) and stimulus-evoked (exocytotic) release of transmitter: a simple method to separate the two types of release. *Neurochem Int* 33: 359–366, 1998.
31. Yamazaki T, Akiyama T, Kitagawa H, Takauchi Y, Kawada T, Sunagawa K. A new, concise dialysis approach to assessment of cardiac sympathetic nerve terminal abnormalities. *Am J Physiol Heart Circ Physiol* 272: H1182–H1187, 1997.
32. Zheng F, Kidd C, Bowser-Riley F. Effects of moderate hypothermia on baroreflex and pulmonary chemoreflex heart rate response in decerebrate ferrets. *Exp Physiol* 81: 409–420, 1996.

## Angiotensin II attenuates myocardial interstitial acetylcholine release in response to vagal stimulation

Toru Kawada,<sup>1</sup> Toji Yamazaki,<sup>2</sup> Tsuyoshi Akiyama,<sup>2</sup> Meihua Li,<sup>1,3</sup> Can Zheng,<sup>1,3</sup> Toshiaki Shishido,<sup>1</sup> Hidezo Mori,<sup>2</sup> and Masaru Sugimachi<sup>1</sup>

<sup>1</sup>Department of Cardiovascular Dynamics, Advanced Medical Engineering Center and <sup>2</sup>Department of Cardiac Physiology, National Cardiovascular Center Research Institute, Osaka; and <sup>3</sup>Japan Association for the Advancement of Medical Equipment, Tokyo, Japan

Submitted 5 April 2007; accepted in final form 19 July 2007

**Kawada T, Yamazaki T, Akiyama T, Li M, Zheng C, Shishido T, Mori H, Sugimachi M.** Angiotensin II attenuates myocardial interstitial acetylcholine release in response to vagal stimulation. *Am J Physiol Heart Circ Physiol* 293: H2516–H2522, 2007. First published July 20, 2007; doi:10.1152/ajpheart.00424.2007.—Although ANG II exerts a variety of effects on the cardiovascular system, its effects on the peripheral parasympathetic neurotransmission have only been evaluated by changes in heart rate (an effect on the sinus node). To elucidate the effect of ANG II on the parasympathetic neurotransmission in the left ventricle, we measured myocardial interstitial ACh release in response to vagal stimulation (1 ms, 10 V, 20 Hz) using cardiac microdialysis in anesthetized cats. In a control group ( $n = 6$ ), vagal stimulation increased the ACh level from  $0.85 \pm 0.03$  to  $10.7 \pm 1.0$  (SE) nM. Intravenous administration of ANG II at  $10 \mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  suppressed the stimulation-induced ACh release to  $7.5 \pm 0.6$  nM ( $P < 0.01$ ). In a group with pretreatment of intravenous ANG II receptor subtype 1 (AT<sub>1</sub> receptor) blocker losartan (10 mg/kg,  $n = 6$ ), ANG II was unable to inhibit the stimulation-induced ACh release ( $8.6 \pm 1.5$  vs.  $8.4 \pm 1.7$  nM). In contrast, in a group with local administration of losartan (10 mM,  $n = 6$ ) through the dialysis probe, ANG II inhibited the stimulation-induced ACh release ( $8.0 \pm 0.8$  vs.  $5.8 \pm 1.0$  nM,  $P < 0.05$ ). In conclusion, intravenous ANG II significantly inhibited the parasympathetic neurotransmission through AT<sub>1</sub> receptors. The failure of local losartan administration to nullify the inhibitory effect of ANG II on the stimulation-induced ACh release indicates that the site of this inhibitory action is likely at parasympathetic ganglia rather than at postganglionic vagal nerve terminals.

cardiac microdialysis; cats; losartan

ANG II HAS a variety of effects on the cardiovascular system (22): it acts on the vascular beds to increase peripheral vascular resistance and also on the adrenal cortex to cause volume retention. These direct effects of ANG II contribute to the maintenance of arterial pressure (AP). Aside from these direct effects, ANG II has been shown to modulate the sympathetic nervous system both centrally (7, 9) and peripherally (10). With respect to the sympathetic regulation in the heart, however, exogenous ANG II does not facilitate stimulation- and ischemia-induced norepinephrine release in the porcine left ventricle (18). Compared with a number of reports on the sympathetic system, only a few reports are available as to the effects of ANG II on the parasympathetic system. In 1982, Potter (23) demonstrated that ANG II (5–10  $\mu\text{g}$  iv, body wt not

reported) inhibited bradycardia induced by vagal stimulation in dogs. In that study, administration of ACh reduced the heart rate to an identical degree in the presence or absence of ANG II, suggesting that the inhibition of bradycardia by ANG II was attributable to the inhibition of the ACh release from the vagal nerve terminals. In contrast, Andrews et al. (3) reported that ANG II (500 ng/kg iv) did not inhibit bradycardia induced by vagal stimulation in ferrets. In a rat heart failure model, ANG II receptor subtype 1 (AT<sub>1</sub> receptor) antagonist losartan enhanced the bradycardic response to vagal stimulation (5). In pithed rats, an angiotensin-converting enzyme (ACE) inhibitor captopril also enhanced the bradycardic response to vagal stimulation (25, 26). In all of these studies, changes in the heart rate were used as a functional measurement of peripheral vagal function because of the difficulty in measuring the ACh release in the in vivo heart. Accordingly, whether ANG II affects the vagal control over the ventricle remains unknown. The aim of the present study was to examine the effect of ANG II on the vagal stimulation-induced ACh release in the left ventricular myocardium by measuring the interstitial ACh levels directly using a cardiac microdialysis technique (1, 13–15). We also explored the possible sites of action for the effect of ANG II on the stimulation-induced ACh release by administering losartan systemically from the femoral vein or locally through the dialysis fiber. Because ACh has a protective effect on the ischemic myocardium (12, 24, 29), elucidating the effect of ANG II on the ACh release in the ventricle would be helpful to understand the mechanism of ACE inhibitor or AT<sub>1</sub> receptor antagonist for the treatment of heart diseases (16, 17).

### MATERIALS AND METHODS

#### Surgical Preparation

Animal care was provided in strict accordance with the *Guiding Principles for the Care and Use of Animals in the Field of Physiological Sciences* approved by the Physiological Society of Japan. All protocols were approved by the Animal Subject Committee of the National Cardiovascular Center. Twenty eight adult cats weighing from 1.9 to 4.9 kg were anesthetized using an intraperitoneal injection of pentobarbital sodium (30–35 mg/kg) and were then ventilated mechanically with room air mixed with oxygen. The depth of anesthesia was maintained by a continuous intravenous infusion of pentobarbital sodium ( $1\text{--}2 \text{ mg}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ ) through a catheter inserted in the right femoral vein. Systemic AP was monitored by a catheter inserted in the right femoral artery. Heart rate was determined from an

Address for reprint requests and other correspondence: T. Kawada, Dept. of Cardiovascular Dynamics, National Cardiovascular Center Research Institute, 5-7-1 Fujishirodai, Suita, Osaka 565-8565, Japan (e-mail: torukawa@res.nccv.go.jp).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

electrocardiogram using a cardi tachometer. Esophageal temperature of the animal, measured using a thermometer (CTM-303; TERUMO), was maintained at  $\sim 37^{\circ}\text{C}$  using a heating pad and a lamp. Both vagal nerves were exposed and sectioned bilaterally through a midline cervical incision. With the animal in the lateral position, we resected the left fifth and sixth ribs to approach the heart. After the incision of the pericardium, the heart was suspended in a pericardial cradle. Stainless steel wires were attached to the apex and the posterior wall of the left ventricle to pace the heart. Using a fine guiding needle, we implanted a dialysis probe transversely through the anterolateral free wall of the left ventricle. Next, we attached a pair of bipolar platinum electrodes to the cardiac end of each sectioned vagal nerve. The nerves and electrodes were covered in warmed mineral oil for insulation. We gave heparin sodium (100 U/kg) intravenously to prevent blood coagulation. At the end of the experiment, postmortem examination confirmed that the semipermeable membrane of the dialysis probe had been implanted in the left ventricular myocardium.

#### Dialysis Technique

The materials and properties of the dialysis probe have been described previously (1). Briefly, we designed a transverse dialysis probe in which a dialysis fiber of semipermeable membrane (13 mm length,  $310\ \mu\text{m}$  outer diameter,  $200\ \mu\text{m}$  inner diameter; PAN-1200, 50,000 mol wt cutoff; Asahi Chemical) was attached at both ends to

polyethylene tubes (25 cm length,  $500\ \mu\text{m}$  outer diameter,  $200\ \mu\text{m}$  inner diameter). The dialysis probe was perfused at a rate of  $2\ \mu\text{l}/\text{min}$  with Ringer solution containing the cholinesterase inhibitor physostigmine ( $100\ \mu\text{M}$ ). Experimental protocols were started 2 h after implanting the dialysis probe when the ACh concentration in the dialysate reached a steady state. ACh concentrations in the dialysate were measured by an HPLC system with electrochemical detection (Eicom, Kyoto, Japan).

Figure 1 schematizes the three original protocols and two supplemental protocols utilized in the present study. The hatched rectangle indicates the baseline sampling, whereas the solid rectangles indicate the sampling during the 10-min vagal stimulation period (1 ms, 10 V, 20 Hz) in each protocol. The stimulus was set supramaximal to most easily delineate the possible effect of ANG II on myocardial interstitial ACh release. In all of the vagal stimulation periods, we paced the heart at 200 beats/min to avoid the difference in heart rate affecting the vagal stimulation-induced ACh release (14). For baseline sampling periods, we paced the heart at 200 beats/min when spontaneous heart rate was  $< 200$  beats/min.

**Protocol 1** ( $n = 6$ ). We examined the effects of intravenous administration of ANG II on vagal stimulation-induced myocardial ACh release. We collected a dialysate sample under baseline conditions. We then stimulated the vagal nerve and paced the heart for 10 min and collected a dialysate sample during the stimulation period




Fig. 1. Schematic representation of the protocols used in the present study. After implantation of the dialysis probe (2 h), we obtained a baseline dialysate sample (hatched rectangles) for 10 min. Thereafter, we obtained 4 dialysate samples during vagal stimulation with fixed-rate pacing for 10 min (filled rectangles) at intervening intervals of 15 min. In *protocols 1* through *3*, after obtaining 2 control trials ( $VS_1$  and  $VS_2$ ), we initiated intravenous administration of ANG II at  $10\ \mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  and waited for 15 min to obtain a dialysate sample during vagal stimulation with fixed-rate pacing (A10-VS). We then increased the dose of ANG II to  $100\ \mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  and waited for an additional 15 min before obtaining a dialysate sample during vagal stimulation with fixed-rate pacing (A100-VS). In *protocol 2*, the ANG II receptor subtype 1 blocker losartan was administered by bolus injection (10 mg/kg) before obtaining a baseline dialysate sample and also immediately before the beginning of each dose of ANG II administration ( $\blacktriangledown$ ). In *protocol 3*, we administered losartan (10 mM) through the dialysis probe throughout the protocol. In *protocol 4*, we first collected a dialysate sample using perfusate free of physostigmine. We then replaced the perfusate with Ringer solution containing physostigmine and collected dialysate samples of baseline and vagal stimulation (VS). Approximately 15 min after the onset of iv ANG II administration at  $10\ \mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ , we collected dialysate samples of baseline (A10-baseline) and vagal stimulation (A10-VS). In *protocol 5*, we collected dialysate samples during a control vagal stimulation (VS) and during the 2 doses of iv ANG II administration (A10-VS and A100-VS). The pressor effect of ANG II was counteracted by simultaneous iv infusion of the L-type  $\text{Ca}^{2+}$  channel blocker nifedipine.

(VS<sub>1</sub>). After an intervening interval of 15 min, we repeated the 10-min vagal stimulation with fixed-rate pacing and collected another dialysate sample (VS<sub>2</sub>). After performing these two control trials, we began intravenous administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . Approximately 15 min after the onset of the ANG II administration, we collected a dialysate sample (A10-VS) during 10-min vagal stimulation with fixed-rate pacing. We then increased the dose of ANG II at 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . Approximately 15 min after the onset of the higher-dose ANG II administration, we collected a final dialysate sample (A100-VS) during 10-min vagal stimulation with fixed-rate pacing.

**Protocol 2** ( $n = 6$ ). We examined whether the intravenous AT<sub>1</sub> receptor antagonist losartan would block the effects of ANG II on the vagal stimulation-induced myocardial ACh release. We infused losartan potassium intravenously at 10 mg/kg and waited for ~15 min. We then collected baseline, VS<sub>1</sub>, and VS<sub>2</sub> samples with an intervening interval of 15 min, as described in *protocol 1*. Next, after an additional bolus injection of losartan potassium at 10 mg/kg, we began intravenous infusion of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . After ~15 min, we obtained a dialysate sample of A10-VS. Finally, after another bolus injection of losartan potassium at 10 mg/kg, we began intravenous infusion of ANG II at 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . After an additional 15 min, we obtained a dialysate sample of A100-VS.

**Protocol 3** ( $n = 6$ ). We examined whether local administration of losartan would block the effects of ANG II on the vagal stimulation-induced myocardial ACh release. We perfused the dialysis probe with Ringer solution containing 10 mM of losartan potassium. Taking into account the distribution across the semipermeable membrane, we administered losartan at a concentration >400 times higher than that for intravenous administration in *protocol 2*. Because local administrations of larger molecules such as  $\omega$ -conotoxin GVIA (molecular weight 3037) and  $\omega$ -conotoxin MVIIC (mol wt 2,749) were able to suppress vagal stimulation-induced ACh release in our previous study (15), it would be reasonable to assume that losartan potassium (mol wt 461) should have spread in the vicinity of the dialysis fiber, from which the dialysate was collected. Using the same procedures as described in *protocol 1*, we obtained dialysate samples for baseline, VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS. A previous study indicated that ACh measured by cardiac microdialysis in the left ventricle mainly reflected ACh released from the postganglionic nerve terminals and not from the parasympathetic ganglia (1 and see DISCUSSION for details).

**Protocol 4** ( $n = 5$ ). To examine the effects of ANG II on the baseline ACh level, we performed an additional protocol where the baseline ACh level was measured during intravenous infusion of




Fig. 2. Changes in dialysate ACh concentrations obtained from *protocol 1*. Vagal stimulation significantly increased the ACh levels. There was no significant difference in the ACh level between the 2 control trials (VS<sub>1</sub> and VS<sub>2</sub>). The ACh level was significantly lower in A10-VS and A100-VS compared with that measured in VS<sub>1</sub> and VS<sub>2</sub>. There was no significant difference in the ACh level between A10-VS and A100-VS. Values are presented as mean and SE. \* $P < 0.01$  by Tukey's test.




Fig. 3. Changes in dialysate ACh concentrations obtained from *protocol 2*. Vagal stimulation significantly increased the ACh levels. There was no significant difference in the ACh level among the 4 dialysate samples during vagal stimulation (VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS). Values are presented as means and SE. \* $P < 0.01$  by Tukey's test.

ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  (A10-baseline). In this protocol, we also obtained a dialysate sample using the perfusate without the cholinesterase inhibitor physostigmine before the usual dialysate sampling using the perfusate containing physostigmine.

**Protocol 5** ( $n = 5$ ). To avoid the pressor effect of ANG II, we administered an L-type Ca<sup>2+</sup> channel blocker nifedipine (0.5–2.0  $\text{mg}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ ) simultaneously with ANG II and obtained dialysate samples for VS, A10-VS, and A100-VS. In a previous study, intravenous administration of an L-type Ca<sup>2+</sup> channel blocker alone did not affect the vagal stimulation-induced myocardial ACh release significantly (15).

#### Statistical Analysis

All data are presented as mean  $\pm$  SE values. In *protocols 1* through *3*, myocardial interstitial ACh levels were compared among baseline, VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS samples using a repeated-measures ANOVA (8). When there was a significant difference, Tukey's test for all-pairwise comparisons was applied to identify the differences between any two of the samples. Differences were considered significant at  $P < 0.05$ . The mean AP value in the last 1 min of the 10-min vagal stimulation period was treated as the AP value during vagal stimulation. The AP data were compared using a repeated-measures ANOVA among baseline, during the two control stimulations (VS<sub>1</sub> and VS<sub>2</sub>), and before and during vagal stimulation under the two different doses of intravenous ANG II administrations. When there was a significant difference, Dunnett's test for comparison against a single control was applied to identify differences from the baseline value. Differences were considered significant at  $P < 0.05$ . In




Fig. 4. Changes in dialysate ACh concentrations obtained from *protocol 3*. Vagal stimulation significantly increased the ACh levels. There was no significant difference in the ACh level between the 2 control trials (VS<sub>1</sub> and VS<sub>2</sub>). The ACh level was significantly lower in A10-VS and A100-VS compared with that measured in VS<sub>1</sub> and VS<sub>2</sub>. There was no significant difference in the ACh level between A10-VS and A100-VS. Values are presented as means and SE. † $P < 0.05$  and \* $P < 0.01$  by Tukey's test.

Table 1. Mean arterial pressure values before vagal stimulation and during the last 1 min of stimulation

	Baseline	VS <sub>1</sub>	VS <sub>2</sub>	A10	A10-VS	A100	A100-VS
Protocol 1	102 ± 11	93 ± 17	91 ± 17	132 ± 9†	105 ± 19	129 ± 13†	105 ± 21
Protocol 2	102 ± 17	71 ± 16*	69 ± 16*	80 ± 15	68 ± 17*	86 ± 19	72 ± 18*
Protocol 3	102 ± 13	100 ± 17	92 ± 17	139 ± 11*	120 ± 19	147 ± 11*	122 ± 21

Data are means ± SE obtained from baseline, two control trials (VS<sub>1</sub> and VS<sub>2</sub>), before (A10) and during (A10-VS) vagal stimulation under iv administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ , and before (A100) and during (A100-VS) vagal stimulation under iv administration of ANG II at 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . The heart was paced at 200 beats/min whenever vagal stimulation was applied. † $P < 0.05$  and \* $P < 0.01$  from the respective baseline values by Dunnett's test.

protocol 4, the baseline ACh levels were compared before and during the ANG II administration using a paired *t*-test. The ACh levels during vagal stimulation were also compared before and during ANG II administration using a paired *t*-test. In protocol 5, the ACh levels and the mean AP values were compared among VS, A10-VS, and A100-VS using a repeated-measures ANOVA followed by Tukey's test.

## RESULTS

In protocol 1, vagal stimulation significantly increased myocardial interstitial ACh levels (Fig. 2). There was no significant difference between two control trials with an intervening interval of 15 min [VS<sub>1</sub>: 10.7 ± 1.0 (SE) nM and VS<sub>2</sub>: 9.9 ± 0.9 (SE) nM]. Intravenous administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  significantly attenuated the vagal stimulation-induced ACh release (A10-VS: 7.5 ± 0.6 nM) to ~71% of VS<sub>1</sub>. Although the intravenous administration of ANG II at 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  also significantly attenuated the vagal stimulation-induced ACh release (A100-VS: 7.3 ± 0.9 nM) to ~68% of VS<sub>1</sub>, the ACh levels were not different from those of A10-VS.

In protocol 2, vagal stimulation significantly increased myocardial interstitial ACh levels under control stimulations (VS<sub>1</sub>: 8.6 ± 1.5 nM and VS<sub>2</sub>: 8.7 ± 2.0 nM; Fig. 3). With a pretreatment of intravenous losartan, intravenous ANG II was unable to suppress the vagal stimulation-induced ACh release (A10-VS: 8.4 ± 1.7 nM and A100-VS: 7.1 ± 1.4 nM). Although the mean level of ACh tended to be lower in A100-VS compared with VS<sub>1</sub> or VS<sub>2</sub>, the differences were not statistically significant.

In protocol 3, vagal stimulation significantly increased myocardial interstitial ACh levels under control stimulations (VS<sub>1</sub>: 8.0 ± 0.8 nM and VS<sub>2</sub>: 7.9 ± 0.8 nM; Fig. 4). Intravenous

ANG II at either 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  or 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  significantly suppressed the vagal stimulation-induced ACh release to ~72% (A10-VS: 5.8 ± 1.0 nM) and 62% (A100-VS: 5.0 ± 0.7 nM of that seen in VS<sub>1</sub>), respectively.

In protocol 1, the AP values before the vagal stimulation during the intravenous ANG II administrations (A10 and A100) were significantly higher than the baseline AP value (Table 1). The AP values during vagal stimulation (VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS) were not different from the baseline AP value. In protocol 2, the AP value before the first administration of losartan was 126 ± 14 mmHg. The AP values before the vagal stimulation during the intravenous ANG II administrations (A10 and A100) were not significantly different from the baseline AP value. The AP values during vagal stimulation (VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS) were significantly lower than the baseline AP value. In protocol 3, the AP values before vagal stimulation during the intravenous ANG II administrations (A10 and A100) were significantly higher than the baseline AP value. The AP values during vagal stimulation (VS<sub>1</sub>, VS<sub>2</sub>, A10-VS, and A100-VS) did not differ statistically from the baseline AP value.

Figure 5 illustrates typical chromatograms obtained from one animal in protocol 4. The baseline ACh level was below the limit of determination (0.5 nM) when the perfusate did not contain physostigmine (physostigmine-free baseline). The baseline ACh level was above the limit of determination when perfusate contained 100  $\mu\text{M}$  physostigmine. (This perfusate was usually used for the ACh measurements.) Vagal stimulation increased the ACh level (VS). The intravenous administration of ANG II did not affect the baseline ACh level (A10-baseline) but significantly attenuated the ACh level during vagal stimulation (A10-VS).




Fig. 5. Typical chromatograms for the ACh measurements obtained from protocol 4. ACh was less than the limit of determination when perfusate did not contain physostigmine (physostigmine-free baseline). The baseline ACh level was above the limit of determination when perfusate contained 100  $\mu\text{M}$  physostigmine. (This perfusate was usually used for the ACh measurements.) Vagal stimulation increased the ACh level (VS). The administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  did not affect baseline ACh level (A10-baseline) but significantly attenuated the vagal stimulation-induced ACh release (A10-VS). See Table 2 for pooled data of ACh levels. ND, not detected.

Table 2. Mean arterial pressure values and ACh concentrations obtained in protocol 4

	Physostigmine-free Baseline	Baseline	VS	A10-Baseline	A10-VS
ACh, nM	Not detected	1.6±0.4	10.6±2.4	1.7±0.5	7.8±2.1*
Mean arterial pressure, mmHg	111±11	109±12	103±6	148±3*	118±6

Data are means ± SE obtained from physostigmine-free baseline, baseline, control vagal stimulation (VS), and baseline (A10-baseline) and vagal stimulation (A10-VS) under iv administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ . There was no significant difference in the ACh level between baseline and A10-baseline by a paired-*t*-test. The ACh level was significantly lower in A10-VS than in VS by a paired-*t*-test. Mean arterial pressure was significantly higher in A10-baseline compared with the physostigmine-free baseline value by Dunnett's test. \**P* < 0.01.

In protocol 5, the pressor effect of ANG II was counteracted by the simultaneous intravenous infusion of nifedipine (Table 3). Under this condition, the intravenous administration of ANG II significantly attenuated the stimulation-induced ACh level to ~83% (A10-VS) and 72% (A100-VS) of that seen in VS.

## DISCUSSION

The present study demonstrated that intravenous ANG II significantly inhibited the vagal stimulation-induced myocardial interstitial ACh release in the left ventricle in anesthetized cats. Intravenous administration of losartan abolished the inhibitory effect of ANG II on the stimulation-induced ACh release, suggesting that the inhibitory action of ANG II was mediated by AT<sub>1</sub> receptors.

### Inhibitory Effect of ANG II on Myocardial Interstitial ACh Release

Only a few reports have focused on the modulatory effects of ANG II on the parasympathetic nervous system (3, 5, 25, 26), all of which have used the heart rate reduction in response to vagal stimulation as a functional measurement to assess the peripheral vagal function. Although ANG II has been shown to inhibit the ACh release in the rat entorhinal cortex in vitro (4), the direct evidence for the inhibitory effect of ANG II on the ACh release in the peripheral vagal neurotransmission in vivo has been lacking. The present study demonstrated that intravenous ANG II inhibited the vagal nerve stimulation-induced ACh release in the left ventricle in vivo (Fig. 2). As for the sympathetic system in the heart, Lameris et al. (18) have previously demonstrated that ANG II does not affect the sympathetic nerve stimulation-induced norepinephrine release. The in-

significant effect of ANG II on the sympathetic neurotransmission and the inhibitory effect of ANG II on the parasympathetic neurotransmission may provide the basis for a study by Takata et al. (26) in which ACE inhibitor enhanced cardiac vagal but not sympathetic neurotransmission.

An increased activity of the renin-angiotensin system is common in chronic heart failure and has been considered to be a stimulus for aggravation of the disease. Inhibition of the renin-angiotensin system by ACE inhibitors or by AT<sub>1</sub> receptor blockers can prevent the ventricular remodeling and improve the survival rate (16, 17), suggesting that ANG II is indeed involved in the aggravation of heart failure. ACh, on the other hand, can exert a cardioprotective effect against myocardial ischemia in several experimental settings (12, 24, 29). If ANG II inhibits the peripheral vagal neurotransmission, blockade of ANG II would increase the vagal effect on the heart. Actually, Du et al. (5) demonstrated that losartan enhanced bradycardia induced by vagal stimulation in rats with chronic myocardial infarction. In that study, however, the ventricular effect of vagal stimulation was not assessed. The results of the present study indicate that ANG II inhibited the vagal neurotransmission in the ventricle. Blockade of ANG II is therefore expected to increase the vagal effect on the ventricular myocardium when the vagal outflow from the central nervous system is unchanged. Although no literatures appear to be available as to the chronic effect of ACh on the prognosis of heart failure, electrical vagal stimulation was able to improve the survival rate of chronic heart failure in rats (19). In that study, the magnitude of the vagal stimulation was such that the heart rate decreased only by 20–30 beats/min in rats, suggesting that a modest increase in vagal tone would be sufficient to produce a cardioprotective effect. It is plausible that blockade of ANG II yields beneficial effects on chronic heart failure not only by antagonizing the sympathetic effects but also by enhancing the vagal effects on the ventricle.

Vagal stimulation was able to reduce the left ventricular contractility as assessed by end-systolic elastance only when sympathetic stimulation coexisted (20), suggesting that the effect of vagal stimulation on ventricular contractility would be secondary to sympathoinhibition. Accordingly, contribution of the inhibitory effect of ANG II on the stimulation-induced ACh release to the physiological regulation of ventricular contractility might be marginal. We think that the finding is important as a peripheral mechanism of vagal withdrawal in heart diseases accompanying the activation of the renin-angiotensin system.

Table 3. Mean arterial pressure values and ACh concentrations obtained in protocol 5

	VS	A10-VS	A100-VS
ACh, nM	12.7±1.1	10.6±1.1†	9.2±1.5*
Mean arterial pressure, mmHg	83.4±12.2	68.4±6.3	70.4±9.5

Data are means ± SE from a control vagal stimulation trial (VS), during vagal stimulation under iv administration of ANG II at 10  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  (A10-VS) and during vagal stimulation under iv administration of ANG II at 100  $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$  (A100-VS). The heart was paced at 200 beats/min during vagal stimulation. In this protocol, the pressor effect of ANG II was counteracted by simultaneous iv administration of the L-type Ca<sup>2+</sup> channel blocker nifedipine (0.5–2  $\text{mg}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ ). †*P* < 0.05 and \**P* < 0.01 from the VS group by Tukey's test. There was no significant difference between A10-VS and A100-VS in the ACh level. There were no significant differences in mean arterial pressure among the three trials.



*Possible Site of the Inhibitory Action of ANG II on ACh Release*

In *protocol 3*, we examined whether local administration of losartan was able to nullify the inhibitory effect of ANG II on the vagal stimulation-induced ACh release. The utility of local administration of pharmacological agents through the dialysis fiber has been confirmed previously. As an example, local administration the Na<sup>+</sup> channel inhibitor tetrodotoxin through the dialysis fiber completely blocked the nerve stimulation-induced ACh release (14). With respect to the source for ACh, intravenous administration of the nicotinic antagonist hexamethonium bromide completely blocked the stimulation-induced ACh release, whereas local administration of hexamethonium bromide did not, suggesting the lack of parasympathetic ganglia in the vicinity of dialysis fiber (1). In support of our interpretation, a neuroanatomic finding indicates that three ganglia, away from the left anterior free wall targeted by the dialysis probe, provide the major source of left ventricular postganglionic innervation in cats (11). Therefore, the myocardial interstitial ACh measured by cardiac microdialysis in the left ventricle mainly reflects the ACh release from the postganglionic vagal nerve terminals. The results of *protocol 3* indicate that losartan spread around the postganglionic vagal nerve terminals failed to abolish the inhibitory effect of ANG II on the stimulation-induced ACh release. Because intravenous administration of losartan was able to abolish the inhibitory effect of ANG II on the stimulation-induced ACh release (*protocol 2*), the site of this inhibitory action is likely at parasympathetic ganglia rather than at postganglionic vagal nerve terminals. The fact that AT<sub>1</sub> receptors are rich in parasympathetic ganglia (2) would support our interpretation.

ANG II has a direct vasoconstrictive effect on the coronary artery (30). At the same time, however, the intravenous administration of ANG II tended to increase mean AP during vagal stimulation by ~15 mmHg in *protocol 1* (Table 1). Although it was statistically insignificant, if this increase in mean AP increased cardiac oxygen demand, the coronary blood flow might have been increased (27), resulting in an increased rate of washout in the myocardial tissue. The possibility cannot be ruled out that such a washout mechanism contributed to the reduction of stimulation-induced ACh release during ANG II administration. However, the baseline ACh level was not decreased by ANG II in *protocol 4*, suggesting that the washout rate did not increase significantly. In addition, even when the pressor effect of ANG II was counteracted by nifedipine, ANG II was still able to inhibit the vagal stimulation-induced ACh release in *protocol 5*. Therefore, we think that the change in washout rate was not a principal mechanism for the reduction of stimulation-induced ACh release by ANG II.

The mechanisms for the baseline ACh release under the vagotomized condition were not identified in the present study. In the motor nerve terminals, a so-called nonquantal release of ACh is documented, which is independent of nerve activity (6). Incorporation of the vesicular transport system in the membrane of the nerve terminals during an exocytosis process is considered to be responsible for the mechanism of nonquantal ACh release. A similar mechanism might contribute to the baseline ACh release in the vagal nerve terminals.

Several limitations need to be addressed. First, the dose of ANG II might have increased the plasma ANG II concentration

beyond the physiological range. In this regard, the observed effect might be rather pharmacological or pathological than physiological. Nevertheless, because there are local synthesis and degradation of ANG II in the heart (21, 28), the inhibition of ACh release by ANG II could operate locally in the heart. Second, whether ANG II inhibited the ACh release from the preganglionic nerve terminals or it suppressed the excitability of the postganglionic nerve fibers to ACh was not identified in the present study. Third, the involvement of ANG II receptor subtype 2 (AT<sub>2</sub> receptor) in the modulation of peripheral parasympathetic neurotransmission was not examined in the present study because intravenous losartan was able to abolish the inhibitory effect of ANG II on the stimulation-induced ACh release. However, if coactivation of AT<sub>1</sub> and AT<sub>2</sub> receptors is required for the inhibitory effect of ANG II, blockade of AT<sub>2</sub> receptors would also abolish the inhibitory effect. Fourth, we tested just one level of vagal stimulation. Whether the effect of ANG II on the stimulation-induced ACh release depends on the vagal stimulation intensity remains to be resolved.

In conclusion, intravenous ANG II reduced the vagal nerve stimulation-induced ACh release in the left ventricle. Intravenous losartan abolished the inhibitory effect of ANG II on the stimulation-induced ACh release, suggesting that this inhibition was mediated by AT<sub>1</sub> receptors. Because local administration of losartan via dialysis fiber was unable to nullify the inhibitory effect of ANG II on the stimulation-induced ACh release, the site of this inhibitory action is likely parasympathetic ganglia. The present results imply that the beneficial effects of ACE inhibitors and AT<sub>1</sub> receptor antagonists in the treatment of heart diseases may include not only the suppression of sympathetic activity but also the enhancement of vagal activity to the ventricle.

#### GRANTS

This study was supported by a Health and Labour Sciences Research Grant for Research on Advanced Medical Technology, a Health and Labour Sciences Research Grant for Research on Medical Devices for Analyzing, Supporting, and Substituting the Function of Human Body, and Health and Labour Sciences Research Grant H18-Iryo-Ippan-023 from the Ministry of Health, Labour, and Welfare of Japan.

#### REFERENCES

1. Akiyama T, Yamazaki T, Ninomiya I. In vivo detection of endogenous acetylcholine release in cat ventricles. *Am J Physiol Heart Circ Physiol* 266: H854–H860, 1994.
2. Allen AM, Zhuo J, Mendelsohn FA. Localization and function of angiotensin AT<sub>1</sub> receptors. *Am J Hypertens* 13: 31S–38S, 2000.
3. Andrews PL, Dutia MB, Harris PJ. Angiotensin II does not inhibit vagally-induced bradycardia or gastric contractions in the anaesthetized ferret. *Br J Pharmacol* 82: 833–837, 1984.
4. Barnes JM, Barnes NM, Costall B, Horovitz ZP, Naylor RJ. Angiotensin II inhibits the release of [<sup>3</sup>H]acetylcholine from rat entorhinal cortex in vitro. *Brain Res* 491: 136–143, 1989.
5. Du XJ, Cox HS, Dart AM, Esler MD. Depression of efferent parasympathetic control of heart rate in rats with myocardial infarction: effect of losartan. *J Cardiovasc Pharmacol* 31: 937–944, 1998.
6. Edwards C, Doležal V, Tuček S, Zemková H, Vyskočil F. Is an acetylcholine transport system responsible for nonquantal release of acetylcholine at the rodent myoneuronal junction? *Proc Natl Acad Sci USA* 82: 3354–3358, 1985.
7. Gao L, Wang W, Li Y, Schultz HD, Liu D, Cornish KG, Zucker IH. Sympathoexcitation by central ANG II: roles for AT<sub>1</sub> receptor upregulation and NAD(P)H oxidase in RVLM. *Am J Physiol Heart Circ Physiol* 288: H2271–H2279, 2005.
8. Glantz SA. *Primer of Biostatistics* (5th ed.). New York: McGraw-Hill, 2002.

9. Hirooka Y, Head GA, Potts PD, Godwin SJ, Bendle RD, Dampney RA. Medullary neurons activated by angiotensin II in the conscious rabbit. *Hypertension* 27: 287–296, 1996.
10. Hughes J, Roth RH. Evidence that angiotensin enhances transmitter release during sympathetic nerve stimulation. *Br J Pharmacol* 41: 239–255, 1971.
11. Johnson TA, Gray AL, Lauenstein JM, Newton SS, Massari VJ. Parasympathetic control of the heart. I. An interventriculo-septal ganglion is the major source of the vagal intracardiac innervation of the ventricles. *J Appl Physiol* 96: 2265–2272, 2004.
12. Kakinuma Y, Ando M, Kuwabara M, Kafare RG, Okudela K, Kobayashi M, Sato T. Acetylcholine from vagal stimulation protects cardiomyocytes against ischemia and hypoxia involving additive non-hypoxic induction of HIF-1 $\alpha$ . *FEBS Lett* 579: 2111–2118, 2005.
13. Kawada T, Yamazaki T, Akiyama T, Sato T, Shishido T, Inagaki M, Takaki H, Sugimachi M, Sunagawa K. Differential acetylcholine release mechanisms in the ischemic and non-ischemic myocardium. *J Mol Cell Cardiol* 32: 405–414, 2000.
14. Kawada T, Yamazaki T, Akiyama T, Shishido T, Inagaki M, Uemura K, Miyamoto T, Sugimachi M, Takaki H, Sunagawa K. In vivo assessment of acetylcholine-releasing function at cardiac vagal nerve terminals. *Am J Physiol Heart Circ Physiol* 281: H139–H145, 2001.
15. Kawada T, Yamazaki T, Akiyama T, Uemura K, Kamiya A, Shishido T, Mori H, Sugimachi M. Effects of Ca<sup>2+</sup> channel antagonists on nerve stimulation-induced and ischemia-induced myocardial interstitial acetylcholine release in cats. *Am J Physiol Heart Circ Physiol* 291: H2187–H2191, 2006.
16. Konstam MA, Neaton JD, Poole-Wilson PA, Pitt B, Segal R, Sharma D, Dashbach EJ, Carides GW, Dickstein K, Riegger G, Camm AJ, Martinez FA, Bradstreet DC, Ikeda LS, Santoro EP, investigators ELITEII. Comparison of losartan and captopril on heart failure-related outcomes and symptoms from the losartan heart failure survival study (ELITE II). *Am Heart J* 150: 123–131, 2005.
17. Konstam MA, Rousseau MF, Kronenberg MW, Udelson JE, Melin J, Stewart D, Dolan N, Edens TR, Ahn S, Kinan D, Howe DM, Kilcoyne L, Metherall J, Benedict C, Yusuf S, Pouleur H, investigators SOLVD. Effects of the angiotensin converting enzyme inhibitor enalapril on the long-term progression of left ventricular dysfunction in patients with heart failure. *Circulation* 86: 431–438, 1992.
18. Lameris TW, de Zeeuw S, Duncker DJ, Alberts G, Boomsma F, Verdouw PD, van den Meiracker AH. Exogenous angiotensin II does not facilitate norepinephrine release in the heart. *Hypertension* 40: 491–497, 2002.
19. Li M, Zheng C, Sato T, Kawada T, Sugimachi M, Sunagawa K. Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation* 109: 120–124, 2004.
20. Nakayama Y, Miyano H, Shishido T, Inagaki M, Kawada T, Sugimachi M, Sunagawa K. Heart rate-independent vagal effect on end-systolic elastance of the canine left ventricle under various levels of sympathetic tone. *Circulation* 104: 2277–2279, 2001.
21. Paul M, Mehr AP, Kreutz R. Physiology of local renin-angiotensin systems. *Physiol Rev* 86: 747–803, 2006.
22. Peach MJ. Renin-angiotensin system: Biochemistry and mechanisms of action. *Physiol Rev* 57: 313–370, 1977.
23. Potter EK. Angiotensin inhibits action of vagus nerve at the heart. *Br J Pharmacol* 75: 9–11, 1982.
24. Qin Q, Downey JM, Cohen MV. Acetylcholine but not adenosine triggers preconditioning through PI3-kinase and a tyrosine kinase. *Am J Physiol Heart Circ Physiol* 284: H727–H734, 2003.
25. Rechtman M, Majewski H. A facilitatory effect of anti-angiotensin drugs on vagal bradycardia in the pithed rat and guinea-pig. *Br J Pharmacol* 110: 289–296, 1993.
26. Takata Y, Arai T, Suzuki S, Kurihara J, Uezono T, Okubo Y, Kato H. Captopril enhances cardiac vagal but not sympathetic neurotransmission in pithed rats. *J Pharmacol Sci* 95: 390–393, 2004.
27. Tune JD, Gorman MW, Feigl EO. Matching coronary blood flow to myocardial oxygen consumption. *J Appl Physiol* 97: 404–415, 2004.
28. van Kats JP, Danser AH, van Meegen JR, Sassen LM, Verdouw PD, Schalekamp MA. Angiotensin production by the heart. A quantitative study in pigs with the use of radiolabeled angiotensin infusions. *Circulation* 98: 73–81, 1998.
29. Yao Z, Gross GJ. Acetylcholine mimics ischemic preconditioning via a glibenclamide-sensitive mechanism in dogs. *Am J Physiol Heart Circ Physiol* 264: H2221–H2225, 1993.
30. Zhang C, Knudson JD, Setty S, Araiza A, Dincer üD, Kuo L, Tune JD. Coronary arteriolar vasoconstriction to angiotensin II is augmented in prediabetic metabolic syndrome via activation of AT<sub>1</sub> receptors. *Am J Physiol Heart Circ Physiol* 288: H2154–H2162, 2005.

## Regional difference in ischaemia-induced myocardial interstitial noradrenaline and acetylcholine releases

Toru Kawada<sup>a,\*</sup>, Toji Yamazaki<sup>b</sup>, Tsuyoshi Akiyama<sup>b</sup>, Toshiaki Shishido<sup>a</sup>, Shuji Shimizu<sup>a</sup>, Masaki Mizuno<sup>a</sup>, Hidezo Mori<sup>b</sup>, Masaru Sugimachi<sup>a</sup>

<sup>a</sup> Department of Cardiovascular Dynamics, Advanced Medical Engineering Center, National Cardiovascular Center Research Institute, 5-7-1 Fujishirodai, Suita, Osaka 565-8565, Japan

<sup>b</sup> Department of Cardiac Physiology, National Cardiovascular Center Research Institute, Japan

Received 18 May 2007; received in revised form 28 June 2007

### Abstract

Knowledge of the regional differences in myocardial interstitial noradrenaline (NA) and acetylcholine (ACh) levels during ischaemia would be important to understand the abnormality of neuronal environment surrounding the ischaemic heart. Using a cardiac microdialysis technique, we compared ischaemia-induced changes in the myocardial interstitial NA and ACh levels among three groups of anesthetized cats: the anterior free wall of the left ventricle (ANT group,  $n=7$ ; the left anterior descending coronary artery was occluded), the posterior free wall of the left ventricle (POST group,  $n=6$ ; the left circumflex coronary artery was occluded), and the right ventricle (RV group,  $n=6$ ; the right coronary artery was occluded). The maximum NA level was not different between the ANT and POST groups but was significantly lower in the RV group ( $P<0.01$ ) [70 nM (SD 37), 106 nM (SD 99), and 7 nM (SD 10), respectively]. The maximum ACh level was not different between the ANT and POST groups but was significantly lower in the RV group ( $P<0.05$ ) [16 nM (SD 7), 20 nM (SD 15), and 6 nM (SD 2), respectively]. In contrast, there were no significant differences in NA or ACh release in response to a local administration of ouabain (10 mM) among the ANT, POST, and RV groups ( $n=6$  each). In conclusion, the regional difference of the ischaemic effects, rather than the regional difference in the functional distributions of sympathetic and vagal efferent nerve terminals, might contribute to the lower levels of ischaemia-induced NA and ACh releases in the RV group.

© 2007 Elsevier B.V. All rights reserved.

**Keywords:** Cardiac microdialysis; Coronary artery occlusion; Ouabain; Cats

The heart is under an incessant control by the autonomic nervous system. The principal molecule that affects the myocardium is noradrenaline (NA) for the sympathetic nerve and acetylcholine (ACh) for the parasympathetic nerve. Acute myocardial ischaemia causes abnormality of the neural regulation via mechanisms such as pathological cardio-cardiac reflexes and disruption of the nerves traversing the ischaemic region (Zipes 1990; Hainsworth, 1991; Elvan and Zipes, 1998; Armour, 1999; Kawada et al., 2002). Although occlusion of the left anterior descending coronary artery

(LAD) has been shown to increase myocardial interstitial NA and ACh levels in the ischaemic region (Shindo et al., 1996; Kawada et al., 2000; Lameris et al., 2000), whether the effects of ischaemia on the myocardial interstitial NA and ACh levels are homogeneous within the left ventricle and between the left and right ventricles remains unknown. The cardio-depressor reflex similar to that induced by veratridine (the Bezold–Jarisch reflex) is frequently observed in inferoposterior but not in anterior myocardial infarction, suggesting the regional difference in the vagal afferent fibre distribution within the left ventricle (Thames et al., 1978; Walker et al., 1978). Regional differences are also reported in the distributions of sympathetic and vagal efferent nerves within the left ventricle and between the left and right ventricles

\* Corresponding author. Tel.: +81 6 6833 5012x2427; fax: +81 6 6835 5403.

E-mail address: torukawa@res.ncvc.go.jp (T. Kawada).

(Pierpont et al., 1984; Schmid et al., 1978). We hypothesized that the effects of ischaemia on the myocardial interstitial NA and ACh levels would also show regional differences.

We used a cardiac microdialysis technique and measured dialysate NA and ACh concentrations as indices of myocardial interstitial NA and ACh levels in anesthetized cats (Akiyama et al., 1991, 1994; Yamazaki et al., 1997; Kawada et al., 2001). We compared ischaemia-induced changes in the myocardial interstitial NA and ACh levels among the following regions: the anterior free wall of the left ventricle (ANT group) perfused by the LAD, the posterior free wall of the left ventricle (POST group) perfused by the left circumflex coronary artery (LCX), and the right ventricle (RV group) perfused by the right coronary artery (RCA). In addition, we compared changes in the myocardial interstitial NA and ACh levels in response to a local administration of ouabain through the dialysis probe (Yamazaki et al., 1999; Kawada et al., 2001). The advantage of the local administration of ouabain might be that we can assess the transmitter releasing function of the sympathetic and vagal efferent nerve terminals in the working heart without significant effects on the systemic haemodynamics.

## 1. Materials and methods

### 1.1. Surgical preparation

Animal care was conducted in accordance with the *Guiding Principles for the Care and Use of Animals in the Field of Physiological Sciences* approved by the Physiological Society of Japan. All protocols were approved by the Animal Subjects Committee of the National Cardiovascular Center. Adult cats weighing 2.2 to 5.0 kg were anaesthetized via an intraperitoneal injection of pentobarbital sodium (30–35 mg/kg) and ventilated mechanically with room air mixed with oxygen. The depth of anaesthesia was maintained with a continuous intravenous infusion of pentobarbital sodium ( $1\text{--}2\text{ mg kg}^{-1}\text{ h}^{-1}$ ) through a catheter inserted into the right femoral vein. Mean systemic arterial pressure was measured from a catheter inserted into the right femoral artery. The heart rate was determined using an electrocardiogram.

We performed an ischaemia protocol and a local ouabain protocol in different animals. In each protocol, the experimental animals were divided into ANT, POST, and RV groups. In the ANT and POST groups, the left fifth and/or sixth ribs were resected to allow access to the heart. In the ANT group, a 3-0 silk suture was passed around the LAD just distal to the first diagonal branch for later occlusion. Using a fine guiding needle, a dialysis probe was implanted transversely into the anterior free wall of the left ventricle perfused by the LAD. In the POST group, a 3-0 silk suture was passed around the LCX for later occlusion, and a dialysis probe was implanted transversely into the posterior free wall of the left ventricle perfused by the LCX. In the RV group, the right fifth and/or sixth ribs were resected to expose the heart. A 3-0 silk suture was passed around the RCA for later occlusion. The right ventricular wall was picked up with

a pair of forceps, and a dialysis probe was threaded transversely through the myocardium using a fine guiding needle. Heparin sodium (100 U/kg) was administered intravenously to prevent blood coagulation. A postmortem examination confirmed that the dialysis probe did not penetrate into the ventricular cavity. In the local ouabain protocol, similar experimental settings without a coronary snare were prepared for the three groups of animals.

### 1.2. Dialysis technique

The materials and properties of the dialysis probe have been described previously (Akiyama et al., 1991, 1994). Briefly, a dialysis fibre (13 mm length, 310  $\mu\text{m}$  O.D., 200  $\mu\text{m}$  I.D.; PAN-1200, 50,000 molecular weight cutoff, Asahi Chemical, Osaka, Japan) was glued at both ends to polyethylene tubes (25 cm length, 500  $\mu\text{m}$  O.D., 200  $\mu\text{m}$  I.D.). The dialysis probe was perfused at a rate of 2  $\mu\text{l}/\text{min}$  with Ringer solution containing a cholinesterase inhibitor eserine ( $10^{-4}\text{ M}$ ). Two hours elapsed before the dialysate sampling was started to allow dialysate NA and ACh concentrations reached steady states. One sampling period was set at 15 min, which yielded a sample volume of 30  $\mu\text{l}$ . The actual dialysate sampling lagged by 5 min behind a given collection period taking into account the dead space volume between the dialysis membrane and the sample tube. Each sample was collected in a microtube containing 3  $\mu\text{l}$  of phosphate buffer




Fig. 1. Changes in myocardial interstitial noradrenaline (NA) and acetylcholine (ACh) levels during coronary occlusion. ANT: region of the anterior free wall of the left ventricle perfused by the left anterior descending coronary artery, POST: region of the posterior free wall of the left ventricle perfused by the left circumflex coronary artery, RV: region of the right ventricle perfused by the right coronary artery. The coronary occlusion significantly increased myocardial interstitial NA levels in the ANT and POST groups (top panel). Changes in the NA levels were not statistically significant in the RV group. The coronary occlusion significantly increased myocardial interstitial ACh levels in all of the ANT, POST, and RV groups (bottom panel). Data are mean and SD values. \* $P < 0.05$  and † $P < 0.01$  from the corresponding baseline value by Dunnett's test.




Fig. 2. Maximum levels of myocardial interstitial NA and ACh induced by coronary occlusions. The maximum NA level did not differ between the ANT and POST groups, but it was significantly lower in the RV group (left panel). The maximum ACh level did not differ between the ANT and POST groups, but it was significantly lower in the RV group than in the POST group (right panel). Data are mean and SD values. Exact *P* values determined by Tukey test are supplied.

(0.1 M, pH 3.5) to prevent amine oxidation. The ACh concentration in the dialysate was measured directly by high performance liquid chromatography with electrochemical detection (HPLC-ECD) system (Eicom, Kyoto, Japan). The NA concentration in the dialysate was measured by another HPLC-ECD system after removing interfering compounds with an alumina procedure.

### 1.3. Protocols

#### 1.3.1. Ischaemia protocol

In each animal in the ANT ( $n=7$ ), POST ( $n=6$ ), and RV ( $n=6$ ) groups, a 15-min dialysate sample was collected under baseline conditions. Thereafter, each corresponding coronary artery was occluded for 60 min, and four consecutive 15-min dialysate samples were obtained during the occlusion period.

#### 1.3.2. Local ouabain protocol

In each animal in the ANT ( $n=6$ ), POST ( $n=6$ ), and RV ( $n=6$ ) groups, a 15-min dialysate sample was collected under baseline conditions. The perfusate for the dialysis probe was then replaced with Ringer solution containing 10 mM of ouabain. The local administration of ouabain has been shown to evoke myocardial interstitial NA and ACh releases without significant effects on the systemic haemodynamics (Yamazaki et al., 1999; Kawada et al., 2001). Four consecutive 15-min dialysate samples were obtained during the local administration of ouabain.

Table 1

Changes in mean arterial pressure by occlusions of the anterior descending coronary artery (ANT), the left circumflex coronary artery (POST), and the right coronary artery (RV)

	Baseline	5	15	30	45	60
ANT	116 (31)	114 (27)	108 (24)	108 (25)	108 (23)	110 (25)
POST	139 (31)	119 (31) <sup>†</sup>	125 (31) <sup>†</sup>	130 (31) <sup>*</sup>	132 (28)	135 (33)
RV	101 (20)	101 (26)	99 (22)	101 (23)	103 (25)	103 (23)

Baseline and values after 5, 15, 30, 45, and 60 min of the occlusion are presented (in mm Hg). Data are mean (SD) values. <sup>\*</sup> $P<0.05$  and <sup>†</sup> $P<0.01$  from the corresponding baseline value.

### 1.4. Statistical analysis

All data are presented as mean (SD) values. Because the increase in NA reached more than 100 times the baseline value in the ANT and POST groups, NA data were compared after logarithmic transformation. To examine the effects of ischaemia or the local administration of ouabain on the NA or ACh level in each group, we used one-way repeated-measures analysis of variance (Glantz, 2002). When there was a significant difference in measured values among the collection periods, Dunnett's test was applied to identify the difference from the baseline value. To compare the maximum NA or ACh response among the ANT, POST, and RV groups, we used one-way analysis of variance. When there was a significant difference among the three groups, Tukey test was applied for simultaneous all pairwise comparisons. The differences were considered significant at  $P<0.05$ .

## 2. Results

Coronary artery occlusion increased the NA levels to more than 100 times the respective baseline levels in the ANT and POST groups (Fig. 1, top). In the RV group, although the mean NA level increased to nearly 10 times the baseline level, the change was not statistically significant, possibly due to the large variance of the NA responses across the animals. The coronary occlusion increased the ACh levels to approximately 15 times the respective baseline levels in the ANT and POST groups whereas it increased the

Table 2

Changes in the heart rate by occlusions of the anterior descending coronary artery (ANT), the left circumflex coronary artery (POST), and the right coronary artery (RV)

	Baseline	5	15	30	45	60
ANT	193 (18)	166 (13) <sup>†</sup>	174 (15) <sup>†</sup>	175 (18) <sup>†</sup>	173 (15) <sup>†</sup>	171 (15) <sup>†</sup>
POST	183 (25)	159 (21) <sup>†</sup>	167 (20) <sup>†</sup>	167 (25) <sup>†</sup>	170 (25) <sup>†</sup>	169 (27) <sup>†</sup>
RV	188 (34)	175 (35)	176 (33)	179 (35)	181 (36)	183 (38)

Baseline and values after 5, 15, 30, 45, and 60 min of the occlusion are presented (in beats min<sup>-1</sup>). Data are mean (SD) values. <sup>†</sup> $P<0.01$  from the corresponding baseline value.




Fig. 3. Changes in myocardial interstitial NA and ACh levels induced by a local administration of ouabain through the dialysis probe. Ouabain significantly increased myocardial interstitial NA levels in all of the ANT, POST, and RV groups (top panel). The maximum NA level did not differ among the three groups. Ouabain significantly increased myocardial interstitial ACh levels in all of the ANT, POST, and RV groups (bottom panel). The maximum ACh level did not differ among the three groups. Data are mean and SD values. \* $P < 0.05$  and † $P < 0.01$  from the corresponding baseline value by Dunnett's test.

ACh levels to approximately 8 times the baseline level in the RV group (Fig. 1, bottom).

The maximum NA level observed during ischaemia was not different between the ANT and POST groups, but it was significantly lower in the RV group than in the ANT and POST groups (Fig. 2, left). The maximum ACh level observed during ischaemia was not different between the ANT and POST groups, but it was significantly lower in the RV group than in the POST group (Fig. 2, right).

Changes in mean arterial pressure obtained from the ischaemia protocol are summarized in Table 1. Occlusion of the LAD did not change mean arterial pressure significantly (the ANT group), whereas occlusion of the LCX significantly decreased mean arterial pressure at 5, 15, and 30 min of the ischaemic period (the POST group). Occlusion of the RCA did not change mean arterial pressure (the RV group).

Changes in the heart rate obtained from the ischaemia protocol are summarized in Table 2. Occlusion of the LAD significantly decreased the heart rate throughout the occlusion period (the ANT group). Occlusion of the LCX also significantly decreased the heart rate throughout the occlusion period (the POST group). In contrast, occlusion of the RCA did not change the heart rate significantly (the RV group).

A local administration of ouabain increased the NA levels in all of the ANT, POST, and RV groups (Fig. 3, top). There were no significant differences in the maximum NA levels among the three groups ( $P = 0.40$  by ANOVA). The local administration of ouabain also increased the ACh levels in all of the ANT, POST, and RV groups (Fig. 3, bottom). The ACh levels reached their maximum levels during 15–30 min

of the administration period, then slightly decreased from the maximum levels during 30–45 and 45–60 min of the administration period but remained higher than the respective baseline levels. There were no significant differences in the maximum ACh levels among the three groups ( $P = 0.35$  by ANOVA).

### 3. Discussion

The level of ischaemia-induced NA release was significantly lower in the RV group than in the ANT and POST groups (Fig. 2, left). The level of ischaemia-induced ACh release was significantly lower in the RV group than in the POST group (Fig. 2, right). To our knowledge, this is the first report showing the differential effects of ischaemia on the myocardial interstitial NA and ACh levels among the main coronary arteries.

#### 3.1. Regional difference in the ischaemia-induced myocardial interstitial NA release

Mechanisms responsible for the ischaemia-induced NA release have been extensively studied (Schömig A et al., 1984, 1988). An exocytotic release mechanism participates in the NA release in the early phase of ischaemia (within approximately 20 min from the onset of ischaemia) (Akiyama and Yamazaki, 1999). Energy depletion in the ischaemic region impairs the  $\text{Na}^+ - \text{K}^+$  ATPase activity and induces axoplasmic  $\text{Na}^+$  accumulation. Because the NA uptake carrier is driven by the  $\text{Na}^+$  gradient across the plasma membrane (Schwartz, 2000), the accumulation of intracellular  $\text{Na}^+$  causes the reverse transport of axoplasmic NA to extracellular space. This nonexocytotic release mechanism becomes predominant as the ischaemic period is prolonged (Akiyama and Yamazaki, 1999; Lameris et al., 2000). In the present study, there was no significant difference in the levels of ischaemia-induced NA release between the ANT and POST groups (Fig. 2, left), suggesting that the extent of energy depletion caused by the coronary artery occlusion might be similar between the anterior and posterior regions of the left ventricle. In contrast, the level of ischaemia-induced NA release was much lower in the RV group (Fig. 2, left). Much lower oxygen consumption in the right ventricle than in the left ventricle (Weiss et al., 1978; Kusachi et al. 1982) may have delayed the progression of ischaemia and/or mitigated the severity of ischaemia.

To examine whether the observed difference in the ischaemia-induced NA release was attributable to the regional difference in the functional distribution of sympathetic efferent nerve terminals, we measured the amount of myocardial interstitial NA release in response to a local administration of ouabain (Yamazaki et al., 1999). Because locally administered ouabain spreads in the vicinity of the semipermeable membrane of the dialysis fibre and evokes the NA release, the NA concentration in the dialysate thus measured is considered to reflect the density of sympathetic

nerve terminals around the dialysis fibre. There were no significant differences in the maximum NA levels in response to the ouabain administration among the ANT, POST, and RV groups (Fig. 3, top), suggesting that the functional distribution of sympathetic efferent nerve terminals did not account for the lower level of ischaemia-induced NA release in the RV group.

The results of local administration of ouabain, showing no significant regional differences in the functional distribution of sympathetic efferent nerve terminals, are comparable to histological studies. In the human heart (Kawano et al., 2003), the number of tyrosine hydroxylase (TH)-positive nerves is similar between the left and right ventricles, and the number of TH-positive nerves in the anterior wall is 1.2 times greater than that in the posterior wall of the left ventricle. Although Pierpont et al. (Pierpont et al., 1984) reported a regional difference in the NA content in the canine left ventricle, the major difference observed in their study is an increasing gradient of NA from the apex to the base of the ventricle. In addition, the NA content was similar between the left and right ventricles in their study [495 ng/g (SD 267) vs. 503 ng/g (SD 123)].

### 3.2. Regional difference in the ischaemia-induced myocardial interstitial ACh release

In previous studies, we have demonstrated that acute myocardial ischaemia causes myocardial interstitial ACh release in the ischaemic region (Kawada et al., 2000; 2006b). An exocytotic release mechanism may be involved in the ischaemia-induced ACh release within approximately 15 min from the onset of ischaemia (Kawada et al., 2000, 2006a). Thereafter, a local ACh release mechanism independent of vagal nerve discharge may play a dominant role in the ischaemia-induced ACh release. In the present study, there was no significant difference in the levels of ischaemia-induced ACh release between the ANT and POST groups (Fig. 2, right). In contrast, the level of ischaemia-induced ACh release was much lower in the RV group than in the POST group. These results suggest that the energy depletion during ischaemia might be less severe or delayed in the right ventricle compared to that in the left ventricle.

Similar to the ouabain-induced NA release, there were no significant differences in the levels of ouabain-induced ACh release among the ANT, POST, and RV groups (Fig. 3, bottom). In a histological study of the human heart (Kawano et al., 2003), the number of acetylcholine esterase (AChE)-positive nerves in the right ventricle is 1.2 times greater than that in the left ventricle, and the number of AChE-positive nerves in the posterior wall is 1.4 times greater than that in the anterior wall of the left ventricle. In the guinea pig heart, the level of choline acetyltransferase activity was approximately two times higher in the right ventricle than in the left ventricle (Schmid et al., 1978). Notwithstanding the discrepancies among reports, these histochemical studies indicate that the number of the vagal nerve terminals in the right

ventricle is not less than that in the left ventricle. In other words, the regional difference in the vagal efferent nerve distribution may not account for the lower level of ischaemia-induced ACh release in the RV group as compared with the POST group.

### 3.3. Pathological significance

The pathological significance of the NA and ACh releases in the ischaemic region is still to be explored. Although high levels of NA reveal cardiotoxicity (Rona, 1985), depletion of catecholamine in reserpinized animals fails to reduce the myocardial infarct size (Toombs et al., 1993; Vander Heide et al., 1995). However, in the reserpinized animals, not only the ischaemic area but also the non-ischaemic area is subjected to catecholamine depletion, making the interpretation of the results difficult. Acetylcholine, when administered prior to coronary occlusion, induces an ischaemic preconditioning mimetic effect (Qin et al., 2002). Acetylcholine also exerts protection on myocytes against hypoxia (Kakinuma et al., 2005). Generally speaking, the excess NA might be harmful whereas the presence of ACh might be beneficial to the myocardium.

One possible feature of the cardiac microdialysis may be that it can monitor the time course of changes in myocardial interstitial NA and ACh levels (Shindo et al., 1996; Kawada et al., 2000; Lameris et al., 2000). Although myocardial ischaemia evokes both the NA and ACh releases, the ACh release is more prompt compared to the NA release. When calculating the percentage against the respective maximum level, the mean NA levels were only 3 and 2% whereas the mean ACh levels reached 79 and 53% in the ANT and POST groups, respectively, during the 0–15 min of the ischaemic period (Fig. 1, note the logarithmic scaling in the top panel). In the RV group, the mean NA level was 28% whereas the mean ACh level reached 72% during the 0–15 min of the ischaemic period. It seems that the ACh release is a protective mechanism against a forthcoming excess of NA in the ischaemic region. Further studies are required to elucidate the significance of a local neuronal environment in modifying the severity of myocardial ischaemia.

### 3.4. Regional difference in the reflex effects

In the ischaemia protocol, the reflexes from the heart and the arterial baroreflex might have modified the efferent nerve activities. Both the ANT and POST groups showed a significant decrease in the heart rate during ischaemia (Table 2), suggesting an increase in the vagal tone and/or a decrease in the sympathetic tone. Because mean arterial pressure was either unchanged (the ANT group) or decreased (the POST group) (Table 1), the baroreflex cannot account for the decrease in the heart rate during ischaemia. The decreased mean arterial pressure in the POST group compared to the unchanged mean arterial pressure in the ANT group suggests that the cardiodepressor reflex was stronger in the POST than

the ANT group. These differences, however, did not cause the difference in the maximum levels of ischaemia-induced NA and ACh releases between the ANT and POST groups (Fig. 2). Because local release mechanisms became predominant as the ischaemic period is prolonged (Akiyama and Yamazaki, 1999; Kawada et al., 2000, 2006a; Lameris et al., 2000), the difference in the efferent nerve activities might not have affected the maximum levels of NA and ACh releases significantly. In contrast to the ANT and POST groups, the RV group did not show significant changes in mean arterial pressure or the heart rate, suggesting that the effect of RCA occlusion on the systemic haemodynamics was minimal in the present study. Despite the absence of significant bradycardia, the myocardial interstitial ACh level was significantly increased in the RV group during ischaemia, suggesting the involvement of a local release mechanism.

There are limitations to the present study. First, we could not examine regional differences in the NA and ACh releases along the transmural axis from the epicardial layer to the endocardial layer, because we could not control the exact depth of the dialysis fibre implanted transversely in the ventricular wall. Second, species differences should be taken into account when interpreting the present data. As an example, we usually observed a bradycardic response not only during LCX occlusion but also during LAD occlusion in cats (Table 2). In contrast, LAD occlusion in dogs more frequently evokes tachycardia and hypertension (Thames et al., 1978; Zipes, 1990).

#### 4. Conclusion

The maximum levels of ischaemia-induced NA and ACh releases did not differ between the ANT and POST groups but were significantly lower in the RV group. In contrast, myocardial interstitial NA and ACh releases in response to a local administration of ouabain did not show regional differences among the ANT, POST, and RV groups. The regional difference in the ischaemic effects, rather than the regional difference in the functional distributions of sympathetic and vagal efferent nerve terminals, might contribute to the lower levels of ischaemia-induced NA and ACh releases in the RV group.

#### Acknowledgments

This study was supported by “Health and Labour Sciences Research Grant for Research on Advanced Medical Technology”, “Health and Labour Sciences Research Grant for Research on Medical Devices for Analyzing, Supporting and Substituting the Function of Human Body”, and “Health and Labour Sciences Research Grant H18-Iryo-Ippan-023” from the Ministry of Health, Labour and Welfare of Japan.

#### References

- Akiyama, T., Yamazaki, T., 1999. Norepinephrine release from cardiac sympathetic nerve endings in the in vivo ischemic region. *J. Cardiovasc. Pharmacol.* 34, S11–S14.
- Akiyama, T., Yamazaki, T., Ninomiya, I., 1991. In vivo monitoring of myocardial interstitial norepinephrine by dialysis technique. *Am. J. Physiol. Heart Circ. Physiol.* 261, H1643–H1647.
- Akiyama, T., Yamazaki, T., Ninomiya, I., 1994. In vivo detection of endogenous acetylcholine release in cat ventricles. *Am. J. Physiol. Heart Circ. Physiol.* 266, H854–H860.
- Armour, J.A., 1999. Myocardial ischaemia and the cardiac nervous system. *Cardiovasc. Res.* 41, 41–54.
- Elvan, A., Zipes, D.P., 1998. Right ventricular infarction causes heterogeneous autonomic denervation of the viable peri-infarct area. *Circulation* 97, 484–492.
- Glantz, S.A., 2002. *Primer of Biostatistics*, 5th ed. McGraw-Hill, New York.
- Hainsworth, R., 1991. Reflexes from the heart. *Physiol. Rev.* 71, 617–658.
- Kakinuma, Y., Ando, M., Kuwabara, M., Katare, R.G., Okudela, K., Kobayashi, M., Sato, T., 2005. Acetylcholine from vagal stimulation protects cardiomyocytes against ischemia and hypoxia involving additive non-hypoxic induction of HIF-1 $\alpha$ . *FEBS Lett.* 579, 2111–2118.
- Kawada, T., Yamazaki, T., Akiyama, T., Sato, T., Shishido, T., Inagaki, M., Takaki, M., Sugimachi, M., Sunagawa, K., 2000. Differential acetylcholine release mechanisms in the ischemic and non-ischemic myocardium. *J. Mol. Cell Cardiol.* 32, 405–414.
- Kawada, T., Yamazaki, T., Akiyama, T., Shishido, T., Inagaki, M., Uemura, K., Miyamoto, T., Sugimachi, M., Takaki, H., Sunagawa, K., 2001. In vivo assessment of acetylcholine-releasing function at cardiac vagal nerve terminals. *Am. J. Physiol. Heart Circ. Physiol.* 281, H139–H145.
- Kawada, T., Yamazaki, T., Akiyama, T., Mori, H., Uemura, K., Miyamoto, T., Sugimachi, M., Sunagawa, K., 2002. Disruption of vagal efferent axon and nerve terminal function in the posts ischemic myocardium. *Am. J. Physiol. Heart Circ. Physiol.* 283, H2687–H2691.
- Kawada, T., Yamazaki, T., Akiyama, T., Li, M., Ariumi, H., Mori, H., Sunagawa, K., Sugimachi, M., 2006a. Vagal stimulation suppresses ischemia-induced myocardial interstitial norepinephrine release. *Life Sci.* 78, 882–887.
- Kawada, T., Yamazaki, T., Akiyama, T., Uemura, K., Kamiya, A., Shishido, T., Mori, H., Sugimachi, M., 2006b. Effects of Ca<sup>2+</sup> channel antagonists on nerve stimulation-induced and ischemia-induced myocardial interstitial acetylcholine release in cats. *Am. J. Physiol. Heart Circ. Physiol.* 291, H2187–H2191.
- Kawano, H., Okada, R., Yano, K., 2003. Histological study on the distribution of autonomic nerves in the human heart. *Heart Vessels* 18, 32–39.
- Kusachi, S., Nishiyama, O., Yasuhara, K., Saito, D., Haraoka, S., Nagashima, H., 1982. Right and left ventricular oxygen metabolism in open-chest dogs. *Am. J. Physiol. Heart Circ. Physiol.* 243, H761–H766.
- Lameris, T.W., de Zeeuw, S., Alberts, G., Boomsma, F., Duncker, D.J., Verdouw, P.D., Veld, A.J., van Den Meiracker, A.H., 2000. Time course and mechanism of myocardial catecholamine release during transient ischemia in vivo. *Circulation* 101, 2645–2650.
- Pierpont, G.L., DeMaster, E.G., Cohn, J.N., 1984. Regional differences in adrenergic function within the left ventricle. *Am. J. Physiol. Heart Circ. Physiol.* 246, H824–H829.
- Qin, Q., Downey, J.M., Cohen, M.V., 2002. Acetylcholine but not adenosine triggers preconditioning through PI3-kinase and a tyrosine kinase. *Am. J. Physiol. Heart Circ. Physiol.* 284, H727–H734.
- Rona, G., 1985. Catecholamine cardiotoxicity. *J. Mol. Cell Cardiol.* 17, 291–306.
- Schömig, A., Dart, A.M., Dietz, R., Mayer, E., Kübler, W., 1984. Release of endogenous catecholamines in the ischemic myocardium of the rat Part. A: locally mediated release. *Circ. Res.* 55, 689–701.
- Schömig, A., Kurz, T., Richardt, G., Schömig, E., 1988. Neuronal sodium homeostasis and axoplasmic amine concentration determine calcium-independent noradrenaline release in normoxic and ischemic rat heart. *Circ. Res.* 63, 214–226.
- Schmid, P.G., Greif, B.J., Lund, D.D., Roskoski Jr., R., 1978. Regional choline acetyltransferase activity in the guinea pig heart. *Circ. Res.* 42, 657–660.
- Schwartz, J.H., 2000. Neurotransmitters. In: Kandel, E.R., Schwartz, J.H., Jessell, T.M. (Eds.), *Principles of Neural Science*, 4th ed. McGraw-Hill, New York, pp. 280–297.



- Shindo, T., Akiyama, T., Yamazaki, T., Ninomiya, I., 1996. Regional myocardial interstitial norepinephrine kinetics during coronary occlusion and reperfusion. *Am. J. Physiol., Heart Circ. Physiol.* 270, H245–H251.
- Thames, M.D., Klopfenstein, H.S., Abboud, F.M., Mark, A.L., Walker, J.L., 1978. Preferential distribution of inhibitory cardiac receptors with vagal afferents to the inferoposterior wall of the left ventricle activated during coronary occlusion in the dog. *Circ. Res.* 43, 512–519.
- Toombs, C.F., Wiltse, A.L., Shebuski, R.J., 1993. Ischemic preconditioning fails to limit infarct size in reserpinized rabbit myocardium. Implication of norepinephrine release in the preconditioning effect. *Circulation* 88, 2351–2358.
- Vander Heide, R.S., Schwartz, L.M., Jennings, R.B., Reimer, K.A., 1995. Effect of catecholamine depletion on myocardial infarct size in dogs: role of catecholamines in ischemic preconditioning. *Cardiovasc. Res.* 30, 656–662.
- Walker, J.L., Thames, M.D., Abboud, F.M., Mark, A.L., Klopfenstein, H.S., 1978. Preferential distribution of inhibitory cardiac receptors in left ventricle of the dog. *Am. J. Physiol., Heart Circ. Physiol.* 235, H188–H192.
- Weiss, H.R., Neubauer, J.A., Lipp, J.A., Shinha, A.K., 1978. Quantitative determination of regional oxygen consumption in the dog heart. *Circ. Res.* 42, 394–401.
- Yamazaki, T., Akiyama, T., Kitagawa, H., Takauchi, Y., Kawada, T., Sunagawa, K., 1997. A new, concise dialysis approach to assessment of cardiac sympathetic nerve terminal abnormalities. *Am. J. Physiol., Heart Circ. Physiol.* 272, H1182–H1187.
- Yamazaki, T., Akiyama, T., Kawada, T., 1999. Effects of ouabain on in situ cardiac sympathetic nerve endings. *Neurochem. Int.* 35, 439–445.
- Zipes, D.P., 1990. Influence of myocardial ischemia and infarction on autonomic innervation of heart. *Circulation* 82, 1095–1105.

## Cardiac epinephrine synthesis and ischemia-induced myocardial epinephrine release

Yosuke Kuroko<sup>a</sup>, Toji Yamazaki<sup>b,\*</sup>, Noriyuki Tokunaga<sup>a</sup>, Tsuyoshi Akiyama<sup>b</sup>,  
Hirotoshi Kitagawa<sup>b</sup>, Kozo Ishino<sup>a</sup>, Shunji Sano<sup>a</sup>, Hidezo Mori<sup>b</sup>

<sup>a</sup> Department of Cardiovascular Surgery, Okayama University Graduate School of Medicine and Dentistry, Okayama, 700-8558, Japan

<sup>b</sup> Department of Cardiac Physiology, National Cardiovascular Center Research Institute, 5-7-1 Fujishiro-dai, Suita, Osaka, 565-8565, Japan

Received 4 September 2006; received in revised form 13 February 2007; accepted 14 February 2007

Available online 21 February 2007

Time for primary review 15 days

### Abstract

**Objective:** Phenylethanolamine-*N*-methyltransferase (PNMT), the enzyme that synthesizes epinephrine (EPI) from norepinephrine (NE) in the adrenal gland, is present in extra-adrenal tissues including heart. Ischemia evokes an excessive NE accumulation in the myocardial interstitial spaces. Therefore, cardiac PNMT activity with high NE levels may contribute to cardiac EPI synthesis and release evoked by ischemia.

**Methods:** We measured dialysate EPI levels in the left ventricle of anesthetized rabbits using a cardiac microdialysis technique. The dialysate EPI level served as an index of the myocardial interstitial EPI level. Locally administered NE-induced dialysate EPI responses were measured. The left circumflex coronary artery was occluded for 60 min and the dialysate EPI and NE levels in the ischemic region were measured. Coronary occlusion-induced EPI responses were compared with and without administration of a PNMT inhibitor (SKF29661) in the presence and absence of desipramine (catecholamine transport blocker).

**Results:** Local administration of NE (250, 2500 ng/ml) increased the EPI levels to  $734 \pm 125$  and  $2088 \pm 367$  pg/ml respectively. These increases in dialysate EPI were suppressed by the PNMT inhibitor. Acute myocardial ischemia significantly increased the EPI levels to  $3607 \pm 1069$  pg/ml in the ischemic region, and these were suppressed by the PNMT inhibitor ( $1417 \pm 581$  pg/ml). The pretreatment with desipramine suppressed ischemia-induced EPI release, which did not differ with ( $725 \pm 155$  pg/ml) and without administration of a PNMT inhibitor ( $743 \pm 172$  pg/ml).

**Conclusion:** The cardiac PNMT in the left ventricle is capable of synthesizing EPI with markedly elevated NE levels in the myocardial interstitial space.

© 2007 European Society of Cardiology. Published by Elsevier B.V. All rights reserved.

**Keywords:** Autonomic nervous system; Interstitial space; Ischemia; Reperfusion; Neurotransmitters

### 1. Introduction

It is generally accepted that myocardial ischemia evokes an excessive catecholamine accumulation in the myocardial interstitial space [1,2]. This high catecholamine level in the myocardial interstitium is thought to aggravate the progression of myocardial cell injury and incidence of malignant arrhythmia [3,4]. From *in vitro* and *in vivo* studies, several mechanisms are presently suggested to induce release of norepinephrine (NE) from the nerve endings [1,5,6]. The

outward NE transport through uptake<sub>1</sub> carrier has been proposed as an important mechanism responsible for the ischemia-induced NE release. With respect to epinephrine (EPI), however, it is uncertain whether the mechanism of release differs between EPI and NE.

EPI is synthesized mainly from NE in the adrenal medulla by phenylethanolamine *N*-methyltransferase (PNMT) [7] and released into the bloodstream [8]. Myocardial ischemia evokes an excessive NE and EPI accumulation in the myocardial interstitial space although the blood supply is blocked. Therefore, regional release mechanism has been suggested to induce release of EPI in the ischemic region. Early studies reported that PNMT activity was measured in

\* Corresponding author. Tel.: +81 6 6833 5012x2379; fax: +81 6 6872 8092.  
E-mail address: yamazaki@ri.nccvc.go.jp (T. Yamazaki).

homogenates from the heart [9,10]. Excessive NE level and cardiac PNMT activity may provide the prerequisite for cardiac EPI synthesis evoked by ischemia. We speculate that ischemia may promote EPI synthesis and release by high NE accumulation via cardiac PNMT activity.

We have demonstrated the usefulness of dialysis technique for the *in vivo* monitoring of regional myocardial interstitial catecholamine kinetics [11,12]. In the present study, we extend this approach to assessment of PNMT activity using NE as a substrate of EPI synthesis. We examined the role of PNMT activity in the EPI release evoked by myocardial ischemia. Furthermore, we examined the contribution of neuronal catecholamine transport to EPI release evoked by ischemia.

## 2. Materials and methods

### 2.1. Animal preparation

Animal care proceeded in strict accordance with the *Guide for the Care and Use of Laboratory Animals* published by the US National Institute of Health (NIH Publication No. 85-23, revised 1996). Adult Japanese white rabbits (2.5–3.3 kg) were anesthetized with pentobarbital sodium (30–35 mg/kg *i.v.*). The level of anesthesia was maintained with a continuous intravenous infusion of pentobarbital sodium (1–2 mg/kg/h). The rabbits were intubated and ventilated with room air mixed with oxygen. Heart rate, arterial pressure, and electrocardiogram were simultaneously monitored with a data recorder. The fifth or sixth rib on the left side was partially removed to expose the heart. A snare was placed around the main branch of the left circumflex coronary artery (LCX) to act as the occluder for later coronary occlusion. With a fine guiding needle, a dialysis probe was implanted in the region perfused by LCX of the left ventricular wall. Judging from changes in the color of the ventricular wall during a brief coronary occlusion, the dialysis probe was located in the midst of the ischemic region. Heparin sodium (100 IU/kg) was administered intravenously to prevent blood coagulation.

### 2.2. Dialysis technique and epinephrine measurements

Materials suitable for cardiac dialysis probes have been described in detail elsewhere [13]. Briefly, we designed a handmade long transverse dialysis probe. A dialysis fiber (8 mm length, 0.31 mm o.d., and 0.20 mm i.d.; PAN-1200 50,000 molecular weight cutoff, Asahi Chemical Japan) was glued at both ends of a polyethylene tube. The dialysis probe was perfused with Ringer's solution or Ringer's solution containing pharmacological agents at a perfusion speed of 2  $\mu$ l/min using a microinjection pump (Carnegie Medicine CMA/100). One sampling period was 15 min (1 sampling volume = 30  $\mu$ l). Each sample was collected in a microtube containing 3  $\mu$ l of 0.1 N HCl to prevent amine oxidation. Dialysate EPI and NE level were measured by high-performance liquid chromatography

with electrochemical detection (ECD-300, Eicom, Kyoto, Japan) as previously described [14,15]. Dialysate EPI level served as an index of myocardial interstitial EPI level. We commenced the protocol followed by a stabilization period of 2 h. Taking into consideration the dead space between the dialysis fiber and sample tube, we sampled the dialysate.

### 2.3. Experimental protocols

#### 2.3.1. Dialysate EPI levels during local administration of NE

First, to elucidate cardiac PNMT activity, we locally administered NE. The concentration of NE was chosen to be in the same range as in the myocardial ischemic region based on the results of our previous study [6]. After control sampling, we locally administered NE (250 or 2500 ng/ml) through a dialysis probe for 60 min, with dialysate collected during the last 15 min. The same protocol was followed after administration of a PNMT inhibitor (SKF29661) [16] in separate rabbits. SKF29661 (50 mg/kg) was administered intraperitoneally 60 min before control sampling. To confirm whether PNMT activity was located in sympathetic nerve endings or myocardium, we performed chemical sympathetic denervation with hydroxydopamine (6-OHDA) and examined the dialysate EPI response to local NE infusion. Five days previously, rabbits were given 60 mg/kg 6-OHDA intravenously [17]. Dialysate EPI response to NE infusion was measured. Furthermore, we examined the effect of NE uptake<sub>2</sub> inhibition on the EPI response to local NE infusion. We added corticosterone (1 mM) on the perfusate and measured the dialysate EPI response to local NE infusion.

#### 2.3.2. Time course of dialysate EPI levels during the myocardial ischemia in the presence and absence of SKF29661

After control sampling, we occluded the main branch of LCX for 60 min and then released the occluder. We observed the time course of dialysate EPI levels in the ischemic region in six rabbits. We collected five consecutive 15-minute dialysate samples during coronary occlusion and reperfusion (vehicle group). To examine the involvement of PNMT activity on the EPI level, we intraperitoneally administered a PNMT inhibitor (SKF29661) 60 min before control sampling in separate rabbits (SKF group). We performed LCX occlusion and collected dialysate samples as described in vehicle group. We compared EPI responses to LCX occlusion between vehicle and SKF groups.

#### 2.3.3. Influence of desipramine on dialysate EPI levels during myocardial ischemia with and without SKF29661

EPI can be released via non-exocytotic release at the sympathetic nerve terminals [18]. To elucidate the involvement of catecholamine transporter on EPI levels, we locally administered an inhibitor of NE uptake, carrier desipramine (100  $\mu$ M) through a dialysis probe. One hour thereafter, we performed LCX occlusion and collected dialysate samples in

Table 1  
Time course of heart rate and mean arterial pressure during coronary occlusion and reperfusion

	Control	Coronary occlusion (min)				Reperfusion (min)
		15	30	45	60	15
<i>Heart rate (bpm)</i>						
Vehicle group (n=6)*	297±10	289±8	283±9	280±9	281±7	269±6
SKF29661 group (n=6)	319±11	312±12	311±13	313±13	311±14	306±13
Desipramine group (n=6)	312±10	315±11	317±10	318±12	314±13	309±12
Desipramine+SKF29661 group (n=6)	316±12	310±10	314±8	315±7	317±8	311±8
<i>Mean arterial pressure (mm Hg)</i>						
Vehicle group (n=6)	96±8	87±7	84±5	83±4	81±5	81±3 †
SKF29661 group (n=6)	97±3	93±4	93±4	92±4	92±4	91±4 †
Desipramine group (n=6)	98±6	94±6	91±5	89±5	88±4	87±4 †
Desipramine+SKF29661 group (n=6)	101±2	88±6	90±4	88±4	88±3	83±3 †

†Values are means±SE. \* $P<0.05$  vs. values of pretreatment groups.

the above-described protocol in separate rabbits (desipramine group). Furthermore, we tried to determine the influence of desipramine on PNMT induced EPI release during myocardial ischemia. We co-administered intraperitoneally SKF29661 (50 mg/kg) and desipramine locally through a dialysis probe continuously 60 min before control sampling (desipramine+SKF group). We performed LCX occlusion and collected dialysate samples as described in vehicle group. We compared EPI responses to LCX occlusion between desipramine and desipramine+SKF groups.

At the end of each experiment, the rabbits were killed with an overdose of pentobarbital sodium, and the implant regions were checked to confirm that the dialysis probe had been implanted within the cardiac muscle.

#### 2.4. Statistical methods

The effects of myocardial ischemia (NE infusion) and pretreatment were examined using two-way analysis of variance. When statistical significance was detected between two groups, the dialysate EPI levels with and without a PNMT inhibitor were compared by unpaired *t*-test. The dialysate NE levels were compared among four groups using one-way analysis variance followed by Newman–Keuls test for the multiple comparisons against each other. The data of heart rate and mean arterial pressure were compared among four groups using two-way analysis variance. When statistical significance was detected, the Newman–Keuls test was applied. Statistical significance was defined as  $P<0.05$ . Values are presented as means±SE.

### 3. Results

#### 3.1. Time course of heart rate and mean arterial pressure

Local administration of NE through a dialysis probe did not alter heart rate (HR) or mean arterial pressure (MAP) in vehicle or SKF groups. The time course of HR and MAP during myocardial ischemia and reperfusion is shown in

Table 1. Coronary occlusion tended to cause a fall in HR and MAP, but no statistically significant alterations in HR or MAP were obtained against a baseline value within each group. Basal HR in vehicle group was lower than that in the other three groups.

#### 3.2. Dialysate EPI response during local NE administration through a dialysis probe

Fig. 1 shows data obtained from local administration of exogenous NE through a dialysis probe. Dialysate EPI levels increased significantly with increases in the rate of NE infusion. Dialysate EPI levels reached  $734.5±125$ , and  $2081±367$  pg/ml ( $n=6$ ) at 250 and 2500 ng/ml of NE infusion, respectively. In the presence of SKF29661, dialysate EPI levels were significantly suppressed compared to those of the vehicle group. Dialysate EPI levels were  $68±25$ ,  $282±70$  pg/ml ( $n=6$ ) at 250 and 2500 ng/ml of NE infusion, respectively. SKF29661 attenuated EPI responses to 10% of vehicle group. In sympathetic denervation with 6-OHDA, the dialysate EPI response to NE infusion (250 ng) was preserved ( $n=4$ ,  $760±193$  pg/ml). With the perfusate containing the NE uptake<sub>2</sub> inhibitor, corticosterone (1 mM), the dialysate EPI response to NE infusion (250 ng) was suppressed ( $n=6$ ,  $167±27$  pg/ml).




Fig. 1. Dialysate epinephrine levels evoked by norepinephrine perfusion through dialysis probe. SKF = SKF29661, \* $P<0.05$  vs. value at 0–15 min of norepinephrine perfusion, † $P<0.05$  vs. concurrent value of vehicle group.