

Fig. 1. Conditional expression of *Kras*^{G12V} transgene. (A) The CALNL-HAKras^{G12V} transgene is comprised of a hybrid CMV enhancer/chicken β -actin (CAG) promoter, a cassette for the neomycin resistance gene flanked by loxP sites, and a sequence containing a human *Kras*^{G12V} with a hemagglutinin (HA)-tag. Infection with the Cre recombinase-expressing adenovirus results in Cre-mediated recombination of the transgene and removal of the neo-coding region and its associated mRNA polyadenylation signal, generating a functional HA-*Kras*^{G12V} gene expression unit. GpA, rabbit- β -globin poly(A) site; pA, SV40 early poly(A) site. (B) Cre recombinase with nuclear localization signal expressing adenovirus in which Cre expression is under the control of three different promoters: the amylase promoter and the elastase-1 promoter which are active in acinar cells, and the CAG promoter which is a nonspecific promoter.

Adenopack (Vivascience, Hannover, Germany). The titer of the adenovirus was determined by using the Rapid titer kit (Clontech, Mountain View, CA, USA). The virus stock was concentrated to 1.0×10^{10} pfu/mL.

Induction of active Ras in the pancreas. Adenovirus vectors were injected into the pancreatic ducts of 12-week-old adult male rats through the common duct as previously reported. To induce active Ras specifically in acinar cells, adenoviruses (6×10^8 pfu/rat) in which the expression of Cre recombinase was under the control of acinar cell specific promoters, either the amylase-2 (Ad-Amy-Cre) or elastase-1 (Ad-Ela-Cre) promoter, were used. To induce active Ras non-specifically, adenoviruses (6×10^8 pfu/rat) in which the expression of Cre recombinase was under the control of the non-specific CAG promoter were used.

Western blotting. Western blot analysis and detection of activated Ras protein was performed using a Ras Activation Assay kit (Upstate, Lake Placid, NY, USA) as described previously.^(3,8) Concentrations of the proteins were determined by Bio-Rad Protein assay. Proteins were separated by SDS-PAGE. After transfer to a polyvinylidene difluoride membrane, the membrane was blocked with 5% nonfat milk and then incubated for 1 h at room temperature with primary antibodies. The following antibodies were used: anti-Ras, clone Ras10 (Upstate) diluted 1:4000; HA-probe (Y-11; Santa Cruz Biotechnology, Santa Cruz, CA, USA) diluted 1:1,000; and monoclonal anti- β -actin (A5441; Sigma, St Louis, MO) diluted 1:10 000. The primary antibodies were detected using HRP-conjugated secondary antibodies (Southern Biotechnology Associates, Birmingham, AL, USA) and ECL plus (GE Healthcare UK, Buckinghamshire, UK).

Immunostaining. Tissues were fixed in 10% formalin or 4% paraformaldehyde fixative and embedded in paraffin. For Ki67, proliferating cell nuclear antigen (PCNA), and HA-tag staining, sections were boiled for 10 min in a 10-mM citrate buffer (pH

6.0) and then allowed to cool in PBS for 30 min before incubation with antibodies. For anti- α -amylase staining, section slides were incubated for 10 min in a 0.1% trypsin solution at 37°C and then washed in PBS for 5 min before incubation with antibodies.

Before staining, each section was blocked with 10% goat serum (Nichirei Bio Science, Tokyo, Japan) for 5 min at room temperature. The slides were incubated overnight at 4°C with primary antibodies against Ki67 antigen (NCL-Ki67-p; Novocastra Laboratories, Newcastle, UK), diluted 1:3000; PCNA (clone PC10; DakoCytomation, Glostrup, Denmark), diluted 1:50; HA-Tag (6E2; Cell Signaling, Danvers, MA, USA), diluted 1:100; or anti- α -amylase (A8273; Sigma, St Louis, MO, USA), diluted 1:200. Slides were incubated with secondary antibodies conjugated with Alexa Fluor488, 546, and 647 (Invitrogen, Carlsbad, CA, USA), and images were obtained with a FLUOVIEW FV300 confocal microscope (Olympus, Tokyo, Japan) or a BZ-9000 fluorescence microscope (Keyence, Osaka, Japan).

Results

Targeted activation of HA-*Kras*^{G12V} transgenes in mature acinar cells. Injection of transgenic rats with Cre recombinase expressing adenovirus resulted in excision of the stuffer DNA between the CAG promoter and the transgene and consequent expression of the transgene in infected cells (Fig. 1A). *Kras*301/327 rats were injected with Ad-Amy-Cre or Ad-Ela-Cre. Expression of HA-*Kras*^{G12V} was observed only in amylase-positive acinar cells and not in duct, centroacinar, intercalated duct, or islet cells (Table 2) (Fig. 2A; data for Ad-Ela-Cre is identical to that of Ad-Amy-Cre). Some acinar cells with nuclei with an "owl-eye" or "ground glass" appearance, which are generally used for identification of virus-infected cells,⁽⁹⁾ in rats treated with Ad-Amy-Cre or Ela-Cre were also positive for both amylase and HA (Fig. 2A-d,e). All acinar cells positive for HA were entirely negative for Ki67 (Fig. 2A).

Lack of PDA development by targeted activation of Ras^{G12V} in mature acinar cells. None of the *Kras*301/327 rats injected with Ad-Amy-Cre or Ela-Cre developed pancreatic lesions (Ad-Amy-Cre, 0 out of 5; Ad-Ela-Cre, 0 out of 7) after 8 weeks (Fig. 2B, Table 1). Similarly, none of the *Hras*250 rats injected with Ad-Amy-Cre or Ad-Ela-Cre (6×10^8 pfu/rat) developed pancreatic lesions (Ad-Amy-Cre, 0 out of 7; Ad-Ela-Cre, 0 out of 8) after 8 weeks (Table 1). In addition, *Kras*301/327 rats injected with higher titers of Ad-Amy-Cre (6×10^9 pfu/rat) did not develop pancreatic lesions (data not shown). Finally, tumor induction was not observed in injected *Kras*301/327 rats even after 6 months (data not shown).

Neoplasia development by activation of Ras^{G12V} transgenes in ductular cells. Both *Kras*301/327 and *Hras*250 rats injected with Ad-CAG-Cre (6×10^8 pfu/rat) developed pancreatic neoplasias: 22 of 22 *Kras*301/327 rats and 30 of 35 and *Hras*250 rats after 2 to 4 weeks (Table 1), as observed in our previous report.⁽³⁾ Pancreatic neoplasias were also observed in *Kras*301/327 rats injected with lower titers of Ad-CAG-Cre (6×10^7 pfu/rat) (data not shown). Activation of the transgene in the pancreatic ductal lesions of *Kras*301/327 rats was shown by Western blotting using anti-HA antibody (Fig. 3). The expression of HA-*Kras*^{G12V} was detected in pancreatic intraepithelial neoplasia (PanIN) and neoplastic lesions, but not in normal-looking pancreatic duct cells or stromal cells (Fig. 4A). Ki67 or PCNA and HA were positive in PanIN lesions (Fig. 4B) and in many neoplastic cells (Fig. 4C).

Discussion

The morphological and molecular signatures associated with human pancreas tumors suggests that duct epithelium is




Fig. 2. Acinar cell-specific expression of hemagglutinin (HA)-Kras^{G12V}. (A) Localization of amylase (blue) protein, HA-Kras^{G12V} (red) and Ki67 (green) at 2 days after injection of virus with Ad-Amy-Cre (a, b, c, d) and Ela-Cre (e). All the HA-Kras^{G12V} positive cells (red) were acinar cells; expression was not observed in duct cells (**), centroacinar cells (yellow arrowhead), or small duct cells (*). Most virally infected acinar cells positive for HA-Kras^{G12V} were indistinguishable from non-infected acinar cells by hematoxylin-eosin staining. Some infected acinar cells have nuclei with a so-called "owl-eye" (yellow arrows) or "ground glass" (red arrows) appearance. Ki67 (green) is not present in the nuclei of the cells expressing HA-Kras^{G12V} (red). Bar, 20 μ m. (B) None of the Ad-Amy-Cre (a) or the Ad-Ela-Cre (b) groups displayed any pancreatic lesions, even after 8 weeks. Bar, 50 μ m.

responsible for the development of PDA, but it remains unclear whether other pancreatic cells might also contribute to the cyto-genesis of these lesions. In our previous study using the Hras250 rat, 4 weeks after injection of adenovirus with Cre recombinase under the control of the constitutive CAG promoter, proliferative lesions in the duct epithelium, intercalated ducts, and centroacinar cells were widespread, but we could not detect any proliferative lesions in acinar cells; moreover, subsequent neoplastic lesions expressed only duct cell-specific characteristics and not acinar cell-specific ones.⁽³⁾ We have obtained essentially

identical results with Kras transgenic rats as we did with Hras250 rats (data not shown). These results suggest that PDAs may arise from centroacinar cells, intercalated duct, or pancreatic duct epithelium, but not from acinar cells.

The current study was undertaken to clarify whether mature acinar cells in adult rats could be induced to develop to PDA by targeted activation of oncogenic *ras*. Activation of oncogenic *ras* in acinar cells did not lead to the development of any observable pancreatic lesions, while nonspecific activation of oncogenic *ras* in the pancreas resulted in rapid development of

Table 1. Pancreas tumor induction by activation of Hras^{G12V} or hemagglutinin (HA)-Kras^{G12V} oncogene after Cre-adenovirus injection

Oncogene	Virus vector	Number of rats with tumor (%)
Hras ^{G12V}	Amylase-Cre	0/7 (0)
	Elastase-Cre	0/8 (0)
	CAG-Cre	30/35 (87.5)
HA-Kras ^{G12V}	Amylase-Cre	0/5 (0)
	Elastase-Cre	0/7 (0)
	CAG-Cre	22/22 (100)

Table 2. Target cell and tumor type in Hras^{G12V} and hemagglutinin (HA)-Kras^{G12V} transgenic rats

Virus vector	Target cells			Tumor yield	
	Acinar cells	Centroacinar cells	Duct cells	Acinar cells	Duct cells
Amy-Cre	+	-	-	-	-
Ela-Cre	+	-	-	-	-
CAG-Cre ⁽³⁾	+	+	+	-	+




Fig. 3. Transgene activation in *Kras301* and *327* by Western blotting. A high level of active Ras and hemagglutinin (HA)-*Kras*^{G12V} were detected in the pancreas of the Ad-CAG-Cre-treated rats. The amount of active Ras was analyzed by RBD (Ras-binding domain of Raf-1) pull-down assay followed by Western blotting with anti-pan Ras antibody. HA-*Kras*^{G12V} and endogenous total Ras was detected using anti-Pan Ras antibody. HA-*Kras*^{G12V} was detected using anti-HA antibody. β -Actin was used as a loading control.

pancreatic neoplasias. Our results clearly show that conditional expression of oncogenic *ras* in acinar cells in fully developed pancreas tissue does not result in induction of neoplasia in this model (Table 1).

Previous reports in which *Kras* was activated in immature acinar cells during embryonic development^(10,11) suggested that acinar-ductal metaplasia played a role in the development of PDA. In these models, premalignant acinar-ductal metaplasia and acinar tumor mixed with duct-like lesions developed in transgenic mice. This acinar-ductal metaplasia, however, may have occurred before the pancreas fully developed. Our model, on the other hand, targets mature acinar cells which express digestive enzymes, amylase and/or elastase, and these cells do not undergo acinar-ductal metaplasia in response to *ras* activation.

Our results are in agreement with a recent study in which the distribution of *K-RAS2* gene mutations was extensively examined in surgically resected pancreata from human patients and which concluded that ductal neoplasms of the human pancreas did not appear to arise from acinar cells.⁽¹²⁾

Kras mutations were not observed in pancreatic acinar cell carcinoma (ACC) induced in mature rats by administration of azaserine.⁽¹³⁾ Furthermore, alterations in the APC/ β -catenin pathway were detected in 23.5% of human ACC,⁽¹⁴⁾ but mutation of *Kras* was not observed.^(15,16) Thus, it is possible that APC/ β -catenin or another pathway, but not necessarily *Kras* activation, is involved in ACC development.

While pancreas cancer in the hamster model is also believed to arise from ductal epithelial hyperplasia,^(17,18) several studies using transgenic mice^(11,19-24) suggest that PDA may develop from acinar cells. In most of these studies, however, oncogenic stimuli are present during embryonic development, prior to the development of a mature pancreas. Therefore, the acinar cells which were activated and developed into neoplasias in these models could very well be at a different developmental stage to the acinar cells which are present in a mature pancreas. This is important because the majority of PDA patients are 60 years of age or older. It is highly unlikely that an oncogenic insult occur-

ring in the uterus is the root cause of most of these PDAs. Moreover, epidemiological studies indicate the incidence of PDA is closely related to lifestyle.⁽²⁵⁾ Therefore, PDA most likely develops from cells in the mature pancreas. Consequently, pancreas tumor models in which the oncogenic insult occurs during embryonic development are unlikely to be appropriate for determining the cytogenesis of human PDA.

Two models⁽²²⁻²⁴⁾ use conditional activation of Cre recombinase to activate oncogenic *ras* in adult animals: one model⁽²²⁾ uses the tet-off system to control expression of Cre recombinase and the other model^(23,24) uses the tamoxifen-estrogen receptor system to control nuclear localization of Cre recombinase. These studies had slightly conflicting results. In one study, expression of oncogenic *ras* in adult acinar cells did not by itself induce pancreatic lesions; additional treatment causing chronic pancreatitis was also needed.⁽²²⁾ In the other study, expression of oncogenic *ras* was sufficient to induce PanIN-like lesions.^(23,24) In our model, we clearly showed that while expression of oncogenic *ras* is sufficient to induce duct, intercalated duct, and/or centroacinar cells to develop into pancreatic cancers, it is not sufficient to induce acinar cells to develop into pancreatic cancers. Whether these discrepancies are due to experimental procedures, the nature of the Cre recombinase constructs used, or differences between mice and rats remains to be resolved. There are however, a few readily apparent differences. In our rat system, there is no expression of Cre recombinase in the animal until injection of adenovirus-expressing Cre recombinase, and the expression of Cre recombinase is transient. In the model which uses tamoxifen, on the other hand, Cre recombinase is expressed during embryonic development, but nuclear localization is regulated by tamoxifen.^(23,24) In this model, however, there was a low level of tamoxifen-independent recombination events resulting in expression of oncogenic *ras* in embryonic acinar cells.⁽²⁴⁾ It is possible that embryonic acinar cells expressing oncogenic *ras* did not fully differentiate in the adult pancreas; for example, in the mouse colon expression of *Kras*^{G12V} inhibits differentiation.⁽²⁶⁾ Therefore, it is possible that in the tamoxifen-estrogen regulated model,^(23,24) the acinar cells which were activated to undergo metaplasia to duct-like cells in the adult were not actually mature acinar cells. The other obvious difference is that in the model regulated by the tet-off system, two events were required to induce pancreas cancer: activation of oncogenic *Kras* and chronic pancreatitis.⁽²²⁾ Chronic pancreatitis would very likely result in the death of mature acinar cells and their replacement from a proliferative compartment. It is possible that these replacement cells are not fully mature acinar cells, again suggesting the possibility that the acinar cells which underwent metaplasia to duct-like cells were not actually mature acinar cells.

The primary aim of this study was to determine whether activation of oncogenic *Kras* in mature, digestive enzyme-secreting acinar cells would lead to pancreatic lesions. Our findings support our earlier hypothesis that PDA does not develop from *Kras* activation in mature acinar cells. It is possible, however, that PDA could develop from *Kras* activation in immature acinar cells, and in this regard we would like to emphasize the results of Guerra *et al.*⁽²²⁾ in which activation of *Kras* in the mature pancreas accompanied by chronic pancreatitis resulted in induction of PDA in transgenic mice. Importantly, chronic pancreatitis has been shown to be one of the main risk factors for PDA development in humans.^(27,28)

Other factors which may influence PDA development in our model are inflammation and fibrosis. Shortly after infection of pancreatic tissue with Cre recombinase carrying adenovirus to activate the *Kras* transgene, infiltration of macrophages and lymphocytes could be observed. This infiltration is presumably in response to viral infection. A moderate degree of inflammation, however, was still observed in the stromal tissue surrounding the tumors when PDA developed. These findings suggest that




Fig. 4. Pancreatic ductal adenocarcinoma (PDA) induced by injection of Ad-CAG-Cre in *Kras*^{301/327} rats. (A) The expression of hemagglutinin (HA)-*Kras*^{G12V} (red) was seen only in PDA lesions (on the left of photo), and not in stromal cells, acinar cells (on the right of photo), or normal pancreatic duct cells (*). Bar, 500 μ m. (B) A pancreatic intraepithelial neoplasia (PanIN) lesion was surrounded by fibrous tissue with some infiltration of inflammatory cells. Expression of proliferating cell nuclear antigen (PCNA) (green) and HA protein (red) in a PanIN lesion in rats of the CAG-Cre group. PCNA is preferentially expressed in PanIN cells. Bar, 20 μ m. (C) Expression of Ki67 (green) and HA protein (red) in PDA cells. Many PDA cells (red) are simultaneously positive for Ki67. Bar, 50 μ m.

inflammation may play a role in PDA development in this model. However, interaction between the immune system and tumors is complex and whether inflammation actually promotes PDA development in this model remains to be examined.

A current study has demonstrated that the fibrous element accompanying inflammation can also play an important role in cancer development.⁽²⁹⁾ This aspect of PDA development in our model also remains to be examined.

In summary, while there are discrepancies between different animal models of pancreatic cancer, our results indicate that expression of oncogenic *ras* in fully mature acinar cells does not induce cell proliferation or result in development of any pancreatic lesions. Thus, we conclude that mature acinar cells are not the origin of PanIN or pancreatic neoplasia in this model.

Acknowledgments

We thank Dr T. Shirai (Nagoya City University) for his kind advice for histological examination and assistance in histological specimen preparation. This work was supported in part by a Grant-in-Aid for Scientific Research (C) from Japan Society for the Promotion of Science; a Grant-in-Aid for Cancer Research (15-2, 16-13, 17S-6, 20S-8) from the Ministry of Health, Labour and Welfare, Japan; a Grant-in-Aid for Research on Nanotechnical Medical (H19-Nano-Ippan-014) from the Ministry of Health, Labour, and Welfare of Japan; a Grant-in-Aid for Research on Risk of Chemical Substances (H19-Kagaku-Ippan-006) from the Ministry of Health, Labour, and Welfare of Japan; and a Grant-in-Aid for Nagoya Ohjinkai Young Investigators Medical Research Award from Nagoya City University, Japan.

References

- 1 Beger HG, Rau B, Gansauge F, Poch B, Link KH. Treatment of pancreatic cancer: challenge of the facts. *World J Surg* 2003; 27: 1075–84.
- 2 Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007; 57: 43–66.

- 3 Ueda S, Fukamachi K, Matsuoka Y *et al.* Ductal origin of pancreatic adenocarcinomas induced by conditional activation of a human Ha-ras oncogene in rat pancreas. *Carcinogenesis* 2006; 27: 2497–510.
- 4 Kanegae Y, Lee G, Sato Y *et al.* Efficient gene activation in mammalian cells by using recombinant adenovirus expressing site-specific Cre recombinase. *Nucleic Acids Res* 1995; 23: 3816–21.

- 5 Niwa H, Yamamura K, Miyazaki J. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 1991; **108**: 193–9.
- 6 Asamoto M, Ochiya T, Toriyama-Baba H *et al.* Transgenic rats carrying human c-Ha-ras proto-oncogenes are highly susceptible to N-methyl-N-nitrosourea mammary carcinogenesis. *Carcinogenesis* 2000; **21**: 243–9.
- 7 Minami K, Okuno M, Miyawaki K *et al.* Lineage tracing and characterization of insulin-secreting cells generated from adult pancreatic acinar cells. *Proc Natl Acad Sci U S A* 2005; **102**: 15116–21.
- 8 Fukamachi K, Imada T, Ohshima Y, Xu J, Tsuda H. Purple corn color suppresses Ras protein level and inhibits 7,12-dimethylbenz[a]anthracene-induced mammary carcinogenesis in the rat. *Cancer Sci* 2008; **99**: 1841–6.
- 9 Kobayashi TK, Sato S, Tsubota K, Takamura E. Cytological evaluation of adenoviral follicular conjunctivitis by cytobrush. *Ophthalmologica* 1991; **202**: 156–60.
- 10 Schmid RM. Acinar-to-ductal metaplasia in pancreatic cancer development. *J Clin Invest* 2002; **109**: 1403–4.
- 11 Grippo PJ, Nowlin PS, Demeure MJ, Longnecker DS, Sandgren EP. Preinvasive pancreatic neoplasia of ductal phenotype induced by acinar cell targeting of mutant Kras in transgenic mice. *Cancer Res* 2003; **63**: 2016–9.
- 12 Shi C, Hong SM, Lim P *et al.* KRAS2 mutations in human pancreatic acinar-ductal metaplastic lesions are limited to those with PanIN: implications for the human pancreatic cancer cell of origin. *Mol Cancer Res* 2009; **7**: 230–6.
- 13 Longnecker DS, Curphey TJ. Adenocarcinoma of the pancreas in azaserine-treated rats. *Cancer Res* 1975; **35**: 2249–58.
- 14 Abraham SC, Wu TT, Hruban RH *et al.* Genetic and immunohistochemical analysis of pancreatic acinar cell carcinoma: frequent allelic loss on chromosome 11p and alterations in the APC/beta-catenin pathway. *Am J Pathol* 2002; **160**: 953–62.
- 15 Terhune PG, Memoli VA, Longnecker DS. Evaluation of p53 mutation in pancreatic acinar cell carcinomas of humans and transgenic mice. *Pancreas* 1998; **16**: 6–12.
- 16 Van Kranen HJ, Vermeulen E, Schoren L *et al.* Activation of c-K-ras is frequent in pancreatic carcinomas of Syrian hamsters, but is absent in pancreatic tumors of rats. *Carcinogenesis* 1991; **12**: 1477–82.
- 17 Tsutsumi M, Kondoh S, Noguchi O *et al.* K-ras gene mutation in early ductal lesions induced in a rapid production model for pancreatic carcinomas in Syrian hamsters. *Jpn J Cancer Res* 1993; **84**: 1101–5.
- 18 Tsutsumi M, Konishi Y. Precancerous conditions for pancreatic cancer. *J Hepatobiliary Pancreat Surg* 2000; **7**: 575–9.
- 19 Tuveson DA, Zhu L, Gopinathan A *et al.* Mist1-KrasG12D knock-in mice develop mixed differentiation metastatic exocrine pancreatic carcinoma and hepatocellular carcinoma. *Cancer Res* 2006; **66**: 242–7.
- 20 Hingorani SR, Petricoin EF, Maitra A *et al.* Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell* 2003; **4**: 437–50.
- 21 Wagner M, Luhrs H, Kloppel G, Adler G, Schmid RM. Malignant transformation of duct-like cells originating from acini in transforming growth factor transgenic mice. *Gastroenterology* 1998; **115**: 1254–62.
- 22 Guerra C, Schuhmacher AJ, Canamero M *et al.* Chronic pancreatitis is essential for induction of pancreatic ductal adenocarcinoma by K-Ras oncogenes in adult mice. *Cancer Cell* 2007; **11**: 291–302.
- 23 Habbe N, Shi G, Meguid RA *et al.* Spontaneous induction of murine pancreatic intraepithelial neoplasia (mPanIN) by acinar cell targeting of oncogenic Kras in adult mice. *Proc Natl Acad Sci U S A* 2008; **105**: 18913–8.
- 24 De La OJ, Emerson LL, Goodman JL *et al.* Notch and Kras reprogram pancreatic acinar cells to ductal intraepithelial neoplasia. *Proc Natl Acad Sci U S A* 2008; **105**: 18907–12.
- 25 Jiao L, Mitrou PN, Reedy J *et al.* A combined healthy lifestyle score and risk of pancreatic cancer in a large cohort study. *Arch Intern Med* 2009; **169**: 764–70.
- 26 Haigis KM, Kendall KR, Wang Y *et al.* Differential effects of oncogenic K-Ras and N-Ras on proliferation, differentiation and tumor progression in the colon. *Nat Genet* 2008; **40**: 600–8.
- 27 Lowenfels AB, Maisonneuve P, Cavallini G *et al.* Pancreatitis and the risk of pancreatic cancer. International Pancreatitis Study Group. *N Engl J Med* 1993; **328**: 1433–7.
- 28 Malka D, Hammel P, Maire F *et al.* Risk of pancreatic adenocarcinoma in chronic pancreatitis. *Gut* 2002; **51**: 849–52.
- 29 Sangai T, Ishii G, Kodama K *et al.* Effect of differences in cancer cells and tumor growth sites on recruiting bone marrow-derived endothelial cells and myofibroblasts in cancer-induced stroma. *Int J Cancer* 2005; **115**: 885–92.

