

Table 1 Nodal metastasis, tumor recurrence, and tumor-related deaths according to Allred scores for p53 in tumor-stromal fibroblasts in all patients with IDC who received neoadjuvant therapy

	Cases (%)	Number of patients (%)					
		Nodal metastasis		Tumor recurrence		Tumor-related death	
		Present	Absent	Present	Absent	Yes	No
Model 1			0.680		<0.001		0.035
Allred score	290	166 (57)	124 (43)	75 (26)	215 (74)	21 (7)	269 (93)
0	63 (22)	33 (52)	30 (48)	3 (5)	60 (95)	0	63 (100)
2	24 (8)	14 (58)	10 (42)	4 (17)	20 (83)	0	24 (100)
3	65 (22)	38 (58)	27 (42)	17 (26)	48 (74)	7 (11)	58 (89)
4	64 (22)	40 (63)	24 (37)	23 (36)	41 (64)	8 (13)	56 (87)
5	29 (10)	18 (63)	11 (37)	8 (28)	21 (72)	1 (3)	28 (97)
6	27 (9)	12 (44)	15 (56)	12 (44)	15 (56)	5 (19)	22 (81)
7	16 (6)	10 (63)	6 (37)	7 (44)	9 (56)	0	16 (100)
8	2 (1)	1 (50)	1 (50)	1 (50)	1 (50)	0	2 (100)
Model 2			<0.001		<0.001		<0.001
Allred score	273	176 (65)	97 (35)	80 (29)	193 (71)	24 (9)	249 (91)
0	142 (52)	77 (54)	65 (46)	19 (14)	123 (86)	4 (3)	138 (97)
2	4 (1)	3 (75)	1 (25)	0	4 (100)	0	4 (100)
3	39 (14)	27 (69)	12 (31)	13 (33)	26 (67)	1 (3)	38 (97)
4	44 (16)	37 (84)	7 (16)	23 (52)	21 (48)	7 (16)	37 (84)
5	33 (12)	26 (79)	7 (21)	19 (58)	14 (42)	9 (27)	24 (73)
6	9 (3)	5 (56)	4 (44)	5 (56)	4 (44)	2 (22)	7 (78)
7	1 (1)	0	1 (100)	1 (100)	0	1 (100)	0
8	1 (1)	1 (100)	0	0	1 (100)	0	1 (100)

NOTE. Model 1: Allred scores for p53 in tumor-stromal fibroblasts based on biopsy specimens obtained before neoadjuvant therapy. Model 2: Allred scores for p53 in tumor-stromal fibroblasts based on surgical specimens obtained after neoadjuvant therapy.

HER2 category in tumor cells significantly increased the trend values for relative risk for pathologic complete response (Table 2).

3.4. Factors significantly associated with tumor recurrence and tumor death

The univariate analyses of all of the cases as a whole showed that the Allred scores for p53 in tumor-stromal fibroblasts in the biopsy specimens and the surgical specimens were significantly associated with tumor recurrence and tumor-related death (Table 1, Fig. 2). In the multivariate analyses using model 1, UICC pTNM-pN category and the presence of lymph vessel invasion significantly increased the

trend values for the hazard rates (HRs) for tumor recurrence and tumor-related death (data not shown). The Allred scores for p53 in tumor-stromal fibroblasts and the presence of an FF significantly increased the trend values for the HRs for tumor recurrence, and the Allred scores for ER in tumor cells significantly increased the trend value for the HR for tumor-related death in the multivariate analyses (data not shown). When model 2 was used, the Allred scores for p53 in tumor-stromal fibroblasts, the Allred scores for ER in tumor cells, UICC pTNM-pN category, and histologic grade significantly increased the trend values for the HRs for tumor recurrence and tumor-related death, and the Allred scores for p53 in tumor cells and residual tumor size significantly increased the trend values for the HRs for tumor-related death in the multivariate analyses (data not shown).

Table 2 Multivariate analysis for pathologic complete response in all patients with IDC who received neoadjuvant therapy

	Pathologic complete response	
	Trend RR (trend 95% CI)	P for trend
UICC pTNM-pN category (N0, N1, N2, N3)	0.07 (0.02-0.27)	<.001
HER2 category in tumor cells (0, 1, 2, 3)	1.81 (1.19-2.76)	.005

Abbreviations: RR, relative risk; N0, no nodal metastasis; N1, 1 to 3 nodal metastases; N2, 4 to 9 nodal metastases; N3, 10 or more nodal metastases. NOTE. Pathologic complete responses were adjusted for UICC pTNM-pN category assessed in surgical specimens obtained after neoadjuvant therapy, and HER2 category in tumor cells, Allred scores for ERs, Allred scores for PRs, histologic grade, and tumor necrosis assessed in biopsy specimens obtained before neoadjuvant therapy.




Fig. 2 Disease-free survival curves of patients with IDC according to the Allred scores for p53 in their tumor-stromal fibroblasts assessed in biopsy specimens obtained before neoadjuvant therapy (A) and in surgical specimens obtained after neoadjuvant therapy (B). Disease-free survival of patients with IDC classified by the Allred scores for p53 in tumor-stromal fibroblasts in biopsy and surgical specimens is significantly shortened as the scores increase (A, B: $P < .001$).

In the UICC pTNM stage 0 and I group of patients with IDC who received neoadjuvant therapy, age, lymph vessel invasion, and the Allred scores for p53 in tumor-stromal fibroblasts in the biopsy specimens obtained before neoadjuvant therapy significantly increased the trend values for the HRs for tumor recurrence in the multivariate analyses (Table 3, model 1). Among the factors in the surgical specimens obtained after neoadjuvant therapy, only the Allred scores for p53 in tumor stromal fibroblasts significantly increased the trend values for the HRs for tumor recurrence in the univariate analysis (data not shown).

In the group of UICC pTNM stage II IDC patients who received neoadjuvant therapy, the Allred scores for p53 in tumor-stromal fibroblasts, the Allred scores for PRs in tumor cells, and HER2 category in tumor cells in the biopsy

specimens obtained before neoadjuvant therapy significantly increased the trend values for the HRs for tumor recurrence in the multivariate analyses (Table 4, model 1), and the Allred scores for ERs in tumor cells and the Allred scores for p53 in tumor-stromal fibroblasts in the surgical specimens obtained after neoadjuvant therapy significantly increased the trend values for the HR for tumor recurrence in the multivariate analysis (Table 4, model 2).

Table 3 Multivariate analysis for tumor recurrence in UICC pTNM stage 0 and I patients with IDC who received neoadjuvant therapy

	Tumor recurrence	
	Trend HR (trend 95% CI)	P for trend
Model 1 (n = 92)		
Age (≤ 39 , >39 y)	0.21 (0.06-0.80)	.020
Lymph vessel invasion (absent, present)	4.49 (1.09-18.38)	.036
AS for p53 in tumor-stromal fibroblasts (0, 2-8)	1.45 (1.01-2.21)	.048

Abbreviation: AS, Allred score.

NOTE. Model 1: tumor recurrence was adjusted for age, lymph vessel invasion, and Allred scores for p53 in tumor-stromal fibroblasts assessed in biopsy specimens obtained before neoadjuvant therapy.

Table 4 Multivariate analyses for tumor recurrence in UICC pTNM stage II patients with IDC who received neoadjuvant therapy

	Tumor recurrence	
	Trend HR (Trend 95% CI)	P for trend
Model 1 (n = 86)		
AS for p53 in tumor-stromal fibroblasts (0, 2-8)	1.56 (1.15-2.51)	.007
AS for PRs in tumor cells (0, 2-8)	0.83 (0.71-0.98)	.026
HER2 category in tumor cells (0, 1, 2, 3)	1.61 (1.01-2.59)	.047
Model 2 (n = 89)		
AS for ERs in tumor cells (0, 2-8)	0.87 (0.76-0.99)	.035
AS for p53 in tumor-stromal fibroblasts (0, 2-8)	1.22 (1.00-1.49)	.044

NOTE. Model 1: tumor recurrence was adjusted for the Allred scores for p53 in tumor-stromal fibroblasts, the Allred scores for PRs in tumor cells, HER2 category in tumor cells, and Allred scores for ERs in tumor cells and lymph vessel invasion in the biopsy specimens obtained before neoadjuvant therapy. Model 2: tumor recurrence was adjusted for the Allred scores for ERs in tumor cells and Allred score for p53 in tumor-stromal fibroblasts in surgical specimens obtained after neoadjuvant therapy.

Table 5 Multivariate analyses for tumor recurrence and tumor death in UICC pTNM stage III patients with IDC who received neoadjuvant therapy

	Tumor recurrence		Tumor-related death	
	Trend HR (trend 95% CI)	<i>P</i> for trend	Trend HR (trend 95% CI)	<i>P</i> for trend
Model 1 (n = 112)				
Histologic grade (1, 2, 3)	2.60 (1.51-4.52)	<.001	3.87 (1.25-12.14)	.019
AS for p53 in tumor-stromal fibroblasts (0, 2-8)	1.21 (1.04-1.37)	.010	0.93 (0.70-1.26)	.651
AS for ERs in tumor cells (0, 2-8)	0.95 (0.85-1.07)	.397	0.77 (0.63-0.94)	.014
Adjuvant therapy (no, yes)	—		0.29 (0.09-0.97)	.043
HER 2 category in tumor cells (0, 1, 2, 3)	—		1.71 (1.01-2.91)	.048
Model 2 (n = 120)				
AS for ERs in tumor cells (0, 2-8)	0.87 (0.80-0.94)	<.001	0.77 (0.66-0.90)	<.001
AS for p53 in tumor-stromal fibroblasts (0, 2-8)	1.36 (1.14-1.63)	<.001	1.44 (1.12-1.87)	.005
Histologic grade (1, 2, 3)	1.84 (1.15-2.91)	.013	6.00 (1.96-18.31)	.002
AS for p53 in tumor cells (0, 2-8)	1.14 (1.01-1.28)	.040	1.09 (0.85-1.41)	.492
Residual invasive tumor size (≤20, >20 to ≤50, >50 mm)	2.21 (1.30-3.71)	.003	—	

NOTE. —, not significant in univariate analysis. Model 1: tumor recurrence was adjusted for histologic grade, the Allred scores for p53 in tumor-stromal fibroblasts, the Allred scores for ERs in tumor cells, the Allred scores for p53 in tumor cells, and Allred scores for PRs in tumor cells assessed in biopsy specimens obtained before neoadjuvant therapy, and adjusted for type of neoadjuvant therapy.

Tumor-related death was adjusted for histologic grade, the Allred scores for p53 in tumor-stromal fibroblasts, the Allred scores for ERs in tumor cells, HER 2 category in tumor cells, the Allred scores for PRs in tumor cells assessed in biopsy specimens obtained before neoadjuvant therapy, and adjusted for adjuvant therapy.

Model 2: tumor recurrence was adjusted for the Allred scores for ERs in tumor cells, the Allred scores for p53 in tumor-stromal fibroblasts and tumor cells, histologic grade, residual invasive tumor size, the Allred scores for PRs in tumor cells, tumor necrosis, lymph vessel invasion in the surgical specimens obtained after neoadjuvant therapy, and adjusted for type of neoadjuvant therapy.

Tumor-related death was adjusted for the Allred scores for ERs in tumor cells, the Allred scores for p53 in tumor-stromal fibroblasts and tumor cells, histologic grade, the Allred scores for PRs in tumor cells, HER 2 category in tumor cells, and tumor necrosis in the surgical materials obtained after neoadjuvant therapy, and adjusted for adjuvant therapy.

In the group of UICC pTNM stage III IDC patients who received neoadjuvant therapy, in model 1, histologic grade significantly increased the trend values for the HRs for tumor recurrence and tumor-related death in the multivariate analyses (Table 5). The Allred scores for p53 in tumor-stromal fibroblasts significantly increased the trend values for the HR for tumor recurrence, and the Allred scores for ERs in tumor cells, HER2 category in tumor cells, and adjuvant therapy status significantly increased the trend values for the HR for tumor-related death in the multivariate analysis (Table 5). In model 2, the Allred scores for ERs in tumor cells, the Allred scores for p53 in tumor-stromal fibroblasts, and histologic grade significantly increased the trend values for the HRs for tumor recurrence and tumor-related death in the multivariate analyses (Table 5). The Allred scores for p53 in tumor cells and residual invasive tumor size significantly increased the trend values for tumor recurrence in the multivariate analysis (Table 5).

4. Discussion

This study clearly demonstrated significant correlations between the Allred scores for ER, PR, and p53 in tumor cells and HER2 category in tumor cells assessed in the

biopsy specimens obtained before neoadjuvant therapy and in the surgical specimens obtained after neoadjuvant therapy, and these findings also confirmed the results of other studies [21-23]. It can therefore be concluded that expression of ER, PR, p53, and HER2 in tumor cells is not modified by neoadjuvant therapy. By contrast, although a marginally significant correlation was observed between the Allred scores for p53 in tumor-stromal fibroblasts in the biopsy specimens and in the surgical specimens, the Allred scores for p53 in tumor-stromal fibroblasts tended to be lower in the surgical specimens than in the biopsy specimens. The following 2 explanations for this finding appear to be possible: (1) neoadjuvant therapy down-regulates the status of p53 expression in tumor-stromal fibroblasts and (2) the fixation time interval by 10% formalin suppresses p53 immunoreactivity that reflects some reactive changes within tumor-stromal fibroblasts in biopsy specimens in surgical specimens because the time interval for tissue fixation by 10% formalin is usually shorter in biopsy specimens than in surgical specimens. Because there was a significant correlation between p53 expression in tumor cells in the biopsy specimens before neoadjuvant therapy and in the surgical specimens after neoadjuvant therapy in this study, p53 immunoreactivity in tumor-stromal fibroblasts in biopsy specimens that reflect reactive changes unrelated to nodal metastasis may be

suppressed in tumor-stromal fibroblasts in surgical specimens for the possible reasons mentioned above, thereby resulting in Allred scores for p53 in tumor-stromal fibroblasts in the surgical specimens being probably significantly associated with the presence of nodal metastasis of IDC in this study.

Negative lymph node status, tumor cells being negative for hormone receptor immunoreactivity, and tumor cells being positive for HER2 immunoreactivity have been found to be significantly associated with a pathologic complete response of patients with breast cancer in other studies [24-26], and all these factors were also significantly associated with a pathologic complete response in the univariate analyses or the multivariate analysis in this study. Although tumor fibroblasts play a significant role in regulating tumor sensitivity to a variety of chemotherapeutic agents [27], and p53 activation in tumor-stromal fibroblasts sensitizes tumors to chemotherapy [28,29], no significant association between either p53 expression in tumor-stromal fibroblasts or in tumor cells and pathologic complete response was observed in this study.

This results of this study also clearly demonstrated that Allred scores for p53 in tumor-stromal fibroblasts in biopsy or surgical specimens are a very important outcome predictive factor for IDC patients who have received neoadjuvant therapy independent of UICC pTNM status, but the outcome predictive power of the Allred scores for p53 in tumor-stromal fibroblasts assessed in the surgical specimens obtained after neoadjuvant therapy was superior to that of the Allred scores for p53 in tumor-stromal fibroblasts assessed in the biopsy specimens obtained before neoadjuvant therapy. Thus, we can conclude that the outcome predictive power of the Allred scores for p53 in tumor-stromal fibroblasts should be evaluated in surgical specimens obtained after neoadjuvant therapy.

This study did not investigate Allred scores for p53 for associations with the presence of p53 gene abnormalities in tumor-stromal fibroblasts. Although p53 mutations in tumor-stromal fibroblasts are a common lesion in primary breast cancer and other cancers and have a positive effect on cancer growth [8,30-32], some studies have shown an absence of p53 mutations in the tumor-stroma of breast cancer [33,34], and the possibility of technical problems, for example, polymerase chain reaction artifacts mimicking p53 gene abnormalities, has been pointed out by Campbell et al [35]. Thus, although the mechanism responsible for increasing the malignant potential of IDCs that is related to the expression of p53 in tumor-stromal fibroblasts should be investigated from the standpoint of p53 gene abnormalities, the p53 immunoreactivity in tumor-stromal fibroblasts may in fact reflect specific reactive changes within tumor-stromal fibroblasts that are related to the outcome of patients with IDC.

In previous studies, we and others have reported that an FF, a characteristic histologic feature of the tumor stroma in primary invasive tumors, is a very useful and accurate prognostic histologic tumor-stromal indicator for the out-

come of patients with IDC who have not received neoadjuvant therapy [12,13,36,37]. The present study clearly demonstrated that the presence of an FF (in the biopsy specimens obtained before neoadjuvant therapy, but not in the surgical specimens obtained after neoadjuvant therapy) was a factor that was significantly associated with tumor recurrence. This strongly suggests that neoadjuvant therapy produced FF-like stromal changes in the IDCs of the patients who received neoadjuvant therapy and that the true FFs in the IDCs could not be differentiated from the FF-like stromal changes. Thus, the outcome predictive power of FFs should be assessed in biopsy specimens obtained before neoadjuvant therapy.

In conclusion, this is the first study to clearly demonstrate that p53 expression by tumor-stromal fibroblasts, especially in surgical specimens obtained after neoadjuvant therapy, is strongly associated with the presence of nodal metastasis and the outcome of IDC patients who received neoadjuvant therapy. The modified Allred scoring system is very suitable for accurately assessing p53-expressing tumor-stromal fibroblasts in IDCs independent of the UICC pTNM stage of the IDC. p53 expression in tumor-stromal fibroblasts will probably become a very important target for tumor-gene therapy of IDCs of the breast in patients treated with neoadjuvant therapy.

References

- [1] Finak G, Bertos N, Pepin F, et al. Stromal gene expression predicts clinical outcome in breast cancer. *Nat Med* 2008;14:518-27.
- [2] Sheehan KM, Gulmann C, Eichler GS, et al. Signal pathway profiling of epithelial and stromal compartments of colonic carcinoma reveals epithelial-mesenchymal transition. *Oncogene* 2008;27:323-31.
- [3] Singer CF, Gschwantler-Kaulich D, Fink-Retter A, et al. Differential gene expression profile in breast cancer-derived stromal fibroblasts. *Breast Cancer Res Treat* 2008;110:273-81.
- [4] Loussouarn D, Campion L, Sagan C, et al. Prognostic impact of syndecan-1 expression in invasive ductal carcinomas. *Br J Cancer* 2008;98:1993-8.
- [5] Studebaker AW, Storci G, Werbeck JL, et al. Fibroblasts isolated from common sites of breast cancer metastasis enhance cancer cell growth rates and invasiveness in an interleukin-6-dependent manner. *Cancer Res* 2008;68:9087-95.
- [6] Hasebe T, Sasaki S, Imoto S, et al. Proliferative activity of intratumoral fibroblasts is closely correlated with lymph node and distant organ metastases of invasive ductal carcinoma of the breast. *Am J Pathol* 2000;156:1701-10.
- [7] Hasebe T, Sasaki S, Imoto S, et al. Highly proliferative fibroblasts forming fibrotic focus govern metastasis of invasive ductal carcinoma of the breast. *Mod Pathol* 2001;14:325-37.
- [8] Moinfar F, Man YG, Amould L, et al. Concurrent and independent genetic alterations in the stromal and epithelial cells of mammary carcinoma: implication for tumorigenesis. *Cancer Res* 2000;60:2562-6.
- [9] Patocs A, Zhang L, Xu Y, et al. Breast-cancer stromal cells with TP53 mutations and nodal metastases. *N Engl J Med* 2007;357:2543-51.
- [10] Bloom HJG, Richardson WW. Histological grading and prognosis in breast cancer. *Br J Cancer* 1957;11:359-77.
- [11] Gilchrist KW, Gray R, Fowble B, et al. Tumor necrosis is a prognostic predictor for early recurrence and death in lymph node-positive breast

- cancer: a 10-year follow-up study of 728 eastern cooperative oncology group patients. *J Clin Oncol* 1993;11:1929-35.
- [12] Hasebe T, Tsuda H, Hirohashi S, et al. Fibrotic focus in infiltrating ductal carcinoma of the breast: a significant histopathological prognostic parameter for predicting the long-term survival of the patients. *Breast Cancer Res Treat* 1998;49:195-208.
- [13] Hasebe T, Sasaki S, Imoto S, et al. Prognostic significance of fibrotic focus in invasive ductal carcinoma of the breast: a prospective observational study. *Mod Pathol* 2002;15:502-16.
- [14] Fisher B. Biological and clinical considerations regarding the use of surgery and chemotherapy in the treatment of primary breast cancer. *Cancer* 1977;40:574-87.
- [15] Fisher B. Adjuvant chemotherapy in the primary management of breast cancer. *Med Clin North Am* 1977;61:953-65.
- [16] Allred DC, Clark GM, Elledge R, et al. Association of p53 protein expression with tumor cell proliferation rate and clinical outcome in node-negative breast cancer. *J Natl Cancer Inst* 1993;85:200-6.
- [17] Allred DC, Harvey JM, Berardo MD, et al. Prognostic and predictive factors in breast cancer by immunohistochemical analysis. *Mod Pathol* 1998;11:155-68.
- [18] Mohsin S, Weiss H, Havighurst T, et al. Progesterone receptor by immunohistochemistry and clinical outcome in breast cancer: a validation study. *Mod Pathol* 2004;17:1545-54.
- [19] Badve SS, Baehner FL, Gray RP, et al. Estrogen- and progesterone-receptor status in ECOG2197: comparison of immunohistochemistry by local and central laboratories and quantitative reverse transcription polymerase chain reaction by central laboratory. *J Clin Oncol* 2008;26:2473-81.
- [20] Wolff AC, Hammond ME, Schwartz JN, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *Arch Pathol Lab Med* 2007;131:18-43.
- [21] Faneyte IF, Schrama JG, Peterse JL, et al. Breast cancer response to neoadjuvant chemotherapy: predictive markers and relation with outcome. *Br J Cancer* 2003;88:406-12.
- [22] Taucher S, Rudas M, Mader RM, et al. Influence of neoadjuvant therapy with epirubicin and docetaxel on the expression of HER2/neu in patients with breast cancer. *Breast Cancer Res Treat* 2003;82:207-13.
- [23] Arens N, Bleyl U, Hildenbrand R. HER2/neu, p53, Ki67, and hormone receptors do not change during neoadjuvant chemotherapy in breast cancer. *Virchows Arch* 2005;446:489-96.
- [24] Wang J, Buchholz TA, Middleton LP, et al. Assessment of histologic features and expression of biomarkers in predicting pathologic response to anthracycline-based neoadjuvant chemotherapy in patients with breast cancer. *Cancer* 2002;94:3107-14.
- [25] von Minckwitz G, Sinn HP, Raab G, et al. Clinical response after two cycle compared to HER2, Ki-67, p53, and bcl-2 in independently predicting a pathologic complete response after preoperative chemotherapy in patients with operable carcinoma of the breast. *Breast Cancer Res* 2008;10:R30.
- [26] Sullivan PS, Apple SK. Should histologic type be taken into account when considering neoadjuvant chemotherapy in breast carcinoma? *Breast J* 2009;15:146-54.
- [27] Fraser M, Bai T, Tsang BK. Akt promotes cisplatin resistance in human ovarian cancer cells through inhibition of p53 phosphorylation and nuclear function. *Int J Cancer* 2008;122:534-46.
- [28] Dudley AC, Shin SC, Cliffe AR, et al. Attenuated p53 activation in tumor-associated stromal cells accompanies decreased sensitivity to etoposide and vincristine. *Br J Cancer* 2008;99:118-25.
- [29] Lufkas D, Timis G, Papavassiliou AG, et al. p53 mutations in stromal fibroblasts sensitize tumors against chemotherapy. *Int J Cancer* 2008;123:967-71.
- [30] Bierie B, Moses HL. Under pressure: stromal fibroblasts change their ways. *Cell* 2005;123:985-7.
- [31] Hill R, Song Y, Cardiff RD, et al. Selective evolution of stromal mesenchyme with p53 loss in response to epithelial tumorigenesis. *Cell* 2006;123:1001-11.
- [32] Kiaris H, Chatzistamou I, Trimis G, et al. Evidence for nonautonomous effect of p53 tumor suppressor in carcinogenesis. *Cancer Res* 2005;65:1627-30.
- [33] Allinen M, Beroukhi R, Cai L, et al. Molecular characterization of the tumor microenvironment in breast cancer. *Cancer Cell* 2004;6:17-32.
- [34] Lebreton SC, Newgreen DF, Thompson EW, et al. Induction of epithelial to mesenchymal transition in PMC42-LA human breast carcinoma cells by carcinoma-associated fibroblast secreted factors. *Breast Cancer Res* 2007;9:R19.
- [35] Campbell IG, Qiu W, Polyak K, et al. Breast-cancer stromal cells with TP53 mutations. *N Engl J Med* 2008;10:1634-5.
- [36] Baak JP, Colpaert CG, van Diest PJ, et al. Multivariate prognostic evaluation of the mitotic activity index and fibrotic focus in node-negative invasive breast cancers. *Eur J Cancer* 2005;41:2093-101.
- [37] Colpaert C, Vermeulen PB, van Beest P, et al. Intratumoral hypoxia resulting in the presence of a fibrotic focus is an independent predictor of early distant relapse in lymph node-negative breast cancer patients. *Histopathology* 2001;39:416-25.

Original Articles

Clinical and Pathological Features of Intracystic Papillary Carcinoma of the Breast

TOMONORI AKAGI¹, TAKAYUKI KINOSHITA¹, TADAHIKO SHIEN¹, TAKASHI HOJO¹, SADAKO AKASHI-TANAKA¹, and YUSUKE MURATA²

¹Division of Breast Surgery and ²Pathological Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

Abstract

Purpose. To evaluate the clinicopathological features of intracystic papillary carcinoma (ICPC), which have not been established given its rarity and lack of standard diagnostic criteria.

Methods. We reviewed the clinicopathological findings and treatment outcomes of 14 patients with ICPC diagnosed between 2002 and 2006.

Results. Intracystic papillary carcinoma was diagnosed by fine-needle aspiration biopsy in three patients and by core-needle biopsy in six patients. A preoperative diagnosis was not made in five patients. Three patients underwent magnetic resonance imaging preoperatively, which helped to differentiate benign tumors and maintain free surgical margins. The final pathological diagnosis was invasive carcinoma in 2 (14.2%) of the 14 patients. The patients were followed up for 1–72 months, during which time only one died, of a cancer-unrelated cause.

Conclusion. Our results show that ICPC is more difficult to diagnose than common breast cancer preoperatively. Excisional biopsy was necessary when fine-needle aspiration and core-needle biopsy could not provide a diagnosis. Magnetic resonance imaging is helpful to differentiate a benign tumor from invasive disease.

Key words Intracystic papillary carcinoma · Magnetic resonance imaging

Introduction

Intracystic papillary carcinoma (ICPC) of the breast is a rare malignant tumor, accounting for fewer than 2%

of breast cancers.¹ According to the Japanese Society for Breast Cancer, ICPC includes ductal carcinoma in situ (DCIS). Several reports have described invasive ICPC with synchronous liver metastases.^{2–6} Intracystic papillary carcinoma is more difficult to diagnose than common breast cancer. Because of the lack of standard criteria for diagnosis and treatment, the clinicopathological features and treatments of this type of breast cancer have not been defined. We reviewed the clinical and pathological features of 14 patients who underwent surgery for ICPC between 2002 and 2006.

Patients and Methods

Between 2000 and 2006, 2700 cases of primary breast cancer were diagnosed at the National Cancer Center Hospital, 14 of which were diagnosed as ICPC based on clinicopathological analysis. We reviewed the clinical features, pathological findings, and treatments of these 14 patients. Immunohistochemical evaluation was performed according to the DAKO criteria, with the ABC staining method. Immunohistochemical examinations for ER and PgR were defined as positive when there was positive nuclear reactivity. Positivity was scored as follows: 0, 0% positive cells; 1+, less than 10% positive cells; 2+, 10%–50% positive cells; and 3+, more than 50% positive cells. Immunohistochemical examinations for p53 were also defined as positive when there was positive nuclear reactivity. Positivity was scored as follows: 0, 0% of positive cells; +/-, less than 10%; 1+, 10 to 50%; and 2+, more than 50%. HER2 was defined as positive depending on the cytoplasmic membrane reactivity. The grading system for HER2 was also scored from 0 to 3+ by the immunohistological method reported by Bilous et al.⁷

Table 1. Clinicopathological features of the 14 patients with intracystic papillary carcinoma

No.	Age (years)/sex	Duration from detecting the tumor to operation (months)	Location	US cystic size (mm)	US solid size (mm)	US shape of solid component	MMG shape of mass	MMG calcification	MRI	FNA	CNB	Preoperative diagnosis
1	84/F	2	A	22	5	Irregular	Irregular	None	—	—	—	Not given
2	83/F	2	D	11	6	Regular	Regular	None	—	Class 5	—	DC
3	75/F	3	A	22	7	Irregular	Irregular	A	—	Class 3	—	Not given
4	60/F	4	B	36	10	Regular	Regular	None	—	Class 2	+	Not given
5	43/F	3	A	15	3	Regular	Regular	None	—	—	+	Not given
6	36/F	9	C	34	17	Irregular	No mass	None	—	—	+	ICPC
7	57/F	4	E	10	4	Regular	Regular	None	—	Class 5	—	DC
8	70/M	6	E	50	15	Irregular	Irregular	None	—	—	+	ICPC
9	75/F	2	A	28	20	Regular	Regular	A	—	Class 5	—	DC
10	48/F	3	A	23	5	Regular	Regular	P	—	Class 2	+	Not given
11	74/F	8	A	14	14	Regular	—	—	BCP	—	+	ICPC
12	82/F	24	C	200	30	Regular	—	—	BCP	Class 2	+	ICPC
13	81/F	2	A	170	52	Irregular	Irregular	None	BCP	Class 2	+	ICPC
14	71/F	2	E	60	21	Irregular	Irregular	None	BCP	—	+	ICPC

US, ultrasonography; MMG, mammography; MRI, magnetic resonance imaging; FNA, fine-needle aspiration; CNB, core-needle biopsy; A, amorphous; P, pleomorphic; BCP, breast cancer type in MRI; DC, ductal carcinoma

Results

The clinical data are summarized in Table 1. The patients consisted of one man and 13 women and their ages ranged from 36 to 82 years (median 72.5 years). The initial manifestation was a breast lump in all patients, 13 of whom noticed the breast lump, whereas it was detected by breast cancer screening in 1 patient. The time from tumor detection to treatment ranged from 2 to 24 months (median, 5.2 months). The size of the cystic component ranged from 1 to 20 cm (mean, 4 cm), and the size of the solid component ranged from 3 to 52 mm (median, 12 mm). The tumor was located in areas A, B, C, D, and E in seven, one, two, one, and three patients, respectively.

Ultrasonography showed a multicystic lesion in one patient, and a unicystic lesion in 13 patients. All patients had solid components with intracystic growth. The cystic component ranged from 11 to 220 mm (median, 22.5 mm), and the solid component ranged from 3 to 52 mm (median, 12 mm). The solid components were variable, regular, or irregular in shape.

Twelve patients underwent mammography, which showed a smooth mass in seven, an irregular mass in four, and no mass in one. Four patients had amorphous or pleomorphic calcifications. Magnetic resonance imaging (MRI) showed a breast cancer pattern in all three patients who underwent this examination. It also showed invasion of the cystic wall in one patient.

Fine-needle aspiration was done in 8 of the 14 patients and the tumor was designated as class 5 in 3 (37%) patients, class 3 in 1, and class 2 in 4. Core-needle biopsy was done of five of the tumors designated as class 3 or class 2. Five other patients underwent core-needle biopsy without fine-needle aspiration. A diagnosis of ICPC was made in six (60%) of these ten patients. A diagnosis was not able to be made by core-needle biopsy in four patients, who required excisional biopsy for a definite diagnosis. One patient did not undergo fine-needle aspiration or core-needle biopsy preoperatively.

The pathological features are summarized in Table 2. Thirteen patients underwent mastectomy or partial mastectomy; with axillary lymph node dissection in five, without axillary lymph node dissection in four, and with sentinel lymph node dissection in four. The intracystic fluid was either serous or bloody. Pathological findings revealed invasive ICPC in 2 (14.2%) patients and DCIS was detected around the ICPC in 3 (21.4%) patients. Axillary lymph node metastasis was found in one patient. Estrogen receptor, progesterone receptor, HER2, and p53 were positive in 14 (100%), 13 (92.8%), 3 (21.4%) and 2 (14.2%) patients, respectively.

Thirteen patients were treated with tamoxifen post-operatively, and three of the eight who underwent

Table 2. Pathological findings of intracystic papillary carcinoma (ICPC)

No.	Proposed operation	Invasion of cystic wall	DCIS around ICPC	Lymph node metastasis	ER PgR HER2	p53	Histologic grade	Nuclear grade
1	Bp	–	–	No dissection	ER2 PgR2 HER2 1	–	1	1
2	Bp	–	–	No dissection	ER2 PgR0 HER2 0	–	1	1
3	Bt+sampling	–	–	0/2	ER2 PgR2 HER2 0	+	2	2
4	Bq	–	–	No dissection	ER2 PgR1 HER2 0	–	2	2
5	Bp+Ax	–	–	0/11	ER2 PgR2 HER2 0	–	2	2
6	Bp+Ax	–	+	0/22	ER2 PgR2 HER2 0	–	1	1
7	Bt+Ax	–	–	0/20	ER2 PgR2 HER2 1	–	2	2
8	Bp	–	–	No dissection	ER2 PgR2 HER2 2	2+	2	2
9	Bt+Ax	–	–	0/18	ER1 PgR1 HER2 0	–	2	3
10	Bq+SLN	–	–	0/4	ER1 PgR2 HER2 0	–	1	1
11	Bp	+	–	No dissection	ER2 PgR2 HER2 0	–	1	1
12	Bt+SLN	–	–	1/5	ER3 PgR3 HER2 0	–	1	1
13	Bt+SLN	–	+	0/5	ER3 PgR3 HER2 0	–	1	1
14	Bt+SLN	+	+	0/3	ER3 PgR2 HER2 0	–	1	1

DCIS, ductal carcinoma in situ

partial mastectomy were also treated with radiation therapy. All 14 patients were followed up for 1–72 months. At the time of writing, 13 patients were alive without evidence of recurrence and one had died of a cause unrelated to cancer.

Discussion

Intracystic papillary carcinoma is a rare type of breast cancer characterized by papillary growth within a macroscopic cyst. It accounts for fewer than 2% of all breast cancers.¹ Generally, ICPC shows no invasive growth outside of the cyst and is treated similarly to DCIS. However, there are reports of invasive ICPC with synchronous liver metastases.^{2–6} Yet, because of its rarity and the lack of diagnostic criteria, the clinicopathological features and treatments of ICPC have not been established.

The average age of onset is higher than that for the more common types of breast cancer, at about 65 years old (range, 34–92 years).^{8–10} The average age of onset in this series was 36–82 years old (median, 72.5). Some studies have reported a longer period from tumor detection to treatment for ICPC than for common breast cancer.^{8,9} In our study, it ranged from 1 to 24 months (median, 5 months), which suggests that ICPC grows more slowly than common breast cancer, and that it has a lower pathological grade and a tendency not to form ulcerations.

Intracystic papillomas are difficult to diagnose. Previous studies reported that the average age of onset was 40.7–47 years, and that 81% of intracystic papillomas in patients older than 60 years old were carcinoma.^{8–10} Intracystic papillary carcinoma tumors tend to be larger

than intracystic papillomas, but this does not necessarily help differentiate malignancy from benign growth.¹¹

Ultrasonography was thought to be a useful modality to differentiate malignant from benign tumors, but as seen in this series, the shapes of the solid components can be variable, regular, or irregular in malignant and benign tumors. Thus, several studies have found that ultrasonography is not useful for identifying benign tumors.^{11,12} Although ultrasonography can differentiate malignancy from benign tumors relatively easily when there is invasion, ICPC without invasion is difficult to diagnose with ultrasonography.

Magnetic resonance imaging is one of the most useful diagnostic techniques for common breast cancer, as it shows the patterns of the time–intensity curves of the lesion, allowing us to differentiate cancerous from benign tumors. Naoshige et al. reported that dynamic MRI imaging is very useful in the differential diagnosis of ICPC.⁶ Kusuma et al. also reported that the MRI findings correlated with the pathological findings.¹³ Only three of our patients underwent MRI, which showed malignant patterns in the time–intensity curve in all three. Moreover, in one patient it showed invasive growth outside of the cyst, corresponding to the pathological findings (Fig. 1). This finding demonstrates the strong potential of MRI to differentiate benign tumors from invasive disease.

Fine-needle aspiration or core-needle biopsy is important if the preoperative image indicates a potential malignancy. Fine-needle aspiration should be done initially, followed by core-needle biopsy, unless the fine-needle biopsy reveals class 5. In this series excisional biopsy was necessary when fine-needle aspiration or core-needle biopsy could not provide a diagnosis. It is more difficult to diagnose ICPC than common breast




Fig. 1. Dynamic magnetic resonance imaging showed malignant patterns in the time-intensity curve of the lesion

cancer because nuclear atypicality of ICPC is not prominent. Therefore, a correct diagnosis is dependent on an adequate preoperative biopsy specimen.

The treatments for ICPC and DCIS are generally the same, although cases of invasive ICPC with synchronous liver metastases have been reported.²⁻⁶ According to Yamashita et al., invasive ICPC is no longer rare and intraductal spread beyond 2 cm from the cystic wall is possible.¹⁰ In this series, two patients had invasive ICPC and another patient had axillary lymph node metastases despite no evidence of invasion in any pathologic section. It is likely that this patient had invasive disease that was missed on the available pathologic sections. Thus, it is important to obtain negative pathological surgical margins. Intracystic papillary carcinoma has the potential to be invasive, which can be evaluated by MRI. Standard neoadjuvant and adjuvant treatments have not been established and surgical resection remains the first line of treatment. The frequency of lymph node metastasis of ICPC has been reported as 0%–36%, which is lower than that of common breast cancer.^{9,14} Four patients in this series were treated with sentinel biopsy, which we have been performing in our department since 2004. It is now reasonable not to perform

axillary lymph node dissection, as sentinel biopsy is an accepted indicator of ICPC.

All of the tumors in this series were positive for estrogen and progesterone receptors, and the patients were given tamoxifen as adjuvant therapy. Eight patients who underwent breast-conserving treatment received radiation. Although no definitive conclusions about adjuvant treatments have been made, ICPC should generally be treated like DCIS.

Based on our experience and review of the literature, we conclude that it is critical to evaluate the malignant potential of ICPC and to decide on the most appropriate adjuvant treatment for each individual patient.

References

1. MacGrogan G, Moynihan F, Raju U. Tumors of the breast and female genital organs (WHO/IARC classification of tumors). 1st ed. Lyon: IARC Press; 2003. p. 79–80.
2. Hashimoto Y, Hito Y, Koike M, Itakura M, Yano S. Four cases of intracystic breast cancer (in Japanese). *Geka (J Jpn Surg)* 2006;68:365–70.
3. Okita R, Ohsumi S, Takashima S, Aogi K, Nishimura R. Synchronous liver metastases of intracystic papillary carcinoma with invasion of the breast. *Breast Cancer* 2005;12:327–30.
4. Collins LC, Carlo VP, Hwang H, Barry TS, Grown AM, Schnitt SJ. Intracystic papillary carcinoma of the breast: a reevaluation using a panel of myoepithelial cell markers. *Am J Surg Pathol* 2006;30:1002–7.
5. Ko KH, Kim EK, Park BW. Invasive papillary carcinoma of the breast presenting as posttraumatic recurrent hemorrhagic cysts. *Yonsei Med J* 2006;31:575–7.
6. Tochika N, Takano A, Yoshimoto T, Tanaka J, Sugimoto T, Kobayashi M, et al. Intracystic carcinoma of the male breast: report of a case. *Surg Today* 2001;31:806–9.
7. Bilous M, Doseff M, Hanna W, Isola J, Lebeau A, Moreno A, et al. Current perspectives on HER2 testing: a review of national testing guidelines. *Mod Pathol* 2003;16:173–82.
8. Czernobilsky B. Intracystic carcinoma of the female breast. *Surg Gynecol Obstet* 1967;124:93–8.
9. McKittrick JE, Doane WA, Failing RM. Intracystic papillary carcinoma of the breast. *Am Surg* 1969;35:195–202.
10. Yamashita A, Yoshimoto T, Iwase T, Watanabe S, Kasumi F. Clinicopathological images of intracystic breast carcinoma (in Japanese with English abstract). *Nihon Rinshyo Geka Gakkai-zasshi (J Jpn Soc Clin Surg)* 1994;55:2726–31.
11. Hayashi T, Nishida M, Sato K, Yamasaki T, Takami K, Hiraide H. Study of intracystic tumors lesions of the breast experienced at the department (in Japanese with English abstract). *Nihon Rinshyo Geka Gakkai-zasshi (J Jpn Soc Clin Surg)* 1996;57: 2355–9.
12. Saikawa Y, Kosaka A. Eight cases of intracystic cancer of the breast (in Japanese with English abstract). *Nihon Rinshyo Geka Gakkai-zasshi (J Jpn Soc Clin Surg)* 1991;52:2887–90.
13. Kusuma R, Takayama F, Tsuchiya S. MRI of the breast: comparison of MRI signals and histological characteristics of the same slices. *Med Mol Morphol* 2005;38:204–15.
14. Inayoshi A, Oshiro Y, Machida H. Evaluation of ultrasonography and needle aspiration cytology for intracystic tumors of the breast (in Japanese with English abstract). *Nihon Rinshyo Geka Gakkai-zasshi (J Jpn Soc Clin Surg)* 1999;60: 893–7.

Comparison among different classification systems regarding the pathological response of preoperative chemotherapy in relation to the long-term outcome

Tadahiko Shien · Chikako Shimizu · Kunihiko Seki · Taro Shibata · Takashi Hojo · Masashi Ando · Tsutomu Kohno · Noriyuki Katsumata · Sadako Akashi-Tanaka · Takayuki Kinoshita · Yasuhiro Fujiwara

Received: 31 January 2008 / Accepted: 5 February 2008 / Published online: 20 February 2008
© Springer Science+Business Media, LLC. 2008

Abstract Neoadjuvant chemotherapy (NAC) is increasingly used for operable disease. However there are several pathological response classification systems and the correlation between the pathological response to NAC according to each system and the patient outcome is still under debate. From 1998 to 2006, 370 primary breast cancer patients underwent curative surgical treatment after NAC containing both anthracycline and taxane at the National Cancer Center Hospital. We retrospectively evaluated the clinical and pathological response using the cTMN, Fisher's, Chevallier's, and the Japanese Breast Cancer Society classification systems (JBSC) respectively, and analyzed the correlation between each pathological response and disease free survival (DFS). Ninety-five (26%) patients had tumor recurrence. The five-year DFS according to Fisher's system was pCR, 80% and pINV, 63%. The five-year DFS according to Chevallier's system was Grade 1, 83%, Grade 2, 85%, Grade 3, 62%, and Grade 4, 65%. The five-year DFS according to the JBSC system was Grade 3, 77%, Grade 2, 68%, Grade 1a, 68%, Grade 1b, 58%, and Grade 0,

52%. None of the pathological response systems reached a statistically significant difference. In the classification by the post-treatment number of metastatic axillary lymph nodes, the 5-year DFS was $n = 0$, 86%; $n = 1-3$, 64%; $n = 4-9$, 44%; and $n > 10$ positive: 25% ($P < .0001$). In pathologically node negative patients, there were no significant differences in the DFS among all the classification systems. All three classifications analyzed were considered inadequate as the prognostic marker of the long-term outcome after NAC and further studies are warranted to optimize the prediction.

Keywords Breast cancer · Neoadjuvant · Chemotherapy · Response · Predictor

Introduction

Breast cancer has recently become the most common malignancy among Japanese women. Approximately 40,000 women are annually affected and breast cancer mortality has been increasing. National efforts to establish an early detection system by screening mammography has begun, but many of the primary cases still present with a palpable mass in the breast.

Neoadjuvant chemotherapy (NAC) has been accepted as one of the standards of care not only for locally advanced breast cancer but also for primary operable breast cancer. The disease free survival (DFS) and overall survival (OS) of patients treated with NAC is at least equivalent to those treated with post-operative adjuvant chemotherapy and the chance of breast conservation increases in patients with larger tumors [1, 2]. Although the benefit of the addition of taxane to anthracycline in the preoperative setting in terms of long-term outcome remains controversial, regimens that

T. Shien · T. Hojo · S. Akashi-Tanaka · T. Kinoshita
Department of Breast Surgery, National Cancer Center Hospital,
Tokyo, Japan

T. Shien · C. Shimizu (✉) · M. Ando · T. Kohno ·
N. Katsumata · Y. Fujiwara
Breast and Medical Oncology Division, National Cancer Center
Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan
e-mail: cshimizu@ncc.go.jp

K. Seki
Department of Pathology, National Cancer Center Hospital,
Tokyo, Japan

T. Shibata
Statistics and Cancer Control Division, National Cancer Center,
Tokyo, Japan

combine both anthracycline and taxane, either sequentially or concomitantly, are widely used.

Prognostic factors after primary chemotherapy include the clinical and pathological response to primary chemotherapy, the cTNM stage, and axillary lymph-node status after chemotherapy. “Pathological complete response (pCR)” correlates with an improved DFS and OS and has often been used as the surrogate primary endpoint for NAC. However, the classification systems for pathological response vary among studies and the system that best reflects the long-term outcome remains unidentified. Thus in this study, we applied various pathological response systems in the published literature to the same patient cohort treated with NAC including anthracycline and taxane to compare their usefulness in the prediction of the long-term outcome after NAC.

Patients and methods

Patients and treatments

All breast cancer patients treated with NAC containing both anthracycline and taxane between May 1998 and October 2006 at the National Cancer Center Hospital were extracted from the surgical database to be included in this retrospective study. NAC was indicated in patients with clinical stage II or III primary breast cancer with tumors larger than 3 cm. Core needle biopsy was performed before NAC to obtain a pathological diagnosis. The NAC regimens included (1) four cycles of doxorubicin (DOX, 50 mg/m²) and docetaxel (DOC, 60 mg/m²) (AT) followed by additional adjuvant treatment with two cycles of AT or four cycles of iv CMF (cyclophosphamide, methotrexate and 5FU), (2) four cycles of fluorouracil (500 mg/m²)/epirubicin (100 mg/m²)/cyclophosphamide (500 mg/m²) plus 12 weekly cycles of paclitaxel (80 mg/m²) (FECT), (3) four cycles of doxorubicin (60 mg/m²)/cyclophosphamide (600 mg/m²) plus 12 weekly cycles of paclitaxel (80 mg/m²) (ACT). After November 2002, in patients with HER2-overexpression tumors, trastuzumab (initially 4 mg/kg and 2 mg/kg weekly) was administered with paclitaxel for 12 weeks in the ACT and FECT treated populations (ACTH and FECTH, respectively). Five years of endocrine therapy was scheduled when either the pre-treatment biopsy specimen or surgical specimen post-chemotherapy were positive for the estrogen or progesterone receptor.

Evaluation of pathological factors

Pretreatment diagnosis was established by pathologists from a core needle biopsy specimen. Surgical specimens

were sectioned at about 7–10 mm and the pathological response was evaluated by pathologists. The expression levels of ER (1D5, Dako Cytomation), PgR (1A6, Novocastra) and HER2 (HercepTest[®], Dako Cytomation) were examined with immunohistological staining. ER and PgR were classed as positive when more than 10% of cancer cell nuclei were stained, regardless of the intensity of the staining. HER2 was scored as follows: (0): negative for cells, (1+): slightly positive in more than 10% of cancer cells, (2+): moderately positive in more than 10% of cancer cells, (3+): markedly positive in more than 10% of cancer cells. Additional fluorescent in situ hybridization (FISH) for HER2 amplification (Pathvision, Vysis) was performed and when IHC (3+) or FISH-positive (HER2/CEP17 signal ratio ≥ 2.0) were defined as HER2-positive.

The response criteria used in this study included Fisher’s system [1], Chevalier’s system [3] and the histological response criteria of the Japanese Breast Cancer Society (JBSC) [4, 5]. The key definitions of each response classification system are described in Table 1. To summarize, Fisher’s system evaluated only the histological evidence of invasive disease in the primary tumor, Chevallier’s system incorporated nodal status and the JBSC system measured morphological changes in of the tumor cells and the proportion of histological changes in the primary tumor. The histological effect in both the primary tumor and axillary lymph node should be separately evaluated in the JBSC system, but the standard of how to combine the effect is not mentioned. Therefore we used the pathological response in only the primary lesion in the present study.

In addition, we evaluated pretreatment clinical staging, the clinical response to preoperative chemotherapy and postoperative pathological lymph node status. The clinical response to preoperative chemotherapy was decided from the two greatest perpendicular diameters (before each chemotherapy treatment and before surgery) of tumors in the breast and axillary lymph nodes. No clinical evidence of palpable tumor in the breast and axillary lymph nodes was defined as a clinical complete response (cCR). A reduction in the total tumor size of 30% or greater was graded as clinical partial response (cPR). An increase in total tumor size of more than 20% or appearance of new suspicious ipsilateral axillary adenopathy was considered progressive disease (cPD). Tumors that did not meet the criteria for objective response and progression were considered as stable disease (cSD).

Statistical analysis

Disease free survival (DFS) was calculated from the date that NC was initiated to the date of the first relapse including loco-regional recurrence or the last visit without

Table 1 Pathological response classification systems

Classification system	Key definitions
Fisher et al. [1]	Classification based on microscopic evidence pCR = no histological evidence of invasive tumor cells (specimens with only noninvasive cells included) pINV = histological evidence of invasive disease of any extent
Chevallier et al. [3]	Classification using both microscopic and macroscopic evidence Grade 1 (pCR) = disappearance of all tumor on either macroscopic or microscopic assessment Grade 2 = presence of in situ carcinoma in the breast, no invasive tumor, no tumor in the ALNs Grade 3 = presence of invasive carcinoma with stromal alteration Grade 4 = no/few modifications of the tumor appearance
JBCS	Classification using both microscopic and macroscopic evidence in primary tumor Grade 0 = no therapeutic effect Grade 1 = <66% therapeutic effect, but >33% effect evident Grade 2 = subjectively >66% therapeutic effect, but <near total therapeutic effect Grade 3 = disappearance of all tumor on either macroscopic or microscopic assessment

pCR, pathological complete response; ALN, Axillary lymph node; pINV, pathological invasive disease; JBCS, Japanese breast cancer society

relapse. Kaplan–Meier plots and the log-rank test were used to assess the difference in survival. All comparisons were two-tailed. Cox-proportional hazards models were fitted for OS and DFS and included variables identified a priori as being associated with survival and the ALN status. Other variables not identified a priori were entered into the model one at a time and assessed for statistical significance. All pair-wise interactions were tested. The fit of the model and the proportional hazards assumption were assessed visually with residual plots. The statistical significance level (P) was taken as a measure of the strength of evidence against the null hypothesis, and $P < .05$ was considered statistically significant.

Results

Three hundred and seventy patients with operable breast cancer were included in this study. Table 2 lists the patient and tumor characteristics. The median age was 50 years (26–71) and 192 (52%) patients were over the age of 50. Clinical staging at diagnosis was IIA in 104 (28%), IIB in 114 (31%), IIIA in 75 (20%) and IIIB in 77 (21%). ER and PgR positive patients were respectively 148 (40%) and 152 (41%). 183 (49%) patients were treated with AT, 73 (20%) with ACT and 90 (24%) with FECT. Trastuzumab as administered to four patients among the ACT-treated patients (ACTH) and 20 among the FECT-treated patients. Ten percent of patients with HER2-positive breast cancer received trastuzumab in this study.

Ninety-six patients (26%) had tumor recurrence with a median follow-up of 45 months (range 4–104). Nine patients had only loco-regional recurrence without distant metastasis. Only 42 patients died within this period.

The clinical and pathological response results are shown in Table 3. The overall clinical response rate to NAC was 88% (cCR + cPR) and the cCR rate was 28%. According to Fisher's classification, pCR and pINV was 65 (18%) and 305 (82%). According to Chevallier's classification, 30 (8%) patients achieved a Grade 1 (disappearance of all

Table 2 Patient and tumor characteristics

Parameter	No. of patients (%)
Total	370
Age	
Age <50	179 (48)
Age >51	191 (52)
Pretreatment pathology	
Invasive ductal carcinoma	347 (94)
Invasive lobular carcinoma	13 (4)
Mucinous carcinoma	7 (2)
Others	3 (1)
Hormone receptors	
ER positive	148 (40)
PgR positive	152 (41)
HER2	
Positive (>2+)	132 (36)
Neoadjuvant chemotherapy	
AT	183 (49)
ACT	73 (20)
ACTH	4 (1)
FECT	90 (24)
FECTH	20 (5)
Surgery	
Partial mastectomy	136 (37)
Total mastectomy	234 (63)

Table 3 Response to neoadjuvant chemotherapy, Cox proportional hazards model for disease free survival

Parameter	No. of patients (%)	Hazard ratio (95% CI)
Fisher's classification		
pCR	65 (18)	1.00
pINV	305 (82)	1.07 (0.56–2.73)
Chevallier's classification		
Grade 1	30 (8)	1.00
Grade 2	21 (6)	1.03 (0.18–5.85)
Grade 3	172 (46)	1.00 (0.43–2.26)
Grade 4	147 (40)	1.31 (0.27–5.66)
JBCS classification		
Grade 3	34 (9)	1.00
Grade 2	102 (28)	1.39 (0.54–3.39)
Grade 1b	81 (22)	0.96 (0.36–2.32)
Grade 1a	141 (38)	0.61 (0.21–1.71)
Grade 0	12 (3)	0.50 (0.56–2.73)
Pathological lymph node status		
$n = 0$	174 (47)	1.00
$n = 1–3$	102 (28)	0.78 (0.54–1.10)
$n = 4–9$	57 (15)	1.57* (1.07–2.24)
$n > 10$	37 (10)	2.71* (1.83–3.95)
Clinical stage		
IIA	104 (28)	1.00
IIB	114 (31)	0.68 (0.45–1.01)
IIIA	75 (20)	1.23 (0.86–1.74)
IIIB	77 (21)	1.22 (0.85–1.74)
Clinical response		
CCR + cPR	324 (88)	1.00
CSD + cPD	46 (12)	1.44* (1.10–1.87)

CI, Confidence interval, * $P < .001$

tumors in either breast or lymph node) pathological response. According to the JBCS classification, there were 34 (9%) patients with Grade 3 pathological response (pathologically no residual tumor in the breast). Post-treatment pathological nodal status was negative in 174 (47%), 1–3 positive in 102 (28%), 4–9 positive in 57 (15%), and >10 positive in 37 (10%) patients, respectively. In the Cox proportional hazards model, the classification of pathological lymph node status and clinical response were identified as being independently significantly associated with patient outcomes (Table 3). Pretreatment hormone receptor status was not associated with pathological response or DFS. Inclusion of trastuzumab in NAC was associated with the pathological response in HER2-positive tumors ($P = 0.04$), but there was no statistical difference in the DFS (data not shown).

Figure 1 illustrates the Kaplan–Meier curves of the patient cohort of DFS according to each pathological response classification system (Fisher's, Chevallier's,

JBCS). Among these classification systems, Fisher's tended to show a correlation with DFS, however, it did not reach a statistically significant difference ($P = .067$). The five-year DFS rates in Grade 3, Grade 2, Grade 1a, Grade 1b and Grade according to the JBCS system were 77%, 68%, 68%, 58%, and 52%, respectively ($P = .525$). According to Chevallier's system, the five-year DFS rates for Grade 1, Grade 2, Grade 3 and Grade 4 were 83%, 85%, 62% and 65%, respectively ($P = .16$).

The five-year DFS according to the number of post-treatment axillary node metastases was $n = 0$, 86%; $n = 1–3$, 64%; and $n = 4–9$, 25%. Figure 2 shows the DFS according to the pre-treatment cTMN classification, post-treatment pathological nodal status and clinical response to NAC. The pre-treatment clinical stage, clinical response to NAC and post-treatment pathological nodal status were strong predictors of DFS ($P < .0001$, $P = .0005$, $P < .0001$, respectively).

The pathological response results in post-treatment pathological node negative patients are shown in Fig. 3. Pathological node-negative patients accounted for 174 (47%) out of 370 patients. Since the number of Grade 0 patients according to the JBCS system was only two, they were excluded from the analysis. There were no significant relationships between the three pathological response classification systems and the DFS in pathologically node-negative patients. Neither clinical response ($P = .142$) nor pre-treatment clinical stage ($P = .231$) predicted DFS in node-negative patients.

Discussion

Pathological and biological markers predicting "pCR" in NAC have been evaluated in several studies [6, 7], but there is no consensus on the definition of pathological response. It is particularly unclear whether the classification needs the measurement of the extent of therapeutic effect including the disappearance of tumor cells and decrease of tumor cellularity [1–3, 8, 9]. The frequency distribution of residual tumor size was altered markedly by the inclusion of tumor cellularity, and the accurate pathologic response information may be provided the product of pathologic size and tumor cellularity [10]. The results in our study showed that the evaluation of tumor cellularity and tumor size by both Chevallier's and the JBCS classification systems was not useful for predicting prognosis in both all patients and node-negative patients. This result was in contrast to another study, where the reduction of tumor cellularity significantly correlated with the overall and disease free survival [11]. The negative finding in our study may be due to the small sample size of the study and limited number of events in each category of the

Fig. 1 Kaplan–Meier curves of disease free survival according to pathological response classification systems examined. (a) Fisher’s classification; (b) Chevallier’s classification; (c) JBCS classification





Fig. 2 Kaplan–Meier curves of disease free survival according to (a) Pathologic nodal status; (b) Clinical staging and (c) Clinical response




classification. Moreover the variety of chemotherapy regimens used as NAC may have affected the result. Particularly, trastuzumab was used only in the recent HER2-positive patient cohort.

However, studies including ours indicate the importance of incorporating the pathological nodal status in the prediction of prognosis for patients after NAC [12–15]. Fisher’s classification is the most popular classification

system using major clinical trials such as NSABP trials, but this classification system is diagnosed simply based on the disappearance of invasive tumor cells, regardless of non-invasive tumor cells, only in the primary tumor. Although Fisher’s system is simple, objective and its usefulness as a predictive marker has been validated [1–3, 9, 14], incorporation of the therapeutic effect in axillary lymph nodes may be necessary for more precise outcome prediction.

Fig. 3 Kaplan–Meier curves of disease free survival in node negative patients (a) Fisher's classification; (b) Chevallier classification; (c) JBCS classification; (d) Clinical staging; (e) Clinical response



On the other hand, clinical response was the significant predictor of the disease free survival in this study as reported in several other papers [13, 16–19]. Clinical response reflects the activity of chemotherapeutic agents. Clinical responders had a better prognosis compared with non-responders. The pretreatment clinical stage correlated with disease free survival, but there were good responders among the patients with advanced primary lesions and clinically positive axillary lymph nodes. Although pCR significantly correlated with the clinical response, the importance of the clinical response in outcome prediction may remain in patients with residual tumor or pathologically negative axillary lymph node after NAC.

In conclusion, we think that all three classifications analyzed in this study were not adequate as a prognostic marker of long-term outcome after NAC. The evaluation of the therapeutic effect in primary tumors warrants further study, especially in pathologically node-negative patients after NAC. Given the suggestion that the benefit of certain

chemotherapy regimens might be different depending on the biological tumor characteristics (e.g. hormone responsive, HER2, triple negative), the validity of pCR as a prognostic marker might better be tested independently in each biological subset. Moreover, the validity of pCR with NAC including biologically targeted drugs such as trastuzumab should also be revisited.

References

1. Fisher B, Bryant J, Wolmark N (1998) Effect of preoperative chemotherapy on the outcome of women with operable breast cancer. *J Clin Oncol* 16:2672–2685
2. Bollet MA, Sigal-Zafrani B, Gambotti L et al (2006) Pathological response to preoperative concurrent chemo-radiotherapy for breast cancer: results of a phase II study. *Eur J Cancer* 42:2286–2295
3. Chevallier B, Roche H, Olivier JP et al (1993) Pilot study of intensive induction chemotherapy (FEC-HD) results in a high histologic response rate. *Am J Clin Oncol* 16:223–228

4. Japanese Breast Cancer Society (2005) General rules for clinical and pathological recording of breast cancer. *Breast Cancer* 12:S12–S14
5. Committee for production of histopathological criteria. Japanese breast cancer society (2001) Histopathological criteria for assessment of therapeutic response in breast cancer. *Breast Cancer* 8:1
6. Petit T, Wilt M, Velten M et al (2004) Comparative value of tumour grade, hormonal receptors, Ki-67, HER-2 and topoisomerase II alpha status as predictive markers in breast cancer patients treated with neoadjuvant Anthracycline-based chemotherapy. *Eur J Cancer* 40:205–211
7. Burcombe RJ, Makris A, Richman PI et al (2005) Evaluation of ER, PgR, HER-2 and Ki-67 as predictors of response to neoadjuvant Anthracycline chemotherapy for operable breast cancer. *Br J Cancer* 92:147–155
8. Estevez LG, Gradishar WJ (2004) Evidence-based use of neoadjuvant Taxane in operable and inoperable breast cancer. *Clin Cancer Res* 10:3249–3261
9. Jones RL, Lakhani SR, Ring AE (2006) Pathological complete response and residual DCIS following neoadjuvant chemotherapy for breast carcinoma. *Br J Cancer* 94:358–362
10. Rajan R, Poniecka A, Smith TL et al (2004) Change in tumor cellularity of breast carcinoma after neoadjuvant chemotherapy as a variable in the pathologic assessment of response. *Cancer* 100:1365–1373
11. Ogston KN, Miller ID, Payne S et al (2003) A new histological grading system to assess response of breast cancers to primary chemotherapy: prognostic significance and survival. *Breast* 12:320–327
12. Bear HD, Anderson S, Smith RE et al (2006) Sequential preoperative or postoperative docetaxel added to preoperative doxorubicin plus cyclophosphamide for operable breast cancer: National Surgical Adjuvant Breast and Bowel Project Protocol B-27. *J Clin Oncol* 24:1–9
13. Pierga JY, Mouret E, Laurence V et al (2003) Prognostic factors for survival after neoadjuvant chemotherapy in operable breast cancer: the role of clinical response. *Eur J Cancer* 39:1089–1096
14. Amat S, Abrial C, Penault-Llorca F et al (2005) High prognostic significance of residual disease after neoadjuvant chemotherapy: a retrospective study in 710 patients with operable breast cancer. *Breast Cancer Res Treat* 94:255–263
15. Chollet P, Amat S, Cure H et al (2002) Prognostic significance of a complete pathological response after induction chemotherapy in operable breast cancer. *Br J Cancer* 86:1041–1046
16. Hortobagyi GN, Buzdar AU, Kau SW (1998) Management of stage III primary breast cancer with primary chemotherapy, surgery, and radiation therapy. *Cancer* 62:2507–2516
17. Cameron DA, Anderson ED, Levack P et al (1997) Primary systemic therapy for operable breast cancer-10-year survival data after chemotherapy and hormone therapy. *Br J Cancer* 76:1099–1105
18. Coudert BP, Arnold L, Moreau L et al (2006) Pre-operative systemic (neo-adjuvant) therapy with Trastuzumab and docetaxel for HER2-overexpressing stage II or stage III breast cancer: results of a multicenter phase II trial. *Ann Oncol* 17:409–419
19. Lenert JT, Vlastos G, Mirza NQ et al (1999) Primary tumor response to induction chemotherapy as a predictor of histological status of axillary nodes in operable breast cancer patients. *Ann Surg Oncol* 6:762–767

Clinicopathological Features of Tumors as Predictors of the Efficacy of Primary Neoadjuvant Chemotherapy for Operable Breast Cancer

Tadahiko Shien · Sadako Akashi-Tanaka · Kunihisa Miyakawa · Takashi Hojo · Chikako Shimizu · Kunihiko Seki · Masashi Ando · Tsutomu Kohno · Naruto Taira · Hiroyoshi Doihara · Noriyuki Katsumata · Yasuhiro Fujiwara · Takayuki Kinoshita

Published online: 25 October 2008
© Société Internationale de Chirurgie 2008

Abstract

Background Neoadjuvant chemotherapy (NC) is standard therapy for patients with locally advanced breast cancer and is increasingly used for early-stage operable disease. Clinical and pathological responses are important prognostic parameters for NC, which aims to achieve a pathological complete response or tumor reduction to reduce the volume of subsequent breast resection. Clinicopathological markers that predict patient response to NC are needed to individualize treatment.

Methods From 1998 to 2006, 368 patients with primary breast cancer underwent curative surgical treatment after NC (anthracycline and/or taxane without trastuzumab). We retrospectively evaluated the clinicopathological features and classification of the tumors using computed tomography

(CT) before NC and analyzed the correlation with the pathological complete response (pCR) and reduction of tumor size after treatment.

Results The overall response and pCR rates in these patients were 86% and 17%, respectively. In multivariate analysis, classification as a scirrhous-type tumor was an independent predictor of reduced likelihood of pCR ($p = 0.0115$; odds ratio 0.21). For tumor reduction, histological grade 3 ($p = 0.0002$; odds ratio 3.3) and localized tumors identified by using CT imaging ($p = 0.0126$; odds ratio 2.4) were independent predictors in multivariate analysis.

Conclusions In this study, NC often did not result in pCR for breast cancers classified as scirrhous. Furthermore, tumor type classification using CT imaging and histological grading was effective to predict tumor reduction in response to NC that included an anthracycline and/or a taxane.

T. Shien · S. Akashi-Tanaka · T. Hojo · T. Kinoshita
Department of Surgery, National Cancer Center Hospital,
Tokyo, Japan

T. Shien (✉) · N. Taira · H. Doihara
Department of Cancer and Thoracic Surgery,
Okayama University, 1-5-2 Shikata-cho, Okayama-Shi,
Okayama 700-8558, Japan
e-mail: tshien@md.okayama-u.ac.jp

K. Miyakawa
Nagano PET Imaging and Diagnostic Center,
Nagano, Japan

C. Shimizu · M. Ando · T. Kohno · N. Katsumata ·
Y. Fujiwara
Breast and Medical Oncology Division, National
Cancer Center Hospital, Tokyo, Japan

K. Seki
Department of Pathology, National Cancer Center Hospital,
Tokyo, Japan

Introduction

Neoadjuvant chemotherapy (NC) is used to reduce the size of locally advanced breast cancer tumors, and hence, the area to be resected, or to enable breast conservation for cases in which it was otherwise not possible. In clinical practice, because currently available anticancer drugs are extremely effective, these goals are achieved in many patients and the primary tumors completely disappear (i.e., pathological complete response (pCR)) in some patients by the end of NC. Data from large-scale studies have revealed that the patients who achieved pCR after preoperative administration of anticancer drugs have significantly better prognoses than other patients. These preoperative chemotherapy regimens primarily consist of an anthracycline. A

taxane may be added for some patients and additionally, trastuzumab is included for HER-2-positive patients. Indeed, the percentage of patients who experienced pCR increased when an anthracycline was added to their treatment regimens, and further increased with the addition of a taxane [1, 2]. With NC, limited surgery is assumed to be performed after the volume of the advanced breast cancer tumor is reduced, whereas NC is designed to extend the survival of patients by causing tumors to disappear solely by using anticancer drugs. Therefore, even those patients with breast cancer who have relatively small tumors close to their early-stage are currently treated first with anticancer drugs. Although preoperative chemotherapy has been used in wider range of cases, there are no practical criteria for its indications in terms of the results from clinicopathological examinations. Clinically, some patients show excellent responses to anticancer drugs and NC should be performed proactively, whereas other patients do not significantly benefit from these drugs and NC may not be necessary. Thus, individually predicting the efficacy of NC used for different purposes and deciding whether it should be performed is a current clinical goal.

In recent translational research, the efficacy of anticancer or hormone drugs were predicted by immunologically examining the sensitivity of the patients to these drugs [3]. As the indications of NC continue to expand, it is necessary to precisely select therapeutic methods, including the type of anticancer drugs, based on small tissue samples and laboratory test results that are available before surgeries. In the present study, we retrospectively examined cases treated at our clinic to determine whether it is possible to predict the efficacy of NC used for different purposes based on pretreatment tissue samples and the tumor shape observed using pretreatment CT imaging.

Methods

Patients and treatments

All patients diagnosed with operable breast cancer and treated between May 1998 and July 2006 at the National Cancer Center Hospital (NCCH; Tokyo, Japan) with NC, including an anthracycline and a taxane, were included in this retrospective study. NC was indicated for clinical stage II tumors and tumors >3 cm or stage III breast cancer tumors. Core-needle biopsy was performed before NC to allow a pathological diagnosis. Doxorubicin (DOX, 50 mg/m²) and docetaxel (DOC, 60 mg/m²) (AT regimen) were administered in four cycles every 3 weeks before surgery. Additional adjuvant treatment with DOX/DTX was given if the patients achieved complete or partial remission after preoperative chemotherapy or were otherwise treated with

four cycles of intravenous cyclophosphamide, methotrexate, and 5-fluorouracil. FECT treatment was four cycles of 5-fluorouracil (500 mg/m²)/epirubicin (100 mg/m²)/cyclophosphamide (500 mg/m²) plus 12 weekly cycles of paclitaxel (80 mg/m²) followed by surgery. The ACT regimen was 4 cycles of doxorubicin (60 mg/m²)/cyclophosphamide (600 mg/m²) plus 12 weekly cycles of paclitaxel (80 mg/m²) followed by surgery. The T regimen was 12 weekly cycles of paclitaxel (80 mg/m²) followed by surgery. Recently, patients with breast cancer that showed an HER-2 overexpression phenotype have received trastuzumab as PST. However, in this study we excluded these patients because we have only recently begun to use trastuzumab, and many HER-2-positive patients did not receive this treatment. Tamoxifen (20 mg/day) or anastrozole (10 mg/day) was administered for 5 years when pretreatment biopsy specimens or surgical postchemotherapy specimens were positive for estrogen receptor (ER) or progesterone receptor (PgR). The surgical treatment employed was mastectomy or breast-conserving surgery with axillary lymph node dissection (level 2) and that was decided from both of preoperative general diagnosis (palpation, MMG, US, and MDCT findings) and intraoperative pathological findings.

Evaluation of pathological factors

Pretreatment diagnoses were established by our pathologists using a core-needle biopsy or a surgical resection. The expression levels of hormone receptors and HER-2 were determined by using immunohistological examinations. Surgical specimens were sectioned to an approximately 7–10-mm thickness and pathologically classified by pathologists. Pathologic features were noted and invasive ductal carcinomas (IDCs) were classified as one of three subtypes (papillotubular, solid-tubular, and scirrhous) according to the General and Pathological Recording of Breast Cancer guideline established by the Japanese Breast Cancer Society [4]. The diagnosis of invasive lobular carcinoma was based on tumor histology showing the absence of E-cadherin by immunohistological examination on the pretreatment specimens. The criteria for histological grading of IDCs were based on a modification of those recommended by the World Health Organization [5, 6]. The response criteria used in this study include Fisher's system [7]; pCR means no histological evidence of invasive tumor cells (specimens with only noninvasive cells were included), whereas pINV indicated the presence of invasive tumor cells. The criterion for ER- and PgR-positive samples was specific signals in more than 10% of the cancer cell nuclei, regardless of intensity. HER-2 positivity was defined as 3+, i.e., markedly positive in more than 10% of the cancer cells.

Clinical responses to preoperative chemotherapy were reflected by the two greatest perpendicular diameters (before each chemotherapy treatment and before surgery) of tumors in the breast and an axillary lymph node. No clinical evidence of palpable tumor in the breast and axillary lymph nodes was defined as a clinical complete response (cCR). Reduction in the total tumor size by 30% or more was graded as a clinical partial response (cPR). An increase in the total tumor size of more than 20% or appearance of new suspicious ipsilateral axillary adenopathy was considered progressive disease (cPD). Tumors that did not meet any of the criteria for response or progression were considered unchanged (cNC).

CT imaging

CT examinations were performed with the patient in the supine position using a helical CT scanner (X-Vigor; Toshiba Medical Systems, Japan) between January and June 2000 or using an MDCT scanner (Aquilion, Toshiba) beginning in July 2000. The first noncontrast-enhanced CT scan served as the baseline with scanning performed from the cranial end of the sternum to the inframammary fold. Subsequently, an enhanced zoomed scan was obtained to visualize the entire breast. A bolus of 100 ml of nonionic contrast material (300 mgI/ml) was injected intravenously at a rate of 3 ml per second via an antecubital vein on the side opposite the affected breast using an automated injector. Image acquisition was started 40 s after the start of the bolus injection. The reconstruction interval was 5 mm.

The tumor shape was classified into two types: localized tumors visualized as single lesions and nonlocalized tumors, including those with surrounding lesions, multiple lesions, or glandular spreading (Fig. 1). CT imaging was used before both NC and surgery. The maximum tumor size measurements and the tumor shape classification were obtained using the CT images and compared with the size measured during the pathological examination.

Results

From May 1998 to July 2006, 403 patients were administered an anthracycline and/or a taxane as NC at the NCCH. Excluding the patients who received trastuzumab, the indication of which was not clear at the time of the study, concomitantly with a taxane, 368 patients who were diagnosed with breast cancer using pretreatment cutting needle biopsies were included in this study. The patient backgrounds are shown in Table 1. Among the patients, 194 (53%) were aged 50 years or younger and 174 (47%) were aged 51 years or older. The clinical stages of the patients at the first visit were IIA, IIB, IIIA, and IIIB for 29%, 31%, 13%, and 20%, respectively. According to the histological examinations of pretreatment cutting needle biopsies, 333 patients (90%) had an IDC, 19%, 36%, and 36% of which were classified as papillotubular type, solid-tubular type and scirrhous type, respectively. Other than IDC, 14 patients (4%) had an invasive lobular carcinoma (ILC) and 7 patients (2%) had a mucinous carcinoma.

Fig. 1 Classification of tumor by CT imaging. **a** Localized type. **b-1** Nonlocalized type: glandular spreading. **b-2** Nonlocalized type: tumor with surrounding lesions




Table 1 Patient and disease characteristics (*N* = 368)

Parameter	No. of patients	%
Age (years)		
≤50	194	53
≥51	174	47
Clinical stage		
IIA	105	29
IIB	114	31
IIIA	74	13
IIIB	75	20
Pretreatment pathology		
Invasive ductal carcinoma	333	90
Papillotubular type	68	19
Solid-tubular type	131	36
Scirrhous type	134	36
Invasive lobular carcinoma	14	4
Mucinous carcinoma	7	2
Other	14	4
Hormone receptors		
ER positive	150	41
PgR positive	218	59
HER2		
Positive	57	15
Histological grade		
G1	18	5
G2	169	46
G3	181	49
Neoadjuvant chemotherapy		
AC	3	1
ACT	75	20
AT	185	50
FECT	92	25
T	13	4
Surgery		
Partial mastectomy	136	37
Total mastectomy	232	63
Clinical response		
CR	99	27
PR	218	59
NC	46	13
PD	5	1
Pathological response		
pCR	64	17
pINV	304	83
Postoperative pathological tumor size (mm)		
Median	24	
Range	0–130	
No. of pathological LN metastases		
0	164	45
1–3	108	29

Table 1 continued

Parameter	No. of patients	%
4–9	58	16
≥10	38	10

PgR, progesterone receptor; *ER*, estrogen receptor; *CR*, complete response; *PR*, partial response; *NC*, neoadjuvant chemotherapy; *pCR*, pathological complete response; *LN*, lymph node

Immunohistological examinations revealed that 41%, 59%, and 15% of the patients were positive for ER, PgR, and HER-2, respectively. The histological grade was G2 and G3 in 46% and 49% of the patients, respectively, indicating that many patients had relatively high-grade disease. As NC regimens, AC, ACT, AT, FECT, and T were used in 1%, 20%, 50%, 25%, and 4% of the patients, respectively. The clinical response rate to NC was 86% (27% for cCR and 59% for cPR), and 64 patients (17%) achieved a pCR pathological response. The median postoperative pathological tumor size was 24 (range, 0–130) mm. Whereas 45% of the patients were node-negative, 16% of the patients had four or more and approximately 10% of the patients had ten or more metastatic lymph nodes. Among the 368 patients, we further examined 267 patients who underwent CT imaging before treatment (Table 2). Classification of the tumor shape based on CT imaging showed localized tumors in 65 patients (24%). The median maximum tumor size measured using pretreatment CT was 40 (range, 15–120) mm. When we compared pretreatment maximum tumor size and the postoperative pathological tumor size in these patients, the treatment reduced the maximum tumor size by 30% or more in 146 patients (55%).

Table 3 shows the results of univariate analysis performed to evaluate the relationship between the efficacy of

Table 2 Tumor characteristics in CT images (*N* = 267)

Parameter	No. of patients	%
Tumor type		
Localized type	65	24
Nonlocalized type	202	76
Pretreatment tumor size (mm)		
Median	40	
Range	15–120	
Tumor reduction rate ^a		
>30%	146	55
<30%	121	33

^a $\times 100$ (Pretreatment tumor size – pathological tumor size)/pretreatment tumor size); pretreatment tumor sizes were measured in imaging from computed tomography