

3. 16週目 検査値比較

		遅延型	即効型
16W BMI	N	60.00	64.00
	Mean	23.643	23.224
	SD	3.128	3.020
	MEDIAN	23.700	23.000
	MIN	16.40	17.90
	MAX	33.90	35.20
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.4491
	P-VALUE Wilcoxon	.	0.3497
16W 筋力 (全身)	N	60.00	64.00
	Mean	3.655	4.125
	SD	2.358	2.387
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.2725
	P-VALUE Wilcoxon	.	0.3031
16W 筋力 (腕)	N	60.00	64.00
	Mean	3.917	4.672
	SD	2.302	2.303
	MEDIAN	4.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.0704
	P-VALUE Wilcoxon	.	0.0687
16W 筋力 (脚)	N	60.00	64.00
	Mean	3.217	3.828
	SD	2.285	2.286
	MEDIAN	2.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.1391
	P-VALUE Wilcoxon	.	0.1332

(Continued)

3. 16週目 検査値比較

		遅延型	即効型
16W 握力 右	N	60.00	63.00
	Mean	33.917	35.484
	SD	7.873	7.797
	MEDIAN	32.000	35.000
	MIN	20.00	7.50
	MAX	61.00	51.50
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.2696
	P-VALUE Wilcoxon	.	0.1202
16W 握力 左	N	60.00	64.00
	Mean	34.480	34.969
	SD	14.611	8.200
	MEDIAN	31.000	35.500
	MIN	20.00	10.50
	MAX	128.00	55.50
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.8173
	P-VALUE Wilcoxon	.	0.1308
16W HBA1c	N	60.00	67.00
	Mean	6.227	6.369
	SD	1.075	1.059
	MEDIAN	5.900	6.200
	MIN	4.80	4.80
	MAX	9.20	9.90
	NMISS	2	0
	P-VALUE STUDENT T-TEST	.	0.4553
	P-VALUE Wilcoxon	.	0.3738

3. 検査値比較 差の変化量 (16W-開始時)

		遅延型	即効型
16W-開始時 BMI	N	60.00	63.00
	Mean	-0.032	0.147
	SD	0.891	0.612
	MEDIAN	0.000	0.100
	MIN	-5.00	-1.40
	MAX	1.40	2.00
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.1963
	P-VALUE Wilcoxon	.	0.3256
16W-開始時 筋力 (全身)	N	60.00	63.00
	Mean	0.088	0.016
	SD	1.221	1.238
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	4.00	5.00
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.7445
	P-VALUE Wilcoxon	.	0.7297
16W-開始時 筋力 (腕)	N	60.00	63.00
	Mean	0.017	0.079
	SD	1.790	1.274
	MEDIAN	0.000	0.000
	MIN	-4.00	-3.00
	MAX	7.00	4.00
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.8226
	P-VALUE Wilcoxon	.	0.6146
16W-開始時 筋力 (脚)	N	60.00	63.00
	Mean	0.072	0.095
	SD	1.088	1.304
	MEDIAN	0.000	0.000
	MIN	-3.00	-4.00
	MAX	4.00	5.00
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.9137
	P-VALUE Wilcoxon	.	0.5799

(Continued)

3. 検査値比較 差の変化量 (16W-開始時)

		遅延型	即効型
16W-開始時 握力 右	N	59.00	62.00
	Mean	0.436	0.595
	SD	3.323	4.038
	MEDIAN	0.200	0.500
	MIN	-6.50	-8.50
	MAX	13.50	12.50
	NMISS	3	5
	P-VALUE STUDENT T-TEST	.	0.8133
	P-VALUE Wilcoxon	.	0.9420
16W-開始時 握力 左	N	59.00	63.00
	Mean	2.717	0.673
	SD	13.517	4.073
	MEDIAN	1.000	0.000
	MIN	-9.00	-15.00
	MAX	98.00	9.00
	NMISS	3	4
	P-VALUE STUDENT T-TEST	.	0.2540
	P-VALUE Wilcoxon	.	0.8234
16W-開始時 HBA1c	N	45.00	52.00
	Mean	0.051	-0.031
	SD	0.490	0.475
	MEDIAN	0.000	0.000
	MIN	-1.40	-1.60
	MAX	2.20	1.00
	NMISS	17	15
	P-VALUE STUDENT T-TEST	.	0.4063
	P-VALUE Wilcoxon	.	0.8012

3. 28週目 検査値比較

		遅延型	即効型
28W BMI	N	60.00	65.00
	Mean	23.683	23.288
	SD	3.254	3.362
	MEDIAN	23.550	23.000
	MIN	16.00	14.80
	MAX	34.70	35.50
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.5057
	P-VALUE Wilcoxon	.	0.3618
28W 筋力 (全身)	N	59.00	65.00
	Mean	3.593	4.154
	SD	2.298	2.245
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	3	2
	P-VALUE STUDENT T-TEST	.	0.1721
	P-VALUE Wilcoxon	.	0.1596
28W 筋力 (腕)	N	59.00	65.00
	Mean	4.085	4.831
	SD	2.380	2.336
	MEDIAN	3.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	3	2
	P-VALUE STUDENT T-TEST	.	0.0809
	P-VALUE Wilcoxon	.	0.0766
28W 筋力 (脚)	N	59.00	65.00
	Mean	3.169	4.015
	SD	2.175	2.315
	MEDIAN	2.000	4.000
	MIN	1.00	1.00
	MAX	8.00	9.00
	NMISS	3	2
	P-VALUE STUDENT T-TEST	.	0.0386
	P-VALUE Wilcoxon	.	0.0470

(Continued)

3. 28週目 検査値比較

		遅延型	即効型
28W 握力・右	N	60.00	64.00
	Mean	33.942	35.875
	SD	7.830	8.866
	MEDIAN	32.750	36.000
	MIN	20.00	3.50
	MAX	59.00	52.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.2017
	P-VALUE Wilcoxon	.	0.0804
28W 握力・左	N	60.00	65.00
	Mean	32.742	35.269
	SD	7.841	7.690
	MEDIAN	30.750	36.000
	MIN	18.00	10.50
	MAX	59.00	51.00
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.0714
	P-VALUE Wilcoxon	.	0.0201
28W HbA1c	N	61.00	66.00
	Mean	6.256	6.320
	SD	1.120	0.948
	MEDIAN	6.200	6.300
	MIN	4.80	4.50
	MAX	10.60	8.70
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.7283
	P-VALUE Wilcoxon	.	0.4213
28W PSA	N	51.00	63.00
	Mean	1.071	1.250
	SD	1.095	0.892
	MEDIAN	0.788	1.050
	MIN	0.19	0.09
	MAX	6.96	5.05
	NMISS	11	4
	P-VALUE STUDENT T-TEST	.	0.3394
	P-VALUE Wilcoxon	.	0.0513

3. 検査値比較 差の変化量 28W比較-開始時

		遅延型	即効型
28W-開始時 BMI	N	60.00	64.00
	Mean	-0.048	0.050
	SD	1.134	1.115
	MEDIAN	0.050	0.000
	MIN	-5.50	-7.00
	MAX	1.80	2.50
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.6273
	P-VALUE Wilcoxon	.	0.6759
28W-開始時 筋力 (全身)	N	59.00	64.00
	Mean	-0.102	0.016
	SD	1.255	1.241
	MEDIAN	0.000	0.000
	MIN	-4.00	-4.00
	MAX	5.00	3.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.6033
	P-VALUE Wilcoxon	.	0.3692
28W-開始時 筋力 (腕)	N	59.00	64.00
	Mean	0.034	0.281
	SD	1.903	1.972
	MEDIAN	0.000	0.000
	MIN	-4.00	-6.00
	MAX	7.00	7.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.4810
	P-VALUE Wilcoxon	.	0.2327
28W-開始時 筋力 (脚)	N	59.00	64.00
	Mean	-0.097	0.250
	SD	1.200	1.024
	MEDIAN	0.000	0.000
	MIN	-4.00	-4.00
	MAX	4.00	3.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.0866
	P-VALUE Wilcoxon	.	0.0163

(Continued)

3. 検査値比較 差の変化量 28W比較-開始時

		遅延型	即効型
28W-開始時 握力・右	N	59.00	63.00
	Mean	0.300	1.244
	SD	3.033	6.025
	MEDIAN	0.500	1.500
	MIN	-7.50	-31.00
	MAX	7.00	13.50
	NMISS	3	4
	P-VALUE STUDENT T-TEST	.	0.2813
	P-VALUE Wilcoxon	.	0.2031
28W-開始時 握力・左	N	59.00	64.00
	Mean	0.669	1.350
	SD	4.085	3.266
	MEDIAN	0.000	1.750
	MIN	-6.00	-5.50
	MAX	24.00	9.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.3078
	P-VALUE Wilcoxon	.	0.0972
28W-開始時 HbA1c	N	45.00	51.00
	Mean	0.053	-0.125
	SD	0.484	0.541
	MEDIAN	0.000	0.000
	MIN	-1.30	-1.70
	MAX	1.40	0.90
	NMISS	17	16
	P-VALUE STUDENT T-TEST	.	0.0928
	P-VALUE Wilcoxon	.	0.1457
28W-開始時 PSA	N	50.00	62.00
	Mean	0.242	0.380
	SD	0.826	0.684
	MEDIAN	0.051	0.198
	MIN	-0.74	-0.99
	MAX	5.46	3.76
	NMISS	12	5
	P-VALUE STUDENT T-TEST	.	0.3363
	P-VALUE Wilcoxon	.	0.0094

3. 52週目 検査値比較

		遅延型	即効型
52W PSA	N	53.00	60.00
	Mean	1.032	1.090
	SD	0.722	0.652
	MEDIAN	0.793	1.023
	MIN	0.26	0.11
	MAX	3.30	3.00
	NMISS	9	7
	P-VALUE STUDENT T-TEST	.	0.6579
	P-VALUE Wilcoxon	.	0.3710
52W HbA1c	N	57.00	62.00
	Mean	6.172	6.279
	SD	0.951	0.920
	MEDIAN	6.100	6.200
	MIN	5.00	5.00
	MAX	8.90	9.50
	NMISS	5	5
	P-VALUE STUDENT T-TEST	.	0.5337
	P-VALUE Wilcoxon	.	0.3971
52W BMI	N	61.00	66.00
	Mean	23.531	23.194
	SD	3.411	3.368
	MEDIAN	23.700	23.000
	MIN	16.60	16.00
	MAX	34.60	35.70
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.5763
	P-VALUE Wilcoxon	.	0.4617
52W 筋力 (全身)	N	61.00	66.00
	Mean	3.656	4.394
	SD	2.128	2.183
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0563
	P-VALUE Wilcoxon	.	0.0572

(Continued)

3. 52週目 検査値比較

		遅延型	即効型
52W 筋力 (腕)	N	61.00	66.00
	Mean	3.738	4.439
	SD	2.040	2.099
	MEDIAN	3.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0587
	P-VALUE Wilcoxon	.	0.0465
52W 筋力 (脚)	N	61.00	66.00
	Mean	3.328	4.015
	SD	2.143	2.215
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0784
	P-VALUE Wilcoxon	.	0.0854
52W 握力 右	N	61.00	65.00
	Mean	34.008	36.269
	SD	8.132	7.406
	MEDIAN	33.000	36.000
	MIN	19.50	16.50
	MAX	58.50	52.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.1050
	P-VALUE Wilcoxon	.	0.0628
52W 握力 左	N	61.00	66.00
	Mean	32.415	35.136
	SD	7.629	8.262
	MEDIAN	31.000	37.000
	MIN	19.00	10.00
	MAX	55.00	54.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0566
	P-VALUE Wilcoxon	.	0.0249

(Continued)

3. 52週目 検査値比較

		遅延型	即効型
52W 骨塩定量	N	52.00	60.00
	Mean	102.98	100.35
	SD	19.589	16.614
	MEDIAN	106.00	100.50
	MIN	60.00	67.00
	MAX	147.00	143.00
	NMISS	10	7
	P-VALUE STUDENT T-TEST	.	0.4435
	P-VALUE Wilcoxon	.	0.3782

3. 検査値比較 変化量比較 (52W-開始時)

		遅延型	即効型
52W-開始時 BMI	N	61.00	65.00
	Mean	-0.121	-0.028
	SD	1.090	0.864
	MEDIAN	0.000	-0.200
	MIN	-4.50	-1.60
	MAX	1.90	2.30
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.5930
	P-VALUE Wilcoxon	.	0.6408
52W-開始時 筋力 (全身)	N	61.00	65.00
	Mean	0.016	0.262
	SD	1.408	1.314
	MEDIAN	0.000	0.000
	MIN	-3.00	-4.00
	MAX	5.00	4.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.3141
	P-VALUE Wilcoxon	.	0.0975
52W-開始時 筋力 (腕)	N	61.00	65.00
	Mean	-0.279	-0.092
	SD	1.496	1.783
	MEDIAN	0.000	0.000
	MIN	-4.00	-5.00
	MAX	3.00	4.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.5275
	P-VALUE Wilcoxon	.	0.5556
52W-開始時 筋力 (脚)	N	61.00	65.00
	Mean	0.103	0.262
	SD	1.179	1.189
	MEDIAN	0.000	0.000
	MIN	-3.00	-3.00
	MAX	4.00	4.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.4550
	P-VALUE Wilcoxon	.	0.1739

(Continued)

3. 検査値比較 変化量比較 (52W-開始時)

		遅延型	即効型
52W-開始時 握力 右	N	60.00	64.00
	Mean	0.545	1.592
	SD	3.044	4.564
	MEDIAN	0.500	1.000
	MIN	-9.00	-12.00
	MAX	8.00	16.50
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.1380
	P-VALUE Wilcoxon	.	0.2012
52W-開始時 握力 左	N	60.00	65.00
	Mean	0.722	1.022
	SD	4.402	4.922
	MEDIAN	-0.250	1.500
	MIN	-8.00	-30.00
	MAX	27.00	9.50
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.7210
	P-VALUE Wilcoxon	.	0.0391
52W-開始時 HbA1c	N	44.00	52.00
	Mean	0.020	-0.152
	SD	0.532	0.562
	MEDIAN	0.050	0.000
	MIN	-1.60	-2.10
	MAX	1.00	0.90
	NMISS	18	15
	P-VALUE STUDENT T-TEST	.	0.1282
	P-VALUE Wilcoxon	.	0.0771
52W-開始時 PSA	N	51.00	59.00
	Mean	0.160	0.240
	SD	0.392	0.396
	MEDIAN	0.076	0.160
	MIN	-0.77	-0.96
	MAX	1.80	1.33
	NMISS	11	8
	P-VALUE STUDENT T-TEST	.	0.2890
	P-VALUE Wilcoxon	.	0.1177

(Continued)

3. 検査値比較 変化量比較 (52W-開始時)

		遅延型	即効型
52W-開始時 骨塩定量	N	52.00	57.00
	Mean	4.712	3.488
	SD	8.338	4.645
	MEDIAN	2.300	3.000
	MIN	-9.00	-8.00
	MAX	45.94	18.00
	NMISS	10	10
	P-VALUE STUDENT T-TEST	.	0.3402
	P-VALUE Wilcoxon	.	0.8744

4. QOL比較 52週目

		遅延型	即効型
PF	N	55.00	55.00
	Mean	83.505	91.091
	SD	16.438	9.264
	MEDIAN	90.000	95.000
	MIN	20.00	60.00
	MAX	100.00	100.00
	NMISS	7	12
	P-VALUE STUDENT T-TEST	.	0.0035
	P-VALUE Wilcoxon	.	0.0019
PR	N	55.00	55.00
	Mean	83.977	92.045
	SD	26.455	13.091
	MEDIAN	100.00	100.00
	MIN	0.00	50.00
	MAX	100.00	100.00
	NMISS	7	12
	P-VALUE STUDENT T-TEST	.	0.0451
	P-VALUE Wilcoxon	.	0.3484
BP	N	55.00	54.00
	Mean	74.473	74.870
	SD	25.014	19.406
	MEDIAN	84.000	73.000
	MIN	21.00	31.00
	MAX	100.00	100.00
	NMISS	7	13
	P-VALUE STUDENT T-TEST	.	0.9264
	P-VALUE Wilcoxon	.	0.8168
GH	N	55.00	54.00
	Mean	60.514	60.213
	SD	19.091	15.726
	MEDIAN	57.000	57.000
	MIN	17.00	20.00
	MAX	100.00	100.00
	NMISS	7	13
	P-VALUE STUDENT T-TEST	.	0.9287
	P-VALUE Wilcoxon	.	0.8645

(Continued)

4. QOL比較 52週目

		遅延型	即効型
VT	N	55.00	54.00
	Mean	70.379	69.676
	SD	20.955	18.066
	MEDIAN	75.000	71.875
	MIN	6.25	31.25
	MAX	100.00	100.00
	NMISS	7	13
	P-VALUE STUDENT T-TEST	.	0.8517
	P-VALUE Wilcoxon	.	0.6130
SF	N	56.00	54.00
	Mean	82.143	85.880
	SD	27.163	20.753
	MEDIAN	100.00	100.00
	MIN	12.50	37.50
	MAX	100.00	100.00
	NMISS	6	13
	P-VALUE STUDENT T-TEST	.	0.4205
	P-VALUE Wilcoxon	.	0.6577
RE	N	53.00	55.00
	Mean	88.050	91.667
	SD	22.018	12.108
	MEDIAN	100.00	100.00
	MIN	0.00	50.00
	MAX	100.00	100.00
	NMISS	9	12
	P-VALUE STUDENT T-TEST	.	0.2903
	P-VALUE Wilcoxon	.	0.8048
MH	N	55.00	54.00
	Mean	78.545	76.111
	SD	18.173	16.981
	MEDIAN	85.000	80.000
	MIN	25.00	35.00
	MAX	100.00	100.00
	NMISS	7	13
	P-VALUE STUDENT T-TEST	.	0.4717
	P-VALUE Wilcoxon	.	0.3459

(Continued)

4. QOL比較 52週目

		遅延型	即効型
AMS (心理的)	N	62.00	67.00
	Mean	6.855	6.537
	SD	3.715	4.020
	MEDIAN	7.000	7.000
	MIN	0.00	0.00
	MAX	19.00	15.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.6429
	P-VALUE Wilcoxon	.	0.9753
AMS (身体的)	N	62.00	67.00
	Mean	12.065	12.045
	SD	5.906	7.233
	MEDIAN	12.000	14.000
	MIN	0.00	0.00
	MAX	25.00	26.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9865
	P-VALUE Wilcoxon	.	0.6893
AMS (性機能)	N	62.00	67.00
	Mean	12.048	10.821
	SD	5.846	6.367
	MEDIAN	13.000	13.000
	MIN	0.00	0.00
	MAX	22.00	21.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.2574
	P-VALUE Wilcoxon	.	0.3813
国際前立腺症状スコア計	N	62.00	67.00
	Mean	10.435	8.597
	SD	9.248	8.907
	MEDIAN	8.500	6.000
	MIN	0.00	0.00
	MAX	34.00	33.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.2523
	P-VALUE Wilcoxon	.	0.1950

(Continued)

4. QOL比較 52週目

		遅延型	即効型
国際勃起能スコア5	N	62.00	67.00
	Mean	6.032	8.761
	SD	6.435	7.552
	MEDIAN	5.000	8.000
	MIN	0.00	0.00
	MAX	21.00	24.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.0296
	P-VALUE Wilcoxon	.	0.0363

4. QOL比較 差の変化量 (52W-開始時)

			遅延型	即効型
52W-開始時	PF	N	54.00	53.00
		Mean	0.231	-0.451
		SD	10.952	7.692
		MEDIAN	0.000	0.000
		MIN	-40.00	-35.00
		MAX	20.00	25.00
		NMISS	8	14
		P-VALUE STUDENT T-TEST	.	0.7105
		P-VALUE Wilcoxon	.	0.4333
52W-開始時	PR	N	53.00	53.00
		Mean	-3.774	2.005
		SD	21.278	16.706
		MEDIAN	0.000	0.000
		MIN	-100.00	-43.75
		MAX	37.50	50.00
		NMISS	9	14
		P-VALUE STUDENT T-TEST	.	0.1230
		P-VALUE Wilcoxon	.	0.1914
52W-開始時	BP	N	54.00	53.00
		Mean	0.593	-0.415
		SD	23.728	22.468
		MEDIAN	0.000	0.000
		MIN	-59.00	-48.00
		MAX	49.00	69.00
		NMISS	8	14
		P-VALUE STUDENT T-TEST	.	0.8220
		P-VALUE Wilcoxon	.	0.7103
52W-開始時	GH	N	54.00	54.00
		Mean	0.264	1.764
		SD	13.820	13.670
		MEDIAN	0.000	3.250
		MIN	-37.00	-40.00
		MAX	35.00	40.00
		NMISS	8	13
		P-VALUE STUDENT T-TEST	.	0.5719
		P-VALUE Wilcoxon	.	0.5410

(Continued)

4. QOL比較 差の変化量 (52W-開始時)

		遅延型	即効型	
52W-開始時	VT	N	54.00	54.00
		Mean	-1.350	2.083
		SD	19.669	19.197
		MEDIAN	0.000	0.000
		MIN	-50.00	-43.75
		MAX	43.75	50.00
		NMISS	8	13
		P-VALUE STUDENT T-TEST	.	0.3607
		P-VALUE Wilcoxon	.	0.4917
52W-開始時	SF	N	55.00	54.00
		Mean	-1.364	-1.389
		SD	26.860	21.122
		MEDIAN	0.000	0.000
		MIN	-75.00	-50.00
		MAX	50.00	50.00
		NMISS	7	13
		P-VALUE STUDENT T-TEST	.	0.9957
		P-VALUE Wilcoxon	.	0.7097
52W-開始時	RE	N	52.00	54.00
		Mean	-2.564	3.704
		SD	17.661	19.200
		MEDIAN	0.000	0.000
		MIN	-58.33	-33.33
		MAX	58.33	75.00
		NMISS	10	13
		P-VALUE STUDENT T-TEST	.	0.0835
		P-VALUE Wilcoxon	.	0.1522
52W-開始時	MH	N	54.00	54.00
		Mean	0.741	0.532
		SD	17.277	17.197
		MEDIAN	0.000	0.000
		MIN	-40.00	-45.00
		MAX	50.00	60.00
		NMISS	8	13
		P-VALUE STUDENT T-TEST	.	0.9500
		P-VALUE Wilcoxon	.	0.9285

(Continued)