

3. 16週目 検査値比較 完全型

		遅延型	即効型
16W 握力 左	MAX	128.00	48.50
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.8891
	P-VALUE Wilcoxon	.	0.0900
16W HBA1 c	N	42.00	43.00
	Mean	6.100	6.200
	SD	0.952	1.015
	MEDIAN	5.800	6.000
	MIN	4.80	4.80
	MAX	8.10	8.70
	NMISS	1	0
	P-VALUE STUDENT T-TEST	.	0.6408
	P-VALUE Wilcoxon	.	0.7051

3. 16週目 検査値比較 境界型

		遅延型	即効型
16W BMI	N	16.00	22.00
	Mean	24.481	22.955
	SD	3.619	2.869
	MEDIAN	24.500	22.900
	MIN	19.30	18.70
	MAX	33.90	27.70
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.1555
	P-VALUE Wilcoxon	.	0.2611
16W 筋力 (全身)	N	16.00	22.00
	Mean	4.375	4.364
	SD	2.156	2.321
	MEDIAN	4.000	4.500
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.9878
	P-VALUE Wilcoxon	.	0.9762
16W 筋力 (腕)	N	16.00	22.00
	Mean	4.375	5.409
	SD	2.630	2.384
	MEDIAN	4.500	5.500
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1

(Continued)

3. 16週目 検査値比較 境界型

		遅延型	即効型
16W 筋力 (腕)	P-VALUE STUDENT T-TEST	.	0.2142
	P-VALUE Wilcoxon	.	0.2254
16W 筋力 (脚)	N	16.00	22.00
	Mean	3.938	4.045
	SD	2.294	2.193
	MEDIAN	4.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.8840
16W 握力 右	P-VALUE Wilcoxon	.	0.8460
	N	16.00	22.00
16W 握力 右	Mean	36.594	36.114
	SD	9.990	9.639
	MEDIAN	37.500	37.000
	MIN	21.50	7.50
	MAX	61.00	51.50
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.8821
	P-VALUE Wilcoxon	.	0.7448
16W 握力 左	N	16.00	22.00
	Mean	37.675	36.500
	SD	9.928	9.261
	MEDIAN	36.150	38.000
	MIN	20.00	17.50

(Continued)

3. 16週目 検査値比較 境界型

		遅延型	即効型
16W 握力 左	MAX	58.00	55.50
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.7101
	P-VALUE Wilcoxon	.	0.8244
16W HbA1c	N	16.00	23.00
	Mean	6.519	6.730
	SD	1.350	1.074
	MEDIAN	6.100	6.500
	MIN	4.90	5.30
	MAX	9.20	9.90
	NMISS	1	0
	P-VALUE STUDENT T-TEST	.	0.5892
	P-VALUE Wilcoxon	.	0.4574

3. 検査値比較 差の変化量 (16W-開始時) 完全型

		遅延型	即効型
16W-開始時 BMI	N	42.00	40.00
	Mean	-0.124	0.111
	SD	0.965	0.568
	MEDIAN	-0.100	0.100
	MIN	-5.00	-1.40
	MAX	1.20	1.30
	NMISS	1	3
	P-VALUE STUDENT T-TEST	.	0.1860
	P-VALUE Wilcoxon	.	0.2512
16W-開始時 筋力 (全身)	N	42.00	40.00
	Mean	0.102	0.125
	SD	1.316	1.343
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	4.00	5.00
	NMISS	1	3
	P-VALUE STUDENT T-TEST	.	0.9388
	P-VALUE Wilcoxon	.	0.4444
16W-開始時 筋力 (腕)	N	42.00	40.00
	Mean	0.095	0.125
	SD	2.081	1.324
	MEDIAN	0.000	0.000
	MIN	-4.00	-3.00
	MAX	7.00	4.00
	NMISS	1	3

(Continued)

3. 検査値比較 差の変化量 (16W-開始時) 完全型

		遅延型	即効型
16W-開始時 筋力 (腕)	P-VALUE STUDENT T-TEST	.	0.9390
	P-VALUE Wilcoxon	.	0.8118
16W-開始時 筋力 (脚)	N	42.00	40.00
	Mean	0.214	0.200
	SD	1.159	1.305
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	4.00	5.00
	NMISS	1	3
	P-VALUE STUDENT T-TEST	.	0.9583
	P-VALUE Wilcoxon	.	0.8534
16W-開始時 握力 右	N	41.00	39.00
	Mean	0.317	0.367
	SD	3.144	3.748
	MEDIAN	0.500	0.500
	MIN	-6.50	-7.00
	MAX	6.00	8.50
	NMISS	2	4
	P-VALUE STUDENT T-TEST	.	0.9489
	P-VALUE Wilcoxon	.	0.8170
16W-開始時 握力 左	N	41.00	40.00
	Mean	3.232	0.598
	SD	16.117	3.966
	MEDIAN	0.000	-0.500
	MIN	-9.00	-6.50

(Continued)

3. 検査値比較 差の変化量 (16W-開始時) 完全型

		遅延型	即効型
16W-開始時 握力 左	MAX	98.00	9.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.3183
	P-VALUE Wilcoxon	.	0.7621
16W-開始時 HBA1c	N	33.00	33.00
	Mean	0.015	-0.030
	SD	0.411	0.425
	MEDIAN	0.000	0.000
	MIN	-1.40	-1.40
	MAX	1.30	1.00
	NMISS	10	10
	P-VALUE STUDENT T-TEST	.	0.6603
	P-VALUE Wilcoxon	.	0.7170

3. 検査値比較 差の変化量 (16W-開始時) 境界型

		遅延型	即効型
16W-開始時 BMI	N	16.00	22.00
	Mean	0.119	0.168
	SD	0.627	0.677
	MEDIAN	0.250	0.150
	MIN	-1.10	-1.30
	MAX	1.00	2.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.8201
	P-VALUE Wilcoxon	.	0.8939
16W-開始時 筋力 (全身)	N	16.00	22.00
	Mean	0.188	-0.227
	SD	0.911	1.020
	MEDIAN	0.000	0.000
	MIN	-1.00	-3.00
	MAX	2.00	1.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.2042
	P-VALUE Wilcoxon	.	0.3635
16W-開始時 筋力 (腕)	N	16.00	22.00
	Mean	-0.125	0.000
	SD	0.806	1.234
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	1.00	3.00
	NMISS	1	1

(Continued)

3. 検査値比較 差の変化量 (16W-開始時) 境界型

		遅延型	即効型
16W-開始時 筋力 (腕)	P-VALUE STUDENT T-TEST	.	0.7259
	P-VALUE Wilcoxon	.	0.6581
16W-開始時 筋力 (脚)	N	16.00	22.00
	Mean	-0.106	-0.091
	SD	0.511	1.342
	MEDIAN	0.000	0.000
	MIN	-1.00	-4.00
	MAX	1.00	2.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.9656
	P-VALUE Wilcoxon	.	0.5836
16W-開始時 握力 右	N	16.00	22.00
	Mean	0.763	0.914
	SD	4.014	4.644
	MEDIAN	-0.150	0.500
	MIN	-4.50	-8.50
	MAX	13.50	12.50
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.9172
	P-VALUE Wilcoxon	.	0.9055
16W-開始時 握力 左	N	16.00	22.00
	Mean	1.831	0.614
	SD	3.236	4.342
	MEDIAN	2.750	1.000
	MIN	-5.00	-15.00

(Continued)

3. 検査値比較 差の変化量 (16W-開始時) 境界型

		遅延型	即効型
16W-開始時 握力 左	MAX	7.50	7.50
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.3507
	P-VALUE Wilcoxon	.	0.2301
16W-開始時 HBA)c	N	12.00	19.00
	Mean	0.150	-0.032
	SD	0.674	0.565
	MEDIAN	0.000	0.000
	MIN	-0.50	-1.60
	MAX	2.20	0.70
	NMISS	5	4
	P-VALUE STUDENT T-TEST	.	0.4248
	P-VALUE Wilcoxon	.	0.9837

3. 28週目 検査値比較 完全型

		遅延型	即効型
28W BMI	N	41.00	41.00
	Mean	23.139	23.639
	SD	2.928	3.397
	MEDIAN	23.000	23.000
	MIN	16.00	17.50
	MAX	30.10	35.50
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.4774
	P-VALUE Wilcoxon	.	0.8419
28W 筋力(全身)	N	40.00	41.00
	Mean	3.475	4.122
	SD	2.353	2.337
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	3	2
	P-VALUE STUDENT T-TEST	.	0.2181
	P-VALUE Wilcoxon	.	0.2171
28W 筋力(腕)	N	40.00	41.00
	Mean	4.125	4.780
	SD	2.345	2.393
	MEDIAN	3.500	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	3	2

(Continued)

3. 28週目 検査値比較 完全型

		遅延型	即効型
28W 筋力(腕)	P-VALUE STUDENT T-TEST	.	0.2168
	P-VALUE Wilcoxon	.	0.2111
28W 筋力(脚)	N	40.00	41.00
	Mean	3.100	3.976
	SD	2.182	2.382
	MEDIAN	2.000	4.000
	MIN	1.00	1.00
	MAX	7.00	9.00
	NMISS	3	2
	P-VALUE STUDENT T-TEST	.	0.0886
	P-VALUE Wilcoxon	.	0.1220
28W 握力・右	N	41.00	40.00
	Mean	32.793	35.163
	SD	6.719	9.023
	MEDIAN	32.500	35.000
	MIN	20.00	3.50
	MAX	48.00	52.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.1832
	P-VALUE Wilcoxon	.	0.1145
28W 握力・左	N	41.00	41.00
	Mean	31.207	34.439
	SD	6.116	7.388
	MEDIAN	30.000	33.500
	MIN	20.50	10.50

(Continued)

3. 28週目 検査値比較 完全型

		遅延型	即効型
28W 握力・左	MAX	47.50	51.00
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.0340
	P-VALUE Wilcoxon	.	0.0145
28W HbA1c	N	42.00	42.00
	Mean	6.114	6.200
	SD	1.026	0.946
	MEDIAN	5.800	6.050
	MIN	4.80	4.50
	MAX	9.40	8.20
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.6917
P-VALUE Wilcoxon	.	0.4820	
28W PSA	N	35.00	40.00
	Mean	1.119	1.183
	SD	1.212	0.707
	MEDIAN	0.806	1.066
	MIN	0.19	0.09
	MAX	6.96	2.78
	NMISS	8	3
	P-VALUE STUDENT T-TEST	.	0.7765
	P-VALUE Wilcoxon	.	0.1517

3. 28週目 検査値比較 境界型

		遅延型	即効型
28W BMI	N	17.00	23.00
	Mean	24.818	22.557
	SD	3.803	3.295
	MEDIAN	24.400	22.300
	MIN	19.40	14.80
	MAX	34.70	27.70
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.0516
P-VALUE Wilcoxon	.	0.0977	
28W 筋力 (全身)	N	17.00	23.00
	Mean	4.000	4.217
	SD	2.236	2.173
	MEDIAN	4.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.7590
P-VALUE Wilcoxon	.	0.7503	
28W 筋力 (腕)	N	17.00	23.00
	Mean	4.294	4.957
	SD	2.494	2.325
	MEDIAN	4.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
NMISS	0	0	

(Continued)

3. 28週目 検査値比較 境界型

		遅延型	即効型
28W 筋力(腕)	P-VALUE STUDENT T-TEST	.	0.3932
	P-VALUE Wilcoxon	.	0.3918
28W 筋力(脚)	N	17.00	23.00
	Mean	3.529	4.087
	SD	2.239	2.295
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.4476
	P-VALUE Wilcoxon	.	0.3985
28W 握力・右	N	17.00	23.00
	Mean	36.882	36.804
	SD	10.018	8.754
	MEDIAN	37.500	36.500
	MIN	21.00	14.00
	MAX	59.00	52.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9792
	P-VALUE Wilcoxon	.	0.8695
28W 握力・左	N	17.00	23.00
	Mean	36.618	36.326
	SD	10.531	8.128
	MEDIAN	34.500	37.000
	MIN	18.00	21.00

(Continued)

3. 28週目 検査値比較 境界型

		遅延型	即効型
28W 握力・左	MAX	59.00	50.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9217
	P-VALUE Wilcoxon	.	0.8588
28W HbA1c	N	17.00	23.00
	Mean	6.624	6.583
	SD	1.333	0.917
	MEDIAN	6.500	6.600
	MIN	5.00	5.30
	MAX	10.60	8.70
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9089
	P-VALUE Wilcoxon	.	0.7735
28W PSA	N	15.00	22.00
	Mean	1.015	1.386
	SD	0.810	1.178
	MEDIAN	0.788	1.039
	MIN	0.30	0.27
	MAX	2.91	5.05
	NMISS	2	1
	P-VALUE STUDENT T-TEST	.	0.2967
	P-VALUE Wilcoxon	.	0.3534

3. 検査値比較 差の変化量 28W比較-開始時 完全型

		遅延型	即効型
28W-開始時 BMI	N	41.00	40.00
	Mean	-0.200	0.207
	SD	1.174	0.677
	MEDIAN	-0.100	0.100
	MIN	-5.50	-1.60
	MAX	1.60	1.70
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.0601
	P-VALUE Wilcoxon	.	0.0677
28W-開始時 筋力 (全身)	N	40.00	40.00
	Mean	-0.075	0.200
	SD	0.888	1.159
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	3.00	3.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.2373
	P-VALUE Wilcoxon	.	0.1455
28W-開始時 筋力 (腕)	N	40.00	40.00
	Mean	0.200	0.650
	SD	1.977	1.875
	MEDIAN	0.000	0.000
	MIN	-2.00	-2.00
	MAX	7.00	7.00
	NMISS	3	3

(Continued)

3. 検査値比較 差の変化量 28W比較-開始時 完全型

		遅延型	即効型
28W-開始時 筋力 (腕)	P-VALUE STUDENT T-TEST	.	0.2994
	P-VALUE Wilcoxon	.	0.0915
28W-開始時 筋力 (脚)	N	40.00	40.00
	Mean	0.125	0.375
	SD	0.791	0.807
	MEDIAN	0.000	0.000
	MIN	-1.00	-1.00
	MAX	3.00	3.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.1655
	P-VALUE Wilcoxon	.	0.0837
28W-開始時 握力・右	N	40.00	39.00
	Mean	-0.163	0.905
	SD	3.245	7.081
	MEDIAN	0.250	2.000
	MIN	-7.50	-31.00
	MAX	7.00	13.50
	NMISS	3	4
	P-VALUE STUDENT T-TEST	.	0.3897
P-VALUE Wilcoxon	.	0.2000	
28W-開始時 握力・左	N	40.00	40.00
	Mean	0.375	1.660
	SD	4.739	3.533
	MEDIAN	-0.250	1.250
	MIN	-6.00	-5.50

(Continued)

3. 検査値比較 差の変化量 28W比較-開始時 完全型

		遅延型	即効型
28W-開始時 握力・左	MAX	24.00	9.00
	NMISS	3	3
	P-VALUE STUDENT T-TEST	.	0.1731
	P-VALUE Wilcoxon	.	0.0270
28W-開始時 HbA1c	N	32.00	32.00
	Mean	0.034	-0.053
	SD	0.451	0.525
	MEDIAN	0.000	0.000
	MIN	-1.10	-1.60
	MAX	1.40	0.90
	NMISS	11	11
	P-VALUE STUDENT T-TEST	.	0.4769
	P-VALUE Wilcoxon	.	0.9193
28W-開始時 PSA	N	35.00	40.00
	Mean	0.286	0.341
	SD	0.952	0.471
	MEDIAN	0.058	0.199
	MIN	-0.39	-0.32
	MAX	5.46	1.74
	NMISS	8	3
	P-VALUE STUDENT T-TEST	.	0.7482
	P-VALUE Wilcoxon	.	0.0179

3. 検査値比較 差の変化量 28W比較-開始時 境界型

		遅延型	即効型
28W-開始時 BMI	N	17.00	23.00
	Mean	0.259	-0.209
	SD	0.937	1.621
	MEDIAN	0.400	0.000
	MIN	-2.40	-7.00
	MAX	1.40	2.50
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.2944
28W-開始時 筋力(全身)	N	17.00	23.00
	Mean	-0.176	-0.304
	SD	1.944	1.363
	MEDIAN	0.000	0.000
	MIN	-4.00	-4.00
	MAX	5.00	3.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.8079
28W-開始時 筋力(腕)	N	17.00	23.00
	Mean	-0.235	-0.348
	SD	1.678	2.058
	MEDIAN	0.000	0.000
	MIN	-4.00	-6.00
	MAX	2.00	3.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.8716

(Continued)

3. 検査値比較 差の変化量 28W比較-開始時 境界型

		遅延型	即効型
28W-開始時 筋力(腕)	P-VALUE STUDENT T-TEST	.	0.8547
	P-VALUE Wilcoxon	.	0.7881
28W-開始時 筋力(脚)	N	17.00	23.00
	Mean	-0.512	0.043
	SD	1.779	1.331
	MEDIAN	0.000	0.000
	MIN	-4.00	-4.00
	MAX	4.00	2.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.2653
	P-VALUE Wilcoxon	.	0.1066
28W-開始時 握力・右	N	17.00	23.00
	Mean	1.335	1.830
	SD	2.403	3.891
	MEDIAN	1.000	1.000
	MIN	-2.00	-4.40
	MAX	7.00	11.50
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.6462
	P-VALUE Wilcoxon	.	0.8692
28W-開始時 握力・左	N	17.00	23.00
	Mean	1.353	0.696
	SD	2.227	2.733
	MEDIAN	1.500	2.000
	MIN	-2.50	-5.00

(Continued)

3. 検査値比較 差の変化量 28W比較-開始時 境界型

		遅延型	即効型
28W-開始時 握力・左	MAX	5.00	5.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.4222
	P-VALUE Wilcoxon	.	0.5536
28W-開始時 HbA1c	N	13.00	19.00
	Mean	0.100	-0.247
	SD	0.574	0.560
	MEDIAN	0.100	-0.100
	MIN	-1.30	-1.70
	MAX	1.10	0.80
	NMISS	4	4
	P-VALUE STUDENT T-TEST	.	0.0985
	P-VALUE Wilcoxon	.	0.0355
28W-開始時 PSA	N	15.00	22.00
	Mean	0.138	0.449
	SD	0.416	0.969
	MEDIAN	0.048	0.154
	MIN	-0.74	-0.99
	MAX	1.15	3.76
	NMISS	2	1
	P-VALUE STUDENT T-TEST	.	0.2497
	P-VALUE Wilcoxon	.	0.2654

3. 52週目 検査値比較 完全型

		遅延型	即効型
52W PSA	N	37.00	37.00
	Mean	1.044	1.117
	SD	0.726	0.686
	MEDIAN	0.793	1.045
	MIN	0.26	0.11
	MAX	3.30	3.00
	NMISS	6	6
	P-VALUE STUDENT T-TEST	.	0.6568
	P-VALUE Wilcoxon	.	0.4051
52W HbA1c	N	40.00	39.00
	Mean	5.990	6.154
	SD	0.787	0.922
	MEDIAN	5.850	6.200
	MIN	5.00	5.00
	MAX	7.80	9.50
	NMISS	3	4
	P-VALUE STUDENT T-TEST	.	0.3977
	P-VALUE Wilcoxon	.	0.4734
52W BMI	N	42.00	42.00
	Mean	22.979	23.390
	SD	3.034	3.630
	MEDIAN	23.050	23.000
	MIN	16.60	16.00
	MAX	30.60	35.70
	NMISS	1	1

(Continued)

3. 52週目 検査値比較 完全型

		遅延型	即効型
52W BMI	P-VALUE STUDENT T-TEST	.	0.5742
	P-VALUE Wilcoxon	.	0.9038
52W 筋力(全身)	N	42.00	42.00
	Mean	3.476	4.310
	SD	2.133	2.247
	MEDIAN	3.000	4.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0850
	P-VALUE Wilcoxon	.	0.0862
52W 筋力(腕)	N	42.00	42.00
	Mean	3.476	4.286
	SD	1.928	2.040
	MEDIAN	3.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0652
	P-VALUE Wilcoxon	.	0.0550
52W 筋力(脚)	N	42.00	42.00
	Mean	3.119	4.000
	SD	2.144	2.327
	MEDIAN	2.500	4.000
	MIN	1.00	1.00

(Continued)

3. 52週目 検査値比較 完全型

		遅延型	即効型
52W 筋力 (脚)	MAX	8.00	8.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0748
	P-VALUE Wilcoxon	.	0.0840
52W 握力 右	N	42.00	41.00
	Mean	32.917	35.659
	SD	6.806	6.901
	MEDIAN	32.500	35.000
	MIN	20.00	23.00
	MAX	49.00	52.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.0721
	P-VALUE Wilcoxon	.	0.0892
52W 握力 左	N	42.00	42.00
	Mean	30.774	34.952
	SD	6.296	7.606
	MEDIAN	30.000	37.000
	MIN	19.00	10.50
	MAX	43.00	54.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.0075
	P-VALUE Wilcoxon	.	0.0067

(Continued)

3. 52週目 検査値比較 完全型

		遅延型	即効型
52W 骨塩定量	N	34.00	37.00
	Mean	102.74	102.65
	SD	17.053	17.182
	MEDIAN	105.50	98.000
	MIN	69.00	74.00
	MAX	131.00	143.00
	NMISS	9	6
	P-VALUE STUDENT T-TEST	.	0.9831
	P-VALUE Wilcoxon	.	0.8674

3. 52週目 検査値比較 境界型

		遅延型	即効型
52W PSA	N	14.00	22.00
	Mean	1.103	1.056
	SD	0.734	0.617
	MEDIAN	0.932	1.090
	MIN	0.32	0.29
	MAX	2.70	2.30
	NMISS	3	1
	P-VALUE STUDENT T-TEST	.	0.8378
	P-VALUE Wilcoxon	.	0.9483
52W HbA1c	N	15.00	22.00
	Mean	6.607	6.545
	SD	1.204	0.878
	MEDIAN	6.300	6.700
	MIN	5.10	5.20
	MAX	8.90	8.90
	NMISS	2	1
	P-VALUE STUDENT T-TEST	.	0.8589
	P-VALUE Wilcoxon	.	0.9753
52W BMI	N	17.00	23.00
	Mean	24.671	22.700
	SD	3.956	2.860
	MEDIAN	24.400	22.800
	MIN	18.80	17.80
	MAX	34.60	27.00
	NMISS	0	0

(Continued)

3. 52週目 検査値比較 境界型

		遅延型	即効型
52W BMI	P-VALUE STUDENT T-TEST	.	0.0750
	P-VALUE Wilcoxon	.	0.1587
52W 筋力(全身)	N	17.00	23.00
	Mean	4.353	4.565
	SD	2.029	2.150
	MEDIAN	4.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.7537
52W 筋力(腕)	N	17.00	23.00
	Mean	4.529	4.739
	SD	2.239	2.261
	MEDIAN	4.000	5.000
	MIN	1.00	1.00
	MAX	8.00	8.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.7725
52W 筋力(脚)	P-VALUE Wilcoxon	.	0.7507
	N	17.00	23.00
52W 筋力(脚)	Mean	4.059	4.043
	SD	2.076	2.099
	MEDIAN	4.000	4.000
	MIN	1.00	1.00

(Continued)

3. 52週目 検査値比較 境界型

		遅延型	即効型
52W 筋力 (脚)	MAX	8.00	7.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9818
	P-VALUE Wilcoxon	.	1.0000
52W 握力 右	N	17.00	23.00
	Mean	36.824	37.130
	SD	10.776	8.382
	MEDIAN	39.000	39.500
	MIN	19.50	16.50
	MAX	58.50	49.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9198
	P-VALUE Wilcoxon	.	0.6027
52W 握力 左	N	17.00	23.00
	Mean	36.412	35.304
	SD	9.497	9.648
	MEDIAN	37.000	37.000
	MIN	19.50	10.00
	MAX	55.00	51.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.7199
	P-VALUE Wilcoxon	.	0.8373

(Continued)

3. 52週目 検査値比較 境界型

		遅延型	即効型
52W 骨塩定量	N	16.00	22.00
	Mean	101.94	96.409
	SD	25.347	15.610
	MEDIAN	101.50	100.50
	MIN	60.00	67.00
	MAX	147.00	122.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.4114
	P-VALUE Wilcoxon	.	0.5642

3. 検査値比較 変化量比較 (52W-開始時) 完全型

		遅延型	即効型
52W-開始時 BMI	N	42.00	41.00
	Mean	-0.255	-0.015
	SD	1.164	0.930
	MEDIAN	-0.050	-0.100
	MIN	-4.50	-1.60
	MAX	1.90	2.30
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.3029
	P-VALUE Wilcoxon	.	0.8768
52W-開始時 筋力 (全身)	N	42.00	41.00
	Mean	0.000	0.390
	SD	1.230	1.358
	MEDIAN	0.000	0.000
	MIN	-3.00	-3.00
	MAX	4.00	4.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.1735
	P-VALUE Wilcoxon	.	0.1542
52W-開始時 筋力 (腕)	N	42.00	41.00
	Mean	-0.405	0.171
	SD	1.547	1.815
	MEDIAN	0.000	0.000
	MIN	-4.00	-5.00
	MAX	3.00	4.00
	NMISS	1	2

(Continued)

3. 検査値比較 変化量比較 (52W-開始時) 完全型

		遅延型	即効型
52W-開始時 筋力 (腕)	P-VALUE STUDENT T-TEST	.	0.1236
	P-VALUE Wilcoxon	.	0.1674
52W-開始時 筋力 (脚)	N	42.00	41.00
	Mean	0.190	0.415
	SD	1.042	1.303
	MEDIAN	0.000	0.000
	MIN	-2.00	-3.00
	MAX	3.00	4.00
	NMISS	1	2
	P-VALUE STUDENT T-TEST	.	0.3887
	P-VALUE Wilcoxon	.	0.2558
52W-開始時 握力 右	N	41.00	40.00
	Mean	0.207	1.320
	SD	3.100	4.533
	MEDIAN	0.500	1.000
	MIN	-9.00	-12.00
	MAX	6.50	11.00
	NMISS	2	3
	P-VALUE STUDENT T-TEST	.	0.2001
52W-開始時 握力 左	P-VALUE Wilcoxon	.	0.2180
	N	41.00	41.00
52W-開始時 握力 左	Mean	0.463	1.851
	SD	4.969	3.272
	MEDIAN	-0.500	1.500
	MIN	-8.00	-5.00

(Continued)

3. 検査値比較 変化量比較 (52W-開始時) 完全型

		遅延型	即効型
52W-開始時 握力 左	MAX	27.00	9.50
	NMISS	2	2
	P-VALUE STUDENT T-TEST	.	0.1392
	P-VALUE Wilcoxon	.	0.0064
52W-開始時 HbA1c	N	32.00	33.00
	Mean	-0.053	-0.121
	SD	0.559	0.524
	MEDIAN	0.000	0.000
	MIN	-1.60	-2.10
	MAX	1.00	0.80
	NMISS	11	10
	P-VALUE STUDENT T-TEST	.	0.6141
	P-VALUE Wilcoxon	.	0.4885
52W-開始時 PSA	N	37.00	37.00
	Mean	0.132	0.298
	SD	0.306	0.377
	MEDIAN	0.076	0.210
	MIN	-0.36	-0.11
	MAX	1.33	1.32
	NMISS	6	6
	P-VALUE STUDENT T-TEST	.	0.0411
	P-VALUE Wilcoxon	.	0.0290

(Continued)

3. 検査値比較 変化量比較 (52W-開始時) 完全型

		遅延型	即効型
52W-開始時 骨塩定量	N	34.00	34.00
	Mean	5.445	3.200
	SD	9.207	4.304
	MEDIAN	2.300	3.000
	MIN	-4.00	-8.00
	MAX	45.94	12.00
	NMISS	9	9
	P-VALUE STUDENT T-TEST	.	0.2022
	P-VALUE Wilcoxon	.	0.8441

3. 検査値比較 変化量比較 52W-開始時 境界型

		遅延型	即効型
52W-開始時 BMI	N	17.00	23.00
	Mean	0.112	-0.065
	SD	0.892	0.771
	MEDIAN	0.100	-0.300
	MIN	-1.40	-1.20
	MAX	1.60	2.20
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.5059
	P-VALUE Wilcoxon	.	0.4031
52W-開始時 筋力 (全身)	N	17.00	23.00
	Mean	0.176	0.043
	SD	1.811	1.261
	MEDIAN	0.000	0.000
	MIN	-3.00	-4.00
	MAX	5.00	2.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.7855
	P-VALUE Wilcoxon	.	0.5033
52W-開始時 筋力 (腕)	N	17.00	23.00
	Mean	0.000	-0.565
	SD	1.414	1.701
	MEDIAN	0.000	0.000
	MIN	-3.00	-5.00
	MAX	3.00	2.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.2723
	P-VALUE Wilcoxon	.	0.3459
52W-開始時 筋力 (脚)	N	17.00	23.00
	Mean	0.018	0.000
	SD	1.460	0.953
	MEDIAN	0.000	0.000
	MIN	-3.00	-3.00
	MAX	4.00	1.00
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.9634
	P-VALUE Wilcoxon	.	0.5080

(Continued)

3. 検査値比較 変化量比較 52W-開始時 境界型

		遅延型	即効型
52W-開始時 握力 右	N	17.00	23.00
	Mean	1.276	2.157
	SD	2.971	4.748
	MEDIAN	1.000	1.000
	MIN	-3.50	-6.00
	MAX	8.00	16.50
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.5057
	P-VALUE Wilcoxon	.	0.6117
52W-開始時 握力 左	N	17.00	23.00
	Mean	1.147	-0.326
	SD	2.822	6.886
	MEDIAN	1.500	2.000
	MIN	-5.50	-30.00
	MAX	6.00	3.50
	NMISS	0	0
	P-VALUE STUDENT T-TEST	.	0.4118
	P-VALUE Wilcoxon	.	0.9343
52W-開始時 HbA1c	N	12.00	19.00
	Mean	0.217	-0.205
	SD	0.406	0.635
	MEDIAN	0.200	-0.200
	MIN	-0.40	-1.50
	MAX	1.00	0.90
	NMISS	5	4
	P-VALUE STUDENT T-TEST	.	0.0499
	P-VALUE Wilcoxon	.	0.0533
52W-開始時 PSA	N	14.00	22.00
	Mean	0.233	0.143
	SD	0.569	0.419
	MEDIAN	0.076	0.066
	MIN	-0.77	-0.96
	MAX	1.80	1.33
	NMISS	3	1
	P-VALUE STUDENT T-TEST	.	0.5876
	P-VALUE Wilcoxon	.	0.8839

(Continued)

3. 検査値比較 変化量比較 52W-開始時 境界型

		遅延型	即効型
52W-開始時 骨塩定量	N	16.00	22.00
	Mean	2.556	3.773
	SD	6.249	5.255
	MEDIAN	2.000	2.000
	MIN	-9.00	-3.00
	MAX	15.00	18.00
	NMISS	1	1
	P-VALUE STUDENT T-TEST	.	0.5194
	P-VALUE Wilcoxon	.	0.7101