

For the symptom analysis, changes of LCS from initial score were compared between arms using analysis of covariance with initial score as a covariate.

All analyses were carried out with SAS software release 8.2 (SAS Institute, Cary, NC).

## results

This trial was terminated early due to the unexpected high incidence of interstitial lung disease (ILD) and three treatment-related deaths due to ILD in the combination arm, which were identified by the Adverse Event Reporting system.

### patient characteristics

From January 2002 to September 2003, 130 patients with NSCLC who had failed prior platinum-based chemotherapy from 32 institutions were enrolled (Appendix). These patients were randomly assigned to docetaxel alone ( $n = 65$ ) or docetaxel plus gemcitabine ( $n = 65$ ). One patient died as a result of rapid progressive disease before chemotherapy administration, and one patient did not meet the entry criteria in the docetaxel arm. In addition, one patient did not meet the entry criteria in the docetaxel plus gemcitabine arm. All patients were included in the analysis of survival and PFS, and 64 docetaxel and 65 docetaxel plus gemcitabine patients were assessable for toxicity. Fifty-nine patients with measurable lesions by RECIST

in the docetaxel arm and 57 eligible patients in docetaxel plus gemcitabine arm were assessable for response (Figure 1). Table 1 presents baseline patient characteristics.

The median number of cycles was 3 (range 0–6) and 2 (range 1–8) in the docetaxel and docetaxel plus gemcitabine arms, respectively. The median interval between cycles was 22 days for both arms.

### toxicity

This trial was terminated early due to the unexpected high incidence of ILD and three treatment-related deaths (4.6%) due to ILD in the docetaxel plus gemcitabine arm. These events were identified by the Adverse Event Reporting system. Thirteen (20.0%) patients receiving combination treatment suffered from all grades of ILD, whereas only two (3.1%) patients receiving docetaxel alone suffered from grades 1–2 ILD. Grades 2–4 ILD occurred in 16.9% of docetaxel plus gemcitabine patients, an unexpected high incidence rate. No risk factors were identified contributing to these pulmonary adverse events.

Toxicity was assessed in all patients who received at least one treatment cycle and in all cycles (Table 2). Overall, grades 3–4 neutropenia occurred in 55 docetaxel patients (85.9%) and 53 docetaxel plus gemcitabine patients (81.5%). Grades 3–4 anemia occurred in two patients (3.1%) and 12 patients (18.5%) treated with docetaxel alone and docetaxel plus




Figure 1. CONSORT diagram for the study.

gemcitabine, respectively. Sixteen patients treated with docetaxel (25.0%) and 11 patients with docetaxel plus gemcitabine (16.9%) developed febrile neutropenia. All

Table 1. Patient characteristics

	D arm		DG arm	
	No. of patients	%	No. of patients	%
Patients enrolled	65		65	
Age, years				
Median	62		60	
Range	34–75		34–74	
Gender				
Male	48	73.8	51	78.5
Female	17	26.2	14	21.5
ECOG PS				
0	20	30.8	21	32.3
1	45	69.2	44	67.7
Histology				
Squamous	19	29.2	22	33.8
Adenocarcinoma	40	61.5	40	61.5
Large cell	4	6.2	3	4.6
Others	2	3.1	0	0
Best response of prior chemotherapy				
CR	2	3.1	0	0
PR	38	58.5	40	61.5
SD	20	30.8	19	29.2
PD	5	7.7	6	9.2

D, docetaxel; DG, docetaxel plus gemcitabine; ECOG PS, Eastern Cooperative Oncology Group performance status; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease.

Table 2. Hematological and non-hematological toxicity

	D arm (n = 64)					DG arm (n = 65)				
	NCI-CTC grade					NCI-CTC grade				
Hematological	0–1	2	3	4	3–4%	0–1	2	3	4	3–4%
Anemia	27	35	2	0	3.1	21	32	9	3	18.5
Leukopenia	9	14	29	12	64.1	11	12	32	10	64.6
Neutropenia	7	2	15	40	85.9	8	4	19	34	81.5
Thrombocytopenia	64	0	0	0	0	43	14	8	0	12.3
Non-hematological	0–1	2	3	4	2–4%	0–1	2	3	4	2–4%
Allergic reaction	64	0	0	0	0	59	5	1	0	9.2
Alopecia	45	18	–	–	28.1	49	14	–	–	21.5
ALT	61	2	1	0	4.7	52	10	3	0	20.0
Diarrhea	61	3	0	0	4.7	60	3	2	0	7.7
Edema	63	1	0	0	1.6	64	1	0	0	1.5
Fatigue	56	5	2	1	12.5	56	7	1	1	13.8
Febrile neutropenia	48	–	16	0	25.0	54	–	11	0	16.9
Infection with grades 3–4 neutropenia	59	–	5	0	7.8	56	–	9	0	13.8
Infection without neutropenia	54	8	2	0	15.6	51	4	9	1	21.5
Nausea	55	7	2	–	14.1	55	6	4	–	15.4
Neuropathy	62	2	0	0	3.1	62	2	0	1	4.6
Pneumonitis (ILD)	63	1	0	0	1.6	54	3	7	1	16.9
Stomatitis	61	3	0	0	4.7	60	5	0	0	7.7

D, docetaxel; DG, docetaxel plus gemcitabine; NCI-CTC, National Cancer Institute—Cancer Common Toxicity Criteria; ALT, alanine aminotransferase; ILD, interstitial lung disease.

required antibiotic treatment and G-CSF; however, no patient died. One patient in the docetaxel plus gemcitabine arm developed anaphylactic shock immediately after administration of docetaxel at the second cycle. Grades 2–4 ALT elevation was more frequent with docetaxel plus gemcitabine than with docetaxel (20.0% versus 4.7%). Grades 2–4 non-neutropenic infection occurred more often with docetaxel plus gemcitabine than with docetaxel (21.5% versus 15.6%). Grades 2–4 ILD was more frequent with docetaxel plus gemcitabine than with docetaxel (16.9% versus 1.6%). Other toxic effects were relatively mild (Table 2). Overall, docetaxel plus gemcitabine was more toxic than docetaxel, however, well tolerated except for ILD in docetaxel plus gemcitabine arm.

### treatment efficacy

The overall response rate for docetaxel alone was 6.8% [95% confidence interval (CI) 1.9% to 16.5%] and 7.0% for docetaxel plus gemcitabine (95% CI 2.0% to 17.0%). There was no significant difference between treatment arms ( $P = 0.71$ ; Fisher's exact test).

At the time of this analysis, 50 docetaxel patients (76.9%) and 48 docetaxel plus gemcitabine patients (73.8%) had died. The median survival time was 10.1 months for docetaxel alone and 10.3 months for docetaxel plus gemcitabine (one-sided  $P = 0.36$  stratified log-rank test; Figure 2A). The respective 1-year survival rate was 43.1% (95% CI 31.0% to 55.1%) for docetaxel and 46.0% (95% CI 33.8% to 58.1%) for docetaxel plus gemcitabine.

The median PFS time was 2.1 and 2.8 months for docetaxel and docetaxel plus gemcitabine, respectively (one-sided  $P = 0.028$  stratified log-rank test; Figure 2B).




Figure 2. Overall survival (A) and progression-free survival (B) by treatment arm.

### disease-related symptom assessment

Patients' compliance with disease-related symptom assessment was 100% at baseline and 95.4% at 6 weeks later. Compliance rates were not different between the arms ( $P = 1.00$ ). LCS data were missing in four surveys due to death or severe impairment of the patient's general condition; this accounted for 1.5% of the total number of surveys scheduled. Mean LCS at baseline and 6 weeks were shown in Table 3. There were no significant differences in the LCS changes from baseline to 6 weeks between docetaxel and docetaxel plus gemcitabine arms ( $P = 0.61$ ).

### discussion

This trial was terminated early due to the unexpected high incidence of ILD and three treatment-related deaths due to ILD in the docetaxel plus gemcitabine arm. Our findings seem to indicate that the combination of docetaxel and gemcitabine may be associated with a higher incidence of pulmonary adverse events compared with docetaxel alone, especially in patients with previously treated NSCLC.

Pulmonary toxicity following chemotherapeutic agents, including ILD, has been well recognized for many years. In most cases, this toxicity is mild and self-limiting. However, the mechanism of developing drug-induced ILD is uncertain, and risk factors for developing this disorder have not been identified. In terms of combination therapy with docetaxel and gemcitabine for advanced NSCLC, there were few reports about the incidences of ILD at the time this study was planned. A phase I study of patients with transitional cell carcinoma evaluated thrice-weekly doses of docetaxel given on day 1 plus gemcitabine given on days 1 and 15 and showed that pulmonary toxicity occurred in three of five patients and was

Table 3. Disease-related symptom assessment

Lung Cancer Subscale	D arm	DG arm
Baseline		
Number	$n = 65$	$n = 65$
Mean $\pm$ SD	$19.0 \pm 5.48$	$19.7 \pm 5.25$
6 weeks later		
Number	$n = 62$	$n = 62$
Mean $\pm$ SD	$18.1 \pm 5.56$	$18.9 \pm 5.05$
Difference		
Mean $\pm$ SD	$-1.11 \pm 3.81$	$-0.99 \pm 4.49$

D, docetaxel; DG, docetaxel plus gemcitabine; SD, standard deviation.

the cause of death in one [25]. Recently, some reports have been published about the high incidence of ILD due to the combination regimen of docetaxel and gemcitabine in patients with NSCLC [13, 26, 27], including the present study (Table 4). In Japanese population, ILD is a very complex issue in treatment of patients with lung cancer. Epidermal growth factor tyrosine kinase inhibitor gefitinib is developing ILD significantly in Japanese patients with NSCLC [28]. It is uncertain why ILD is developing more in Japanese patients with NSCLC than the Western patients. Ethnic difference may be one of the explanations for this occurrence. The combination of gemcitabine and docetaxel is associated with a high incidence of severe pulmonary toxicity. The regimen should not be used outside a clinical trial.

The median survival times of 10.1 and 10.3 months and estimated 1-year survival rates of 43.1% and 46.0% with docetaxel alone and docetaxel plus gemcitabine, respectively, suggest that adding gemcitabine to docetaxel did not provide any increased efficacy in patients with previously treated NSCLC. Interestingly, the combination regimen of docetaxel plus gemcitabine significantly improved the median PFS time ( $P = 0.028$ ). Possible reasons for failing to detect a significant difference between survival curves may include an insufficient occurrence of documented events as a result of the study population comprising patients with relatively good prognosis, in addition to a high proportion of patients subsequently receiving third-line therapy. During this study, gefitinib treatment was commonly used for patients with recurrent NSCLC in Japan [29]. Asian ethnicity is a well-known predictive factor for a response for gefitinib [30].

Two randomized phase II trials compared docetaxel alone with docetaxel plus irinotecan in second-line chemotherapy for NSCLC [31, 32]. No significant treatment differences in survival were observed in either trial; however, the trials were phase II study and were not powered or designed to compare survival. This study was not powered to compare survival when it was terminated early due to the unexpected high incidence of ILD in the docetaxel plus gemcitabine arm. However, based on previous studies, as well as the present results, combination chemotherapy with docetaxel and another chemotherapeutic agent has not improved survival in patients with previously treated NSCLC.

In conclusion, docetaxel alone is still the standard second-line treatment for advanced NSCLC. The combination of docetaxel and gemcitabine was too toxic to obtain any survival

**Table 4.** Reports of interstitial lung disease due to docetaxel plus gemcitabine regimen

Author	Year	Study type	Treatment schedule	n	Grades 3-4 ILD (%)	TRD (%)
Rebattu et al. [13]	2001	Phase I/II	Docetaxel (60, 75, 85, 100 mg/m <sup>2</sup> ) day 8; gemcitabine (1000 mg/m <sup>2</sup> ), days 1 and 8, every 3 weeks	49	3 (6.1)	0
Kouroussis et al. [25]	2004	Phase I	Docetaxel (30, 35, 40 mg/m <sup>2</sup> ), days 1, 8 and 15; gemcitabine (700, 800, 900, 1000 mg/m <sup>2</sup> ), days 1, 8 and 15, every 4 weeks	26	6 (23)	2 (7.7)
Matsui et al. [21]	2005	Phase I/II	Docetaxel (50, 60 mg/m <sup>2</sup> ) day 1 or 8; gemcitabine (800, 1000 mg/m <sup>2</sup> ), days 1 and 8, every 3 weeks	59	3 (5.1)	0
Pujor et al. [27]	2005	Phase III	Docetaxel (85 mg/m <sup>2</sup> ) day 8; gemcitabine (1000 mg/m <sup>2</sup> ), days 1 and 8, every 3 weeks	155	8 (5.2)	1 (0.6)
Takeda (present study)	2008	Phase III	Cisplatin (100 mg/m <sup>2</sup> ) day 1; vinorelbine (30 mg/m <sup>2</sup> ), days 1, 8, 15 and 22, every 4 weeks	156	1 (0.6)	0
			Docetaxel (60 mg/m <sup>2</sup> ) day 8; gemcitabine (800 mg/m <sup>2</sup> ), days 1 and 8, every 3 weeks	65	8 (12.3)	3 (4.6)
			Docetaxel (60 mg/m <sup>2</sup> ) day 1, every 3 weeks	64	0 (0)	0

ILD, interstitial lung disease; TRD, treatment-related death.

benefit in patients with recurrent advanced NSCLC. The development of less toxic and more effective chemotherapeutic agents, including molecular targeted drugs, is warranted for the second-line treatment of NSCLC.

## funding

Grant-in-Aid for Cancer Research from the Ministry of Health, Labour and Welfare of Japan.

## acknowledgements

We thank Ms Mieko Imai for data management, Mr Takashi Asakawa, Dr Naoki Ishizuka for statistical analyses in the interim monitoring, and Dr Haruhiko Fukuda for valuable contributions to this study. This study is registered with UMIN-CTR [<http://www.umin.ac.jp/ctr/index.htm> umin.ac.jp/ctr], identification number C000000027].

## appendix

The following institutions participated in the study: Hokkaido Cancer Center (Sapporo), Ibaragi Prefectural Central Hospital (Kasama), Tochigi Cancer Center (Utsunomiya), Nishiginma National Hospital (Shibukawa), Gunma Prefectural Cancer Center Hospital (Ohta), Saitama Cancer Center Hospital (Ina), National Cancer Center Hospital East (Kashiwa), National Cancer Center Hospital (Tokyo), International Medical Center of Japan (Tokyo), Cancer Institute Hospital (Tokyo), Toranomon Hospital (Tokyo), Kanagawa Cancer Center Hospital (Yokohama), Yokohama Municipal Hospital (Yokohama), Niigata Cancer Center Niigata Hospital (Niigata), Gifu Municipal Hospital (Gifu), Aichi Cancer Center Hospital (Nagoya), Nagoya National Hospital (Nagoya), Prefectural Aichi Hospital (Okazaki), Osaka City University Medical School (Osaka), Kinki University School of Medicine (Osaka-Sayama), Osaka Medical Center for Cancer and Cardiovascular Disease (Osaka), Osaka Prefectural Medical Center for

Respiratory and Allergic disease (Habikino), Kinki-Chuo Chest Medical Center (Sakai), Toneyama National Hospital (Toyonaka), Osaka Prefectural General Hospital (Osaka), Osaka City General Hospital (Osaka), Kobe City General Hospital (Kobe), Hyogo Collage of Medicine (Nishinomiya), Hyogo Cancer Center (Akashi), Shikoku Cancer Center Hospital (Matsuyama), Kyusyu University Hospital (Fukuoka), and Kumamoto Regional Medical Center (Kumamoto).

## references

- Parkin DM. Global cancer statistics in the year 2000. *Lancet Oncol* 2001; 2: 533-543.
- Dancey J, Shepherd FA, Gralla RJ et al. Quality of life assessment of second-line docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy: results of a prospective, randomized phase III trial. *Lung Cancer* 2004; 43: 183-194.
- Socinski MA, Morris DE, Masters GA et al. Chemotherapeutic management of stage IV non-small cell lung cancer. *Chest* 2003; 123 (Suppl 1): 226S-243S.
- Shepherd FA, Dancey J, Ramlau R et al. Prospective randomized trial of docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy. *J Clin Oncol* 2000; 18: 2095-2103.
- Fossella FV, DeVore R, Kerr RN et al. Randomized phase III trial of docetaxel versus vinorelbine or ifosfamide in patients with advanced non-small-cell lung cancer previously treated with platinum-containing chemotherapy regimens: the TAX 320 Non-Small Cell Lung Cancer Study Group. *J Clin Oncol* 2000; 18: 2354-2362.
- Crino L, Mosconi AM, Scagliotti G et al. Gemcitabine as second-line treatment for advanced non-small-cell lung cancer: a phase II trial. *J Clin Oncol* 1999; 17: 2081-2085.
- Georgoulas V, Kouroussis C, Androulakis N et al. Front-line treatment of advanced non-small-cell lung cancer with docetaxel and gemcitabine: a multicenter phase II trial. *J Clin Oncol* 1999; 17: 914-920.
- Georgoulas V, Papadakis E, Alexopoulos A et al. Platinum-based and non-platinum-based chemotherapy in advanced non-small-cell lung cancer: a randomised multicentre trial. *Lancet* 2001; 357: 1478-1484.
- Hainsworth JD, Burris HA III, Billings FT III et al. Weekly docetaxel with either gemcitabine or vinorelbine as second-line treatment in patients with advanced non-small cell lung carcinoma: phase II trials of the Minnie Pearl Cancer Research Network. *Cancer* 2001; 92: 2391-2398.

10. Hejna M, Kornek GV, Raderer M et al. Treatment of patients with advanced nonsmall cell lung carcinoma using docetaxel and gemcitabine plus granulocyte-colony stimulating factor. *Cancer* 2000; 89: 516–522.
11. Kakolyris S, Papadakis E, Tsiafaki X et al. Docetaxel in combination with gemcitabine plus rhG-CSF support as second-line treatment in non-small cell lung cancer. A multicenter phase II study. *Lung Cancer* 2001; 32: 179–187.
12. Kosmas C, Tsavaris N, Vadiaka M et al. Gemcitabine and docetaxel as second-line chemotherapy for patients with non-small cell lung carcinoma who fail prior paclitaxel plus platinum-based regimens. *Cancer* 2001; 92: 2902–2910.
13. Rebattu P, Quantin X, Ardiet C et al. Dose-finding, pharmacokinetic and phase II study of docetaxel in combination with gemcitabine in patients with inoperable non-small cell lung cancer. *Lung Cancer* 2001; 33: 277–287.
14. Rischin D, Boyer M, Smith J et al. A phase I trial of docetaxel and gemcitabine in patients with advanced cancer. *Ann Oncol* 2000; 11: 421–426.
15. Spiridonidis CH, Laufman LR, Jones J et al. Phase I study of docetaxel dose escalation in combination with fixed weekly gemcitabine in patients with advanced malignancies. *J Clin Oncol* 1998; 16: 3866–3873.
16. Spiridonidis CH, Laufman LR, Carman L et al. Second-line chemotherapy for non-small-cell lung cancer with monthly docetaxel and weekly gemcitabine: a phase II trial. *Ann Oncol* 2001; 12: 89–94.
17. Kunitoh H, Watanabe K, Onoshi T et al. Phase II trial of docetaxel in previously untreated advanced non-small-cell lung cancer: a Japanese Cooperative Study. *J Clin Oncol* 1996; 14: 1649–1655.
18. Taguchi T, Furue H, Niitani H et al. Phase I clinical trial of RP 56976 (docetaxel) a new anticancer drug. *Gan To Kagaku Ryoho* 1994; 21: 1997–2005.
19. Miyazaki M, Takeda K, Ichimaru Y et al. A phase I/II study of docetaxel (D) and gemcitabine (G) combination chemotherapy for advanced non-small cell lung cancer (NSCLC). *J Clin Oncol* 2001; 20 (Suppl): (Abstr 2812).
20. Niho S, Kubota K, Goto K et al. Combination second-line chemotherapy with gemcitabine and docetaxel for recurrent non-small-cell lung cancer after platinum containing chemotherapy: a phase I/II trial. *Cancer Chemother Pharmacol* 2003; 52: 19–24.
21. Matsui K, Hirashima T, Nitta T et al. A phase I/II study comparing regimen schedules of gemcitabine and docetaxel in Japanese patients with stage IIIB/IV non-small cell lung cancer. *Jpn J Clin Oncol* 2005; 35: 181–187.
22. Arbusk SG, Ivy SP, Setser A et al. The Revised Common Toxicity Criteria: Version 2.0 CTEP. <http://ctep.info.nih.gov> (30 April 1999, date last accessed).
23. Therasse P, Arbusk SG, Eisenhauer EA et al. New guidelines to evaluate the response to treatment in solid tumors. *J Natl Cancer Inst* 2000; 92: 205–216.
24. Cella DF, Bonomi AE, Lloyd SR et al. Reliability and validity of the Functional Assessment of Cancer Therapy-Lung (FACT-L) quality of life instrument. *Lung Cancer* 1995; 12: 199–220.
25. Dunsford LM, Mead MG, Bateman CA et al. Severe pulmonary toxicity in patients treated with a combination of docetaxel and gemcitabine for metastatic transitional cell carcinoma. *Ann Oncol* 1999; 10: 943–947.
26. Kouroussis C, Tsavaris N, Syrigos K et al. High incidence of pulmonary toxicity of weekly docetaxel and gemcitabine in patients with non-small cell lung cancer: results of a dose-finding study. *Lung Cancer* 2004; 44: 363–368.
27. Pujor JL, Breton JL, Gervais R et al. Gemcitabine-docetaxel versus cisplatin-vinorelbine in advanced or metastatic non-small-cell lung cancer: a phase III study addressing the case for cisplatin. *Ann Oncol* 2005; 16: 602–610.
28. Ando M, Okamoto I, Yamamoto N et al. Predictive factors for interstitial lung disease, antitumor response, and survival in non-small-cell lung cancer patients treated with gefitinib. *J Clin Oncol* 2006; 24: 2549–2556.
29. Fukuoka M, Yano S, Giaccone G et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2003; 21: 2237–2246.
30. Thatcher N, Chang A, Parikh P et al. Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell lung cancer: results from a randomized, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer). *Lancet* 2005; 366: 1527–1537.
31. Pectasides D, Pectasides M, Farmakis D et al. Comparison of docetaxel and docetaxel-irinotecan combination as second-line chemotherapy in advanced non-small cell lung cancer: a randomized phase II trial. *Ann Oncol* 2005; 16: 294–299.
32. Wouters FM, Groen HJ, Riesma H et al. Phase II randomized trial of docetaxel vs docetaxel and irinotecan in patients with stage IIIB-IV non-small-cell lung cancer who failed first-line treatment. *Br J Cancer* 2005; 92: 15–20.

Original Article

## A Phase I Study of Gemcitabine and Carboplatin in Patients with Advanced Non-small Cell Lung Cancer and a Performance Status of 2

Young Hak Kim<sup>1,2</sup>, Kaoru Kubota<sup>1</sup>, Koichi Goto<sup>1</sup>, Kiyotaka Yoh<sup>1</sup>, Seiji Niho<sup>1</sup>, Hironobu Ohmatsu<sup>1</sup>, Nagahiro Saijo<sup>1</sup> and Yutaka Nishiwaki<sup>1</sup>

<sup>1</sup>Division of Thoracic Oncology, National Cancer Center Hospital East, Chiba and <sup>2</sup>Department of Respiratory Medicine, Kyoto University Hospital, Kyoto, Japan

Received April 16, 2009; accepted May 7, 2009

**Objective:** The aim of this study was to determine the maximum-tolerated dose (MTD) and the recommended dose of combination chemotherapy with gemcitabine (GEM) and carboplatin (CBDCA) in non-small cell lung cancer (NSCLC) patients with a performance status (PS) of 2.

**Methods:** Chemotherapy-naïve NSCLC patients with PS 2 were enrolled. Chemotherapy consisted of an escalated dose of GEM on days 1 and 8 and CBDCA on day 1 every 3 weeks. Patients were scheduled to receive GEM (mg/m<sup>2</sup>)/CBDCA (area under the curve: AUC) at four dose levels: 800/4 (level 1), 1000/4 (level 2), 1000/4.5 (level 3) and 1000/5 (level 4), respectively.

**Results:** Between February 2004 and August 2006, 13 patients were enrolled in this study. Dose-limiting toxicities (DLTs) were thrombocytopenia, febrile neutropenia and hyponatremia. DLTs were observed in two of six patients at dose level 1 and in three of six patients at dose level 2. Dose level 2 was thus determined to be the MTD. Among 12 evaluable patients, 7 patients had stable diseases and 5 patients had progressive diseases, and the median survival time was 3.8 months.

**Conclusions:** The MTD and the recommended dose for Phase II studies of this regimen were determined to be GEM 1000 mg/m<sup>2</sup> and CBDCA AUC of 4. Additional objective measures are needed to evaluate patients' risk and benefit in future clinical trials for PS 2 patients.

*Key words:* non-small cell lung cancer – performance status 2 – gemcitabine – carboplatin – Phase I

### INTRODUCTION

Platinum-based combination chemotherapy has been shown to improve survival and quality-of-life (QOL) in patients with advanced non-small cell lung cancer (NSCLC) (1,2). In the 1990s, new chemotherapeutic agents, such as gemcitabine (GEM), vinorelbine, docetaxel, paclitaxel (PTX) and irinotecan, were developed. Currently, platinum-based

chemotherapy employing these new agents is accepted as the standard chemotherapy worldwide (3,4). In addition, a meta-analysis demonstrated significant longer progression-free survival of GEM and platinum combination compared with other new agents and platinum combinations (5). Thus, combination chemotherapy with GEM and platinum is now considered as one of the most active regimens for advanced NSCLC.

Like in other types of cancers, performance status (PS) has been shown to be one of the most important prognostic factors for survival in advanced NSCLC (6–8). Patients with

For reprints and all correspondence: Young Hak Kim, Department of Respiratory Medicine, Kyoto University Hospital, 54 Shogoin-Kawaharacho, Sakyo-ku, Kyoto 606-8507, Japan. E-mail: ekim@kubp.kyoto-u.ac.jp

impaired PS generally have lower response rate and shorter survival in spite of high risk for severe toxicities (9,10). Historically, clinical trials have excluded patients with Eastern Cooperative Oncology Group (ECOG) PS of 2 or worse. To date, it has not been fully elucidated whether platinum-based combination chemotherapy is feasible and effective in patients with PS 2.

Carboplatin (CBDCA), an analog of cisplatin (CDDP), has lower nephro- and gastrointestinal toxicity and has been widely used as a substitution of CDDP. Several randomized trials have shown the equivalence between GEM + CBDCA (GC) and GEM + CDDP (GP) in terms of response rate and survival (11,12). In those trials, toxicities, such as emesis, nephropathy and neuropathy were significantly mild in GC. Although recent meta-analysis disclosed slightly but significant survival advantage of CDDP (13,14), GC can be one of the treatment options, especially for patients who are not suitable to receive CDDP. In a randomized Phase III trial comparing GC with vinblastine + CDDP, GC showed better response rate and survival, and toxicities were similar between the two arms (15). Although 70% of all enrolled patients in the study had PS 2, overall response rate and median survival time (MST) were 27% and 11.6 months in GC arm. These survival data were comparable to those in patients with PS 0 or 1 who treated with platinum-based chemotherapy.

These results suggest the potential benefit of GC in patients with PS 2; however, the optimal dose of GC has not been investigated in patients with impaired PS. Therefore, we conducted a Phase I study to determine the maximum-tolerated dose (MTD) and the recommended dose for Phase II studies of GC in advanced NSCLC patients with PS 2.

## PATIENTS AND METHODS

### ELIGIBILITY

Patients with histologically or cytologically proven advanced NSCLC were eligible for the study. Each patient was required to meet the following criteria: (i) clinical stage IIIB or IV; (ii) ECOG PS of 2; (iii) aged 20–75 years; (iv) measurable lesion; (v) no prior chemotherapy; (vi) adequate hematological function (white blood cell  $\geq 3500/\text{mm}^3$ , hemoglobin  $\geq 9.5$  g/dl and platelets  $\geq 100\,000/\text{mm}^3$ ); (vii) adequate hepatic and renal function (total bilirubin  $\leq 1.5$  mg/dl, AST and ALT  $< 100$  IU/l and creatinine  $\leq 1.5$  mg/dl); (viii)  $\text{PaO}_2 \geq 60$  mmHg; and (ix) written informed consent. Patients with active concomitant malignancy, radiologically apparent interstitial pneumonia or pulmonary fibrosis, serious concurrent illness (e.g. uncontrolled diabetes mellitus, hypertension, angina pectoris, myocardial infarction within 3 months after onset or severe infection), history of severe drug allergy or pregnant/lactating women were excluded. The study protocol was approved by the institutional review board of the National Cancer Center.

### TREATMENT SCHEDULE

This was a Phase I, dose-escalation study planned for GEM on days 1 and 8 and CBDCA on day 1 of a 21-day course. The initial dose level of GEM was  $800\text{ mg/m}^2$  and CBDCA was an area under the concentration–time curve (AUC) of  $4\text{ mg min/ml}$ . The actual dose of CBDCA was calculated based on Cockcroft–Gault equation (16) and Calvert formula (17) every course. CBDCA was infused over 60 min, and 60 min after the completion of CBDCA infusion, GEM was administered over 30 min. Prophylactic administration of granulocyte colony-stimulating factor (G-CSF) was not permitted. Administration of G-CSF was permitted for patients with grade 4 neutropenia and/or leukopenia and grade 3 febrile neutropenia. The administration of GEM was omitted on day 8 if patients met one of the following criteria: white blood cell  $< 2000/\text{mm}^3$ , neutrophil  $< 1000/\text{mm}^3$ , platelets  $< 50\,000/\text{mm}^3$  and PS  $\geq 3$ . No dose modification of GEM was permitted on day 8. If dose-limiting toxicity (DLT) was observed, the dose of each drug was reduced to 80% in the next course of chemotherapy. Treatment was to be performed for at least two courses, unless unacceptable toxicity or disease progression occurred.

The DLT was defined as follows: grade 4 thrombocytopenia, grade 3 or grade 4 febrile neutropenia, grade 3 non-hematological toxicity (except for nausea/vomiting and alopecia) and omission of the treatment on day 8. Dose-escalation schedule is shown in Table 1. Initially, three patients were treated at each dose level. If DLT was not observed in any of three patients, dose escalation was made. If DLT was observed in one or two of three patients, an additional three patients were entered in the same dose level. If DLT was observed in three or more of six patients or all of the initial three patients, we considered that the dose was the MTD. If DLT was observed in one or two of six patients, dose escalation was also made. Dose escalation was decided by the toxic data only in the first course of chemotherapy.

### BASELINE AND TREATMENT ASSESSMENT

Pre-treatment evaluation consisted of complete medical history and physical examination, complete blood cell counts, blood chemistry studies, electrocardiograph, arterial blood gas analysis, chest radiography, computed tomography (CT) of the chest, CT or ultrasound study of the abdomen,

Table 1. Dose-escalation schedule

Dose level	Gemcitabine ( $\text{mg/m}^2$ )	Carboplatin (AUC)	No. of patients
1	800	4	3–6
2	1000	4	3–6
3	1000	4.5	3–6
4	1000	5	3–6

AUC, area under the curve.

CT or magnetic resonance imaging of the brain, and bone scintigraphy. Complete blood cell counts, blood chemistry studies and chest radiography were repeated every week. Creatinine clearance was estimated by the Cockcroft–Gault equation every course. Tumor response was assessed with the Response Evaluation Criteria in Solid Tumor (RECIST) criteria (18). Toxicity was evaluated according to the National Cancer Institute-Common Toxicity Criteria (version 2.0).

**RESULTS**

**PATIENT CHARACTERISTICS**

Between February 2004 and August 2006, 13 patients were enrolled in this study. However, one patient was excluded from the analysis because of the error in dose calculation. Table 2 shows the characteristics of 12 evaluable patients. Eleven patients were male and one was female. The median age of the patients was 68 years (range, 51–72 years). There were five adenocarcinomas, four squamous cell carcinomas, two large cell carcinomas and one pleomorphic carcinoma. Stage IIIB and IV patients were five and six, respectively, and one patient was a relapse after surgical resection.

**DOSE ESCALATION**

At the dose level 1, DLT was observed in two of the first three patients; one experienced grade 3 hyponatremia and the other experienced grade 3 febrile neutropenia. Thereafter, we amended the protocol, and grade 3 hyponatremia was excluded from DLT criteria after that. Another three patients were treated at the same dose. Since these patients did not

**Table 2.** Characteristics of evaluable patients (n = 12)

Characteristics	No. of patients
Gender	
Male	11
Female	1
Age (years)	
Median	68
Range	51–72
Histology	
Adenocarcinoma	5
Squamous cell carcinoma	4
Large cell carcinoma	2
Pleomorphic carcinoma	1
Stage	
IIIB	5
IV	6
Relapse after surgery	1

show any additional DLT, the dosage was then escalated to the next step. At the dose level 2, DLT was observed in two of the first three patients: one experienced grade 3 nausea/vomiting and omission on day 8 and the other experienced grade 3 febrile neutropenia and anorexia. Therefore, another three patients were assigned to receive the treatment at the same dose. Out of those three patients, one patient developed grade 4 febrile neutropenia and grade 3 anorexia. Thus, DLT was observed in three of six patients at the dose level 2. As a result, the dose level 2 (GEM, 1000 mg/m<sup>2</sup> and CBDCA, AUC of 4) was determined to be the MTD.

**TOXICITY**

The worst grades for each patient in the first cycle are listed in Table 3. Grade 3/4 leukopenia or neutropenia was observed in one patient at level 1 and two patients at level 2. Febrile neutropenia was observed in one patient at level 1 and two patients at level 2. Two patients had grade 3/4 anemia at level 1 and one patient required red blood cell

**Table 3.** Toxicities during the first cycle

NCI-CTC grade	Level 1 (n = 6)		Level 2 (n = 6)	
	G1/2	G3/4	G1/2	G3/4
<b>Hematologic</b>				
Leukopenia	1/2	0/1	2/1	2/0
Neutropenia	1/1	1/0	1/1	2/0
Febrile neutropenia	0/0	1/0	0/0	1/1
Anemia	1/3	1/1	2/3	0/0
Thrombocytopenia	1/2	2/1	1/1	4/0
Transaminase	2/0	0/0	4/2	0/0
Bilirubin	0/0	0/0	0/0	1/0
Creatinine	0/0	0/0	0/0	0/0
Hyponatremia	4/0	2/0	5/0	0/0
<b>Non-hematologic</b>				
Nausea/vomiting	2/0	0/0	3/1	1/0
Anorexia	4/1	0/0	2/1	3/0
Fatigue	1/0	0/0	1/2	1/0
Diarrhea	0/0	0/0	2/0	0/0
Constipation	0/0	0/0	0/1	0/0
Mucositis	0/0	0/0	0/0	0/0
Pneumonitis	0/0	0/0	0/0	0/0
Infection	0/0	0/0	0/0	0/0
Skin rash	1/0	0/0	1/0	0/0
Omission on day 8	0		1	
No. of patients with DLT	2		3	

NCI-CTC, National Cancer Institute-Common Toxicity Criteria; DLT, dose-limiting toxicity.



transfusion. No grade 3/4 anemia occurred at level 2. Thrombocytopenia was the principal toxicity of this combination chemotherapy. At level 1, grade 3/4 thrombocytopenias were observed in three patients, and two patients received platelet transfusion. At level 2, two patients experienced grade 3/4 thrombocytopenia requiring no platelet transfusions. Non-hematologic toxicities were generally mild at level 1, however, one patient experienced grade 3 nausea/vomiting and omission of day 8 at level 2. This patient also presented grade 3 hyperbilirubinemia suspected to be drug-induced hepatitis, and died 16 days after the start of the treatment. The worst value of his laboratory data was 6.6 mg/dl in total bilirubin on day 12, 40 IU/l in AST on day 7 and 103 IU/l in ALT on day 7. He had a past history of drug-induced hepatitis related to aspirin. The excluded patient was administered GEM at 800 mg/body. Despite the dose was approximately two-thirds of the planned dose, he experienced grade 3 nausea/vomiting and the treatment was discontinued. The median number of administered cycle was 1. The actual administered cycles were one in seven patients, two in one patient, three in two patients and four in two patients. The reasons for the discontinuation in seven patients who terminated the treatment at one cycle were toxicity for three patients, patient refusal for two patients, treatment delay for one patient and both toxicity and disease progression was for one patient.

ANTI-TUMOR ACTIVITY

There were seven stable diseases and five progressive diseases (PD). No partial or complete response was observed (Table 4). Four patients received second-line chemotherapy after GC: docetaxel for two patients and gefitinib for two patients. One patient received gefitinib experienced partial response; however, remaining three patients had PD also in the second-line treatment. The MST was 3.8 months (Fig. 1).

DISCUSSION

This is the first PS 2-specific Phase I study of GC in Japanese patients, and the MTD and recommended dose were determined to be GEM 1000 mg/m<sup>2</sup> and CBDCA AUC of 4.

Table 4. Drug delivery and anti-tumor efficacy

Dose level	Number of patients	Median course (range)	Overall response			
			CR	PR	SD	PD
1	6	1 (1-4)	0	0	4	2
2	6	2 (1-4)	0	0	3	3

CR, complete response; PR, partial response; SD, stable diseases; PD, progressive diseases.




Figure 1. Kaplan-Meier curve of overall survival. Median overall survival time was 3.8 months.

The recommended dose of CBDCA was lower than other studies conducted in the USA (19,20). With respect to the dose of CBDCA, the method of measuring serum creatinine values is critical. In Japan, most institutions use the enzymatic method, whereas the Jaffe method remains the mainstream in the USA (21). According to the study comparing these two methods, serum creatinine values are higher in the Jaffe method than in the enzymatic method by ~0.2 mg/dl (21). Therefore, at the same AUC, higher CBDCA dose is administered in Japan than in the USA. Incidentally, based on the Calvert formula, a difference of 0.2 mg/dl of creatinine leads to the difference of AUC = 1. In short, the AUC = 4 in Japan roughly corresponds to the AUC = 5 in the USA. For global clinical trials, the difference of methods for measurement of laboratory data also should be paid attention to.

PS is one of the most powerful and reliable prognostic factors in advanced NSCLC (6-8), and a worse PS is characterized by lower response rate to chemotherapy and shorter survival (9,10). Median survival of patients with PS 2 is substantially shorter than that of patients with PS 0 or 1. Moreover, patients with PS 2 are at higher risk for severe toxicity than those with better PS. According to the population-based surveys, up to 30-40% of all advanced NSCLC is characterized PS 2 (22,23). Namely, patients with PS 2 constitute a distinctive, non-trivial subgroup in NSCLC. However, little attention has been paid to this special patient population until recently.

The guidelines from the American Society of Clinical Oncology support the single use of third-generation non-platinum agents for patients with PS 2 (24). This recommendation is mainly based on the results of Phase III trials comparing single-agent chemotherapy with best supportive care alone, in which good tolerability and significant survival benefit or improvement of QOL with single-agent chemotherapy have been demonstrated (25-28). However, PS 2 patients accounted for a small proportion of patients in those trials and any conclusive evidence cannot be drawn for the treatment of patients with PS 2. At present, available

data from PS 2-specific clinical trials are quite limited. In this context, no consensus has been developed on the standard chemotherapy in patients with PS 2.

The role of adding platinum to third-generation single agents is still unclear. Recently, the Norwegian Lung Cancer Study Group reported the results of a retrospective study that compared the outcome of patients with PS 2 to that of patients with PS 0 or 1 who had participated in randomized trials comparing two third-generation, CBDCA-based regimens (29). According to the retrospective study, although MST of patients with PS 2 was significantly shorter than that of patients with PS 0 or 1 (4.5 vs. 8.9 months;  $P < 0.01$ ), toxicity was acceptable for patients with PS 2 and they achieved better symptom improvement compared with patients with PS 0 or 1. ECOG conducted the first PS 2-specific randomized trial (19). In the randomized Phase II trial, two platinum-based chemotherapy regimens, PTX + CBDCA (PC) and GP, have been compared, and both regimens were proved feasible with acceptable toxicity. However, survival time was quite limited in both treatment arms: MST was 6.2 months for PC and 6.9 months for GP, respectively. A Greece Group performed a randomized Phase II trial comparing non-platinum single-agent chemotherapy with CBDCA-based chemotherapy (30). In the study, patients were randomly assigned to either GC or GEM alone and MST was 6.7 months for GC and 4.8 months for GEM alone, respectively ( $P = 0.49$ ), whereas neutropenia ( $P = 0.007$ ) and thrombocytopenia ( $P < 0.001$ ) were more common in GC arm. In contrast, according to a subgroup analysis of the Cancer and Leukemia Group B study 9730 comparing PC with PTX alone, patients with PS 2 (107 patients, 18% of the population) achieved significantly better survival when they were treated with PC than those treated with PTX alone (20). Thus, the role of platinum-based chemotherapy for patients with PS 2 is still controversial.

The results could vary even between PS 2-specific trials due to two major reasons. First, determining PS score is inevitably subjective, there is considerable inter-observer variation even between healthcare professionals (31). Second, there can be significant heterogeneity in the PS 2 patient population: the reasons for impaired PS may be due to tumor-related (such as pain, fatigue and weight loss), to pre-existing co-morbidities (such as chronic obstructive pulmonary disease, cardiovascular disease and age-related decline in functional status) or both, furthermore (32). There is a clear need for a more objective classification system that takes into account the individual effects of disease-related symptoms and co-morbidities. The common co-morbidity scales are the Cumulative Illness Rating Scale-Geriatric (CIRS-G) and the Charlson scale. Their prognostic impacts have been validated prospectively (33,34). Moreover, they are more objective than PS. Although our study did not, all future studies for PS 2 patients should use such co-morbidity scales to stratify patients more accurately.

Recently, molecular-targeted agents, especially epidermal growth factor receptor tyrosine kinase inhibitors such as

gefitinib or erlotinib, have been tested in clinical trials for patients with poor PS. Inoue et al. (35) conducted a Phase II trial of gefitinib in patients with NSCLC whose tumor harboring EGFR gene mutation. In the study, all patients were not feasible for cytotoxic chemotherapy due to poor PS: 26 of 29 patients were PS 2–4. Overall response rate and MST were 66% and 6.5 months, respectively. In addition, PS improvement rate was 79%, and no treatment-related deaths were observed. These excellent results strongly suggest that stratification with molecular status should be required in the future trial of PS 2 or more.

In this study, we determined the MTD and the recommended dose of GC in Japanese patients with PS 2. Response rate and overall survival of the regimen were disappointing. However, some previous studies clearly support the use of platinum agent in PS 2 patients (19,20). Future clinical trials for PS 2 patients should use more objective criterion such as co-morbidity scales in addition to PS in order to measure patients' risk more accurately. Such studies may reveal that which patients should be treated and not be treated with platinum-based chemotherapy among PS 2 patients.

## Funding

This study was supported in part by a Grant-in Aid for Cancer Research from the Japanese Ministry of Health and Welfare.

## Conflict of interest statement

None declared.

## References

1. Souquet PJ, Chauvin F, Boissel JP, Cellerino R, Cormier Y, Ganz PA, et al. Polychemotherapy in advanced non small cell lung cancer: a meta-analysis. *Lancet* 1993;342:19–21.
2. Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *BMJ* 1995;311:899–909.
3. Schiller JH, Harrington D, Belani CP, Langer C, Sandler A, Krook J, et al. Comparison of four chemotherapy regimens for advanced non-small-cell lung cancer. *N Engl J Med* 2002;346:92–8.
4. Bunn PA, Jr. Chemotherapy for advanced non-small-cell lung cancer: who, what, when, why? *J Clin Oncol* 2002;20:23–33S.
5. Le Chevalier T, Scagliotti G, Natale R, Danson S, Rosell R, Stabel R, et al. Efficacy of gemcitabine plus platinum chemotherapy compared with other platinum containing regimens in advanced non-small-cell lung cancer: a meta-analysis of survival outcomes. *Lung Cancer* 2005;47:69–80.
6. Stanley KE. Prognostic factors for survival in patients with inoperable lung cancer. *J Natl Cancer Inst* 1980;65:25–32.
7. Albain KS, Crowley JJ, LeBlanc M, Livingston RB. Survival determinants in extensive-stage non-small-cell lung cancer: the Southwest Oncology Group experience. *J Clin Oncol* 1991;9:1618–26.
8. Brundage MD, Davies D, Mackillop WJ. Prognostic factors in non-small cell lung cancer: a decade of progress. *Chest* 2002;122:1037–57.

9. Gridelli C, Ardizzoni A, Le Chevalier T, Manegold C, Ferrone F, Thatcher N, et al. Treatment of advanced non-small-cell lung cancer patients with ECOG performance status 2: results of an European Experts Panel. *Ann Oncol* 2004;15:419–26.
10. Gridelli C, Maione P, Rossi A, Guerriero C, Ferrara C, Del Gaizo F, et al. Chemotherapy of advanced NSCLC in special patient population. *Ann Oncol* 2006;17(Suppl 5):v72–8.
11. Mazzanti P, Massacesi C, Rocchi MB, Mattioli R, Lippe P, Trivisonne R, et al. Randomized, multicenter, phase II study of gemcitabine plus cisplatin versus gemcitabine plus carboplatin in patients with advanced non-small cell lung cancer. *Lung Cancer* 2003;41:81–9.
12. Zatloukal P, Petruzelka L, Zemanova M, Kolek V, Skrickova J, Pesek M, et al. Gemcitabine plus cisplatin vs. gemcitabine plus carboplatin in stage IIb and IV non-small cell lung cancer: a phase III randomized trial. *Lung Cancer* 2003;41:321–31.
13. Hotta K, Matsuo K, Ueoka H, Kiura K, Tabata M, Tanimoto M. Meta-analysis of randomized clinical trials comparing cisplatin to carboplatin in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2004;22:3852–9.
14. Ardizzoni A, Boni L, Tiseo M, Fossella FV, Schiller JH, Paesmans M, et al. Cisplatin- versus carboplatin-based chemotherapy in first-line treatment of advanced non-small-cell lung cancer: an individual patient data meta-analysis. *J Natl Cancer Inst* 2007;99:847–57.
15. Grigorescu AC, Draghici IN, Nitipir C, Gutulescu N, Corlan E. Gemcitabine (GEM) and carboplatin (CBDCA) versus cisplatin (CDDP) and vinblastine (VLB) in advanced non-small-cell lung cancer (NSCLC) stages III and IV: a phase III randomised trial. *Lung Cancer* 2002;37:9–14.
16. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron* 1976;16:31–41.
17. Calvert AH, Newell DR, Gumbrell LA, O'Reilly S, Burnell M, Boxall FE, et al. Carboplatin dosage: prospective evaluation of a simple formula based on renal function. *J Clin Oncol* 1989;7:1748–56.
18. Therasse P, Arbusk SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000;92:205–16.
19. Langer C, Li S, Schiller J, Tester W, Rapoport BL, Johnson DH, et al. Randomized phase II trial of paclitaxel plus carboplatin or gemcitabine plus cisplatin in Eastern Cooperative Oncology Group performance status 2 non-small-cell lung cancer patients: ECOG 1599. *J Clin Oncol* 2007;25:418–23.
20. Lilenbaum RC, Herndon JE, 2nd, List MA, Desch C, Watson DM, Miller AA, et al. Single-agent versus combination chemotherapy in advanced non-small-cell lung cancer: the cancer and leukemia group B (study 9730). *J Clin Oncol* 2005;23:190–6.
21. Ando Y, Minami H, Saka H, Ando M, Sakai S, Shimokata K. Adjustment of creatinine clearance improves accuracy of Calvert's formula for carboplatin dosing. *Br J Cancer* 1997;76:1067–71.
22. Buccheri G, Ferrigno D, Tamburini M. Karnofsky and ECOG performance status scoring in lung cancer: a prospective, longitudinal study of 536 patients from a single institution. *Eur J Cancer* 1996;32A:1135–41.
23. Radzikowska E, Glaz P, Roszkowski K. Lung cancer in women: age, smoking, histology, performance status, stage, initial treatment and survival. Population-based study of 20 561 cases. *Ann Oncol* 2002;13:1087–93.
24. Pfister DG, Johnson DH, Azzoli CG, Sause W, Smith TJ, Baker S, Jr. American Society of Clinical Oncology treatment of unresectable non-small-cell lung cancer guideline: update 2003. *J Clin Oncol* 2004;22:330–53.
25. The Elderly Lung Cancer Vinorelbine Italian Study Group. Effects of vinorelbine on quality of life and survival of elderly patients with advanced non-small-cell lung cancer. *J Natl Cancer Inst* 1999;91:66–72.
26. Ranson M, Davidson N, Nicolson M, Falk S, Carmichael J, Lopez P, et al. Randomized trial of paclitaxel plus supportive care versus supportive care for patients with advanced non-small-cell lung cancer. *J Natl Cancer Inst* 2000;92:1074–80.
27. Roszkowski K, Pluzanska A, Krzakowski M, Smith AP, Saigi E, Aasebo U, et al. A multicenter, randomized, phase III study of docetaxel plus best supportive care versus best supportive care in chemotherapy-naïve patients with metastatic or non-resectable localized non-small cell lung cancer (NSCLC). *Lung Cancer* 2000;27:145–57.
28. Anderson H, Hopwood P, Stephens RJ, Thatcher N, Cottier B, Nicholson M, et al. Gemcitabine plus best supportive care (BSC) vs BSC in inoperable non-small cell lung cancer—a randomized trial with quality of life as the primary outcome. UK NSCLC Gemcitabine Group. Non-Small Cell Lung Cancer. *Br J Cancer* 2000;83:447–53.
29. Helbekkmo N, Aasebo U, Sundstrom SH, von Plessen C, Brunsvig PF, Bremnes RM; Norwegian Lung Cancer Study Group. Treatment outcome in performance status 2 advanced NSCLC patients administered platinum-based combination chemotherapy. *Lung Cancer* 2008;62:253–60.
30. Kosmidis PA, Dimopoulos MA, Syrigos K, Nicolaidis C, Aravantinos G, Boukovinas I, et al. Gemcitabine versus gemcitabine-carboplatin for patients with advanced non-small cell lung cancer and a performance status of 2: a prospective randomized phase II study of the Hellenic Cooperative Oncology Group. *J Thorac Oncol* 2007;2:135–40.
31. Taylor AE, Oliver DN, Sivanthan T, Chi M, Purnell C. Observer error in grading performance status in cancer patients. *Support Care Cancer* 1999;7:332–5.
32. Gebbia V, Galetta D, De Marinis F. Non small cell lung cancer patients with ECOG PS2: unsolved questions and lessons from clinical trials. *Ann Oncol* 2005;16(Suppl 4):iv123–31.
33. Firat S, Byhardt RW, Gore E. Comorbidity and Karnofsky performance score are independent prognostic factors in stage III non-small-cell lung cancer: an institutional analysis of patients treated on four RTOG studies. Radiation Therapy Oncology Group. *Int J Radiat Oncol Biol Phys* 2002;54:357–64.
34. Frasci G, Lorusso V, Panza N, Comella P, Nicoletta G, Bianco A, et al. Gemcitabine plus vinorelbine versus vinorelbine alone in elderly patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2000;18:2529–36.
35. Inoue A, Kobayashi K, Usui K, Maemondo M, Okinaga S, Mikami I, et al. First-line gefitinib for patients with advanced non-small-cell lung cancer harboring epidermal growth factor receptor mutations without indication for chemotherapy. *J Clin Oncol* 2009;27:1394–400.

# Gefitinib or Carboplatin–Paclitaxel in Pulmonary Adenocarcinoma

Tony S. Mok, M.D., Yi-Long Wu, M.D., F.A.C.S., Sumitra Thongprasert, M.D., Chih-Hsin Yang, M.D., Ph.D., Da-Tong Chu, M.D., Nagahiro Saijo, M.D., Ph.D., Patrapim Sunpaweravong, M.D., Baohui Han, M.D., Benjamin Margono, M.D., Ph.D., F.C.C.P., Yukito Ichinose, M.D., Yutaka Nishiwaki, M.D., Ph.D., Yuichiro Ohe, M.D., Ph.D., Jin-Ji Yang, M.D., Busyamas Chewaskulyong, M.D., Haiyi Jiang, M.D., Emma L. Duffield, M.Sc., Claire L. Watkins, M.Sc., Alison A. Armour, F.R.C.R., and Masahiro Fukuoka, M.D., Ph.D.

## ABSTRACT

### BACKGROUND

Previous, uncontrolled studies have suggested that first-line treatment with gefitinib would be efficacious in selected patients with non–small-cell lung cancer.

### METHODS

In this phase 3, open-label study, we randomly assigned previously untreated patients in East Asia who had advanced pulmonary adenocarcinoma and who were nonsmokers or former light smokers to receive gefitinib (250 mg per day) (609 patients) or carboplatin (at a dose calculated to produce an area under the curve of 5 or 6 mg per milliliter per minute) plus paclitaxel (200 mg per square meter of body-surface area) (608 patients). The primary end point was progression-free survival.

### RESULTS

The 12-month rates of progression-free survival were 24.9% with gefitinib and 6.7% with carboplatin–paclitaxel. The study met its primary objective of showing the noninferiority of gefitinib and also showed its superiority, as compared with carboplatin–paclitaxel, with respect to progression-free survival in the intention-to-treat population (hazard ratio for progression or death, 0.74; 95% confidence interval [CI], 0.65 to 0.85;  $P < 0.001$ ). In the subgroup of 261 patients who were positive for the epidermal growth factor receptor gene (*EGFR*) mutation, progression-free survival was significantly longer among those who received gefitinib than among those who received carboplatin–paclitaxel (hazard ratio for progression or death, 0.48; 95% CI, 0.36 to 0.64;  $P < 0.001$ ), whereas in the subgroup of 176 patients who were negative for the mutation, progression-free survival was significantly longer among those who received carboplatin–paclitaxel (hazard ratio for progression or death with gefitinib, 2.85; 95% CI, 2.05 to 3.98;  $P < 0.001$ ). The most common adverse events were rash or acne (in 66.2% of patients) and diarrhea (46.6%) in the gefitinib group and neurotoxic effects (69.9%), neutropenia (67.1%), and alopecia (58.4%) in the carboplatin–paclitaxel group.

### CONCLUSIONS

Gefitinib is superior to carboplatin–paclitaxel as an initial treatment for pulmonary adenocarcinoma among nonsmokers or former light smokers in East Asia. The presence in the tumor of a mutation of the *EGFR* gene is a strong predictor of a better outcome with gefitinib. (ClinicalTrials.gov number, NCT00322452.)

From the State Key Laboratory in Oncology in South China, Sir YK Pao Centre for Cancer, Department of Clinical Oncology, Chinese University of Hong Kong, Hong Kong (T.S.M.), Guangdong General Hospital, Guangzhou (Y.L.W., J.-J.Y.), Cancer Hospital, Chinese Academy of Medical Sciences, Beijing (D.-T.C.), and Shanghai Chest Hospital, Shanghai (B.H.) — all in China; Maharaj Nakorn Chiang Mai Hospital, Chiang Mai University, Chiang Mai (S.T., B.C.), and Prince of Songkla University, Songkla (P.S.) — both in Thailand; National Taiwan University Hospital, Taipei, Taiwan (C.-H.Y.); National Cancer Center Hospital East, Chiba (N.S., Y.N.), National Kyushu Cancer Center, Fukuoka (Y.I.), National Cancer Center Hospital, Tokyo (Y.O.), AstraZeneca, Osaka (H.J.), and Kinki University School of Medicine, Osaka (M.F.) — all in Japan; Dr. Soetomo Hospital, Surabaya, Indonesia (B.M.); and AstraZeneca, Macclesfield, United Kingdom (E.L.D., C.L.W., A.A.A.). Address reprint requests to Dr. Mok at the Department of Clinical Oncology, Chinese University of Hong Kong, Prince of Wales Hospital, Shatin, HKSAR, China, or at [tony@clo.cuhk.edu.hk](mailto:tony@clo.cuhk.edu.hk).

This article (10.1056/NEJMoa0810699) was published on August 19, 2009, at NEJM.org.

N Engl J Med 2009;361:947-57.

Copyright © 2009 Massachusetts Medical Society.

Reprinted from THE NEW ENGLAND JOURNAL OF MEDICINE (ISSN 0028-4793) Vol. 361:947-957 (September 3, 2009). Copyright © 2009 Massachusetts Medical Society. All rights reserved. Printed in the U.S.A. Fax: (781) 893-8103 [www.nejm.org](http://www.nejm.org)

**I**NHIBITORS OF THE EPIDERMAL GROWTH factor receptor (EGFR) tyrosine kinase have shown clinical efficacy, as compared with the best supportive care<sup>1</sup> or standard chemotherapy,<sup>2</sup> when given as second-line or third-line therapy for advanced non-small-cell lung cancer. Treatment with EGFR tyrosine kinase inhibitors is most effective in women, patients who have never smoked, patients with pulmonary adenocarcinomas, and patients of Asian origin. In these populations, such treatment is associated with favorable rates of objective responses, progression-free survival, and overall survival.<sup>1,3,4</sup> These populations also have a relatively high incidence of somatic mutations in the region of the EGFR gene that encodes the tyrosine kinase domain.<sup>5,6</sup> Studies have shown that in patients with pulmonary adenocarcinoma who had a base-pair deletion at exon 19 (del746\_A750) or a point mutation at exon 21 (L858R), the tumors were highly responsive to EGFR tyrosine kinase inhibitors,<sup>7-9</sup> and subsequent studies of first-line therapy with these agents showed objective response rates of 54.8 to 81.6% and progression-free survival of 9.7 to 13.3 months among patients with these mutations.<sup>10-12</sup>

On the basis of these and other studies,<sup>1,4,13-16</sup> we hypothesized that in a selected population, first-line therapy with an oral EGFR tyrosine kinase inhibitor would be at least as effective as chemotherapy with carboplatin-paclitaxel. In this study, we compared the efficacy, safety, and adverse-event profile of gefitinib with those of carboplatin-paclitaxel when these drugs were used as first-line treatment in nonsmokers or former light smokers in East Asia who had adenocarcinoma of the lung. We also examined the role of an EGFR mutation as a predictor of the efficacy of gefitinib or carboplatin-paclitaxel.

## METHODS

### STUDY DESIGN AND PATIENTS

The First Line Iressa versus Carboplatin/Paclitaxel in Asia (Iressa Pan-Asia Study [IPASS]) study was a phase 3, multicenter, randomized, open-label, parallel-group study comparing gefitinib (Iressa, AstraZeneca) with carboplatin (Paraplatin, Bristol-Myers Squibb) plus paclitaxel (Taxol, Bristol-Myers Squibb) as first-line treatment in clinically selected patients in East Asia who had advanced non-small-cell lung cancer. The primary end point was pro-

gression-free survival. Secondary end points included overall survival (an early analysis, since follow-up is ongoing), the objective response rate, quality of life, reduction in symptoms, safety, and the adverse-event profile. Evaluations of efficacy according to the baseline biomarker status of EGFR were planned exploratory objectives.

Patients were eligible for inclusion in the study if they were 18 years of age or older, had histologically or cytologically confirmed stage IIIB or IV non-small-cell lung cancer with histologic features of adenocarcinoma (including bronchoalveolar carcinoma), were nonsmokers (defined as patients who had smoked <100 cigarettes in their lifetime) or former light smokers (those who had stopped smoking at least 15 years previously and had a total of ≤10 pack-years of smoking), and had had no previous chemotherapy or biologic or immunologic therapy. Other eligibility criteria are described in the Supplementary Appendix, available with the full text of this article at NEJM.org.

The principal investigators and the members of the steering committee (see the Appendix at the end of this article) designed the study in collaboration with the sponsor (AstraZeneca) and supervised the conduct of the trial. The sponsor collected and analyzed the data. The lead academic author had unrestricted access to the data and vouches for the validity and completeness of the results of the trial (see the Supplementary Appendix for further details). All patients provided written informed consent; separate consent was provided for the assessment of EGFR biomarkers. An independent ethics committee at each participating institution approved the study protocol. The study was conducted in accordance with the Declaration of Helsinki, the International Conference on Harmonization Guidelines for Good Clinical Practice, applicable regulatory requirements, and AstraZeneca's policy on bioethics. One planned interim analysis was performed by an independent statistician and reviewed by an independent data and safety monitoring committee (see the Supplementary Appendix).

### STUDY TREATMENT

Patients were randomly assigned, in a 1:1 ratio, to receive gefitinib (250 mg per day, administered orally) or paclitaxel (200 mg per square meter of body-surface area, administered intravenously over a 3-hour period on the first day of the cycle) fol-

lowed immediately by carboplatin (at a dose calculated to produce an area under the concentration-time curve of 5.0 or 6.0 mg per milliliter per minute, administered intravenously over a period of 15 to 60 minutes) in cycles of once every 3 weeks for up to 6 cycles. Randomization was performed with the use of dynamic balancing<sup>17</sup> with respect to performance status, as assessed by the World Health Organization (WHO) performance scale measuring activity (0 or 1, or 2 on a scale of 0 to 4, with lower numbers indicating a higher degree of activity); smoking status (nonsmoker or former light smoker); sex; and center. Treatment continued until progression of the disease, development of unacceptable toxic effects, a request by the patient or physician to discontinue treatment, serious non-compliance with the protocol, or completion of six chemotherapy cycles. Among patients assigned to gefitinib therapy, those whose tumor progressed were offered the opportunity to switch to treatment with carboplatin-paclitaxel; however, if the patient declined or was not a good candidate for that treatment, he or she could receive another approved therapy of the physician's choice. Among patients who were receiving carboplatin-paclitaxel, further therapy after progression of the disease was at the physician's discretion.

#### ASSESSMENTS

Progression-free survival was assessed from the date of randomization to the earliest sign of disease progression, as determined by means of the Response Evaluation Criteria in Solid Tumors (RECIST),<sup>18</sup> or death from any cause. Overall survival was assessed from the date of randomization until death from any cause. Tumor response was assessed every 6 weeks until disease progression. Quality of life was assessed with the use of the Functional Assessment of Cancer Therapy-Lung (FACT-L) questionnaire (in which scores range from 0 to 136, with higher scores indicating better quality of life) and the Trial Outcome Index (TOI, which is the sum of the physical well-being, functional well-being, and lung-cancer subscale [LCS] scores of FACT-L; scores range from 0 to 84, with higher scores indicating better quality of life), and symptoms were assessed with the use of the LCS score (scores range from 0 to 28, with higher scores indicating fewer symptoms). The FACT-L questionnaire<sup>19</sup> was administered at randomization and at week 1, once every 3 weeks

until day 127, once every 6 weeks from day 128 until disease progression, and when the study drug was discontinued. Clinically relevant improvement was predefined as an improvement of six points or more in FACT-L and TOI scores or an improvement of two points or more in LCS scores, with the higher scores maintained for at least 21 days.<sup>20</sup> Safety and tolerability were assessed according to National Cancer Institute Common Terminology Criteria for Adverse Events, version 3.0. Tumor samples from patients who consented to have biomarkers assessed were analyzed at two central laboratories to determine biomarker status, with EGFR mutation status the first priority. Patients were considered to be positive for the EGFR mutation if 1 of 29 EGFR mutations was detected with the use of the amplification refractory mutation system (ARMS) and the DxS EGFR29 mutation-detection kit.<sup>21,22</sup>

#### STATISTICAL ANALYSIS

The primary end point (progression-free survival) was analyzed with the use of a Cox proportional-hazards model in the intention-to-treat population (all randomly assigned patients) to assess the non-inferiority of gefitinib as compared with carboplatin-paclitaxel, with the WHO performance status (0 or 1, or 2), smoking status (nonsmoker or former light smoker), and sex as covariates. For noninferiority to be demonstrated, the 95% confidence interval for the hazard ratio had to lie entirely below the predefined noninferiority limit of 1.2. We estimated that with a total of 944 progression events, the study would have 80% power to demonstrate noninferiority if the treatments were truly equal, with a two-sided 5% probability of an erroneous demonstration of noninferiority. If the 95% confidence interval for the hazard ratio was also below 1, the P value would be less than 0.05 and superiority could be concluded from the same analysis without statistical penalty (closed test procedure).<sup>23</sup> Supportive secondary analyses are described in the Supplementary Appendix. Planned subgroup analyses were performed to compare progression-free survival between treatments in groups defined according to WHO performance status (0 or 1, or 2), smoking status (nonsmoker or former light smoker), sex, age at randomization (<65 years or ≥65 years), disease stage at screening (stage IIIB or IV), and presence or absence of biomarkers. Tests to determine in-

teractions of treatment with covariates were used to identify predictive factors by assessing whether there was a significant difference in the treatment effect for progression-free survival (hazard ratio for progression or death) between subgroups.

Overall survival was analyzed with the use of methods that were similar to those used for the analysis of progression-free survival. The results of an early analysis are presented; follow-up with respect to overall survival is ongoing. The objective response rate (in the intention-to-treat population) and quality of life and rates of symptom reduction (among all patients with a baseline and at least one post-baseline quality-of-life assessment that could be evaluated) were assessed with the use of a logistic-regression model with the same covariates as those considered for progression-free survival to calculate odds ratios and 95% confidence intervals. Planned subgroup analyses of the objective response rate were performed with the use of methods that were similar to those used for the analysis of progression-free survival.

Adverse events were summarized for all patients who received at least one dose of the assigned study treatment. The incidence rates of 10 specified safety events (5 that were possibly associated with each study treatment) were compared with the use of Fisher's exact test; adjustment for multiple comparisons was performed with the use of the method of Westfall and Young.<sup>24</sup>

## RESULTS

### PATIENTS AND TREATMENT


From March 2006 through October 2007, a total of 1217 patients from 87 centers in Hong Kong, elsewhere in China, Indonesia, Japan, Malaysia, the Philippines, Singapore, Taiwan, and Thailand were randomly assigned to a study group (Fig. 1). The two groups were well balanced with respect to demographic and baseline characteristics (Table 1). The mean duration of treatment was 6.4 months (median, 5.6; range, 0.1 to 22.8) for gefitinib and 3.4 months (median, 4.1; range, 0.7 to 5.8) for carboplatin-paclitaxel. The median number of treatment cycles in the carboplatin-paclitaxel group was six. At the cutoff date for collection of data (April 14, 2008), a total of 24.5% of the patients in the gefitinib group were continuing to receive the study treatment; all patients in the carboplatin-paclitaxel group had discontinued the drugs. After discontinuation of the assigned treatment at

any time during the study, 38.9% of the patients in the gefitinib group received carboplatin-paclitaxel, and 39.5% of the patients in the carboplatin-paclitaxel group received an EGFR tyrosine kinase inhibitor; 10.5% of the patients in the gefitinib group and 14.0% of those in the carboplatin-paclitaxel group received other anticancer treatments.

### EFFICACY

The median follow-up period for the analysis of progression-free survival was 5.6 months. The median progression-free survival was 5.7 months in the gefitinib group and 5.8 months in the carboplatin-paclitaxel group, approximately coinciding with crossing of the Kaplan-Meier curves. The 12-month rates of progression-free survival were 24.9% with gefitinib and 6.7% with carboplatin-paclitaxel; a total of 950 patients had progression of disease. The study met its primary objective of demonstrating noninferiority and showed the superiority of gefitinib as compared with carboplatin-paclitaxel for progression-free survival (hazard ratio for progression or death, 0.74; 95% confidence interval [CI], 0.65 to 0.85;  $P < 0.001$ ). The probability that a patient would be free of disease progression was greater with carboplatin-paclitaxel in the first 6 months and greater with gefitinib in the following 16 months (Fig. 2A). Progression-free survival was longer in the gefitinib group than in the carboplatin-paclitaxel group in all clinical subgroups; the only clinical factor that affected progression-free survival was age (<65 years: hazard ratio, 0.81; 95% CI, 0.70 to 0.95;  $P = 0.007$ ;  $\geq 65$  years: hazard ratio, 0.58; 95% CI, 0.45 to 0.76;  $P < 0.001$ ;  $P = 0.03$  for the interaction of treatment with age) (Fig. 1 in the Supplementary Appendix).

A total of 1038 patients (85.3%) gave their consent for biomarker analyses, and 683 patients (56.1%) provided samples. EGFR mutation data for 437 patients (35.9%) could be evaluated. Patients with a tissue sample that could be evaluated had demographic characteristics that were similar to those of the overall population (Table 1 in the Supplementary Appendix). Of the 437 samples, 261 (59.7%) were positive for a mutation. Of these 261 samples, 140 (53.6%) had exon 19 deletions, 111 (42.5%) had a mutation at exon 21 (L858R), 11 (4.2%) had a mutation at exon 20 (T790M), and 10 (3.8%) had other mutations; 11 patients had multiple mutations. The proportions of mutations



**Figure 1. Screening, Group Assignment, and Inclusion in Analyses.**

All patients who were randomly assigned to a study group were included in the intention-to-treat analysis; all patients with a baseline and at least one post-baseline quality-of-life assessment that could be evaluated were included in the quality-of-life analysis; patients who did not deviate substantially from the inclusion and exclusion criteria at entry or from the protocol were included in the per-protocol analysis; and all patients who received at least one dose of study treatment were included in the safety analysis. Among the 112 patients who were assessed for eligibility but were not assigned to a study group, the main reasons for exclusion were a serum creatinine level that was higher than 1.5 times the upper limit of the reference range or a creatinine clearance of 60 ml per minute or less; newly diagnosed central nervous system metastases that had not yet been definitively treated with surgery or radiation; or an absolute neutrophil count of less than  $2.0 \times 10^9$  per liter, a platelet count of less than  $100 \times 10^9$  per liter, or a hemoglobin level of less than 10 g per deciliter. A total of 63 patients who were treated with gefitinib continued to receive gefitinib after disease progression, and 1 patient who was treated with carboplatin-paclitaxel continued to receive carboplatin-paclitaxel after disease progression because the investigator believed that the treatment was providing a benefit.

were well balanced between the two groups (Table 2 in the Supplementary Appendix).

There was a significant interaction between treatment and *EGFR* mutation with respect to progression-free survival ( $P < 0.001$ ). Progression-free survival was significantly longer among patients receiving gefitinib than among those receiving carboplatin-paclitaxel in the mutation-positive sub-

group (hazard ratio for progression, 0.48; 95% CI, 0.36 to 0.64;  $P < 0.001$ ) (Fig. 2B) and significantly shorter among patients receiving gefitinib than among those receiving carboplatin-paclitaxel in the mutation-negative subgroup (hazard ratio, 2.85; 95% CI, 2.05 to 3.98;  $P < 0.001$ ) (Fig. 2C). Results in the subgroup with unknown *EGFR*-mutation status (hazard ratio with gefitinib, 0.68; 95%



Table 1. Demographic and Baseline Characteristics in the Intention-to-Treat Population.\*

Characteristic	Gefitinib (N=609)	Carboplatin- Paclitaxel (N=608)
Age — yr		
Median	57	57
Range	24–84	25–84
Sex — no. (%)		
Male	125 (20.5)	127 (20.9)
Female	484 (79.5)	481 (79.1)
Ethnic group — no. (%)†		
Chinese	314 (51.6)	304 (50.0)
Japanese	114 (18.7)	119 (19.6)
Other East Asian‡	179 (29.4)	184 (30.3)
Other	2 (0.3)	1 (0.2)
Smoking history — no. (%)		
Never smoked	571 (93.8)	569 (93.6)
Former light smoker	37 (6.1)	38 (6.2)
Former non-light smoker	1 (0.2)	1 (0.2)
WHO performance status — no. (%)§		
0	157 (25.8)	161 (26.5)
1	391 (64.2)	382 (62.8)
2	61 (10.0)	65 (10.7)
Histologic feature of tumor — no. (%)		
Adenocarcinoma	581 (95.4)	591 (97.2)
Bronchoalveolar carcinoma	27 (4.4)	15 (2.5)
Unknown	1 (0.2)	2 (0.3)
Disease stage at entry — no. (%)		
IIIB	150 (24.6)	144 (23.7)
IV	459 (75.4)	463 (76.2)
Unknown	0	1 (0.2)
Time from diagnosis to randomization — no. (%)		
<6 mo	582 (95.6)	573 (94.2)
≥6 mo	27 (4.4)	34 (5.6)
Unknown	0	1 (0.2)
Disease stage at diagnosis — no. (%)¶		
IA	7 (1.1)	12 (2.0)
IB	2 (0.3)	9 (1.5)
IIA	2 (0.3)	1 (0.2)
IIB	1 (0.2)	6 (1.0)
IIIA	6 (1.0)	3 (0.5)
IIIB	166 (27.3)	163 (26.8)
IV	424 (69.6)	413 (67.9)
Unknown	1 (0.2)	1 (0.2)

\* Percentages may not sum to 100 because of rounding.  
 † Ethnic group was self-reported.  
 ‡ Other East Asian refers to patients who belong to East Asian ethnic groups other than Chinese and Japanese.  
 § The World Health Organization (WHO) performance status measures level of activity and is assessed on a scale of 0 to 4, with lower numbers indicating a higher degree of activity.  
 ¶ All patients had Stage IIIB or IV disease at entry.

CI, 0.58 to 0.81;  $P < 0.001$ ) (Fig. 2D) were similar to those for the overall population.

The objective response rate in the overall population was significantly higher with gefitinib than with carboplatin–paclitaxel (43.0% vs. 32.2%; odds ratio, 1.59; 95% CI, 1.25 to 2.01;  $P < 0.001$ ) (Table 3 in the Supplementary Appendix) and numerically or statistically greater with gefitinib in all clinical subgroups. The objective response rate was 71.2% with gefitinib versus 47.3% with carboplatin–paclitaxel in the mutation-positive subgroup ( $P < 0.001$ ) and 1.1% (one patient) versus 23.5%, respectively, in the mutation-negative subgroup ( $P = 0.001$ ) (Table 3 in the Supplementary Appendix).

Overall survival in this early analysis (450 patients [37.0%] died, with follow-up ongoing) was similar between the two groups in the overall population (hazard ratio for death in the gefitinib group, 0.91; 95% CI, 0.76 to 1.10) (Fig. 2A in the Supplementary Appendix). Median survival was 18.6 months among patients receiving gefitinib and 17.3 months among patients receiving carboplatin–paclitaxel. After observing the results with respect to progression-free survival, we performed an analysis of overall survival according to mutation status, although this analysis included only 81 deaths in the mutation-positive subgroup and 94 in the mutation-negative subgroup. The hazard ratios with gefitinib were 0.78 (95% CI, 0.50 to 1.20) in the mutation-positive subgroup and 1.38 (95% CI, 0.92 to 2.09) in the mutation-negative subgroup (Fig. 2B and 2C in the Supplementary Appendix).

Significantly more patients in the gefitinib group than in the carboplatin–paclitaxel group had a clinically relevant improvement in quality of life, as assessed by scores on the FACT-L questionnaire (odds ratio, 1.34; 95% CI, 1.06 to 1.69;  $P = 0.01$ ) and by scores on the TOI (odds ratio, 1.78; 95% CI, 1.40 to 2.26;  $P < 0.001$ ) (Fig. 3). Rates of reduction in symptoms, as assessed on the basis of the LCS scores, were similar between patients who received gefitinib and those who received carboplatin–paclitaxel (odds ratio with gefitinib, 1.13; 95% CI, 0.90 to 1.42;  $P = 0.30$ ) (Fig. 3). Results according to mutation status are provided in Figure 3 in the Supplementary Appendix.

**SAFETY AND ADVERSE-EVENT PROFILE**

Table 2 lists the most common adverse events. Gefitinib, as compared with carboplatin–paclitaxel, was associated with a lower rate of grade 3 or 4




Figure 2. Kaplan–Meier Curves for Progression-free Survival.

Kaplan–Meier curves for progression-free survival are shown for the overall population (Panel A), patients who were positive for the EGFR mutation (Panel B), patients who were negative for the EGFR mutation (Panel C), and patients with unknown EGFR mutation status (Panel D). Analyses were performed on the basis of the intention-to-treat population. With respect to the overall population, results of the supportive secondary analyses (including a log-rank test, which is valid under the null hypothesis even when hazards are not proportional, and analysis in the per-protocol population) were consistent with the result of the primary analysis. Hazard ratios were calculated with the use of a Cox proportional-hazards model, with the WHO performance status (0 or 1, or 2), smoking history (nonsmoker or former light smoker), and sex as covariates. EGFR denotes epidermal growth factor receptor.

adverse events, as defined according to the Common Terminology Criteria for Adverse Events (28.7% vs. 61.0%), a lower rate of adverse events leading to discontinuation of the drug (6.9% vs. 13.6%), and a lower rate of dose modification due to toxic effects (16.1% vs. 35.2% for carboplatin and 37.5% for paclitaxel). Adverse events leading to death occurred in 3.8% of the patients treated with gefitinib and in 2.7% of the patients treated with paclitaxel–carboplatin; serious adverse events, including death, occurred in 16.3% and 15.6% of patients in the two groups, respectively; and seri-

ous adverse events leading to hospitalization occurred in 13.8% and 13.1% of patients in the two groups, respectively. The incidences of rash or acne, diarrhea, and elevated liver aminotransferase levels were significantly higher with gefitinib than with carboplatin–paclitaxel, whereas the incidences of neurotoxic effects, nausea and vomiting, and hematologic toxic effects were significantly higher with carboplatin–paclitaxel (Table 4 in the Supplementary Appendix). Interstitial-lung-disease events (i.e., the acute respiratory distress syndrome, interstitial lung disease, pneumonitis, or radiation




Figure 3. Rates of Improvement in Scores for Quality for Life and Symptoms.

Calculations were performed on the basis of all patients with a baseline and at least one post-baseline quality-of-life assessment that could be evaluated. P values were calculated with the use of logistic regression, with the WHO performance status (0 or 1, or 2), smoking history (nonsmoker or former light smoker), and sex as covariates. Clinically relevant improvement was predefined as an improvement of six points or more in scores on the Functional Assessment of Cancer Therapy–Lung (FACT–L, in which scores range from 0 to 136, with higher scores indicating better quality of life) and Trial Outcome Index (TOI, in which scores range from 0 to 84, with higher scores indicating better quality of life) or an improvement of two points or more in scores on the lung-cancer subscale (LCS) of the FACT–L (in which scores range from 0 to 28, with higher scores indicating fewer symptoms), with the higher scores maintained for at least 21 days.

pneumonitis) occurred in 16 patients treated with gefitinib (2.6%), 3 of whom died, and in 8 patients treated with carboplatin-paclitaxel (1.4%), 1 of whom died.

#### DISCUSSION

Platinum-based combination chemotherapy, such as carboplatin-paclitaxel, is the standard first-line therapy for advanced non-small-cell lung cancer.<sup>25,26</sup> The results of this trial showed that gefitinib by itself is superior to carboplatin-paclitaxel in a selected population of East Asian patients.

As initial treatment of non-small-cell lung cancer in East Asian nonsmokers or former light smokers with pulmonary adenocarcinoma, gefitinib, as compared with carboplatin-paclitaxel, prolonged progression-free survival, increased the objective response rate, reduced toxic effects, and improved quality of life. The overall benefit was driven primarily by the subgroup of patients with *EGFR* mutations; in this subgroup, patients treated with gefitinib, as compared with those treated

with carboplatin-paclitaxel, had a remarkably high objective response rate (71.2%) and prolonged progression-free survival (hazard ratio for progression or death, 0.48; 95% CI, 0.36 to 0.64;  $P < 0.001$ ). In the subgroup of patients without *EGFR* mutations, the objective response rate with gefitinib was 1.1%, and progression-free survival favored chemotherapy (hazard ratio with gefitinib, 2.85; 95% CI, 2.05 to 3.98;  $P < 0.001$ ). These contrasting outcomes probably explain the change over time in treatment effect for progression-free survival in the overall population. The initial superiority of carboplatin-paclitaxel was attributed to the benefit that the *EGFR*-mutation-negative subgroup received from chemotherapy but not from gefitinib, whereas prolonged progression-free survival in the *EGFR*-mutation-positive subgroup explained the subsequent improvement favoring gefitinib. Crossing of the Kaplan-Meier curves did not occur in the mutation-positive subgroup or the mutation-negative subgroup.

Lynch et al. found specific *EGFR* mutations that correlated with tumor response to gefitinib.<sup>7</sup> In the Iressa Survival Evaluation in Lung Cancer trial (ISEL; ClinicalTrials.gov number, NCT00242801), the objective response rate for gefitinib-treated patients was 37.5% among the 16 patients with a tumor bearing an *EGFR* mutation as compared with 2.6% among the 116 patients without a mutation.<sup>27</sup> Our trial confirms the predictive value of *EGFR* mutations for the responsiveness of pulmonary adenocarcinoma to gefitinib as compared with carboplatin-paclitaxel. The difference in the rates of objective response between gefitinib-treated patients with an *EGFR* mutation and those without an *EGFR* mutation (71.2% vs. 1.1%) was remarkable. The rate of an objective response to first-line gefitinib in our study is similar to rates reported in other studies in which patients were selected according to *EGFR*-mutation status, including patients in Western countries.<sup>10,12,28</sup> Sequist et al. screened patients (who were selected on the basis of clinical characteristics) for an *EGFR* mutation and reported an objective response rate of 54.8% among 31 gefitinib-treated patients who were positive for an *EGFR* mutation, only 2 of whom were Asian.<sup>12</sup> However, in our study, objective response rates among patients without an *EGFR* mutation were lower than expected, given the results of previous studies.<sup>16,29</sup> One possible explanation is our use of ARMS, a more sensitive technique for detecting *EGFR* mutations.<sup>21,22</sup> When Zhu et al. used ARMS to reanalyze 148 samples

**Table 2. Adverse Events.\***

Adverse Event	Gefitinib (N=607)		Carboplatin-Paclitaxel (N=589)	
	All Adverse Events	CTC Grade 3, 4, or 5	All Adverse Events	CTC Grade 3, 4, or 5
	<i>number (percent)</i>			
Rash or acne†	402 (66.2)	19 (3.1)	132 (22.4)	5 (0.8)
Diarrhea	283 (46.6)	23 (3.8)	128 (21.7)	8 (1.4)
Dry skin	145 (23.9)	0	17 (2.9)	0
Anorexia†	133 (21.9)	9 (1.5)	251 (42.6)	16 (2.7)
Pruritus†	118 (19.4)	4 (0.7)	74 (12.6)	1 (0.2)
Stomatitis†	103 (17.0)	1 (0.2)	51 (8.7)	1 (0.2)
Asthenic conditions†	102 (16.8)	2 (0.3)	259 (44.0)	11 (1.9)
Nausea	101 (16.6)	2 (0.3)	261 (44.3)	9 (1.5)
Paronychia	82 (13.5)	2 (0.3)	0	0
Vomiting	78 (12.9)	1 (0.2)	196 (33.3)	16 (2.7)
Constipation	73 (12.0)	0	173 (29.4)	1 (0.2)
Alopecia	67 (11.0)	0	344 (58.4)	0
Neurotoxic effects†	66 (10.9)	2 (0.3)	412 (69.9)	29 (4.9)
Myalgia	47 (7.7)	3 (0.5)	186 (31.6)	10 (1.7)
Arthralgia	39 (6.4)	1 (0.2)	113 (19.2)	6 (1.0)
Neutropenia‡				
Any	NA	22 (3.7)	NA	387 (67.1)
Febrile	1 (0.2)	1 (0.2)	17 (2.9)	17 (2.9)
Anemia‡	NA	13 (2.2)	NA	61 (10.6)
Leukopenia‡	NA	9 (1.5)	NA	202 (35.0)

\* Calculations were based on 1196 patients who received at least one dose of the study treatment. The Common Terminology Criteria (CTC) grade is defined on the basis of the National Cancer Institute Common Terminology Criteria for Adverse Events, version 3.0. Events are included if they occurred in at least 10% of patients in either treatment group, either while the patients were receiving treatment or during the 28-day follow-up, and if there was at least a 5% difference between groups. There were other adverse events that occurred in few patients and that may or may not have been related to the study drug. NA denotes not available.

† This is a group term (sum of high-level and preferred terms, according to the definitions in the *Medical Dictionary for Regulatory Activities*).

‡ Data are from the laboratory reports of 599 patients who were taking gefitinib and 577 who were taking carboplatin-paclitaxel. Events were included if there was a worsening in the laboratory value (absolute neutrophil count in the case of neutropenia, hemoglobin in the case of anemia, and white-cell count in the case of leukopenia) from baseline to CTC grade 3 or 4.

that had previously been classified as negative for an EGFR mutation, they found 11 new samples with exon 19 mutations.<sup>30</sup> Another possible explanation is that studies that showed higher response rates among mutation-negative patients were not always conducted in previously untreated patients. Mutation-negative status that is determined in a diagnostic sample obtained at the time of the initial presentation may change during subsequent tumor progression or during the course of chemotherapy.<sup>31</sup>

Our findings suggest that, whenever possible, EGFR-mutation status should be determined before the initial treatment of pulmonary adenocarcino-

ma. Ethnic origin, smoking status, and histologic findings help to identify patients who have a high likelihood of having an EGFR mutation; in this study, 59.7% of the tumors in a clinically selected population had EGFR mutations, as compared with 12.1% and 14.8% in the unselected populations in the ISEL and Iressa in NSCLC Trial Evaluating Response and Survival versus Taxotere (INTEREST; NCT00076388) studies, respectively.<sup>2,27</sup>

The efficacy of gefitinib seen in this study was coupled with lower incidences of alopecia, nausea, vomiting, neurotoxic symptoms, and myelosuppression than those seen with carboplatin-paclitaxel. Among 607 patients who received gefitinib