

# HEMODYNAMIC ALTERATIONS IN THE TRANSFERRED TISSUE TO LOWER EXTREMITIES

HIROYUKI SAKURAI, M.D., Ph.D.,\* TAKASHI YAMAKI, M.D., MASAKI TAKEUCHI, M.D., Ph.D., KAZUTAKA SOEJIMA, M.D., Ph.D., TARO KONO, M.D., and MOTOHIRO NOZAKI, M.D., Ph.D.

A higher incidence of failure has been reported for free flaps transplanted to the lower extremities. However, the physiological background of this phenomenon has not been elucidated. We reviewed the 3-day postoperative hemodynamic data for 103 free flaps, including the in situ venous pressure ( $N = 103$ ), arterial pressure ( $N = 53$ ), and surface blood flow ( $N = 42$ ). The cases were divided into two groups based on the recipient site, i.e., lower extremity (the LE group;  $N = 29$ ) and the other (non-LE group;  $N = 74$ ). The venous pressure was significantly higher in the LE group ( $26.6 \pm 2.2$  vs.  $14.8 \pm 1.2$  mmHg), whereas the arterial pressure immediately after surgery was lower than the non-LE group. The hemodynamic data within the transferred tissues demonstrated significant differences between groups, especially in the early postoperative period. There is a possibility that the high venous pressure may aggravate the poor perfusion in tissues transferred to the lower extremities. © 2008 Wiley-Liss, Inc. *Microsurgery* 29:000–000, 2009.

Lower extremity reconstruction has enjoyed renewed interest with the development of free tissue transfers.<sup>1–3</sup> However, thrombosis and flap failure remain major complications of this method. Despite improved overall success rates approaching 99% in several retrospective analyses of free flaps,<sup>4–6</sup> transfers to the lower extremities still show failure rates of more than 10%, especially under posttraumatic conditions.<sup>2,7,8</sup> Although a multitude of factors determine the success or failure of a microvascular free tissue transfer,<sup>9</sup> the actual reason for the higher risk of flap failure in lower extremity reconstruction has not been elucidated. Infection has been suggested as a predominant cause of morbidity in free flaps transferred to the lower legs.<sup>10</sup> However, other factors must be considered in the majority of cases of failure in which aggressive wound care and antisepsis were properly undertaken. Hypercoagulability associated with thrombocytosis may be related to the higher incidence of vascular thrombosis in lower leg reconstructions,<sup>11</sup> although this scenario also is applicable only to limited cases in limited situations.

The vascular system in the legs is unique because it has a high hydrostatic pressure during orthostasis,<sup>12,13</sup> and it is easily affected by several disorders on both the arterial and venous sides. We hypothesized that tissues transferred to the lower extremities may present different hemodynamics compared with other parts of the body in the early postoperative period. Currently, we have developed a novel method for monitoring the flap circulation

with regard to the intraparenchymatous vascular pressure.<sup>14</sup> This study is based upon data collected in patients who underwent microvascular free tissue transfer with this novel monitoring system. The patient profiles such as sex, etiology, and preferred tissue for reconstruction and recipient vessels were totally different among patient groups. In spite of these inherent differences, we decided that it would be informative to determine the hemodynamic characteristics in tissues transferred to different recipient sites.

## PATIENTS AND METHODS

### Patients

This study included 103 consecutive patients who underwent free tissue transfer with venous pressure monitoring between 2003 and 2007. This monitoring system, approved by the ethical committee of Tokyo Women's Medical University, included a continuous data acquisition system,<sup>14</sup> therefore, we retrospectively reviewed hemodynamic data in the transferred flap. The patients with a history of cardiopulmonary disorder or hypertension were excluded in the study. All patients underwent surgery under general anesthesia. There were 38 female and 65 male patients. The patients ranged in age from 9 to 82 years, with a mean age of 54 years. The patients were divided into two groups depending on the recipient site, lower extremity (LE group;  $N = 29$ ), and other sites (non-LE group;  $N = 74$ ). The specific donor sites employed in the two groups are shown in the Table 1. In the LE group, we preferentially used the deep venous system rather than the superficial venous system as the recipient vein (Table 2). We included four cases of stasis ulcer because of superficial venous insufficiency, for which superficial venous stripping was performed prior to the free tissue transfer.

Department of Plastic and Reconstructive Surgery, Tokyo Women's Medical University, Shinjuku-ku, Tokyo 162-8666, Japan

\*Correspondence to: Hiroyuki Sakurai, M.D., Ph.D., 8-1 Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan. E-mail: bighiro@prs.twmu.ac.jp

Received 13 July 2008; Accepted 19 August 2008

Published online in Wiley InterScience(www.interscience.wiley.com). DOI 10.1002/micr.20570

### Venous Pressure Measurements

The patients were fully informed concerning the reasons, the technique of catheter placement, the details of

the monitoring procedure, and the consequences as well as possible complications associated with catheter placement.

After completion of vascular anastomoses, an intravenous catheter (3 Fr, 1.0-mm OD, Atom Medical, Tokyo, Japan) was inserted into a side branch of the anastomosed vein. The catheter was then connected to a fluid pressure transducer (P231D, Statham Gould, Oxnard, CA) and to a physiological recorder (BSS-9800, Nihon Kohden, Tokyo, Japan). The line solution of the venous catheter contained heparin at 10 U/ml in 0.9% sodium chloride, and the flow rate of the flush solution was ~3 ml/hr, resulting in continuous infusion of the heparin at 720 U/day. Zero calibrations were taken at the level of the right atrium.

Three days after the operation, the catheter was locked and disconnected from the transducer, but left in position for a further 4 days. Thereafter, the catheter was gently tugged at and withdrawn from the flap.

Table 1. Patient Profiles: Gender, Age, and Donor Site

	LE group (N = 29)	Non-LE group (N = 74)
Male/female	24/5	41/33
Age (year)	49 ± 4	52 ± 2
Type of tissues		
Anterolateral thigh flap	15	16
Groin flap	5	9
LD musculocutaneous flap	3	11
Omentum	2	
Fibula osteocutaneous flap	2	4
Rectus abdominis MC flap		12
Jejunum		7
Forearm flap		6
DIEP flap		3
DP		3
Others	2	3

Table 2. Patient Profiles of the LE Group

No.	Sex	Age	Etiology	Flap	Recipient vessel		Measurement			Comment
					Artery	Vein	VP	AP	LDP	
1	M	29	Tibial bone open fracture	FOCF	PTA	PTV	+			
2	M	64	Intractable ulcer on the lower leg	ALT	ATA	ATV	+			
3	M	74	Intractable ulcer on the lower leg	Omentum	ATA	ATV	+	+		
4	M	71	Tibial bone open fracture	FOCF	ATA	ATV	+			
5	M	35	Traumatized lower leg	ALT	ATA	ATV	+	+		Arterial thrombosis
6	M	59	Intractable stasis ulcer on the lower leg	Omentum	ATA	ATV	+	+		
7	M	17	Scar contracture of the dorsal foot	Groin flap	DPA	GSV	+			
8	M	68	Chronic osteomyelitis of tibial bone	ALT	ATA	ATV	+	+	+	
9	M	71	MM at the heel	ALT	PTA	PTV	+		+	
10	M	46	Intractable stasis ulcer on the lower leg	ALT	PTA	PTV	+		+	
11	M	43	SCAR contracture around knee joint	Groin flap	DGA	GSV, DGV	+	+	+	
12	M	80	SCC at the medial foot	ALT	PTA	PTV	+	+	+	
13	F	63	Intractable stasis ulcer on the lower leg	ALT	PTA	PTV	+	+	+	
14	M	55	Chronic osteomyelitis of the femur	ALT	DGA	GSV	+	+	+	
15	M	9	Degloving injury of the foot	LD	ATA	ATV	+	+	+	
16	M	28	Mangled lower extremity	Fillet flap	FA	FV	+			
17	M	33	Intractable ulcer on the foot	ALT	ATA	ATV	+	+		Hematoma
18	F	64	Intractable ulcer on the lower leg	LD	ATA	ATV	+			
19	F	13	Postburn contracture on the thigh	Groin flap	SIEA	GSV	+		+	
20	M	68	SCC on the lower leg	ALT	ATA	ATV	+	+	+	
21	F	58	AVM on the foot	Groin flap	LTA	SSV	+	+	+	
22	M	49	Postburn contracture on the lower leg	Scapular flap	DGA	SSV	+		+	
23	M	49	Diabetic gangrene	ALT	ATA	ATV	+	+		
24	M	59	Intractable ulcer on the lower leg	LD	FA	ATV	+	+		
25	M	64	Traumatized foot	Groin flap	DPA	GSV	+	+	+	
26	M	20	Traumatized foot	ALT	PTA	PTV	+	+	+	
27	F	68	Intractable stasis ulcer on the lower leg	ALT	PTA	PTV	+	+	+	Venous thrombosis
28	M	48	Intractable ulcer on the foot	ALT	ATA	ATV	+	+	+	
29	M	17	Intractable ulcer on the foot	ALT	PTA	PTV	+	+	+	

M, male; F, female; FOCF, fibula osteocutaneous flap; ALT, anterolateral thigh flap; LD, latissimus dorsi MC flap; PTA (V), posterior tibial artery (vein); ATA (V), anterior tibial artery (vein); DPA, dorsalis pedis artery; DGA(V), descending genicular artery (vein); LTA, lateral tarsal artery; GSV, great saphenous vein; SSV, small saphenous vein.

### Arterial Pressure Monitoring

Fifty-three of the patients (LE group: 19, non-LE group: 34) also underwent in situ arterial catheterization for further hemodynamic analysis. The method was essentially the same as that for the venous catheterization. However, the line solution of the arterial catheter did not contain heparin, avoiding the risk of bleeding tendency in the transferred tissue. In some case, the arterial catheterization was impossible because of the small size of the diameter. In those cases, we used Doppler auscultation for the monitoring of arterial thrombosis.

### Laser Doppler Flowmetry

In addition, the surface blood flow was continuously monitored in 42 patients (LE group: 17, non-LE group: 25) using a laser Doppler flowmeter (Laserflo BPM 403, TSI, St. Paul, MN) with a standard right angle probe having a head diameter of 19 mm (Model P-430). This probe is applicable only to the exteriorized skin island. This apparatus provided flow rates in millimeters per minute per 100 g of tissue.

### Data Acquisition

All parameters were continuously recorded for 3 days at a sampling frequency of 250 Hz via a commercial data acquisition system (PowerLab<sup>®</sup>, AD Instruments Pty., Australia). During this postoperative period, all patients were confined to bed rest. Because all parameters were influenced by the patient's positioning, posture, and movements, analysis was performed of sections showing stable hemodynamics for more than 30 minutes while the patients were at rest and in the supine position (immediately after surgery and 12 and 72 hours after surgery). The mean values of the venous pressure, arterial pressure, and blood flow were obtained using the customized program included in the data acquisition system (PowerLab).

### Data Analysis and Statistics

Data are expressed as means  $\pm$  SEM. The effects of different recipient sites (i.e., LE, non-LE groups) on the hemodynamic data were tested using ANOVA with repeated measures across time. Statistical significance was defined as a *P* value of 0.05 or less. All statistical analyses were performed with SAS statistical package version 8.2.

## RESULTS

There were three cases with postoperative anastomotic failure (two cases of venous thrombosis and one case of arterial thrombosis) in the 103 cases. One of the cases with venous thrombosis received a mandibular reconstruction with free fibular osteocutaneous flap (in the non-LE group). In this case, cutaneous flap was located

in the oral cavity, therefore, the flow probe of the LDF was not readily applicable. The other case who developed venous thrombosis presented with an intractable stasis ulcer. In this case, we applied both venous pressure monitoring and LDF monitoring. Both methods were quite sensitive responding the compromised venous outflow. The arterial thrombosis in another case was detected with the continuous arterial pressure monitoring. All three cases were successfully salvaged by the early detection with the monitoring system. There was no statistically significant difference in the incidence of the vascular thrombosis between two groups.

Free flaps transplanted to the lower extremities demonstrated a significantly higher venous pressure compared with the other sites (LE:  $26.6 \pm 2.2$  mmHg, non-LE:  $14.8 \pm 1.2$  mmHg) (see Fig. 1). The high venous pressure in the LE group declined gradually, whereas the venous pressure in the non-LE group remained virtually unchanged during the 3-day postoperative period.

The arterial pressure in the LE group was significantly lower than that in the non-LE group (see Fig. 2). Therefore, the arteriovenous pressure gradients, defined as the driving pressure for perfusion of transferred tissues, showed significant differences between groups (see Fig. 3). The driving pressure in the LE group was  $52.6 \pm 3.7$  mmHg immediately after surgery, whereas those in the non-LE group was  $75.0 \pm 2.4$  mmHg. Thereafter, the lower driving pressure in the LE group gradually increased to  $59.8 \pm 3.6$  mmHg at 72 hours after surgery; however, it was still significantly lower than the non-LE group.

The surface blood flow measured by the laser Doppler flowmeter in the LE group was  $2.2 \pm 0.6$  ml/min/100 g immediately after surgery, whereas that in the non-LE group was  $3.7 \pm 0.5$  ml/min/100 g (see Fig. 4). There was no statistically significant difference between groups. As the driving pressure increased during the 3-day postoperative period, the blood flow increased significantly in the LE group ( $4.1 \pm 0.9$  ml/min/100 g on the 3rd postoperative day). On the other hand, the blood flow in the non-LE group was relatively constant during the 3-day postoperative period.

## DISCUSSION

Free tissue transfer using a microvascular technique has been firmly established since its introduction more than three decades ago. However, little is known about the physiological aspects within the transferred tissue. The monitoring technique currently developed in our institution,<sup>14</sup> in which the in situ vascular pressure can be continuously measured, provides us a great opportunity to elucidate the hemodynamic changes after free tissue transfer. Although this method required catheterization of




Figure 1. Changes in venous pressure within the tissues transferred to the lower extremity (LE group) and the other sites (non-LE group). Values are given as the mean  $\pm$  SEM. \*: Significant difference ( $P < 0.05$ ) from the value immediately after surgery. †: Significant difference ( $P < 0.05$ ) between groups.




Figure 3. Changes in arteriovenous pressure gradient within the tissues transferred to the lower extremity (LE group) and the other sites (non-LE group). Values are given as the mean  $\pm$  SEM. \*: Significant difference ( $P < 0.05$ ) from the value immediately after surgery. †: Significant difference ( $P < 0.05$ ) between groups.




Figure 2. Changes in arterial pressure within the tissues transferred to the lower extremity (LE group) and the other sites (non-LE group). Values are given as the mean  $\pm$  SEM. †: Significant difference ( $P < 0.05$ ) between groups.




Figure 4. Changes in the value of LDF after free tissue transfer. Values are given as the mean  $\pm$  SEM. \*Significant difference ( $P < 0.05$ ) from the value immediately after surgery.

the vessels within the transferred flap, the safety and efficacy of such catheterization has been well documented in previous reports in which anticoagulant agents were continuously infused through a catheter.<sup>15-17</sup>

In this study, the hemodynamic data within the transferred tissue were strongly influenced by the recipient site. The venous pressure in the LE group was significantly higher compared with the values in the non-LE group. It is conceivable that the positioning of the lower extremities may affect the venous pressure. Because we analyzed the venous pressure during the initial 3 days after surgery while the patients rested in bed the whole day, the gravitational effects on the venous pressure were

considered to be at a minimum. Additionally, the site of the arterial catheterization was identical to the venous catheterization; therefore, the effect of the position of the recipient sites on the hemodynamic data was negligible when one looked at the pressure gradient between the arterial and venous sides. Because the arterial pressure in the LE group was significantly lower than that in the non-LE group, the arteriovenous pressure gradient showed clear differences between the groups. According to the concept of Burton,<sup>18</sup> a decreased driving pressure results in reduced blood flow. Although we could not detect significant difference in the incidence of anastomotic failure between the two groups, there is a possibil-

ity that the hemodynamic differences might be related to the reported high failure rates in lower leg reconstruction using free-tissue transfer.<sup>2,7,8</sup>

Lower limb hemodynamic impairment associated with surgery and general anesthesia has been well documented. Foate et al.<sup>19</sup> reported that the mean arterial leg blood flow in patients under general anesthesia was only 28% of the control at 1 hour after surgery. This compromised arterial inflow has been linked to a higher rate of early postoperative graft failure in infrainguinal arterial bypass surgery, whereas epidural anesthesia did not cause this adverse effect of general anesthesia.<sup>20</sup> Knaggs et al.<sup>21</sup> reported that the maximal venous flow attenuation was seen during the early recovery period. By disrupting the laminar flow of the venous circulation, stasis brings platelets into contact with the endothelium, leads to the build-up of thrombi and promotes the activation of endothelial cells, setting off a vicious circle of hypercoagulability. Although the mechanisms involved in the susceptibility of the lower leg hemodynamics to general anesthesia and surgical interventions are not fully understood, the early postoperative period is the most critical for vascular compromise even after free tissue transfer.

Despite the lower driving pressure in the LE group, we could not detect significant differences in the LDF data. This inability to detect differences may be attributed to the method we used for assessment of the blood flow. It is well known that the flow values with LDF vary greatly depending on the patient and the type of tissue.<sup>22-24</sup> Therefore, one must not rely on absolute values, and recent reports emphasize the importance of observing the trend of the perfusion rather than the absolute value.<sup>22,23</sup> It is of interest that the decrease in the venous pressure during a 3-day postoperative period was associated with an increased blood flow in the LE group.

When the venous pressure in a limb is elevated to more than 25 mmHg, the cutaneous, subcutaneous, and muscle vascular resistances increase within that region, resulting in a reduction in blood flow of about 40%.<sup>13</sup> This reflex has been termed the venoarteriolar response, because stretch receptors located in small veins induce changes in the arteriolar vascular tone "upstream" of the veins.<sup>12</sup> A local mechanism, not the central neurogenic response, seems to mediate the venoarteriolar response. Therefore, the high venous pressure may aggravate the poor perfusion in the transferred tissue through this response, leading to the high incidence of failure in lower limb reconstruction.

This study is limited by its small number of patients. Further investigations are warranted to determine the effects of injury-related variables, the type of transferred tissue, the selection of the recipient vessels, local heparinization, and epidural anesthesia. Nonetheless, monitoring the vascular pressure through in situ catheterization

provided us further insight into the pathophysiology after free tissue transfer.


## CONCLUSIONS

In summary, tissues transferred to the lower extremities demonstrated a higher venous pressure and poorer perfusion immediately after surgery compared with tissues transferred to other parts of the body. Thereafter, the venous pressure gradually decreased in association with an increase in the blood perfusion of the flap. There is a possibility that the impaired hemodynamics in tissues transferred to the lower extremities might be a contributing factor to the higher incidence of flap failure during the early postoperative period.

## REFERENCES

- Pollak AN, McCarthy ML, Burgess AR. Short-term wound complications after application of flaps for coverage of traumatic soft-tissue defects about the tibia. The Lower Extremity Assessment Project (LEAP) Study Group. *J Bone Joint Surg Am* 2000;82A:1681-1691.
- Gonzalez MH, Tarandy DI, Troy D, Phillips D, Weinzwieg N. Free tissue coverage of chronic traumatic wounds of the lower leg. *Plast Reconstr Surg* 2002;109:592-600.
- Godina M. Early microsurgical reconstruction of complex trauma of the extremities. *Plast Reconstr Surg* 1986;78:285-292.
- Hidalgo DA, Disa JJ, Cordeiro PG, Hu QY. A review of 716 consecutive free flaps for oncologic surgical defects: Refinement in donor-site selection and technique. *Plast Reconstr Surg* 1998;102:722-732.
- Kroll SS, Schusterman MA, Reece GP, Miller MJ, Evans GR, Robb GL, Baldwin BJ. Timing of pedicle thrombosis and flap loss after free-tissue transfer. *Plast Reconstr Surg* 1996;98:1230-1233.
- Disa JJ, Cordeiro PG, Hidalgo DA. Efficacy of conventional monitoring techniques in free tissue transfer: An 11-year experience in 750 consecutive cases. *Plast Reconstr Surg* 1999;104:97-101.
- Benacquista T, Kasabian AK, Karp NS. The fate of lower extremities with failed free flaps. *Plast Reconstr Surg* 1996;98:834-840.
- Langstein HN, Chang DW, Miller MJ, Evans GR, Reece GP, Kroll SS, Robb GL. Limb salvage for soft-tissue malignancies of the foot: An evaluation of free-tissue transfer. *Plast Reconstr Surg* 2002;109:152-159.
- Nahabedian MY, Momen B, Manson PN. Factors associated with anastomotic failure after microvascular reconstruction of the breast. *Plast Reconstr Surg* 2004;114:74-82.
- Swartz WM, Mears DC. The role of free-tissue transfers in lower-extremity reconstruction. *Plast Reconstr Surg* 1985;76:364-373.
- Choe EI, Kasabian AK, Kolker AR, Karp NS, Zhang L, Bass LS, Nardi M, Josephson G, Karpatkin M. Thrombocytosis after major lower extremity trauma: Mechanism and possible role in free flap failure. *Ann Plast Surg* 1996;36:489-494.
- Hassan AA, Tooke JE. Mechanism of the postural vasoconstrictor response in the human foot. *Clin Sci (Lond)* 1988;75:379-387.
- Okazaki K, Fu Q, Martini ER, Shook R, Conner C, Zhang R, Crandall CG, Levine BD. Vasoconstriction during venous congestion: Effects of venoarteriolar response, myogenic reflexes, and hemodynamics of changing perfusion pressure. *Am J Physiol Regul Integr Comp Physiol* 2005;289:R1354-R1359.
- Sakurai H, Nozaki M, Takeuchi M, Soejima K, Yamaki T, Kono T, Fukaya E, Yamamoto Y. Monitoring the changes in intraparenchymatous venous pressure to ascertain flap viability. *Plast Reconstr Surg* 2007;119:2111-2117.
- May JW Jr, Rothkopf DM. Salvage of a failing microvascular free muscle flap by direct continuous intravascular infusion of heparin: A case report. *Plast Reconstr Surg* 1989;83:1045-1048.

16. Harris NR II, Webb MS, May JW Jr. Intraoperative physiologic blood flow studies in the TRAM flap. *Plast Reconstr Surg* 1992;90:553-558.
17. Hudson DA, Engelbrecht G, Duminy FJ. Another method to prevent venous thrombosis in microsurgery: An in situ venous catheter. *Plast Reconstr Surg* 2000;105:999-1003.
18. Burton AC. Resistance to flow in vascular beds. *Nature* 1958;182:1456-1457.
19. Foate JA, Horton H, Davis FM. Lower limb blood flow during transurethral resection of the prostate under spinal or general anaesthesia. *Anaesth Intensive Care* 1985;13:383-386.
20. Christopherson R, Beattie C, Frank SM, Norris EJ, Meinert CL, Gottlieb SO, Yates H, Rock P, Parker SD, Perler BA. Perioperative morbidity in patients randomized to epidural or general anesthesia for lower extremity vascular surgery. Perioperative Ischemia Randomized Anesthesia Trial Study Group. *Anesthesiology* 1993;79:422-434.
21. Knaggs AL, Delis KT, Mason P, Macleod K. Perioperative lower limb venous haemodynamics in patients under general anaesthesia. *Br J Anaesth* 2005;94:292-295.
22. Yuen JC, Feng Z. Monitoring free flaps using the laser Doppler flowmeter: Five-year experience. *Plast Reconstr Surg* 2000;105:55-61.
23. Heller L, Levin LS, Klitzman B. Laser Doppler flowmeter monitoring of free-tissue transfers: Blood flow in normal and complicated cases. *Plast Reconstr Surg* 2001;107:1739-1745.
24. Mailaender P, Machens HG, Waurick R, Rieck B, Berger A. Routine monitoring in patients with free tissue transfer by laser-Doppler flowmetry. *Microsurgery* 1994;15:196-202.

available at [www.sciencedirect.com](http://www.sciencedirect.com)journal homepage: [www.elsevier.com/locate/burns](http://www.elsevier.com/locate/burns)

## Case reports

# Prefabricated flap for multiple facial units reconstruction using a jejunal seromuscular patch as a vascular carrier

Hiroyuki Sakurai\*, Masaki Takeuchi, Daiki Nakamori, Taro Kono, Kazutaka Soejima, Takashi Yamaki, Motohiro Nozaki

Department of Plastic and Reconstructive Surgery, Tokyo Women's Medical University, 8-1 Kawada-cho Shinjuku-ku, Tokyo 162-8666, Japan

## ARTICLE INFO

### Article history:

Accepted 6 September 2008

## 1. Introduction

Prefabricated flap is a versatile reconstructive option especially in patients with severe burns, because it can create new donor sites without limitation of natural vascular territories. In this method, a vascular carrier is implanted to a new skin territory. Following a period of maturation and neovascularization, the prefabricated flap can be transferred, based on the implanted pedicle. The very first attempt of prefabricated flap was reported by Washio [1] in a canine model using an intestinal seromuscular patch as the vascular carrier. However, the development of various types of tissue transfer, such as muscle [2], fascia [3], and vascular pedicle alone, has precluded the usage of intestine in this clinical setting.

For reconstruction of the facial skin, the concept of the aesthetic units [4] has been well accepted since its introduction more than five decades ago. Prefabricated flaps have been utilized for reconstruction of individual aesthetic unit such as nose, ear, cheek, lip and neck [5–7]. However, in a case of composite units defect, a single vascular carrier may not be sufficient to perfuse all units to be reconstructed. In these cases, separated vascular carriers that can nourish each unit independently are desirable. We experienced a case in which the units of nose, upper lip, and lower lip were simultaneously reconstructed with a prefabricated flap. In this case, we used jejunal seromuscular patches as “vascular carriers”, because

the jejunum could be divided in multiple segments owing to the arcade structure of the vessels. Description of the technical details and the experience gained with the patient form the basis of this report.

## 2. Case Report

A 72-year-old man sustained a 37.5 percent body surface area burn, including the whole face, neck, upper and lower extremities. Association of severe inhalation injury resulted in complete laryngeal obstruction necessitating tracheotomy. Sequential debridement and skin grafting were required to primarily close the burn wound. After complete wound closure, the patient presented with severe facial deformities including partial loss of the nose and bilateral ears (Fig. 1). The lower lip was severely pulled down, resulting in poor oral hygiene and interference with the oral sphincter function. The left anterior chest was left intact for the facial reconstruction, and we initiated a plan to develop a prefabricated flap in this area for the simultaneous reconstruction of the composite aesthetic units including nose, upper lip and lower lip.

The jejunal segment was followed to approximately 40 cm below the ligament of Treitz, where the vascular arcades were evaluated by transillumination. Approximately 25 cm of jejunum was harvested to obtain the long pedicle. The

\* Corresponding author. Tel.: +81 3 3353 8111; fax: +81 3 3225 0940.  
E-mail address: [bighiro@prs.twmu.ac.jp](mailto:bighiro@prs.twmu.ac.jp) (H. Sakurai).  
0305-4179/\$36.00 © 2008 Elsevier Ltd and ISBI. All rights reserved.  
doi:10.1016/j.burns.2008.09.011



Fig. 1 – Preoperative findings.

methods to elongate the vascular pedicle were described elsewhere [8,9]. Briefly, the oral side of the mesentery was dissected transversely leaving the most distal arcade, vasa recta, while the anal side of the arcade was connected to the mesenteric artery and vein. After completion of the revascularization with microvascular anastomoses of jejunal vessels to the left facial artery and vein, the jejunal flap was divided into three segments. After the conduits of the jejunum were opened, the mucosa was hydrodissected with an epinephrine solution in order to decrease cumbersome oozing and to facilitate dissection [10]. The mucosa was sharply dissected away leaving a thin seromuscular layers (Fig. 2). These three seromuscular flaps were implanted to the subcutaneous layer of the anterior chest through the incision of the suppositional oral orifice, for prefabrication of nose, upper lip, and lower lip,



Fig. 2 – Jejunal seromuscular flaps after dissection of the mucosal layer.

independently. At the same time, costal cartilage was transplanted to the nasal portion sandwiched between skin and transferred seromuscular layer. (Fig. 3)


Five weeks after surgery, the prefabricated flap was elevated. The vascular pedicle and implanted jejunal segment were easily dissected with minimum adhesion to the underlying tissue (Fig. 4). Even after fenestration for the mouth and nostrils, three segments of the prefabricated flap demonstrated stable perfusion. After transplantation of the prefabricated flap, the donor site was closed with mesh skin grafting (Fig. 5). The Postoperative course was uneventful, and free from complications. Three years after surgery, the shape of the nose was well maintained without any shrinkage of the prefabricated flap (Fig. 6).

### 3. Discussion

Severe post-burn scarring of the face presents a formidable challenge to the reconstructive surgeon. Skin grafts are generally used during acute burn management, but they almost invariably contract over time. Because full supplementation of dermis and soft tissue is required to correct a fundamental problem, various types of flaps have been used to reconstruct the facial skin [11,12]. In addition to functional considerations, this area needs to be reconstructed with the concept of aesthetic unit [4,11].

The clinical applications of flap prefabrication have increased reconstructive capabilities and overcome problems not solved satisfactorily by the available classical free flaps. Because this method can also modify the shape of skin flaps, reconstruction of individual units such as nose, ear, cheek, lip and neck have been previously reported [5-7]. The prerequisites for an ideal vascular carrier for prefabricated flap are high





**Fig. 3 – Prefabricated flap created in the left anterior chest. Segment 1: prefabrication for nsasal reconstruction. The costal cartilage was transplanted between skin and transferred seromuscular layer. Segment 2: prefabrication for upper lip reconstruction. Segment 3: prefabrication for lower lip reconstruction.**



**Fig. 4 – Dissection of the prefabricated flap. The vascular pedicle and implanted jejunal segments were easily dissected with minimum adhesion to the underlining tissue.**

intrinsic vascularity, a vascular pedicle of adequate length and caliber, limited bulk, acceptable donor site morbidity, and predictable ability to form new vascular connections. In the previous study, Tark et al. [13] demonstrated that the musculovascular pedicle surpasses the other carriers in revascularization of new tissues, with its abundant vascular beds and a vascular pedicle of adequate length and caliber. However, the bulkiness and donor site morbidity may limit the use of muscles as carriers. On the other hand, a fasciovascular pedicle is a thin pliable tissue that allows wider attachments to the area of prefabrication, although it is lagging in quantitative skin-surface perfusion because of less intrinsic vascularity [13]. In this regard, the jejunal seromuscular patch met all properties expected for the ideal vascular carrier.

Small intestinal submucosa (SIS) is a type I collagen-based biomaterial extracted from the porcine intestine in a manner that removes all cells. This material can attract host cell migration [14,15], therefore, it has been clinically used as a scaffold for tissue engineering in a variety of systems such as the bladder [16], urethra [17], blood vessels [18], tendon [19], and the abdominal wall [20]. In addition, the SIS has been reported to induce the proliferation of human dermal microvascular endothelial cells, producing several angiogenic cytokines [21,22] such as fibroblast growth factor (FGF-2), transforming growth factor- $\beta$  (TGF- $\beta$ ), and vascular endothelial growth factor (VEGF). These favorable properties of the SIS in tissue engineering supported the superiority of the jejunal seromuscular patch as the vascular carrier of prefabricated flaps.

The unique architecture of the mesenteric vessels was the major reason why the jejunal seromuscular patch was used in this particular patient who required composite facial units reconstruction. After fenestration for the mouth and nostril, a single vascular carrier may not be sufficient to perfuse all units. The jejunal patch could be easily divided into three



Fig. 5 – Simultaneous reconstruction of the composite facial units. (Left) elevated prefabricated flap. (Right) immediately after operation.



Fig. 6 – Three years after reconstruction.

segments, facilitating simultaneous reconstruction of three different facial units.

To the best of our knowledge, this is the first report of clinical application using intestinal seromuscular patch as a vascular carrier for prefabrication. Although necessity of a laparotomy is a drawback of this method, it is worthy to be considered especially when the composite facial units need to be restored with limited donor site.

REFERENCES

- [1] Washio H. An intestinal conduit for free transplantation of other tissues. *Plast Reconstr Surg* 1971;48:48-51.
- [2] Shintomi Y, Ohura T. The use of muscle vascularized pedicle flaps. *Plast Reconstr Surg* 1982;70:725-35.

Please cite this article in press as: Sakurai H, et al. Prefabricated flap for multiple facial units reconstruction using a jejunal seromuscular patch as a vascular carrier. *Burns* (2009), doi:10.1016/j.burns.2008.09.011

- [3] Khouri RK, Ozbek MR, Hruza GJ, Young VL. Facial reconstruction with prefabricated induced expanded (PIE) supraclavicular skin flaps. *Plast Reconstr Surg* 1995;95:1007-15.
- [4] Gonzalez-Ulloa M. Restoration of the face covering by means of selected skin in regional aesthetic units. *Br J Plast Surg* 1956;9:212-21.
- [5] Teot L, Cherenfant E, Otman S, Giovannini UM. Prefabricated vascularised supraclavicular flaps for face resurfacing after postburns scarring. *Lancet* 2000;355:1695-6.
- [6] Pribaz JJ, Fine N, Orgill DP. Flap prefabrication in the head and neck: a 10-year experience. *Plast Reconstr Surg* 1999;103:808-20.
- [7] Evans DM. Facial reconstruction after a burn injury using two circumferential radial forearm flaps, and a dorsalis pedis flap for the nose. *Br J Plast Surg* 1995;48:471-6.
- [8] Sakurai H, Nozaki M, Sasaki K, Nakazawa H. The use of free jejunal autograft for the treatment of vaginal agenesis: surgical methods and long-term results. *Br J Plast Surg* 2000;53:319-23.
- [9] Sakurai H, Nozaki M. Reconstruction of the pharyngoesophagus with voice restoration. *Int J Clin Oncol* 2005;10:243-6.
- [10] Zubowicz VN, Mitchell CW. Seromuscular patch as an adjunct in reconstruction of the cervical esophagus with free jejunal transfer. *Plast Reconstr Surg* 1995;96:1709-12.
- [11] Sakurai H, Takeuchi M, Fujiwara O, Soejima K, Yamaki T, Kono T, Nozaki M. Total face reconstruction with one expanded free flap. *Surg Technol Int* 2005;14:329-33.
- [12] Sakurai H, Soejima K, Takeuchi M, Kono T, Yamaki T, Nozaki M. Reconstruction of perioral burn deformities in male patients by using the expanded frontal scalp. *Burns* 2007;33:1059-64.
- [13] Tark KC, Tuchler RE, Shaw WW. Flap prefabrication: effectiveness of different vascular carriers. *Ann Plast Surg* 1996;37:298-304.
- [14] Lindberg K, Badylak SF. Porcine small intestinal submucosa (SIS): a bioscaffold supporting in vitro primary human epidermal cell differentiation and synthesis of basement membrane proteins. *Burns* 2001;27:254-66.
- [15] MacLeod TM, Sarathchandra P, Williams G, Sanders R, Green CJ. Evaluation of a porcine origin acellular dermal matrix and small intestinal submucosa as dermal replacements in preventing secondary skin graft contraction. *Burns* 2004;30:431-7.
- [16] Misseri R, Cain MP, Casale AJ, Kaefer M, Meldrum KK, Rink RC. Small intestinal submucosa bladder neck slings for incontinence associated with neuropathic bladder. *J Urol* 2005;174:1680-2.
- [17] Mantovani F, Trinchieri A, Castelnuovo C, Romano AL, Pisani E. Reconstructive urethroplasty using porcine acellular matrix. *Eur Urol* 2003;44:600-2.
- [18] McCready RA, Hodde J, Irwin RJ, Coffey AC, Divelbiss JL, Bryant MA, et al. Pseudoaneurysm formation in a subset of patients with small intestinal submucosa biologic patches after carotid endarterectomy. *J Vasc Surg* 2005;41:782-8.
- [19] Iannotti JP, Codsì MJ, Kwon YW, Derwin K, Ciccone J, Brems JJ. Porcine small intestine submucosa augmentation of surgical repair of chronic two-tendon rotator cuff tears. A randomized, controlled trial. *J Bone Joint Surg Am* 2006;88:1238-44.
- [20] Gabriel A, Gollin G. Management of complicated gastroschisis with porcine small intestinal submucosa and negative pressure wound therapy. *J Pediatr Surg* 2006;41:1836-40.
- [21] Voytik-Harbin SL, Brightman AO, Kraine MR, Waisner B, Badylak SF. Identification of extractable growth factors from small intestinal submucosa. *J Cell Biochem* 1997;67:478-91.
- [22] Hodde JP, Record RD, Liang HA, Badylak SF. Vascular endothelial growth factor in porcine-derived extracellular matrix. *Endothelium* 2001;8:11-24.

PEPARS

◆特集/四肢のリンパ浮腫の治療

# Untied Stay Suture 法による リンパ管静脈吻合とリンパ管静脈 吻合術の有効性

長谷川健二郎\*<sup>1</sup> 杉山成史\*<sup>2</sup> 難波祐三郎\*<sup>3</sup> 木股敬裕\*<sup>4</sup>

Key Words: リンパ浮腫 (lymphedema), リンパ管静脈吻合術 (lymphaticovenous anastomosis), 微小血管吻合術 (microsurgery), 超微小血管吻合術 (supermicrosurgery), 蛍光リンパ管造影法 (fluorescence lymphography)

**Abstract** 当科では2006年4月よりICG 蛍光リンパ管造影法を用いたリンパ管静脈吻合術を行っている。手術手技では、我々の考案した1st stay suture を untied suture とし、2nd stay suture の糸が通ってから両方の糸を結紮する Untied Stay Suture 法を、特に0.3~0.5 mm 以下のリンパ管静脈吻合に応用している。現在までの手術症例数は上肢13例、下肢63例、合計76症例である。今回術後5か月以上経過観察できた上肢12例・下肢37例を対象に評価した。上肢・下肢症例を合わせた54肢の結果は excellent 17肢(31%), good 21肢(39%), fair 9肢(17%), poor 7肢(13%)であり、70%に効果を示した。

## はじめに

従来、我々は四肢リンパ浮腫に対して、光嶋ら<sup>1)</sup>が報告したスーパーマイクロサージャリーテクニックを用いたリンパ管静脈吻合術(LVA: lymphaticovenous anastomosis)を行ってきた。2006年4月にICG 蛍光リンパ管造影法(FL-ICG: fluorescence lymphography using ICG)を導入することにより、リンパ管の同定が容易となり、また吻合直後にリンパ管から静脈内にリンパ液が入っていることを確認できるためLVAがより確実になってきた。さらに我々の考案したUntied Stay Suture(USS)法は0.5 mm以下のリンパ管や微小血管の吻合には有用であり、LVAの有効性と併せて報告する。

## Untied Stay Suture 法によるリンパ管静脈吻合

まずICG 蛍光リンパ管造影法を用いリンパ管を観察する。手背や足背などの任意の場所にICGを皮内注射し、特殊な赤外線カメラ(Photo Dynamic Eye:PDE, 浜松ホトニクス社製)で観察すると、リンパ管はICGの線状の流れとして造影される(図1)。

リンパの流れに沿ってボールペンで直上の皮膚にマーキングをいれ、同部を切開し顕微鏡下にリンパ管を探して確認し、同時に皮静脈も確保しておく。静脈にはsingle clampを付けるが、リンパ管はリンパ液を流したままのほうが開口部を確認しやすいのでclampを付けない。静脈はできるだけ尾側で、リンパ管は頭側で切離し、縫合部に緊張がかからないようにする(図2)。

口径が0.5 mm以上のリンパ管の吻合は11-0 ナイロン糸で4~6針を目安に従来の微小血管吻合法を用いる。0.3~0.5 mm以下のリンパ管の吻合には12-0 ナイロン糸で我々の考案したUntied Stay Suture法を用いている。

まず静脈側より針を入れる。鑷子の先が血管内

\*1 Kenjiro HASEGAWA, 〒700-8558 岡山市鹿田町2-5-1 岡山大学大学院医歯薬学総合研究科形成再建外科, 講師

\*2 Narushi SUGIYAMA, 同, 助教

\*3 Yuzaburo NAMBA, 同, 准教授

\*4 Yoshihiro KIMATA, 同, 教授




図 1. PDE を使った FL-ICG によるリンパ管の観察  
 星印：ICG を趾間部に皮内注射  
 黄色矢印：造影された足背部のリンパ管



図 2. 吻合直前のリンパ管と皮静脈  
 青色矢印：皮静脈断端  
 黄色矢印：リンパ管断端




図 3. 第1針目の静脈への刺入



図 4. 第1針目のリンパ管への刺入




図 5. untie にされた 1st stay suture




図 6. 第2針目の静脈への刺入




図 7. 第2針目のリンパ管への刺入



図 8. リンパ管・静脈内に通った1st・2nd stay suture



図 9. 第3針目の静脈・リンパ管への刺入




図 10. 第3針目の縫合後




図 11. LVA 終了時  
青色矢印：皮静脈  
黄色矢印：リンパ管




図 12. PDE による LVA 直後の観察  
青色矢印：造影された皮静脈  
黄色矢印：造影されたリンパ管

に入らない場合には軽くそえるようにして刺入する。針が入ったら必ず入ったことを確認する(図3;本症例は静脈の径が約0.7mmと大きく、鑷子が入っている)。

次にリンパ側に針を入れる。注意深く12-0の針先を内膜の下に滑り込ませる(図4)。リンパ管の壁は弱いので、針を抜く時も注意が必要である。

1針目のstay sutureはuntieにし(図5)、糸が抜けないように糸の両端を血管クリップで留め、助手がリンパ管と静脈の断端に適度な緊張がかかるように調節する(図6~8)。一般的には微細血管吻合の際には1針目はstay sutureとして結紮するが、極端に小さなリンパ管の吻合の際には、結紮時に生じる捻れの問題と2針目が入れやすいということより1針目をuntieにして2針目を入れている。1針目の縫合糸を結紮せずにリンパ管-静脈内に残すことによりintravascular stenting(IVaS)法<sup>2)</sup>と類似の効果が得られる(図7)。

2針目はできるだけ1針目から180°の位置に入れるようにする(図6~8)。2針目の刺入方法は1針目と同じであるが、助手がuntieにした1st stay sutureの糸を上手くコントロールし緊張をかけることにより特にリンパ管への刺入が易しくなる(図7)。

2針目が入ったらstay sutureを結紮する。内腔を確認したい場合や鑷子の先を内腔に入れて縫合したい場合には1st・2nd stay sutureをuntieとし、先に3・4針目を結紮した後に2本のstay sutureを結紮することもあるが、LVAの場合には内腔が正しく合っていれば、stay sutureを結紮することによりリンパ管からリンパ液が静脈内に流れ込み、しばらく待機するとリンパ管・静脈が次第に膨らんでくる(図9)。吻合部が膨らむと、次の3・4針目は比較的容易に入れることができる。

3針目つまり表側の縫合であるが、stay sutureを結紮した場合、鑷子が内腔に入らないので注意して針を入れる必要がある(図9, 10)。

最後に血管クリップを反転し表側と同様に裏側

の縫合を施行する。吻合が終了し血管クリップを外し顕微鏡下に確認すると、リンパ流量が十分な場合には静脈血の逆流はなく、静脈がリンパ液で十分に膨らんでいるのが確認できる(図11;本症例では一部逆流した静脈血が吻合部を越えてリンパ管の中にまだ少し残っている)。

吻合後PDEで観察すると、黒く写っているバックグランドシートの上で、リンパ管から静脈内にリンパ液が入り、静脈がリンパ管と同様に造影されていることを確認できる(図12)。

### リンパ管静脈吻合術の有効性

2006年4月にICG蛍光リンパ管造影法を導入した後に、LVAを施行し、術後5か月以上の経過観察が可能であった上肢リンパ浮腫12症例12肢、下肢リンパ浮腫37症例42肢を対象に評価し、有効性について検討した。

上肢症例は全例女性で、乳癌術後に発症していた。年齢は54~87歳で、左上肢7肢、右上肢5肢であった。発症より手術までの期間は1年6か月~29年であった。下肢症例は男性6例、女性31例で、特発性が9症例、2次性が28症例であった。女性は全て女性性器腫瘍術後に発症していた。年齢は35~81歳で、左下肢29肢、右下肢13肢であった。発症から手術までの期間は6か月~30年であった。

評価は図13のごとく、上肢または下肢で、5か所の周囲径を測定し、術前の周囲径に対する術後の周囲径の割合(改善率)をもとめ、いずれかの部位で89%以下に改善したものをexcellent(図14-a)、90~95%の改善をgood(図14-b)とした。96~104%の変化は測定誤差を考慮し不変と考えfair(図14-c)とした。そしていずれの部位にも改善を認めず、105%以上の悪化を認めた場合をpoor(図14-d)と評価した。

上肢、下肢症例を合わせた54肢の結果はexcellent 17肢(31%)、good 21肢(39%)、fair 9肢(17%)、poor 7肢(13%)であった。Excellentとgoodを有効としfairとpoorを無効と考えると、




図 13. 評価方法




図 14. 評価に基づく代表症例

a : excellent, b : good, c : fair, d : poor

a/b  
c/d



表 1. 四肢リンパ浮腫に対する FL-ICG を用いた LVA の術後成績

評価	上肢	下肢	合計 (%)
excellent	5	12	17 (61)
good	7	17	24 (89)
fair	16	8	24 (87)
poor	2	5	7 (23)
	2	12	14 (100)

表 2. 下肢リンパ浮腫に対する FL-ICG を用いた LVA の術後成績

評価	特発性	2次性	合計 (%)
excellent	1	11	12 (29)
good	6	11	17 (40)
fair	1	8	9 (21)
poor	0	5	5 (12)
	10	32	42 (100)

70%が有効で30%が無効であった。上肢症例と下肢症例間には有意差を認めなかった(表1)。下肢症例を特発性と2次性に分けて同様に検討を加えたが両者間にも有意差は認めなかった(表2)。

### 考 察

1st stay suture を untied suture とし 2nd stay suture の糸が通ってから両方の糸を結紮する Untied Stay Suture 法は、壁が薄く透明なリンパ液の流れ込んでくる、特に 0.3~0.5 mm 以下の LVA に有効な手技だと考えられる。また指尖部切断再接着術のような 0.3~0.5 mm 以下の微細血管吻合の場合には、1st・2nd stay suture を untied suture とし先に 3・4 針目を結紮した後に 2本の stay suture を結紮するほうが縫合しやすい。しかし 0.5 mm 以上のリンパ管や微細血管吻合の場合には従来法を用いた方が簡便である。

ICG 蛍光リンパ管造影法が導入される以前の光嶋らのリンパ浮腫に対するスーパーマイクロサージャリーテクニックを用いたリンパ管静脈吻合術の報告によると、上肢リンパ浮腫に対する結果では、12症例を対象に経過観察期間1か月~6年(平均2.2年)で、前腕(肘から10cm遠位)において、0~8cm(平均4.1cm)の減少が得られ、4cm以上の減少がみられたのは7例(58.3%)であった<sup>1)</sup>。また、下肢リンパ浮腫に対する結果では、

13症例に対して術後平均4.6年の経過観察で、下腿(膝から10cm遠位)において平均4.7cmの減少が得られ、4cm以上の減少がみられたのは8例(61.5%)であった<sup>3)</sup>。局麻下の LVA の結果では、52症例を対象に術後平均14.5か月の経過観察で、43症例(82.5%)に効果が得られ、4cm以上の減少がみられたのは17例(32.7%)であった<sup>4)</sup>。これらの報告は評価法に違いがあり、今回の結果と比較することは困難であるが、我々の結果と同様にリンパ管静脈吻合術の有効性を示したものと考えられる。

### 文 献

- 1) Koshima, I, et al. : Supermicrosurgical lymphaticovenular anastomosis for the treatment of lymphedema in the upper extremities. J Reconstr Microsurg. 16 : 437-442, 2000.
- 2) Narushima, M., et al. : Intravascular stenting (IVaS) for safe and precise supermicrosurgery. Ann Plast Surg. 60 : 41-44, 2008.
- 3) Koshima, I, et al. : Long-term follow-up after lymphaticovenular anastomosis for lymphedema in the leg. J Reconstr Microsurg. 19 : 209-215, 2003.
- 4) Koshima, I, et al. : Minimal invasive lymphaticovenular anastomosis under local anesthesia for leg lymphedema. Ann Plast Surg. 53 : 261-266, 2004.

男性下肢リンパ浮腫に対する ICG 蛍光リンパ管造影法を  
用いたリンパ管静脈吻合術

岡山大学大学院医歯薬学総合研究科形成再建外科

長谷川 健二郎, 目谷 雅恵, 雑賀 美帆,  
木股 敬裕

Lymphaticovenous Anastomosis Using Indocyanine Green  
Fluorescence Lymphography for Treatment of Male  
Lymphedema of the Lower Limbs

by

Kenjiro HASEGAWA, Masae METANI, Miho SAIGA,  
Yoshihiro KIMATA

*Department of Plastic and Reconstructive Surgery  
Okayama University Graduate School of Medicine and Dentistry  
Okayama, Japan*

原 著

## 男性下肢リンパ浮腫に対する ICG 蛍光リンパ管造影法を用いたリンパ管静脈吻合術

岡山大学大学院医歯薬学総合研究科形成再建外科

長谷川 健二郎, 目 谷 雅 恵, 雑 賀 美 帆,  
木 股 敬 裕

## Lymphaticovenous Anastomosis Using Indocyanine Green Fluorescence Lymphography for Treatment of Male Lymphedema of the Lower Limbs

by

Kenjiro HASEGAWA, Masae METANI, Miho SAIGA,  
Yoshihiro KIMATA

*Department of Plastic and Reconstructive Surgery  
Okayama University Graduate School of Medicine and Dentistry  
Okayama, Japan*

### Abstract

Since April 2006, we have performed lymphaticovenous anastomosis (LVA) using indocyanine green fluorescence lymphography (FL-ICG). The use of FL-ICG in LVA allowed more precise identification of lymphatics and anastomosis to veins. We report the usefulness of this method for treatment of male lymphedema of the lower limbs.

Among the patients who underwent LVA since we introduced FL-ICG, 6 male patients (7 limbs) with lower-limb lymphedema who were followed postoperatively for 6 months or longer were included in the present study.

Indocyanine green (ICG) was injected intracutaneously at several points on the lower extremity, and the dye distribution was observed using a special infrared camera (Photo Dynamic Eye). ICG is taken up into the lymphatics and its spread along with the lymph flow can be monitored. When a lymphatic vessel was visualized by linear flow of ICG, an incision was made at the site and the lymphatic was searched and identified using an operating microscope, and then anastomosed to the cutaneous vein.

Among a total of 7 limbs, 4 were evaluated as excellent, 3 as good, 0 as fair and

---

Key words : Lymphedema (リンパ浮腫), Lymphaticovenous Anastomosis (リンパ管静脈吻合術),  
Microsurgery (マイクロサージャリー), Supermicrosurgery (超微小血管吻合術),  
Fluorescence Lymphography (蛍光リンパ管造影法)

Good 3肢, Fair 0肢, Poor 0肢であった(表1).

### 代表症例

症例は63歳男性で、6年前に左単径部の滑膜肉腫の切除術・術後放射線治療を受けるも、その後左単径部リンパ節転移を認め、さらに3回のリンパ節切除術と術後放射線治療を受けた。その後左下肢リンパ浮腫を認め、当科に紹介され受診となった。手術は左単径部と左膝内側部の2カ所に端々でLVAを施行した(図3-a)。

術後6ヵ月の状態で、大腿部以外の場所で89%以下に改善しており、評価はExcellentであった。特に、下腿部で7.5cmの改善が認められた(図3-b)。

### 考察

従来、我々は四肢リンパ浮腫に対しKoshimura<sup>2)</sup>が報告したスーパーマイクロサージャリーテクニックを用いたリンパ管静脈吻合術(LVA)を行ってきた。しかし、LVAにおけるリンパ管の同定は術者の経験に左右されるところが大きく、またLVA後のリンパ液が確実に静脈内に入っていることを確認することは困難であった。2006年4月に照喜納ら<sup>7)</sup>が、ICG

蛍光リンパ管造影法を下肢リンパ浮腫症例のLVAに用いた報告をきっかけに、我々もLVAにPDEを用いたICG蛍光リンパ管造影法を導入した。PDEを用いる事により、術中にリアルタイムに流れのあるリンパ管を確認出来るようになり、また吻合直後にリンパ管から静脈内にリンパ液が入っていることを確認できるためLVAがより確実になってきた。特に皮切を加えた後もリンパ管をPDEで観察しながら追いかけることは特記すべき点である。問題点としては深さ約2cmまでしか観察出来ないために、より深部のリンパ管を探す場合にはどうしても経験にもとづいて皮切を加えた後に、PDEで観察することになる点である。

ICG蛍光リンパ管造影法が導入される以前の報告では、光嶋らは、下肢リンパ浮腫13症例に対して、術後平均4.6年の経過観察で、下腿(膝から10cm遠位)において平均4.7cmの減少が得られ、4cm以上の減少がみられたのは8例(61.5%)であった<sup>3)</sup>と報告している。また、局麻下のLVAの結果では、52症例に対して術後平均14.5ヵ月の経過観察で、全体の43症例(82.5%)に効果が得られ、4cm以上の減少がみられたのは17例(32.7%)であった<sup>4)</sup>と報告している。これらの報告は評価法に違いがあり、




図3 症例4: 63歳, 男性, 左下肢リンパ浮腫