

effect. This study investigated preventive effects for healthy people, not curative effects for patients with metabolic syndrome or other diseases. Expanding this study to cover investigation in patient groups is an issue for future consideration.

References

- Abe T, Kawakami Y, Suzuki Y, Gunji A. Prediction equations for body composition of Japanese adults by B-mode ultrasound. *Am J Hum Biol* (1994); **6**: 161–170.
- Ades PA, Savage PD, Brochu M, Tischler MD, Lee NM, Poehlman ET. Resistance training increases total daily energy expenditure in disabled older women with coronary heart disease. *J Appl Physiol* (2005); **98**: 1280–1285.
- Anton MM, Cortez-Cooper MY, DeVan AE, Neidre DB, Cook JN, Tanaka H. Resistance training increases basal limb blood flow and vascular conductance in aging humans. *J Appl Physiol* (2006); **101**: 1351–1355.
- Baechle TR, Earle RW, Wathen D. Resistance training. In: *Essential of strength training and conditioning*. Human Kinetics (2000); 408–409.
- Baron AD, Laakso M, Brechtel G, Hoit B, Watt C, Edelman SV. Reduced postprandial skeletal muscle blood flow contributes to glucose intolerance in human obesity. *J Clin Endocrinol Metab* (1990); **70**: 1525–1533.
- Bonde-Petersen F, Mork AL, Nielsen E. Local muscle blood flow and sustained contractions of human arm and back muscles. *Eur J Appl Physiol Occup Physiol* (1975); **34**: 43–50.
- Borst SE. Interventions for sarcopenia and muscle weakness in older people. *Age Ageing* (2004); **33**: 548–555.
- Cheitlin MD, Armstrong WF, Aurigemma GP, Beller GA, Bierman FZ, Davis JL, Douglas PS, Faxon DP, Gillam LD, Kimball TR, Kusmaul WG, Pearlman AS, Philbrick JT, Rakowski H, Thys DM, Antman EM, Smith SC Jr, Alpert JS, Gregoratos G, Anderson JL, Hiratzka LF, Faxon DP, Hunt SA, Fuster V, Jacobs AK, Gibbons RJ, Russell RO. ACC/AHA/ASE 2003 Guideline Update for the Clinical Application of Echocardiography: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASE Committee to Update the 1997 Guidelines for the Clinical Application of Echocardiography). *J Am Soc Echocardiogr* (2003); **16**: 1091–1110.
- Dela F, Kjaer M. Resistance training, insulin sensitivity and muscle function in the elderly. *Essays Biochem* (2006); **42**: 75–88.
- Dinenno FA, Jones PP, Seals DR, Tanaka H. Limb blood flow and vascular conductance are reduced with age in healthy humans: relation to elevations in sympathetic nerve activity and declines in oxygen demand. *Circulation* (1999); **100**: 164–170.
- Dinenno FA, Seals DR, DeSouza CA, Tanaka H. Age-related decreases in basal limb blood flow in humans: time course, determinants and habitual exercise effects. *J Physiol* (2001a); **531**: 573–579.
- Dinenno FA, Tanaka H, Stauffer BL, Seals DR. Reductions in basal limb blood flow and vascular conductance with human ageing: role for augmented alpha-adrenergic vasoconstriction. *J Physiol* (2001b); **536**: 977–983.
- Evans WJ, Cyr-Campbell D. Nutrition, exercise, and healthy aging. *J Am Diet Assoc* (1997); **97**: 632–638.
- Fleck SJ. Cardiovascular adaptations to resistance training. *Med Sci Sports Exerc* (1988); **20**: S146–S151.
- Hasten DL, Pak-Loduca J, Obert KA, Yarasheski KE. Resistance exercise acutely increases MHC and mixed muscle protein synthesis rates in 78–84 and 23–32 yr olds. *Am J Physiol Endocrinol Metab* (2000); **278**: E620–E626.
- Hunter GR, McCarthy JP, Banman MM. Effects of resistance training on older adults. *Sports Med* (2004); **34**: 329–348.
- Leithe ME, Margorien RD, Hermiller JB, Unverferth DV, Leier CV. Relationship between central hemodynamics and regional blood flow in normal subjects and in patients with congestive heart failure. *Circulation* (1984); **69**: 57–64.
- Lind L, Lithell H. Decreased peripheral blood flow in the pathogenesis of the metabolic syndrome comprising hypertension, hyperlipidemia, and hyperinsulinemia. *Am Heart J* (1993); **125**: 1494–1497.
- MacDougall JD, Ward GR, Sale DG, Sutton JR. Biochemical adaptation of human skeletal muscle to heavy resistance training and immobilization. *J Appl Physiol* (1977); **43**: 700–703.
- MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR. Arterial blood pressure response to heavy resistance exercise. *J Appl Physiol* (1985); **58**: 785–790.
- McDonagh MJ, Davies CT. Adaptive response of mammalian skeletal muscle to exercise with high loads. *Eur J Appl Physiol Occup Physiol* (1984); **52**: 139–155.
- Miyachi M, Kawano H, Sugawara J, Takahashi K, Hayashi K, Yamazaki K, Tabata I, Tanaka H. Unfavorable effects of resistance training on central arterial compliance: a randomized intervention study. *Circulation* (2004); **110**: 2858–2863.
- Miyachi M, Tanaka H, Kawano H, Okajima M, Tabata I. Lack of age-related decreases in basal whole leg blood flow in resistance-trained men. *J Appl Physiol* (2005); **99**: 1384–1390.
- Moreau KL, Donato AJ, Tanaka H, Jones PP, Gates PE, Seals DR. Basal leg blood flow in healthy women is related to age and hormone replacement therapy status. *J Physiol* (2003); **547**: 309–316.
- Ozdemir H, Artas H, Serhadioglu S, Ogur E. Effects of overweight on luminal diameter, flow velocity and intima-media thickness of carotid arteries. *Diagn Interv Radiol* (2006); **12**: 142–146.
- Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC, Kriska A, Leon AS, Marcus BH, Morris J, Paffenbarger RS Jr, Patrick K, Pollock ML, Rippe JM, Sallis J, Wilmore JH. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* (1995); **273**: 402–407.
- Sahn DJ, DeMaria A, Kisslo J, Weyman A. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* (1978); **58**: 1072–1083.
- Shinohara M, Moritani T. Increase in neuromuscular activity and oxygen uptake during heavy exercise. *Ann Physiol Anthropol* (1992); **11**: 257–262.
- Staron RS, Hikida RS, Hagerman FC, Dudley GA, Murray TF. Human skeletal muscle fiber type adaptability to various workloads. *J Histochem Cytochem* (1984); **32**: 146–152.
- Takarada Y, Nakamura Y, Aruga S, Onda T, Miyazaki S, Ishii N. Rapid increase in plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *J Appl Physiol* (2000a); **88**: 61–65.
- Takarada Y, Takazawa H, Ishii N. Applications of vascular occlusion diminish disuse atrophy of knee extensor muscles. *Med Sci Sports Exerc* (2000b); **32**: 2035–2039.
- Tanaka H, Dinenno FA, Monahan KD, Clevenger CM, DeSouza CA, Seals DR. Aging, habitual exercise, and dynamic arterial compliance. *Circulation* (2000); **102**: 1270–1275.
- Tanimoto M, Ishii N. Effects of low-intensity resistance exercise with slow movement and tonic force generation on muscular function in young men. *J Appl Physiol* (2006); **100**: 1150–1157.
- Tanimoto M, Sanada K, Yamamoto K, Kawano H, Gando Y, Tabata I, Ishii N, Miyachi M. Effects of whole-body low-intensity resistance training with slow movement and tonic force generation (LST) on muscular size and strength in young men. *J Strength Cond Res* (2008); **22**: 1926–1938.

Original Article

Attenuated Increases in Blood Pressure by Dynamic Resistance Exercise in Middle-Aged Men

Hiroshi KAWANO^{1),2)}, Hitomi NAKAGAWA³⁾, Sho ONODERA³⁾,
Mitsuru HIGUCHI⁴⁾, and Motohiko MIYACHI²⁾

The present study was performed to test the hypothesis that the blood pressure (BP) response to resistance exercise in middle-aged men with stiffening arteries is greater than that in young men with compliant arteries. The BP responses to acute dynamic resistance exercise (leg press) at individual relative (low, moderate and high) and absolute intensities were investigated in both young and middle-aged men. A total of 21 sedentary healthy normotensive men, 21–25 years of age (young) and 41–59 years of age (middle-aged), were included in the study. At rest, the arterial compliance (simultaneous ultrasound and applanation tonometry) and muscle strength (leg press) were lower, and indices of arterial stiffness and BP were higher in the middle-aged men than in the young men ($p < 0.05$). There were no significant differences in height, body mass, or heart rate between the two groups. During exercise, the systolic BP of the middle-aged men at 80% one-repetition maximum (1RM) was significantly lower than that of the young men for the last half of the exercise period ($p < 0.05$). The amounts of change in systolic and diastolic BP from baseline to the end of resistance exercise were lower in the middle-aged men than in the young men at individual relative intensities ($p < 0.05$) and at individual absolute intensity. In contrast to our hypothesis, these findings indicated that the BP response during dynamic resistance exercise using large muscle groups may be attenuated in middle-aged men relative to young men. (*Hypertens Res* 2008; 31: 1045–1053)

Key Words: aging, resistance exercise, blood pressure, pressor, arterial stiffening

Introduction

Regular physical activity is regarded as an important component of prevention and treatment of age-related increases in cardiovascular disease (1, 2). Aerobic exercise in particular is recommended by major health organizations, including the American Heart Association and American College of Sports Medicine (3, 4), because it shows favorable effects on cardiovascular functions in young, middle-aged, and older men (5–8). In recent years, resistance exercise, another common exer-

cise modality, has gained widespread acceptance in exercise prescription and cardiopulmonary rehabilitation programs and has become an integral component of comprehensive health programs endorsed by the major health organizations (3, 9). However, there is very little information on the potential influence of resistance training on non-musculoskeletal components, in particular the cardiovascular system. Systolic and diastolic blood pressures (BPs) rise rapidly to extremely high values during heavy weight-lifting exercise (10), and BPs are extreme even when exercise is performed with a relatively small muscle mass (11, 12). Most previous studies

From the ¹⁾Graduate School of Human Sciences and ⁴⁾Faculty of Sport Sciences, Waseda University, Tokorozawa, Japan; ²⁾Division of Health Promotion and Exercise, National Institute of Health and Nutrition, Tokyo, Japan; and ³⁾Department of Health and Sports Sciences, Kawasaki University of Medical Welfare, Kurashiki, Japan.

This work was supported by a Grant-in-Aid for Scientific Research (No. 13780041 to M.M.) from the Japan Society for the Promotion of Science and by a Sasagawa Scientific Research Grant (to H.K.) from the Japan Science Society.

Address for Reprints: Hiroshi Kawano, Ph.D., Division of Health Promotion and Exercise, National Institute of Health and Nutrition, 1–23–1 Toyama, Shinjuku-ku, Tokyo 162–8636, Japan. E-mail: hiroshi@aoni.waseda.jp

Received September 11, 2007; Accepted in revised form December 28, 2007.

have focused on the BP responses to resistance exercise in young adults (13–15), and thus have provided little information regarding BP responses in middle-aged and older individuals (12, 16). Furthermore, the interaction between age and BP response for dynamic resistance exercise using large muscle groups has not been reported in middle-aged men with stiffening arteries or young men with compliant arteries.

The stiffness of the large central arteries in the cardiothoracic region increases with advancing age in sedentary humans (6, 17, 18). This physiological alteration reduces the buffering capacity of the arteries, leading to increased pulse pressure, peripheral vessel resistance, and left ventricular wall tension (19, 20), all of which augment the workload of the heart. Generally, it can be assumed that middle-aged and older men with advanced arteriosclerosis would show attenuated pressor buffering function during resistance exercise. Accordingly, we hypothesized that the BP response to dynamic resistance exercise in middle-aged men with stiffening arteries would be larger than that in young men with compliant arteries. The present study was performed to clarify the differences in BP response to acute dynamic resistance exercise at individual relative or absolute intensities between young and middle-aged men.

Methods

Subjects

Twelve young (21–25 years) and nine middle-aged men (41–58 years) were recruited through various forms of advertisement or by posting on bulletin boards at our university, and had not participated in a regular exercise program for at least the previous 2 years. Only male subjects were included in the study to ensure that the interpretation of differences between the two age groups would not be confounded by the possible influence of sex. All subjects were normotensive (<140/90 mmHg), non-obese (body mass index <30 kg/m²), and free of overt chronic diseases as assessed by medical history and physical examination. Subjects taking cardiovascular-acting medications, such as anabolic steroids, or with significant intima-media thickening, plaque formation, and/or other characteristics of atherosclerosis (e.g., ankle-brachial index [ABI] <0.9) were excluded from the study. All subjects gave their written informed consent to participation in the study, and all procedures were reviewed and approved by the Institutional Review Board.

Measurements at Rest

Brachial-Ankle Pulse Wave Velocity and BP

The brachial-ankle pulse wave velocity (baPWV) and BP were assessed using a form PWV/ABI device (Colin Medical Technology, Komaki, Japan). Subjects were examined in the supine position after a rest of at least 5 min. The cuffs were wrapped on both sides of the brachium and ankle, and con-

tained a plethysmographic sensor that determined the waveform data, including BP measurements by the oscillometric method. The baPWV was calculated as distance/time (cm/s) between the brachium and ankle. The time delay between the arrival of the pulse wave at the brachium and ankle was obtained automatically by gating the pulse wave to the peak of the R wave of the electrocardiogram. The distance was estimated from the subject's height as $L = L_a - L_b$ (L_a : path length from the heart to the ankle; L_b : path length from the heart to the brachium). Then, we used the mean of the right and left baPWV values for analysis. The measurements of baPWV and BP were performed three times per day, and the three values were averaged. The reproducibility of baPWV measurement has been validated in our laboratory (coefficient of covariance, $2 \pm 1\%$) and in other studies (21, 22).

Carotid Artery Intima-Media Thickness

The right common carotid artery intima-media thickness (IMT) was measured from images obtained using an ultrasound machine equipped with a high-resolution linear-array broad-band transducer as described previously (23, 24). Ultrasound images were analyzed using computerized image analysis software (NIH Image 1.63). At least 10 measurements of IMT were taken at each segment, and the mean values were used for analysis. This technique has excellent day-to-day reproducibility (coefficient of variance, $3 \pm 1\%$) for the carotid IMT.

Carotid Artery Compliance and Stiffness

A combination of ultrasound imaging of the pulsatile right common carotid artery with simultaneous applanation of tonometrically obtained arterial pressure from the contralateral carotid artery permits noninvasive determination of arterial compliance (23–25). The carotid artery diameter was measured from images obtained using an ultrasound machine equipped with a high-resolution linear-array transducer. A longitudinal image of the cephalic portion of the common carotid artery was acquired 1–2 cm distal to the carotid bulb. All image analyses were performed by the same investigator who was blinded to the group assignments.

Pressure waveforms and amplitudes were obtained from the common carotid artery with a pencil-type probe incorporating a high-fidelity strain-gauge transducer (SPT-301; Millar Instruments, Houston, USA) (26). As baseline levels of BP are subject to hold-down force, the pressure signal obtained by tonometry was calibrated by equating the carotid mean arterial and diastolic BP to the brachial artery value (23–25). In addition to arterial compliance (27), we also calculated the β -stiffness index, which provides an index of arterial compliance adjusted for distending pressure (28). Arterial compliance and the β -stiffness index were calculated using the equations $[(D_1 - D_0)/D_0]/[2(P_1 - P_0)](P_1 - P_0) \times \pi \times (D_0)^2$ and $[\ln(P_1/P_0)]/[(D_1 - D_0)/D_0]$, where D_1 and D_0 are the maximal and minimum diameters, and P_1 and P_0 are the highest and lowest BPs, respectively (23–25). The day-to-day coeffi-

coefficients of variation were $2\pm 1\%$, $7\pm 3\%$, and $5\pm 2\%$ for the carotid artery diameter, pulse pressure, and arterial compliance, respectively.

Measurements during Dynamic Resistance Exercise

Radial BP and ECG

To determine circulatory response, radial BP and ECG were recorded simultaneously in the sitting position at baseline, during exercise, and during the recovery period. ECG and radial BP waveforms were determined using arterial tonometry (JENTOW-7700; Colin Medical Technology) and standard lead electrocardiography (Life Scope 11; Nihon Kohden, Tokyo, Japan), respectively. Both ECG and arterial BP waveforms were sampled at 1,000 samples per second by connecting each device to a computer using an A/D converter (PowerLab; AD Instruments, Colorado Springs, USA). The principle of arterial tonometry is that BP at the radial artery can be obtained by measuring the reaction forces produced by flattening the radial artery. Recently, this method has become preferred over the conventional finger photoplethysmographic method (Finapres), and it has been confirmed that the accuracy and reliability of BP measured by tonometry are greater than those of BP measured by the intra-arterial method. A tonometric sensor was attached to the left wrist, and the wrist was placed on a padded platform at the level of the heart. The oscillometric calibrations were carried out for accurate tonometric measurement before, and sometimes during the main experiment (29).

Strength Testing

Maximal muscular strength was assessed with a leg press machine using air pressure (Keiser; Fitness Apollo Japan Co., Ltd. Tokyo, Japan). A one-repetition maximum (1RM) was determined by having the subjects perform single repetitions with progressively heavier weights, resting 2–3 min between attempts; the heaviest weight that subjects could lift once through a complete range of movement was considered their 1RM. The day-to-day coefficient of variation for 1RM strength in our laboratory is $4\pm 2\%$.

Exercise Protocol I

Subjects rested under quiet conditions before beginning the leg press exercise. After a 60-s baseline period, all subjects randomly performed 10 repetitions of the leg press exercise for 40 s at each of 40%, 60%, and 80% 1RM, followed by an 80-s recovery period. One repetition was performed for 4 s (2 s for concentric and 2 s for eccentric contraction). To measure radial BP accurately, the subject's left arm was supported on an adjustable table during measurement. Subjects were stabilized in the apparatus during exercise using their right hand to hold the support handle on the seat. The left arm was allowed to rest freely by their side to avoid interference with the recording of radial BP from this arm. The subjects were

Table 1. Subject Characteristics

	Young	Middle-aged
<i>N</i>	12	9
Age, years	21.4±0.5	47.8±1.9
Height, cm	170.6±1.7	170.1±2.1
Body mass, kg	65.3±2.1	67.7±2.9
Resting heart rate, bpm	52.3±1.7	60.8±4.4
Brachial systolic BP, mmHg	118±3	123±4
Brachial diastolic BP, mmHg	67±2	80±3*
Brachial mean BP, mmHg	82±4	96±4*
Brachial PP, mmHg	55±4	43±2*
Carotid systolic BP, mmHg	109±2	119±6
Carotid diastolic BP, mmHg	67±2	80±3*
Carotid PP, mmHg	42±2	38±3
Carotid diastolic diameter, mm	5.9±0.1	6.7±0.3*
Carotid intima-media thickness, mm	0.48±0.02	0.63±0.02*
Carotid arterial compliance, mm ² /mmHg	0.17±0.01	0.11±0.01*
Carotid β -stiffness index, a.u.	3.95±0.28	7.30±0.76*
Brachial-ankle PWV, cm/s	1,092±38	1,291±46*
Augmentation index, %	-6.9±5.7	19.6±5.8*
Leg press maximum, kg	350±11	286±19*

Data are mean±SEM. BP, blood pressure; PP, pulse pressure; PWV, pulse wave velocity. Leg press maximum was evaluated by air pressure machine (Keiser; Fitness Apollo Japan Instruments). *Significant at $p < 0.05$ vs. young.

encouraged to avoid deep inhalations while performing the Valsalva maneuver during the exercise. Exercise measurements were performed randomly in a day. The interval time between exercises was controlled at 10 min.

Exercise Protocol II

Eleven young men and nine middle-aged men were studied using protocol II. All subjects performed 10 repetitions of the leg press exercise at individual absolute intensity (145 kgw) 10 min after the end of protocol I. Protocol II was performed using a procedure similar to that in protocol I.

Data Analysis

In the baseline period, during the leg press exercise and recovery periods, waveforms of ECG and radial BP were recorded continuously and simultaneously on a personal computer (iBook G3; Apple Computer, Cupertino, USA). Heart rate and systolic, diastolic, and mean BP were calculated using the Chart5 software package (AD Instruments). Baseline values of BP were taken as the average of the baseline period (1 min) before exercise. Average values every 4 s for one repetition and peak values of BP were obtained during the 40-s exercise period. Eight BP values in the recovery period were averaged at 0–4, 4–8, 16–20, 36–40, 56–60, 76–80, 96–









Fig. 1. Systolic (circles) and diastolic (squares) blood pressure responses during resistance exercises and recovery periods at 40% (bottom), 60% (middle), and 80% (top) of 1RM in young (white) and middle-aged (black) men. Values are means \pm SEM. * $p < 0.05$ vs. young men.

Fig. 2. Heart rate responses during resistance exercises and recovery periods at 40% (bottom), 60% (middle), and 80% (top) of 1RM in young (white) and middle-aged (black) men. Values are means \pm SEM.




Fig. 3. The amounts of change in systolic (SBP; top left) and diastolic (DBP; bottom left) blood pressure, pulse pressure (PP; top right) and heart rate (HR; bottom right) responses to resistance exercises at 40%, 60%, and 80% of 1RM in middle-aged (black bar) and young (white bar) men. Values are means \pm SEM. * $p < 0.05$ vs. young men at the same intensity.

100, and 116–120 s. Heart rate was calculated as the peak value during exercise.

Statistical Analysis

Changes during the leg press exercise were assessed by two-way analysis of variance (group \times time) with repeated measures. In the case of significant F -values, a post hoc test (Newman-Keuls method) was used to identify significant differences among mean values. Peak values were analyzed using the t -test. All data are presented as the means \pm SEM. Statistical significance was set at $p < 0.05$ for all comparisons.

Results

Subject Characteristics

Subjects' characteristics at rest are shown in Table 1. There were no significant differences in height, weight, heart rate, or carotid BP between young and middle-aged men in the present study. Brachial diastolic and mean BP and carotid diastolic diameter were higher in the middle-aged men than in the young group ($p < 0.05$). Brachial systolic BP and carotid pulse pressure were not significantly different between the two groups. β -Stiffness, augmentation index, and baPWV in the middle-aged men were significantly higher than those in the young group ($p < 0.05$). The carotid arterial compliance

and leg press maximum of the middle-aged men were significantly lower than those of the young men ($p < 0.05$).

Exercise Protocol I

Radial systolic and diastolic BP responses during the exercise and recovery periods are shown in Fig. 1. Baseline values of radial diastolic BP under the 40% and 60% 1RM conditions were higher in middle-aged than in young men, and there were no significant differences between groups in other baseline BP values. Systolic BP for 20–40 s during exercise in the middle-aged men at only 80% 1RM was significantly lower than that in the young group ($p < 0.05$). There were no significant differences in heart rate response to resistance exercise between middle-aged and young men at any intensity examined (Fig. 2). Systolic and diastolic BP returned to the baseline values within 60 s during the recovery period. The amounts of change in systolic and diastolic BP and pulse pressure from baseline to the peak response during resistance exercise were significantly lower in the middle-aged men than in the young group at all exercise intensities examined, but there was no significant difference in Δ heart rate (Fig. 3).

Exercise Protocol II

The amounts of change in systolic and diastolic BP and pulse pressure from baseline to the peak response during resistance




Fig. 4. The amounts of change in systolic (SBP; top left) and diastolic (DBP; bottom left) blood pressure, pulse pressure (PP; top right) and heart rate (HR; bottom right) responses to resistance exercise at 145 kgw in middle-aged (black bar) and young (white bar) men. Values are means ± SEM. * $p < 0.05$ vs. young men.

exercise were lower in the middle-aged men than in the young men at individual absolute intensity, but there was no significant difference in Δ heart rate (systole, $p=0.019$; diastole, $p=0.091$; pulse pressure, $p=0.003$; Fig. 4).

Discussion

The major findings of the present study were as follows. 1) The absolute value of BP response to dynamic resistance exercise at 80% 1RM was lower in middle-aged men with stiffening arteries than in young men with compliant arteries. 2) At all relative intensities, the amounts of change in peak BP response to resistance exercise were lower in middle-aged than in young men. 3) At individual absolute intensity, the amounts of change in peak systolic BP response to resistance exercise were lower in middle-aged than in young men. In contrast to our hypothesis, these results suggest that BP responses during dynamic resistance exercise may be attenuated with advancing age at either individual relative or absolute intensity, despite age-related stiffening of the arteries.

Previous studies have suggested that the arteries of middle-aged men appear to be stiffer than those of young men based on measurements of carotid arterial compliance, β -stiffness, augmentation index, and baPWV (5, 6, 22, 23, 30). Our results also showed that arterial stiffening develops at a greater rate in middle-aged men than in young men. However, remarkable hypertrophy (> 1.1 mm) of IMT, which is a char-

acteristic of atherosclerosis, was not observed in any of the subjects in the present study. The maximal muscle strength estimated by the leg press exercise was lower in middle-aged men than in young men. These results indicate that the middle-aged men in the present study had developed arteriosclerosis without atherosclerosis and had lower maximal muscle strength than the young men. Although all BP values at rest in middle-aged men were higher than those in young men, all subjects were normotensive (< 140/90 mmHg). Carotid diastolic diameter in middle-aged men was higher than that in young men. This result was consistent with the results of a previous epidemiological study (31). This alteration may be a physiological adaptation to suppress marked reductions in arterial compliance and vessel diameter induced by hypertrophied IMT with advancing age.

Dynamic resistance exercise is mainly used for health promotion and strength conditioning as it has greater effects on strength and volume of skeletal muscle in comparison with static (isometric) exercise (32). Understanding the pressor response during dynamic resistance exercise using large muscle groups is essential for exercise prescription. However, most previous studies have focused on static resistance exercise (11, 12, 16, 33), and have provided little information regarding the cardiovascular response during dynamic resistance exercise (10), or the interaction between age and BP response to dynamic resistance exercise using large muscle groups. We found that pressor responses during dynamic

resistance exercise at relative intensities were lower in middle-aged than in young men, suggesting that the BP response to dynamic resistance exercise may be attenuated with advancing age despite age-associated arterial stiffening.

From the relative intensities, it is reasonable to hypothesize that the attenuated BP response to resistance exercise in middle-aged men may be induced by the age-related reduction in maximal muscular strength, because of the exercise intensity-associated increase in BP response to resistance exercise in both groups. In the present study, the IRM estimated by the leg press exercise was lower in middle-aged than in young men, suggesting that the absolute intensities during exercise at individual relative intensities were lower in the former than in the latter. Accordingly, we determined BP response during the dynamic leg press exercise at individual absolute intensity (145 kgw) in the middle-aged and young men. The results indicated that the amount of change in BP response to resistance exercise was lower in middle-aged than in young men. These results suggest that age-associated reduction in muscle strength did not contribute to the attenuated pressor response to dynamic resistance exercise in middle-aged as compared with young men.

It is unclear what physiological mechanisms explain the attenuated BP responses during dynamic resistance exercise using large muscle groups in middle-aged as compared with young men. However, we speculate that the mechanism may be as follows. In middle-aged men, the muscle sympathetic nerve activity is higher at rest than in young men (16), whereas during exercise it is lower in the former than the latter (16). This results in attenuation of the increases in cardiac output and peripheral vasoconstriction induced by sympathoexcitation during exercise with advancing age (34–36). The ratio of high-glycolytic muscle fiber type II in skeletal muscle falls from 59% to 48% between the third and sixth decades of life (37), and the transformation to oxidative skeletal muscle fibers results in a lower pressor response evoked by static contraction as compared with glycolytic fibers (38). Therefore, alterations in sympathetic nerve activity and/or skeletal muscle fiber type with advancing age may contribute to the attenuated BP response to resistance exercise in middle-aged as compared with young men.

Sarcopenia and osteoporosis with advancing age are social problems in developed countries with aging populations. The leg press exercise used in this study, as a form of dynamic resistance exercise using predominantly the lower body, is widely accepted in exercise prescription for the prevention and rehabilitation of sarcopenia and osteoporosis, which can lead to falls and femur bone fracture, and may even result in patients becoming bedridden. However, BP rises rapidly and remarkably during high-intensity leg press exercise (10). Indeed, it has been reported the accidents, such as artery dissection and subarachnoid hemorrhage, occur during resistance exercise (39–42). Therefore, care should be taken regarding the rapid and marked increases in BP response to resistance exercise, particularly in middle-aged and older

men. In contrast to our expectations, the results of the present study indicated that pressor responses during dynamic resistance exercise at individual relative and absolute intensities were not higher in middle-aged men with stiffening arteries than in young men with compliant arteries. These results may contribute to our understanding of the cardiovascular responses to resistance exercise at appropriate intensities recommended by the major health organizations in middle-aged men who have developed arterial stiffening.

As it is the simplest parameter of arterial buffering function, pulse pressure was evaluated along with systolic and diastolic BP responses to resistance exercise in the present study. The results indicated that the amounts of change in pulse pressure response to resistance exercise at either relative or absolute intensities were lower in middle-aged men than in young men despite age-related increases in arterial stiffness. The attenuation of pulse pressure response to resistance exercise with advancing age may be affected by systolic function in the left ventricle. Of course, this function is greater in young than in middle-aged men during exercise as well as at rest. Thus, lower pulse pressure response to resistance exercise in middle-aged men may be appropriate. Pulse pressure at rest was also lower in middle-aged than in young men. As pulse pressure at rest increases progressively in normotensive subjects from the fifth decade (43), further studies are needed to determine BP response to resistance exercise in older men with augmented pulse pressure.

The present study had several limitations. Although there have been several reports on BP responses to isometric or aerobic exercise, we did not attempt to compare the BP responses to isometric resistance or aerobic exercise with those to dynamic resistance exercise. Compared to isometric or aerobic exercise, dynamic resistance exercise is more often used for health promotion, strength conditioning and prevention of sarcopenia or osteoporosis in middle-aged and older individuals. Therefore, as a primary approach, it was necessary to clarify the differences in BP response to dynamic resistance training using large muscle groups between young and middle-aged men. Although increases in central arterial BP during exercise may be more important than those in peripheral arterial BP from the standpoint of cardioprotection, we performed noninvasive assessment of only the radial arterial BP response to resistance exercise. Therefore, the results of the present study must be confirmed in future prospective studies focusing on central arterial BP responses to resistance exercise. Finally, the muscular strength maximum was evaluated with a leg press machine using air pressure. The value of muscular strength assessed by this machine may be different from that of muscular strength evaluated using real weights. Although the muscular strength maximum of subjects in the present study was relatively high, our results may not have been affected by this difference.

In conclusion, this study demonstrated that, at either individual relative or absolute intensity, the BP response during dynamic resistance exercise using large muscle groups was

attenuated in middle-aged men as compared with young men despite age-related stiffening of the arteries. These findings may contribute to our understanding of the BP response during dynamic resistance exercise and aid in the safe performance of exercise prescription for prevention and rehabilitation of sarcopenia and osteoporosis in middle-aged and older men.

Acknowledgements

We thank Dr. Kenta Yamamoto and Dr. Izumi Tabata for their advice.

References

- Pate RR, Davis MG, Robinson TN, *et al*: Promoting physical activity in children and youth: a leadership role for schools: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism (Physical Activity Committee) in collaboration with the Councils on Cardiovascular Disease in the Young and Cardiovascular Nursing. *Circulation* 2006; **114**: 1214–1224.
- Thompson PD, Buchner D, Pina IL, *et al*: Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). *Circulation* 2003; **107**: 3109–3116.
- American College of Sports Medicine Position Stand: Exercise and physical activity for older adults. *Med Sci Sports Exerc* 1998; **30**: 992–1008.
- Gordon NF, Scott CB, Wilkinson WJ, Duncan JJ, Blair SN: Exercise and mild essential hypertension. Recommendations for adults. *Sports Med* 1990; **10**: 390–404.
- Tanaka H, Dinenna FA, Monahan KD, *et al*: Aging, habitual exercise, and dynamic arterial compliance. *Circulation* 2000; **102**: 1270–1275.
- Tanaka H, DeSouza CA, Seals DR: Absence of age-related increase in central arterial stiffness in physically active women. *Arterioscler Thromb Vasc Biol* 1998; **18**: 127–132.
- Higashi Y, Sasaki S, Kurisu S, *et al*: Regular aerobic exercise augments endothelium-dependent vascular relaxation in normotensive as well as hypertensive subjects: role of endothelium-derived nitric oxide. *Circulation* 1999; **100**: 1194–1202.
- DeSouza CA, Shapiro LF, Clevenger CM, *et al*: Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 2000; **102**: 1351–1357.
- Williams MA, Haskell WL, Ades PA, *et al*: Resistance exercise in individuals with and without cardiovascular disease: 2007 update: a scientific statement from the American Heart Association Council on Clinical Cardiology and Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2007; **116**: 572–584.
- MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR: Arterial blood pressure response to heavy resistance exercise. *J Appl Physiol* 1985; **58**: 785–790.
- Petrofsky JS, Lind AR: Aging, isometric strength and endurance, and cardiovascular responses to static effort. *J Appl Physiol* 1975; **38**: 91–95.
- Smolander J, Aminoff T, Korhonen I, *et al*: Heart rate and blood pressure responses to isometric exercise in young and older men. *Eur J Appl Physiol Occup Physiol* 1998; **77**: 439–444.
- Fleck SJ, Dean LS: Resistance-training experience and the pressor response during resistance exercise. *J Appl Physiol* 1987; **63**: 116–120.
- Lentini AC, McKelvie RS, McCartney N, Tomlinson CW, MacDougall JD: Left ventricular response in healthy young men during heavy-intensity weight-lifting exercise. *J Appl Physiol* 1993; **75**: 2703–2710.
- Narloch JA, Brandstater ME: Influence of breathing technique on arterial blood pressure during heavy weight lifting. *Arch Phys Med Rehabil* 1995; **76**: 457–462.
- Markel TA, Daley JC 3rd, Hogeman CS, *et al*: Aging and the exercise pressor reflex in humans. *Circulation* 2003; **107**: 675–678.
- Arnett DK, Evans GW, Riley WA: Arterial stiffness: a new cardiovascular risk factor? *Am J Epidemiol* 1994; **140**: 669–682.
- Avolio AP, Deng FQ, Li WQ, *et al*: Effects of aging on arterial distensibility in populations with high and low prevalence of hypertension: comparison between urban and rural communities in China. *Circulation* 1985; **71**: 202–210.
- Nichols WW, O'Rourke MF, Hartley C, McDonald DA: McDonald's Blood Flow in Arteries. London, Edward Arnold, 1998, 564 pp.
- O'Rourke M: Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; **15**: 339–347.
- Iemitsu M, Maeda S, Otsuki T, *et al*: Polymorphism in endothelin-related genes limits exercise-induced decreases in arterial stiffness in older subjects. *Hypertension* 2006; **47**: 928–936.
- Sugawara J, Hayashi K, Yokoi T, *et al*: Brachial-ankle pulse wave velocity: an index of central arterial stiffness? *J Hum Hypertens* 2005; **19**: 401–406.
- Miyachi M, Donato AJ, Yamamoto K, *et al*: Greater age-related reductions in central arterial compliance in resistance-trained men. *Hypertension* 2003; **41**: 130–135.
- Kawano H, Tanaka H, Miyachi M: Resistance training and arterial compliance: keeping the benefits while minimizing the stiffening. *J Hypertens* 2006; **24**: 1753–1759.
- Miyachi M, Kawano H, Sugawara J, *et al*: Unfavorable effects of resistance training on central arterial compliance: a randomized intervention study. *Circulation* 2004; **110**: 2858–2863.
- Kelly R, Hayward C, Avolio A, O'Rourke M: Noninvasive determination of age-related changes in the human arterial pulse. *Circulation* 1989; **80**: 1652–1659.
- Van Merode T, Hick PJ, Hoeks AP, Rahn KH, Reneman RS: Carotid artery wall properties in normotensive and borderline hypertensive subjects of various ages. *Ultrasound Med Biol* 1988; **14**: 563–569.
- Hirai T, Sasayama S, Kawasaki T, Yagi S: Stiffness of systemic arteries in patients with myocardial infarction. A

- noninvasive method to predict severity of coronary atherosclerosis. *Circulation* 1989; **80**: 78–86.
29. Sato T, Nishinaga M, Kawamoto A, Ozawa T, Takatsuji H: Accuracy of a continuous blood pressure monitor based on arterial tonometry. *Hypertension* 1993; **21**: 866–874.
 30. Ronnback M, Fagerudd J, Forsblom C, et al: Altered age-related blood pressure pattern in type 1 diabetes. *Circulation* 2004; **110**: 1076–1082.
 31. Najjar SS, Scuteri A, Lakatta EG: Arterial aging: is it an immutable cardiovascular risk factor? *Hypertension* 2005; **46**: 454–462.
 32. Wernbom M, Augustsson J, Thomee R: The influence of frequency, intensity, volume and mode of strength training on whole muscle cross-sectional area in humans. *Sports Med* 2007; **37**: 225–264.
 33. Boutcher SH, Stocker D: Cardiovascular responses to light isometric and aerobic exercise in 21- and 59-year-old males. *Eur J Appl Physiol Occup Physiol* 1999; **80**: 220–226.
 34. Correia LC, Lakatta EG, O'Connor FC, et al: Attenuated cardiovascular reserve during prolonged submaximal cycle exercise in healthy older subjects. *J Am Coll Cardiol* 2002; **40**: 1290–1297.
 35. Fleg JL, O'Connor F, Gerstenblith G, et al: Impact of age on the cardiovascular response to dynamic upright exercise in healthy men and women. *J Appl Physiol* 1995; **78**: 890–900.
 36. Davy KP, Seals DR, Tanaka H: Augmented cardiopulmonary and integrative sympathetic baroreflexes but attenuated peripheral vasoconstriction with age. *Hypertension* 1998; **32**: 298–304.
 37. Larsson L, Grimby G, Karlsson J: Muscle strength and speed of movement in relation to age and muscle morphology. *J Appl Physiol* 1979; **46**: 451–456.
 38. Wilson LB, Dyke CK, Parsons D, et al: Effect of skeletal muscle fiber type on the pressor response evoked by static contraction in rabbits. *J Appl Physiol* 1995; **79**: 1744–1752.
 39. Elefteriades JA, Hatzaras I, Tranquilli MA, et al: Weight lifting and rupture of silent aortic aneurysms. *JAMA* 2003; **290**: 2803.
 40. Hatzaras I, Tranquilli M, Coady M, et al: Weight lifting and aortic dissection: more evidence for a connection. *Cardiology* 2006; **107**: 103–106.
 41. Haykowsky MJ, Findlay JM, Ignaszewski AP: Aneurysmal subarachnoid hemorrhage associated with weight training: three case reports. *Clin J Sport Med* 1996; **6**: 52–55.
 42. Hogan CJ: An aortic dissection in a young weightlifter with non-Marfan fibrillinopathy. *Emerg Med J* 2005; **22**: 304–305.
 43. Franklin SS, Gustin Wt, Wong ND, et al: Hemodynamic patterns of age-related changes in blood pressure. The Framingham Heart Study. *Circulation* 1997; **96**: 308–315.

体重変化量とウエスト囲変化量との 関連～8年間の変化～

松本 純子, 宮武 伸行, 丸山 弘子, 沼田 健之

岡山県南部健康づくりセンター

Summary

Objectives: We investigated the correlation between changes in body weight and body composition parameters.

Methods: We used the data of 165 Japanese (45.2±11.1 years) at baseline and at 8 year follow-up from a database of 2151 subjects, which is available at the Okayama Southern Institute of Health in Okayama prefecture, Japan. Body weight, waist circumference at the umbilical level, hip circumference, and body fat percentage were used in the analyses.

Results: Body fat percentage was significantly reduced after 8 year. Changes in body weight significantly correlated with changes in waist circumference, changes in hip circumference, and changes in body fat percentage. A decrease in body weight of 1 kg corresponded to a 0.972 cm decrease in waist circumference in men and a 1.069 cm decrease in that in women.

Conclusion: A decrease in body weight of 1 kg corresponded to an almost 1 cm decrease in waist circumference at the umbilical level in Japanese men and women.

Key words

ウエスト囲, 体重, メタボリックシンドローム

はじめに

メタボリックシンドロームは、耐糖能異常や高血圧、脂質代謝異常、肥満という複数の危険因子の重複により、将来心血管疾患の発症に関与するとして注目されている。そして、心血管疾患は危険因子単独の場合より重複することで飛躍的に多くなる¹⁾²⁾。したがって、メタボリックシンドロームを効果的に予防、改善することが重要である。

日本肥満学会 (<http://wwwsoc.nii.ac.jp/jasos/>, accessed on July 13, 2007) では神戸宣言

2006として、メタボリックシンドロームの予防、改善のために、食生活の改善と運動の増加を図り、まず3 kgの減量と3 cmのウエスト囲の減少を実現するサンサン運動を提案している。また、平成18年に厚生労働省から発表された運動指針(エクササイズガイド2006)の中でも、体重1 kgの変化はウエスト囲1 cmの変化に相当すると記載されている³⁾。しかしながら、体重変化量とウエスト囲変化量との関連についてはいまだ明らかでない。

今回私たちは、岡山県南部健康づくりセンター利用者を対象とした8年間の体重変化量とウエスト囲変化量との関連を検討した。

松本 純子

岡山県健康づくり財団岡山県南部健康づくりセンター
岡山市平田408-1
center@okakenko.jp

■対象および方法

対象は、当センターで平成9年度にヘルスチェック（健康度測定）を受けた男性717名（43.8±13.8歳）、女性1434名（43.1±13.6歳）、計2151名のうち、8年後の平成17年度にヘルスチェックを受けた男性51名（41.3±11.5歳）、女性114名（46.9±10.5歳）、計165名であった（表1）。

測定項目は、身長、体重、BMI（Body mass index）、ウエスト囲、ヒップ囲、体脂肪率で、ウエスト囲は両足をそろえて自然に立たせ呼気時臍部で測定し、体脂肪率は空気置換法（BOD POD: Body Composition System, Life Measurement Instruments, Concord, USA）を用いて⁴⁾⁵⁾測定した。

結果はすべて平均値±標準偏差で表し、有意差検定は対応のあるt検定を、体重と各指標との関係はピアソンの相関係数を用い、5%未満

を有意とした。

なお、本検討に関しては、岡山県健康づくり財団倫理委員会の承認を得た。

■結 果

平成17年度のヘルスチェックでは、平成9年度に比較して、男性で体脂肪率が有意に減少し、女性では、体重、BMI、体脂肪率が有意に減少したが、その他の項目では有意な変化は認められなかった（表1）。

体重の変化量とウエスト囲、ヒップ囲、体脂肪率の変化量との間には、男女ともそれぞれ有意な関連が認められた（表2）。回帰式は、男性で、ウエスト囲変化量=0.972×体重変化量-0.169、女性で、ウエスト囲変化量=1.069×体重変化量+1.151となり、体重変化量1kgはウエスト囲変化量1cmにほぼ相当した（表2、図）。

表1 平成9年度と平成17年度での身体組成の変化

	平成9年度	平成17年度	
全 体			
症 例 数	165		
年 齢	45.2 ± 11.1		
身 長 (cm)	159.3 ± 8.6		
体 重 (kg)	58.7 ± 10.5	58.0 ± 10.6	0.0850
BMI (kg / m ²)	23.0 ± 3.0	22.8 ± 3.0	0.1112
ウエスト囲 (cm)	75.2 ± 9.1	75.3 ± 9.1	0.9147
ヒップ囲 (cm)	91.1 ± 5.0	90.8 ± 5.2	0.2076
体脂肪率 (%)	29.6 ± 6.6	27.7 ± 7.4	<0.0001
男 性			
症 例 数	51		
年 齢	41.3 ± 11.5		
身 長 (cm)	169.1 ± 5.1		
体 重 (kg)	68.6 ± 9.0	68.7 ± 8.9	0.9556
BMI (kg / m ²)	24.0 ± 2.9	24.0 ± 3.0	0.9467
ウエスト囲 (cm)	82.6 ± 7.8	82.5 ± 8.0	0.9019
ヒップ囲 (cm)	93.2 ± 4.5	93.0 ± 4.8	0.7240
体脂肪率 (%)	24.3 ± 5.4	22.4 ± 6.6	0.0212
女 性			
症 例 数	114		
年 齢	46.9 ± 10.5		
身 長 (cm)	155.0 ± 5.7		
体 重 (kg)	54.2 ± 7.7	53.2 ± 7.4	0.0084
BMI (kg / m ²)	22.6 ± 2.9	22.2 ± 2.9	0.0246
ウエスト囲 (cm)	71.9 ± 7.6	72.0 ± 7.6	0.7901
ヒップ囲 (cm)	90.2 ± 5.0	89.8 ± 5.1	0.1830
体脂肪率 (%)	32.0 ± 5.7	30.0 ± 6.5	<0.0001

平均値±標準偏差

表2 体重変化量との相関

		r	回帰式
全 体	ウエスト囲変化量 (cm)	0.877	$y=1.002x+0.699$
	ヒップ囲変化量 (cm)	0.844	$y=0.579x+0.051$
	体脂肪率変化量 (%)	0.665	$y=0.748x-1.766$
男 性	ウエスト囲変化量 (cm)	0.917	$y=0.972x-0.169$
	ヒップ囲変化量 (cm)	0.908	$y=0.528x-0.215$
	体脂肪率変化量 (%)	0.692	$y=0.746x-2.824$
女 性	ウエスト囲変化量 (cm)	0.853	$y=1.069x+1.151$
	ヒップ囲変化量 (cm)	0.809	$y=0.652x+0.241$
	体脂肪率変化量 (%)	0.660	$y=0.783x-1.258$




図 体重変化量とウエスト囲変化量との関連

考 察

現在わが国では、メタボリックシンドロームをはじめとした生活習慣病の適切な予防、改善法に注目が集まっている。以前、当センターにおけるメタボリックシンドロームの割合を検討したが、男性30.7%、女性3.6%であった⁹⁾。

さらに、平成20年度からはメタボリックシンドロームの考えをとり入れた新しい健診、保健指導もスタートし、ウエスト囲の計測が必須となる。ウエスト囲は腹部肥満の簡便な指標であ

り、生活習慣病予防の観点からウエスト囲を減少させることは意義深い。

日本肥満学会では、肥満症、メタボリックシンドローム予防、改善のため、まずは体重3 kg、ウエスト囲3 cmの減少を提案しており、また、エクササイズガイド2006の中でも、体重1 kg (7,000kcal) はウエスト囲1 cmに相当すると記載されている。しかし、日本人における体重変化量とウエスト囲変化量に関する検討はわずかである。

以前、私たちが行なった男性肥満者に対する

1年間の運動プログラム（1回90分、週1回で食事のアドバイスは特に行わず、運動中心の介入）を用いた検討では、体重3.3kgの減少が、ウエスト囲4.2cmの減少に相当した⁷⁾。また、ヘルスチェック受診者のうち、1年間隔で2回受診し、生活習慣病などで薬物療法を受けていない男性856名、女性1779名で、体重変化量とウエスト囲変化量との関連についての検討では、回帰式の傾きは男性1.069、女性0.950となり体重変化量1kgはウエスト囲変化量1cmにほぼ相当した⁸⁾。今回の結果もほぼ同様の結果であったことから、8年間にもおよぶ長期にわたっての体重変化量とウエスト囲変化量との関連についても体重変化量1kgはウエスト囲変化量1cmにほぼ相当することがわかった。

今回の検討での問題点は、まず、体重変化量とウエスト囲変化量との関連は検討できたが、メタボリックシンドロームの予防、改善のための体重減少量とウエスト囲減少量とのカットオフ値を検討することはできなかった。また、内臓脂肪面積の測定を直接行なっておらず、内臓脂肪蓄積と身体組成との関連も検討することができなかったため、今後は腹部CTを用いた内臓脂肪面積の測定を行なう必要があると思われる。

以上、当センター利用者を対象とした8年間という比較的長期の体重変化量とウエスト囲変化量との関連を検討した結果、体重変化量1kgは、ウエスト囲変化量1cmにほぼ相当した。今後は、メタボリックシンドロームの予防、改善のために有効なプログラムを計画し、さらなる介入研究が必要であると思われる。

まとめ

当センター利用者を対象とした8年間という比較的長期の体重変化量とウエスト囲変化量との関連を検討した結果、体重変化量1kgは、ウエスト囲変化量1cmにほぼ相当した。

本研究の内容の一部は第38回岡山県医学検査学会で発表予定。

参考文献

- 1) 山田信博: 1. 歴史と疾患概念, I. 疾患概念, *Metabolic syndrome 診断と治療の進歩*, 日本内科学会雑誌2004; 93: 635-641
- 2) 後藤田貴也: メタボリックシンドロームの概念と動脈硬化, *新薬と治療*2004; 54: 29-31
- 3) 運動所要量・運動指針の策定検討会: 健康づくりのための運動指針2006～生活習慣病予防のために～, 平成18年7月
- 4) Miyatake, N. et al.: A new air displacement plethysmograph for the determination of Japanese body composition, *Diabetes Obes Metab* 1999; 1: 347-351
- 5) Miyatake, N. et al.: Comparison of air displacement plethysmograph and bioelectrical impedance for assessing body composition changes during weight loss in Japanese women, *Diabetes Obes Metab* 2005; 7: 268-272
- 6) Miyatake, N. et al.: Prevalence of metabolic syndrome in Okayama prefecture, Japan. *Intern Med* 2006; 45: 107-108
- 7) Miyatake, N. et al.: Changes in serum leptin concentrations in overweight Japanese men after exercise, *Diabetes Obes Metab* 2004; 6: 332-337
- 8) Miyatake, N. et al.: Relationship between changes in body weight and waist circumference in Japanese, *Environ Health Prev Med* 2007; 12: 220-223



Brief report

Comparison of serum uric acid levels between Japanese with and without metabolic syndrome

Takeyuki Numata^a, Nobuyuki Miyatake^{a,*}, Jun Wada^b, Hirofumi Makino^b

^a Okayama Southern Institute of Health, Okayama 700-0952, Japan

^b Department of Medicine and Clinical Science, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan

ARTICLE INFO

Article history:

Accepted 1 October 2007

Published on line 7 March 2008

Keywords:

Metabolic syndrome

Uric acid

Abdominal obesity

ABSTRACT

Objective: The link between uric acid levels and metabolic syndrome was investigated.

Methods: We used data for 997 men and 1290 women. Metabolic syndrome was defined by a new criterion in Japan and uric acid was measured.

Results: Subjects who did not drink alcohol were evaluated. Uric acid was significantly higher in subjects with metabolic syndrome compared with subjects without the syndrome in both sexes. The presence of abdominal obesity and dyslipidemia in men and abdominal obesity in women revealed positive impacts on uric acid levels.

Conclusion: High serum uric acid may be often associated with metabolic syndrome in Japanese people, which accelerates the development of cardiovascular disease from metabolic syndrome.

© 2007 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

We previously reported that 30.7% of men and 3.6% of women were diagnosed as having metabolic syndrome [1]. Metabolic syndrome is closely linked to an increased risk for cardiovascular disease [2], proteinuria [3] and the elevation of hepatic enzymes [4]. Therefore, the prevention and improvement of metabolic syndrome is urgently required. Serum uric acid levels are said to be an independent predictor of cardiovascular disease [5]. However, the link between metabolic syndrome using the new criterion in Japan and uric acid remains to be investigated. In this study, we compared uric

acid between Japanese people with and without metabolic syndrome.

2. Subjects and methods

2.1. Subjects

Using retrospective data from a data base of 14,345 subjects we extracted information from 997 men and 1290 women, aged 20–79 years, who met the following criteria (1) received annual health checkups from June 1997 to May 2006 at Okayama

* Corresponding author at: Okayama Southern Institute of Health, 408-1, Hirata, Okayama 700-0952, Japan. Tel.: +81 862 46 6250; fax: +81 862 46 6330.

E-mail address: center@okakenko.jp (N. Miyatake).

0168-8227/\$ – see front matter © 2007 Elsevier Ireland Ltd. All rights reserved.

doi:10.1016/j.diabres.2007.10.031

Southern Institute of Health, (2) received fasting blood examination, and (3) obtained written informed consent.

2.2. Serum uric acid measurements

Serum uric acid levels were measured by the Uricase-Peroxidase method. The institutional normal range was 2.5–7.0 mg/dl.

2.3. Anthropometric measurements

Anthropometric parameters i.e. height, weight and waist circumference were measured. The waist circumference was measured at the umbilical level.

2.4. Definition of metabolic syndrome

Metabolic syndrome was defined according to the consensus definition of eight Medical Societies in Japan [6]. Men with a waist circumference in excess of 85 cm and women with a waist circumference in excess of 90 cm were diagnosed with metabolic syndrome if they had two or more of the following components: (1) dyslipidemia: triglycerides \geq 150 mg/dl and/or HDL cholesterol $<$ 40 mg/dl; (2) high blood pressure: blood pressure \geq 130/85 mmHg; (3) impaired glucose tolerance: fasting plasma glucose \geq 110 mg/dl.

2.5. Alcohol consumption habits

It is well known that alcohol consumption is closely linked to serum uric acid levels. Data on alcohol consumption habits were obtained at interviews by well-trained staff in a structured way. The subjects were asked if they currently drank alcohol on a regular basis. Subjects answering "yes" were classified as subjects with a drinking habit. Those answering "no" were further asked whether they had ever drunk alcohol on a regular basis. Those answering "no" were classified as subjects without a drinking habit and those answering "yes" were classified as subjects with a previous drinking habit.

2.6. Statistical analysis

Data are expressed as mean \pm standard deviation (S.D.) values. A comparison of parameters between two groups was carried out using χ^2 test, unpaired t test and covariance analysis. A comparison of parameters among four groups was performed by ANOVA and Scheffe's F test: $p < 0.05$ was considered to be statistically significant.

3. Results

A total of 280 men (28.1%) and 46 women (3.6%) were diagnosed as having metabolic syndrome. The results of various parameters in subjects with and without the syndrome are indicated in Table 1. Age was significantly higher in subjects with metabolic syndrome in both sexes and thus, to avoid the influence of age on serum uric acid levels, we compared uric acid by using covariance analysis. In subjects

with metabolic syndrome, uric acid was significantly higher compared with subjects without the syndrome.

We then analyzed the groups with and without each component of metabolic syndrome. Age was significantly higher in subjects with abdominal obesity, impaired glucose tolerance, dyslipidemia, and high blood pressure compared to subjects in both sexes without each component. By comparing uric acid adjusting for age, uric acid was significantly higher in subjects with abdominal obesity and dyslipidemia in men, and also in women with abdominal obesity.

To further evaluate the effect of abdominal obesity and dyslipidemia in men and abdominal obesity in women on uric acid, from enrolled subjects, we used data for 504 men and 624 women. We compared serum uric acid levels by classifying subjects into groups according to a number of different risk factors (Group A: waist circumference (–), dyslipidemia (–), impaired glucose tolerance and/or hypertension (–), Group B: waist circumference (+), dyslipidemia (–), impaired glucose tolerance and/or hypertension (–), Group C: waist circumference (+), dyslipidemia (+), impaired glucose tolerance and/or hypertension (–), Group D: waist circumference (+), dyslipidemia (+), impaired glucose tolerance and/or hypertension (+), (Table 2)). In men, uric acid in subjects in Group C and D was significantly higher than that in subjects in Group A. Uric acid in subjects in Group C was also significantly higher than that in subjects in Group B. Although there was no significant difference between Group A and Group B, uric acid in subjects in Group B was higher than that in subjects in Group A (Group A: 5.7 ± 1.0 mg/dl vs. Group B: 6.1 ± 1.3 mg/dl). In women, uric acid in subjects in Group B and D was significantly higher than that in subjects in Group A.

Although there were no significant differences in drinking habits between the two groups, in order to avoid any influence of alcohol consumption, we performed a separate analysis using only data from subjects with no drinking habit and compared uric acid levels between subjects with and without the syndrome (Table 1). Age was similar between the two male groups. An unpaired t test showed that, uric acid in subjects with metabolic syndrome was significantly higher than that in subjects without. In women, an age difference was noted, and uric acid in subjects with metabolic syndrome was also significantly higher even after adjusting for age by covariance analysis.

4. Discussion

We compared serum uric acid levels in subjects with and without metabolic syndrome using the new criterion in Japan.

Some studies have shown that higher serum uric acid levels are often accompanied by obesity, hypertension [7], dyslipidemia [8], and glucose intolerance [9], all of which play a causal role in the pathogenesis of cardiovascular disease. However, few studies have reported on the association between uric acid and metabolic syndrome using the new criterion in Japan. Kawamoto et al. investigated the relationship between serum uric acid concentrations and metabolic syndrome defined by the modified Japanese criteria using ultrasound image analysis, reporting that the prevalence of metabolic syndrome was significantly increased according to

Table 1 - Comparison of parameters between subjects with and without metabolic syndrome

Men	Metabolic syndrome (+)	Metabolic syndrome (-)	p	p
Number of subjects	280	717	Unpaired t test	Adjusting for age
Age	48.7 ± 10.4	45.4 ± 11.9	<0.0001	
Uric acid (mg/dl)	6.3 ± 1.5	6.0 ± 1.3		0.0133
Men	Waist circumference (+)	Waist circumference (-)	p	p
Number of subjects	508	489		
Age	47.2 ± 10.6	45.5 ± 12.5	<0.0001	
Uric acid (mg/dl)	6.3 ± 1.4	5.8 ± 1.2		0.001
Men	Impaired glucose tolerance (+)	Impaired glucose tolerance (-)	p	p
Number of subjects	249	748		
Age	50.8 ± 10.4	44.8 ± 11.6	<0.0001	
Uric acid (mg/dl)	5.8 ± 1.4	6.2 ± 1.3		0.086
Men	Dyslipidemia (+)	Dyslipidemia (-)	p	p
Number of subjects	485	512		
Age	47.2 ± 11.1	45.5 ± 12.0	0.0249	
Uric acid (mg/dl)	6.3 ± 1.4	5.9 ± 1.3		0.0266
Men	High blood pressure (+)	High blood pressure (-)	p	p
Number of subjects	585	412		
Age	48.3 ± 11.2	43.6 ± 11.6	<0.0001	
Uric acid (mg/dl)	6.2 ± 1.4	5.9 ± 1.3		0.6495
Men	Metabolic syndrome (+)	Metabolic syndrome (-)	p	p
Subjects without drinking habits				
Number of subjects	62	168		
Age	47.0 ± 12.1	45.4 ± 12.7	0.4008	
Uric acid (mg/dl)	6.3 ± 1.4	5.6 ± 1.3	0.0005	
Women	Metabolic syndrome (+)	Metabolic syndrome (-)	p	p
Number of subjects	46	1244		
Age	54.8 ± 10.1	46.9 ± 12.1	<0.0001	
Uric acid (mg/dl)	5.4 ± 1.1	4.4 ± 1.1		0.0412
Women	Waist circumference (+)	Waist circumference (-)	p	p
Number of subjects	89	1201		
Age	51.8 ± 10.8	46.9 ± 12.1	0.0002	
Uric acid (mg/dl)	5.1 ± 1.1	4.3 ± 1.0		0.0113
Women	Impaired glucose tolerance (+)	Impaired glucose tolerance (-)	p	p
Number of subjects	137	1153		
Age	55.6 ± 10.1	46.2 ± 11.9	<0.0001	
Uric acid (mg/dl)	4.7 ± 1.6	4.3 ± 1.0		0.166
Women	Dyslipidemia (+)	Dyslipidemia (-)	p	p
Number of subjects	402	888		
Age	51.5 ± 12.0	45.3 ± 11.6	<0.0001	
Uric acid (mg/dl)	4.6 ± 1.2	4.3 ± 1.0		0.6466
Women	High blood pressure (+)	High blood pressure (-)	p	p
Number of subjects	488	802		
Age	53.3 ± 10.6	43.5 ± 11.5	<0.0001	
Uric acid (mg/dl)	4.7 ± 1.2	4.2 ± 1.0		0.3705
Women	Metabolic syndrome (+)	Metabolic syndrome (-)	p	p
Subjects without drinking habits				
Number of subjects	33	799		
Age	55.9 ± 10.7	48.0 ± 12.1	0.0002	
Uric acid (mg/dl)	5.3 ± 1.2	4.4 ± 1.1		0.0369

The values are expressed as (mean ± S.D.)

Table 2 – Comparison of serum uric acid levels as classified into accumulation of risk factors

	Waist circumference	Dyslipidemia	Impaired glucose tolerance and/or high blood pressure	Number of subjects	Uric acid (mg/dl)	p
Men						
A				140	5.7 ± 1.0	
B				60	6.1 ± 1.3	
C				55	6.8 ± 1.2	ab
D				242	6.3 ± 1.5	a
Women						
A				562	4.2 ± 1.0	
B				11	5.4 ± 1.3	a
C				10	4.4 ± 0.9	
D				35	5.3 ± 1.1	a

a, $p < 0.01$ vs. Group A, b, $p < 0.05$ vs. Group B.

serum uric acid levels in women [10]. Ishizaka et al. also reported that the prevalence of metabolic syndrome showed a graded increase according to serum uric acid values in both sexes using the National Cholesterol Education Program (NCEP) Adult Treatment Panel III (ATPIII) [11]. In this study, serum uric acid levels were significantly higher in subjects with metabolic syndrome than that in subjects without the syndrome, even after strictly considering alcohol consumption. In addition, the contribution of abdominal obesity and dyslipidemia in men and abdominal obesity in women to the development of uric acid was noted. There was a difference in serum uric acid levels according to gender: being lower in women than in men, possibly because of estrogen or other steroid hormones [12]. The prevalence of metabolic syndrome in men was significantly higher than that in women using the criterion in Japan as we previously reported [4]. Although uric acid is related to hypertension and dyslipidemia in some reports [13,14], sex differences and the criterion of metabolic syndrome in Japan might have affected the results in this study.

Potential limitations remain in our study. First, the cross-sectional study design in our study makes it difficult to infer causality between metabolic syndrome and uric acid. Second, we could not prove the mechanism of the link between metabolic syndrome and uric acid. The kidney seems to play an important role in the development of metabolic syndrome [15]. It is possible that a normal kidney response to compensatory hyperinsulinemia associated with insulin resistance in non diabetic subjects contributes to the development of hyperuricemia. This is because hyperinsulinemia causes a significant decrease in the urinary excretion of serum uric acid and sodium [16]. In addition, it is reported that uric acid plays a role in platelet adhesiveness [17] and the formation of free radicals and oxidative stress [18].

In conclusion, uric acid was closely associated with metabolic syndrome in Japanese subjects. Further intervention studies are necessary to test the effect of prevention and treatment of metabolic syndrome on uric acid.

Acknowledgement

This research was supported in part by Research Grants from the Ministry of Health, Labor, and Welfare, Japan.

Conflict of interest

There are no conflicts of interest.

REFERENCES

- [1] N. Miyatake, Y. Kawasaki, H. Nishikawa, S. Takenami, T. Numata, Prevalence of metabolic syndrome in Okayama prefecture, Japan, *Intern. Med.* 45 (2006) 107–108.
- [2] P.W. Wilson, W.B. Kannel, H. Silbershatz, R.B. D'Agostino, Clustering of metabolic factors and coronary heart disease, *Arch. Intern. Med.* 159 (1999) 1104–1109.
- [3] N. Miyatake, J. Wada, Y. Kawasaki, S. Matsumoto, H. Makino, T. Numata, Relationship between metabolic syndrome and proteinuria in the Japanese population, *Intern. Med.* 45 (2006) 599–603.
- [4] N. Miyatake, S. Matsumoto, H. Makino, T. Numata, Comparison of hepatic enzymes between Japanese men with and without metabolic syndrome, *Acta Med. Okayama* 61 (2007) 31–34.
- [5] W.J. Fessel, High uric acid as an indicator of cardiovascular disease, *Am. J. Med.* 68 (1980) 401–404.
- [6] Definition and the diagnostic standard for metabolic syndrome-Committee to evaluate diagnostic standards for metabolic syndrome, *Nippon Naika Gakkai Zasshi*, 94 (2005) 794–809 (Japanese).
- [7] Y. Taniguchi, T. Hayashi, K. Tsumura, G. Endo, S. Fujii, K. Okada, Serum uric acid and the risk for hypertension and Type 2 diabetes in Japanese men: The Osaka Health Survey, *J. Hypertens.* 19 (2001) 1209–1215.
- [8] H.J. Milionis, A.I. Kafika, S.G. Tsouli, V.G. Athyros, E.T. Bairaktari, K.I. Seferiadis, et al., Effects of statin treatment on uric acid homeostasis in patients with primary hyperlipidemia, *Am. Heart J.* 148 (2004) 635–640.
- [9] J. Lee, D. Sparrow, P.S. Vokonas, L. Landsberg, S.T. Weiss, Uric acid and coronary heart disease risk: evidence for a role of uric acid in the obesity–insulin resistance syndrome. The Normative Aging Study, *Am. J. Epidemiol.* 142 (1995) 354–359.
- [10] R. Kawamoto, H. Tomita, Y. Oka, N. Ohtsuka, Relationship between serum uric acid concentration, metabolic syndrome and carotid atherosclerosis, *Intern. Med.* 45 (2006) 605–614.
- [11] N. Ishizaka, Y. Ishizaka, E. Toda, R. Nagai, M. Yamakado, Association between serum uric acid, metabolic syndrome, and carotid atherosclerosis in Japanese individuals, *Arterioscler. Thromb. Vasc. Biol.* 25 (2005) 1038–1044.
- [12] H. Sumino, S. Ichikawa, T. Kanda, T. Nakamura, T. Sakamaki, Reduction of serum uric acid by hormone

- replacement therapy in post menopausal women with hyperuricaemia, *Lancet* 354 (1999) 650.
- [13] R. Klein, B.E. Klein, J.C. Cornoni, J. Maready, J.C. Cassel, H.A. Tyroler, Serum uric acid: its relationship to coronary heart disease risk factors and cardiovascular disease in Evans County, Georgia, *Ann. Intern. Med.* 132 (1973) 401–410.
- [14] T.W. Yoo, K.C. Sung, H.S. Shin, B.J. Kim, B.S. Kim, J.H. Kan, et al., Relationship between serum uric acid concentration and insulin resistance and metabolic syndrome, *Circ. J.* 69 (2005) 928–933.
- [15] G.M. Reaven, The kidney: an unwilling accomplice in syndrome X, *Am. J. Kidney Dis.* 30 (1997) 928–931.
- [16] E. Muscelli, A. Natali, S. Bianchi, R. Bigazzi, A.Q. Galvan, A.M. Sironi, et al., Effect of insulin on renal sodium and uric acid handling in essential hypertension, *Am. J. Hypertens.* 9 (1996) 746–752.
- [17] J.V. Selby, G.D. Friedman, C.P. Quesenberry Jr., Precursors of essential hypertension: pulmonary function, heart rate, uric acid, serum cholesterol, and other serum chemistries, *Am. J. Epidemiol.* 131 (1990) 1017–1027.
- [18] J.L. Zweier, P. Kuppusamy, G.A. Lutty, Measurement of endothelial cell free radical generation: evidence for a central mechanism of free radical injury in postischemic tissues, *Proc. Natl. Acad. Sci. USA* 85 (1988) 4046–4050.

食生活習慣改善を中心とした肥満改善教室「食事ですリムコース」開催の試み

田中晶子 宮武伸行 国橋由美子
Tanaka, Akiko Miyatake, Nobuyuki Kunihashi, Yumiko

林 恵子 久富百合子 山下裕絵
Hayashi, Keiko Hisatomi, Yuriho Yamashita, Hiroe

沼田健之
Numata, Takeyuki

岡山県南部健康づくりセンター

表1 対象

人数	40
年齢 (歳)	57.7±8.0
身長 (cm)	154.8±4.9
体重 (kg)	63.3±9.6
BMI (kg/m ²)	26.5±3.8
腹囲 (cm)	86.2±12.8
ヒップ囲 (cm)	98.4±7.1

平均値±標準偏差

キーワード：肥満, 食生活習慣, 生活習慣

はじめに

平成20年度から、メタボリックシンドロームの概念を取り入れた新しい健診・保健指導がスタートする。岡山県南部健康づくりセンターでは、従来より、運動、食事、休養を3つの柱に、肥満改善教室を開催、評価、報告してきた¹⁻⁵⁾。実際の現場においては運動、食事、休養の生活習慣改善をバランスよく実行するのが理想であるが、運動が得意な人、食生活習慣の改善に意欲のある人などさまざまな参加者がいるのが実情で、肥満改善プログラムもそれに応じて準備されていること

が望ましい。

今回筆者らは、食生活習慣の改善を中心とした肥満改善プログラムを作成し、教室「食事ですリムコース」を実施したので報告する。

対象と方法

対象は、当センターで平成18年度から平成19年度までの間に肥満改善教室「食事ですリムコース」に参加した延べ40人(57.7±8.0歳)(表1)のうち、継続して教室に参加し、前後で測定を行えた男性2名、女性26人の計28人(58.4±8.2歳)である。

図1 プログラムの内容

回数	テーマ	具体的な指導内容
1	オリエンテーション, 肥満について知ろう, 運動の話	BMIの計算, 1日の推定エネルギー必要量の計算, 食事記録の書き方と記録の実践
2	食品のエネルギーやバランスについて, 試食	フードモデルを用いて, 食品の量とエネルギーの説明, 誤った健康情報の是正
3	献立づくり, 試食	1食500kcalの献立を作成, ヘルシーメニューの試食
4	やせるためのコツ, 運動の話	やせるためポイントを説明, 家でできる運動の説明と実技
5	調理実習「おいしいヘルシー料理」10:30~13:00	作成した献立を使っでの調理実習
6	あなたの成果は…?	1カ月後の成果を発表, グループワーク