

$\pm 5.0\%$ of EGFP-positive cells, and $87.3\% \pm 2.6\%$ of Iba1-positive cells coexpressed CD68, which also demonstrated that BM-derived cells expressing Iba1 were of macrophage lineage.

Observation of BM-derived cells expressing Iba1 in the cochlea with a confocal microscope revealed a specific morphological feature that was characterized by a spindle shape with several ramified processes, a characteristic morphological feature of macrophages (Fig. 2a–c). In addition to the immunohistochemistry phenotype, BM-derived cells expressing Iba1 in the cochlea morphologically followed microglia that are referred to as resident tissue macrophages in the CNS.

Systemic Application of M-CSF Increased the Number of Iba1-Positive Cells in the Cochlea

To characterize Iba1-positive cells in the cochlea, we next examined the mobilization of Iba1-positive cells using systemic application of M-CSF, the primary regulator of activation of mononuclear phagocytes in wild-type C57BL/6 mice. Iba1-positive cells in both SL and SG were observed more densely in M-CSF-treated mice (Fig. 3a,b) than in controls (Fig. 3c,d). The density of Iba1-positive cells in SL of the middle turn in the controls was 1.46 ± 0.22 (cells/ $10^4 \mu\text{m}^2$), which increased to 2.54 ± 0.45 after M-CSF treatment (Fig. 3e), although the difference was not significant. By contrast, a significant increase was identified in the density of Iba1-positive cells in SG (from 1.39 ± 0.18 to 2.95 ± 0.3 ; Fig. 3e). These data revealed that Iba1-positive cells in the cochlea are under the control of M-CSF signaling. In addition to the findings of phenotype and morphology, we have demonstrated that BM-derived cells labeled with Iba1 have a quality of resident tissue macrophage in the cochlea.

Cochlear Macrophages Gradually Turn Over for More Than 6 Months

The percentage of chimerism in the peripheral blood at 3 months after BM transplantation was $82.8\% \pm 3.6\%$, indicating that hematopoietic reconstitution was performed successfully at this time point. The chimeric ratio observed in the present study was compatible with previous reports on BM chimeric mice (Yoshimoto et al., 2003; Lang et al., 2006). Sequential observation of Iba1-positive cells in the cochlea of transplanted mice demonstrated that cochlear macrophages labeled with Iba1 survived systemic irradiation and were gradually replaced by EGFP-positive cells derived from transplanted HSCs. One or two weeks after HSC transplantation, no EGFP-positive cells were found within cochlear tissues. By contrast, at 4 weeks after transplantation, expression of EGFP was found in $15.4\% \pm 6.6\%$ of Iba1-positive cells in the cochlea. The ratio for EGFP expression in Iba1-positive cells increased remarkably to $64.9\% \pm 8.1\%$ in cochlear specimens obtained at 3 months (12 weeks) after transplantation, then reached $84.1\% \pm 1.6\%$ at 6 months (24 weeks) after transplanta-




Fig. 3. Systemic application of macrophage colony-stimulating factor increases the density of Iba1-positive cells in the spiral ligament (SL) and spiral ganglion (SG). **a–d**: Several Iba1-positive cells were found in the SL (**c**) and the SG (**d**) following systemic application of macrophage colony-stimulating factor (M-CSF), although few cells expressing Iba1 were observed in control specimens (**a,b**). **e**: Densities of Iba1-positive cells (cells/ $10^4 \mu\text{m}^2$) in SG of M-CSF-treated cochleae were significantly higher than those of control cochleae ($*P = 0.002$, unpaired *t*-test), although no significant difference is found in SL ($P = 0.06$). Bars represent standard errors. Scale bar = $50 \mu\text{m}$.

tion. The differences in the ratio for EGFP expression in Iba1-positive cells were significant at 3 and 6 months compared with 1, 2, or 4 weeks (Fig. 4). The distribution of Iba1-positive cells in the cochlea was identical at each time point (data not shown). The density of Iba1-positive cells in SL was 2.39 ± 0.32 (cells/ $10^4 \mu\text{m}^2$) at




Fig. 4. Sequential analysis of chimeric ratios for Iba1-positive cells following transplantation of hematopoietic stem cells derived from GFP mice. The ratios for EGFP expression in Iba1-positive cells of the whole cochlea are shown at 1, 2, and 4 weeks and 3 (12 weeks) and 6 (24 weeks) months after transplantation. The graph demonstrates a gradual increase in the ratio of EGFP expression in Iba1-positive cells, indicating replacement of native Iba1-positive cells by EGFP-positive cells derived from engrafted hematopoietic stem cells. The ratio at 3 or 6 months is significantly higher than that at 1, 2, or 4 weeks (* $P < 0.001$, ANOVA with Tukey-Kramer's test). Bars show standard errors.

1 week, 2.67 ± 0.07 at 2 weeks, 3.02 ± 0.42 at 4 weeks, 3.25 ± 0.18 at 3 months, and 3.31 ± 0.11 at 6 months after BM transplantation. The density of Iba1-positive cells in SG was 1.35 ± 0.10 (cells/ $10^4 \mu\text{m}^2$) at 1 week, 1.73 ± 0.37 at 2 weeks, 1.22 ± 0.15 at 4 weeks, 1.93 ± 0.32 at 3 months, and 2.26 ± 0.14 at 6 months after BM transplantation. There were no significant differences in the density of Iba1-positive cells in SL among the experimental groups. However, the increase with age in the density of Iba1-positive cells in SG was statistically significant between 1 week and 6 months and between 4 weeks and 6 months. These findings indicate that cochlear macrophages are not eliminated by systemic irradiation and gradually turn over for more than 6 months in nondamaged cochlea.

Cochlear Macrophages Increased in Response to Local Surgical Stress

To examine the reaction of cochlear macrophages to an acute, local, exogenous stress on the cochlea, surgical invasion with application of physiological saline via PSSC was performed on both wild-type and BM chimeric mice. ABR recoding was performed to determine the extent of functional damage following treatment. Significant elevation of ABR thresholds was observed on day 1 after local treatment, whereas no elevation was found on days 7 and 28 at all frequencies (Fig. 5a), indicating that the surgical stress used in the present study caused temporary damage to the cochlea.

Iba1-positive cells were increased temporarily in response to local surgical stress in SL and SG (Fig. 5b–e). The density of Iba1-positive cells in SL exhibited an immediate response to the treatment on day 1 after surgery (3.96 ± 0.16 cells/ $10^4 \mu\text{m}^2$ vs. 1.62 ± 0.18 , preoperative), significantly increased on day 7 after surgery (5.42 ± 0.59), and decreased on day 28 (2.90 ± 0.41 ; Fig. 5f). No significant difference was found in the densities of Iba1-positive cells between before surgery, on day 1, or on day 28. A temporary increase was also found in the density of Iba1-positive cells in SG on day 1 (1.94 ± 0.06) and day 7 (2.66 ± 0.33 ; Fig. 5f) compared with densities of 1.20 ± 0.09 preoperatively and 1.33 ± 0.16 on day 28. The differences between preoperative values and day 7 and between days 7 and 28 were statistically significant.

We also examined the effect of local surgical stress in BM chimeric mice that had been transplanted with HSCs labeled with EGFP 3 months before, to test the mobilization of Iba1-positive cell from BM to the cochlea. The density of cells dually labeled with EGFP and Iba1 studied in SL at day 7 after treatment was 4.01 ± 0.21 (cells/ $10^4 \mu\text{m}^2$) in the operated group and 2.54 ± 0.39 in nonoperated BM chimeric mice, and that in SG was 1.88 ± 0.52 in the operated group and 1.42 ± 0.21 in the nonoperated group. The difference in the density of EGFP-positive cells in SL between the two groups was statistically significant, whereas that in SG was not significant.

Immunohistochemistry for Ki67 was performed before and at days 1 and 7 after surgical treatment, to test whether the increase in cochlear macrophages following surgical treatment was due to proliferation in situ. On day 1 following treatment, 1.75 ± 0.85 cells/section were found to be dually labeled with Ki67 and CD68 ($1.74 \pm 0.71\%$ in total number of cells positive for CD68). On day 7 following treatment, 2.25 ± 0.75 were dually labeled with Ki67 and CD68 ($2.49\% \pm 0.81\%$), in contrast to 1.00 ± 0.40 in untreated mice ($1.53\% \pm 0.51\%$). All Ki67-positive cells in the three groups were found in the lower part of SL. However, no statistically significant difference was observed in the number of Ki67-positive cells among the three groups. These findings indicated that the increase in cochlear macrophages was due mainly to the migration of macrophages from the circulation into the cochlea. However, it was also revealed that the proliferation of macrophages in situ is one of the possible sources for cochlear macrophages.

DISCUSSION

The present study revealed that BM-derived cells are supplied continuously to the cochlea even in the adult mouse and demonstrated that the predominant phenotype of macrophages involved expression of CD68, F4/80, or Iba1 in SL and SG. Our study on the morphology, immunohistochemical phenotype, and reactivity to M-CSF provides evidence that hematopoi-




Fig. 5. Effects of local surgical stress on densities of Iba1-positive cells in the spiral ligament (SL) and spiral ganglion (SG). **a**: Local surgical stress causes significant elevation of ABR thresholds at each frequency on day 1 after treatment ($*P < 0.001$, ANOVA with Tukey-Kramer's test), and the elevation recovered on days 7 and 28 at all the frequencies tested. **b–e**: Immunostaining for Iba1 in cochlear specimens obtained preoperatively (pre-ope; b) and on day 1 (c), day

7 (d), and day 28 (e) after treatment. Increase in Iba1-positive cells was observed in the SL and SG of cochleae obtained on days 1 and 7. **f**: The density of Iba1-positive cells (cells/10⁴ μm²) on day 7 was significantly higher than that in preoperative specimens in both SL ($*P = 0.003$) and SG ($*P = 0.002$) or day 28 in SG. Bars show standard errors. Scale bars = 50 μm.

etic BM-derived cells expressing Iba1 are constitutively present as resident tissue macrophages in the cochlea. The examination of the recruitment of cochlear macrophages demonstrated that these cells exhibited slow turnover for several months during steady-state conditions and quickly increased in response to local surgical stress.

Although earlier studies have demonstrated infiltration of inflammatory cells, including macrophages, into the cochlea following inner ear injury (Fredelius, 1988; Bhawe et al., 1998; Hirose et al., 2005; Tornabene et al., 2006; Ladrech et al., 2007), two recent studies have suggested the presence of resident tissue macrophages in the inner ear.

Lang et al. (2006) reported that BM-derived cells are constitutively present and widely distributed in the cochlea in the same manner demonstrated in the present study. Although they concluded negatively that only 5% of BM-derived cells differentiated into CD45R-positive macrophages, the rate for CD45 expression in BM-derived cells is also compatible with the present study. The authors instead emphasized that histological analysis of the cochleae following HSCs revealed the contribution of BM-derived cells to fibrocytes or mesenchymal cells in the inner ear. However, the specific ratios for immunoreactivity of Na, K-ATPase, or the Na-K-Cl transporter in the BM-derived cells were not determined in their study. Moreover, there is a discrepancy in the ratio for the expression of F4/80 between the study by Lang et al. (2006) and the present study, although both studies used HSC-transplanted mice that exhibited similar chimeric ratios in peripheral blood and SL cells. One possible explanation for this discrepancy is a difference in methods for immunohistochemistry. We used cryostat sections, whereas Lang et al. used paraffin-embedded sections. Our study demonstrated that more than 80% of BM-derived cells in SL and SG show the phenotype of macrophages and that resident tissue macrophages in the cochlea during steady-state conditions are present in a larger number than reported previously.

Another report was made by Hirose et al. (2005) that CD45-positive mononuclear phagocytes are present in the SL of nonnoise-exposed CX3CR1^{GFP/GFP} transgenic mice, and these cells are also labeled with Iba1 or CD68, which is consistent with the results of the present study. However, the density of CD45-positive cells in the cochlea was quite different between the wild-type and CX3CR1^{GFP/GFP} transgenic mice used in their study, although the number of BM-derived cells expressing CD45 in the cochlea demonstrated in the present study was identical to that of CD45-positive cells in the cochlea of nonnoise-exposed wild-type mice presented by Hirose et al. (2005). The most controversial point is whether it is appropriate to assume that macrophages observed in the cochlea can be described as "microglia-like" cells. Although the authors failed to describe the distribution of macrophages in the auditory nervous system including the SG and cochlear modiolus during steady-state conditions, they distinguish the infiltrating macrophages observed after noise exposure from microglia in the CNS by the morphology and the potentiality of exchanging from the vascular space. The proliferation in situ is thought to be one of the main sources of microglia in adults, but some studies have reported that BM-derived cells can enter the CNS and populate the microglial cell compartment (Lawson et al., 1992; Corti et al., 2002; Hess et al., 2004; Simard and Rivest, 2004; Malm et al., 2005). Moreover, previous studies in the CNS using flow cytometry or immunohistochemistry defined a profile of characterization of microglial cells corresponding to the following phenotype: CD68⁺, CD45 low, CD11b⁺, CD11c high, and MHC class II⁺ (Guillemin and Brew, 2004; Floden and Combs, 2007).

Our results on cochlear resident macrophages "CD68⁺, Iba1⁺, and CD45[−]" were compatible with the immunohistochemical definition of microglia as reported above. In addition, the cochlear resident macrophages were observed in the auditory nervous system as well as in the connective tissue of SL in the present study. Because there is a considerable heterogeneity in the phenotypes of the macrophage lineage, and because resident tissue macrophages share several antigens with infiltrating macrophages (Guillemin and Brew, 2004), further studies should be carried out to define a profile of characterization on both resident and infiltrating macrophages in the cochlea.

Many studies on resident tissue macrophages in other organs have been performed with BM chimeric mice. In terms of replacement of resident tissue macrophages by BM, donor-derived cells are reported rapidly to populate the liver with Kupffer cells, resident macrophages in the liver, within 3 weeks, and donor Kupffer cells in liver transplants are replaced with similar kinetics (Naito et al., 1997). Microglia in the CNS exhibit longer turnover period than cochlear resident macrophages. At 12 months following HSC transplantation, 40% of microglia have been replaced by hematopoietic cells (Hess et al., 2004; Simard and Rivest, 2004). In contrast, Xu et al. (2007) reported that EGFP-positive BM-derived cells infiltrate normal retina in significant numbers at 8 weeks after BM transplantation and that by 6 months all retinal microglia/macrophages were replaced by BM-derived EGFP-positive cells, the turnover rate of which is very similar to that of the inner ear in the present findings. From the viewpoint of turnover rates, cochlear resident macrophages have characteristics as resident macrophages in the retina.

Although BM chimeric mice are a powerful tool for the analysis of the dynamics of BM-derived cells, there are some limitations. The results of BM chimeric mice are obtained under extraordinary conditions; systemic lethal irradiation and following BM transplantation. It is also technically difficult to set an ideal negative control, with irradiation but without BM transplantation. Moreover, it is difficult to exclude completely the possibility that the results observed in this study are caused by irradiation-induced damage. However, despite these limitations in studies using BM chimeric mice, our results provided some new insights into the origin and distribution of cochlear resident macrophages and the possible supply of cochlear macrophages by replenishment with BM-derived cells.

Recent studies have demonstrated multiple key functions of resident tissue macrophages not only in phagocytosis of foreign bodies or senescent cells but also in the production and secretion of cytokines and the regulation of specific immune responses (Gordon and Taylor, 2005). In the CNS, microglia have both neurotrophic and neurotoxic properties (Kreutzberg, 1996; Moore and Thanos, 1996; Streit, 1996) and play an effector role in both innate and adaptive immune responses, allowing the CNS to respond rapidly and

efficiently to a wide range of pathogens (Olson and Miller, 2004). Although it has not been elucidated whether the infiltrating macrophages play beneficial or harmful roles in the maintenance of auditory function, pharmacological intervention in the infiltration of macrophages may be a possible strategy for treatment of SNHL. In conclusion, the resident cochlear macrophages have potential as a therapeutic target by means of controlling their ability of phagocytosis, migration, or release of cytokines in the pathology of inner ear immune disorders.

ACKNOWLEDGMENTS

The authors thank Dr. Masaru Okabe (Osaka University) for generously providing GFP transgenic mice and Dr. Junko Okano (Department of Anatomy and Developmental Biology, Graduate School of Medicine, Kyoto University) for critical discussion. We are also very thankful to Dr. Hideaki Ogita, Ms. Yoko Nishiyama, Ms. Rika Sadato, Ms. Akemi Saito, Ms. Keiko Nishio, and Ms. Tomoyo Namura for their excellent technical assistance.

REFERENCES

- Bhave SA, Oesterle EC, Coltrera MD. 1998. Macrophage and microglia-like cells in the avian inner ear. *J Comp Neurol* 398:241–256.
- Corti S, Locatelli F, Donadoni C, Strazzer S, Salani S, Del Bo R, Caccialanza M, Bresolin N, Scarlato G, Comi GP. 2002. Neuroectodermal and microglial differentiation of bone marrow cells in the mouse spinal cord and sensory ganglia. *J Neurosci Res* 70:721–733.
- Floden AM, Combs CK. 2007. Microglia repetitively isolated from in vitro mixed glial cultures retain their initial phenotype. *J Neurosci Methods* 164:218–224.
- Fredelius L. 1988. Time sequence of degeneration pattern of the organ of Corti after acoustic overstimulation. A transmission electron microscopy study. *Acta Otolaryngol* 106:373–385.
- Gordon S, Taylor PR. 2005. Monocyte and macrophage heterogeneity. *Nat Rev Immunol* 5:953–964.
- Guillemin GJ, Brew BJ. 2004. Microglia, macrophages, perivascular macrophages, and pericytes: a review of function and identification. *J Leukocyte Biol* 75:388–397.
- Hess DC, Abe T, Hill WD, Studdard AM, Carothers J, Masuya M, Fleming PA, Drake CJ, Ogawa M. 2004. Hematopoietic origin of microglial and perivascular cells in brain. *Exp Neurol* 186:134–144.
- Hirose K, Discolo CM, Keasler JR, Ransohoff R. 2005. Mononuclear phagocytes migrate into the murine cochlea after acoustic trauma. *J Comp Neurol* 489:180–194.
- Iguchi F, Nakagawa T, Tateya I, Kim TS, Endo T, Taniguchi Z, Naito Y, Ito J. 2003. Trophic support of mouse inner ear by neural stem cell transplantation. *Neuroreport* 14:77–80.
- Imai Y, Ibata I, Ito D, Ohsawa K, Kohsaka S. 1996. A novel gene *iba1* in the major histocompatibility complex class III region encoding an EF hand protein expressed in a monocytic lineage. *Biochem Biophys Res Commun* 224:855–862.
- Kim TS, Nakagawa T, Kitajiri S, Endo T, Takebayashi S, Iguchi F, Kita T, Tamura T, Ito J. 2005. Disruption and restoration of cell-cell junctions in mouse vestibular epithelia following aminoglycoside treatment. *Hear Res* 205:201–209.
- Kreutzberg GW. 1996. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* 19:312–318.
- Ladrech S, Wang J, Simonneau L, Puel JL, Lenoir M. 2007. Macrophage contribution to the response of the rat organ of Corti to amikacin. *J Neurosci Res* 85:1970–1979.
- Lang H, Ebihara Y, Schmiedt RA, Minamiguchi H, Zhou D, Smythe N, Liu L, Ogawa M, Schulte BA. 2006. Contribution of bone marrow hematopoietic stem cells to adult mouse inner ear: mesenchymal cells and fibrocytes. *J Comp Neurol* 496:187–201.
- Lawson LJ, Perry VH, Gordon S. 1992. Turnover of resident microglia in the normal adult mouse brain. *Neuroscience* 48:405–415.
- Lee JE, Nakagawa T, Kim TS, Iguchi F, Endo T, Dong Y, Yuki K, Naito Y, Lee SH, Ito J. 2003. A novel model for rapid induction of apoptosis in spiral ganglions of mice. *Laryngoscope* 113:994–999.
- Malm TM, Koistinaho M, Parepalo M, Vatanen T, Ooka A, Karlsson S, Koistinaho J. 2005. Bone-marrow-derived cells contribute to the recruitment of microglial cells in response to beta-amyloid deposition in APP/PS1 double transgenic Alzheimer mice. *Neurobiol Dis* 18:134–142.
- Moore S, Thanos S. 1996. The concept of microglia in relation to central nervous system disease and regeneration. *Prog Neurobiol* 48:441–460.
- Naito M, Hasegawa G, Takahashi K. 1997. Development, differentiation, and maturation of Kupffer cells. *Microsc Res Techniq* 39:350–364.
- Nakagawa T, Kim TS, Murai N, Endo T, Iguchi F, Tateya I, Yamamoto N, Naito Y, Ito J. 2003. A novel technique for inducing local inner ear damage. *Hear Res* 176:122–127.
- Okabe M, Ikawa M, Kominami K, Nakanishi T, Nishimune Y. 1997. ‘Green mice’ as a source of ubiquitous green cells. *FEBS Lett* 407:313–319.
- Okano T, Nakagawa T, Kita T, Endo T, Ito J. 2006. Cell-gene delivery of brain-derived neurotrophic factor to the mouse inner ear. *Mol Ther* (in press).
- Olson JK, Miller SD. 2004. Microglia initiate central nervous system innate and adaptive immune responses through multiple TLRs. *J Immunol* 173:3916–3924.
- Shiga A, Nakagawa T, Nakayama M, Endo T, Iguchi F, Kim TS, Naito Y, Ito J. 2005. Aging effects on vestibulo-ocular responses in C57BL/6 mice: comparison with alteration in auditory function. *Audiol Neurootol* 10:97–104.
- Shinohara T, Bredberg G, Ulfendahl M, Pykko I, Olivius NP, Kaksonen R, Lindstrom B, Altschuler R, Miller JM. 2002. Neurotrophic factor intervention restores auditory function in deafened animals. *Proc Natl Acad Sci U S A* 99:1657–1660.
- Simard AR, Rivest S. 2004. Bone marrow stem cells have the ability to populate the entire central nervous system into fully differentiated parenchymal microglia. *FASEB J* 18:998–1000.
- Streit WJ. 1996. The role of microglia in brain injury. *Neurotoxicology* 17:671–678.
- Tornabene SV, Sato K, Pham L, Billings P, Keithley EM. 2006. Immune cell recruitment following acoustic trauma. *Hear Res* 222:115–124.
- Xu H, Chen M, Mayer EJ, Forrester JV, Dick AD. 2007. Turnover of resident retinal microglia in the normal adult mouse. *Glia* 55:1189–1198.
- Yoshimoto M, Shinohara T, Heike T, Shiota M, Kanatsu-Shinohara M, Nakahata T. 2003. Direct visualization of transplanted hematopoietic cell reconstitution in intact mouse organs indicates the presence of a niche. *Exp Hematol* 31:733–740.

Potential of Embryonic Stem Cell-Derived Neurons for Synapse Formation With Auditory Hair Cells

Masahiro Matsumoto,¹ Takayuki Nakagawa,^{1*} Ken Kojima,^{1,2}
Tatsunori Sakamoto,¹ Fumino Fujiyama,³ and Juichi Ito¹

¹Department of Otolaryngology-Head and Neck Surgery, Graduate School of Medicine, Kyoto University, Kyoto, Japan

²Establishment of International COE for Integration of Transplantation Therapy and Regenerative Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan

³Department of Morphological Brain Science, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Recent studies have indicated that embryonic stem cells (ESCs) can be a source for the replacement of spiral ganglion neurons (SGNs), auditory primary neurons, and neurite projections from ESC-derived neurons to auditory sensory epithelia. However, the potential of ESC-derived neurons for synapse formation with auditory hair cells (HCs) has not been elucidated. The present study therefore aimed to examine the ability of ESC-derived neurons to form synaptic connections with HCs *in vitro*. Mouse ESC-derived neural progenitors expressing enhanced green fluorescence protein (EGFP) were cocultured with explants of cochlea sensory epithelia obtained from postnatal day 3 mice. After a 7-day culture, neurites of ESC-derived neurons predominantly elongated toward inner hair cells (IHCs), which play a crucial role in sound transmission to SGNs. Immunohistochemical analyses revealed the expression of synapsin 1 and synaptophysin in the nerve endings of ESC-derived neurons adjacent to IHCs, indicating the formation of synaptic connections. Transmission electron microscopy demonstrated synaptic contacts between nerve endings of ESC-derived neurons and IHCs. The present findings show that ESC-derived neurons can make synaptic connections with IHCs. © 2008 Wiley-Liss, Inc.

Key words: embryonic stem cell; hair cell; cochlea; transplantation; spiral ganglion neuron

Sensorineural hearing loss (SNHL) is one of the most common disabilities in our society today, but, because mammalian inner ear cells have limited regenerative activity, treatment options are limited to cochlear implants and the use of hearing aids. Spiral ganglion neurons (SGNs) located in the bony axis (modiolus) of the cochlea are bipolar auditory primary neurons that make synaptic connections with auditory hair cells (HCs) and the cochlear nucleus in the brainstem. They consist of two types of neurons: type I neurons making synaptic

connections with inner hair cells (IHCs) that play a predominant role in the transduction of sound stimuli to neural stimuli, and type II neurons making synaptic connections with outer hair cells (OHCs) that assist in controlling the sensitivity of regions of the auditory epithelia to specific sound stimuli (Spoendlin, 1987; Rubel and Fritzsche, 2002). The loss of SGNs therefore compromises auditory function. SGN degeneration resulting in SNHL occurs during the aging process (Keithley et al., 1989), and can also be caused by several ototoxic agents (Lee et al., 2003) or acoustic trauma (Pujol and Puel, 1999). It compromises the clinical benefits of cochlear implants (Nadol et al., 1989), which rely on direct stimulation of SGNs to improve impaired hearing.

Recent investigations have indicated the potential for SGN regeneration using cell transplantation into the cochlea. Several candidates are available as a source of transplant; previous studies have demonstrated the survival and neurite outgrowth of neural stem cells (Tamura et al., 2004; Hu et al., 2005b) and dorsal ganglion neurons (Hu et al., 2005a) after transplantation into the cochlea. Bone marrow-stromal cells have also shown the capacity for settlement and differentiation into neuronal cells after engraftment into the cochlea (Naito et al., 2004). More recently, the survival of bone marrow stromal cell-derived neurons in the cochlear modiolus has been reported (Matsuoka et al., 2006). Embryonic stem cells (ESCs) are another possible transplant source with the ability to dif-

Contract grant sponsor: Ministry of Education, Science, Sports, Culture and Technology of Japan.

*Correspondence to: Takayuki Nakagawa, MD, PhD, Departments of Otolaryngology-Head and Neck Surgery, Graduate School of Medicine, Kyoto University, Kawaharacho 54, Shogoin, Sakyo-ku, Kyoto, 606-8507 Japan. E-mail: tnakagawa@ent.kuhp.kyoto-u.ac.jp

Received 6 September 2007; Revised 25 January 2008; Accepted 11 March 2008

Published online 9 June 2008 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jnr.21754

ferentiate into neurons, and several neural-inducing methods have been established for this cell type. ESC-derived neural progenitors have a high potential for survival and differentiation into neurons after transplantation into the cochlear modiolus where they were previously shown to massively elongate their neurites toward the peripheral and central auditory systems, reaching host SGNs and the osseous spiral lamina, which is located between SGNs and auditory sensory epithelia (Okano et al., 2005; Corrales et al., 2006). In addition, transplantation of ESC-derived neurons into the cochlear modiolus contributes to the recovery of SGN functions, as monitored by electrically evoked auditory brainstem responses (Okano et al., 2005). In vitro analyses such as coculture studies of ESC-derived neural progenitors with cochlear or vestibular sensory epithelia have demonstrated their differentiation into bipolar neurons (Coleman et al., 2007) and neurite outgrowth that attaches to sensory HCs and expresses synaptophysin at the attachment site (Kim et al., 2005; Matsumoto et al., 2005).

Although there was no direct evidence of synapses establishment, there was functional evidence to indicate the establishment of functioning neural connection (Okano et al., 2005). The aim of the present study was therefore to determine the synapse formation ability of ESC-derived neurons with auditory HCs. In the cochlea, there are the afferent synapses between the peripheral fibers of spiral ganglion neurons and sensory HCs (Sobkowitz, 1992) and the efferent synapses, both axosomatic with the sensory HCs and axodendritic with the peripheral afferents (Sobkowitz et al., 1997, 2002). We cocultured mouse ESC-derived neural progenitors with explants of mouse auditory sensory epithelia and morphologically examined the neural connections that arose between ESC-derived neurons and HCs.

MATERIALS AND METHODS

The Animal Research Committee, Graduate School of Medicine, Kyoto University, approved all experimental protocols. Animal care was under the supervision of the Institute of Laboratory Animals, Graduate School of Medicine, Kyoto University. All experimental procedures were performed in accordance with the NIH *Guide for the care and use of laboratory animals*.

ESCs and Neural Induction

We used the mouse ESCs G4-2 (generously donated by Dr Hitoshi Niwa of Riken CDB, Kobe, Japan), derived from the E14tg2a ESC line (Hooper et al., 1987) and carrying the EGFP gene driven by a ubiquitous strong promoter (CAG promoter), because our previous experiments demonstrated the potential of neurons derived from this cell line for neurite outgrowth toward inner ear HCs (Kim et al., 2005; Matsumoto et al., 2005). Stromal cell-inducing activity (SDIA) was used for the neural induction of ESCs (Kawasaki et al., 2000). Briefly, ESCs were cultured to form differentiated colonies on a feeder layer of PA6 stromal cells derived from newborn mouse calvaria (RCB1127; RIKEN Cell Bank, Japan) in Glas-

gow's modified Eagle's medium (GMEM; Invitrogen, Carlsbad, CA) supplemented with 5% knockout serum replacement (KSR; Invitrogen), 1 mM pyruvate (Sigma, St. Louis, MO), 0.1 mM nonessential amino acids (Invitrogen), and 0.2 mM 2-mercaptoethanol (2-ME; Wako, Osaka, Japan). Colonies that formed on the PA6 monolayer after 6 days of culture were isolated by incubation with collagenase B (Roche Diagnostics, Tokyo, Japan) for 5 min at room temperature (RT).

Isolation of Auditory Epithelia and Coculture

Postnatal day (P) 3 ICR mice were deeply anesthetized with diethyl ether and decapitated. The temporal bones were dissected, and the cochleae were freed from surrounding tissues in 0.01 M phosphate-buffered saline (PBS) at pH 7.4, supplemented with 0.2% glucose. After removal of cochlear lateral walls, auditory epithelia were dissected from the cochlear modiolus. SGNs were carefully removed from the explants using fine forceps as described previously (Matsumoto et al., 2005), resulting in complete removal of neural elements from explant cultures (Fig. 1). Auditory epithelia were explanted intact on a sterile membrane (Falcon BD Biosciences, San Jose, CA) into MEM (Gibco, Grand Island, NY) supplemented with 0.3% glucose in 24-well culture plates. Cultures were incubated at 37°C in a humidified atmosphere of 95% air and 5% CO₂ for 24 hr. The cultured explants were then provided for coculture with ESC-derived neural progenitors. Explants including SGNs were excluded from analysis. Finally, 20 explants were provided for coculture with SDIA-treated ESCs, and an additional four explants cultured alone in the same medium were used as controls.

Cultured auditory epithelia were transferred into GMEM supplemented with 5% KSR, 1 mM pyruvate, 0.1 mM nonessential amino acids, and 0.2 mM 2-ME. Cell suspensions of SDIA-treated ESCs (2 μ l, 1×10^4 cells/ μ l GMEM) were placed inside cultured auditory epithelia, where SGNs are normally located, using a Hamilton syringe (Hamilton, Reno, NV). The medium was changed daily, and cultures were maintained for 7 days.

Tissue Preparation

Histological analysis was performed on whole mounts or cryostat sections. After the culture period, whole mounts (cocultures $n = 5$; controls $n = 2$) were fixed in 4% paraformaldehyde in PBS for 15 min at RT. Cryostat sections (cocultures $n = 15$; controls $n = 2$) were fixed in 4% paraformaldehyde with 0.1% glutaraldehyde in PBS for 30 min and then cryoprotected by immersion in 30% sucrose in PBS at 4°C overnight. Specimens were mounted in OCT compound (Tissue-Tek, Sakura, Tokyo, Japan) and cut at 15 μ m using a cryostat.

Immunohistochemistry

Whole-mount specimens were doubly immunostained for myosin VIIa and β III tubulin to identify the location of HCs in the auditory epithelia and ESC-derived neurons and their neurites, respectively. The latter location was also determined by EGFP expression. We focused on neurite extension




Fig. 1. Explants of auditory epithelia with or without spiral ganglion neurons. On 7 days in vitro (DIV), an explant of the auditory epithelium with spiral ganglion neurons (A) exhibits the presence of neurites and nerve endings expressing β III tubulin (blue) adjacent to hair cells (red; C: xy projection; C': xz projection). Arrows in A indicate the remaining spiral ganglion neurons. An explant of the auditory epithelia after elimination of spiral ganglion neurons (B) demonstrates

three rows of outer hair cells and a single row of inner hair cells expressing myosin VIIa (red) and no neural elements in the auditory epithelium (D: xy projection; D': xz projection). Myo, myosin VIIa; tubulin, β III tubulin; IHC, inner hair cell; OHC, outer hair cell; xy, xy projection image; xz, xz projection image; 7DIV, 7 days in vitro. Scale bars = 200 μ m in A,B; 20 μ m in C,C',D,D'.

of ESC-derived neurons and the relationship between HCs and their extended neurites.

Immunohistochemistry for myosin VIIa, β III tubulin, synapsin 1, synaptophysin, growth-associated protein 43 (GAP43), C-terminal binding protein 2 (ctbp2)/ribeye, and calretinin was carried out on cryostat sections. Synapsin 1 and synaptophysin were used as markers for synaptic vesicles and GAP43 to identify growing neurites. Ctbp2/ribeye was used as a specific marker for synapse ribbons and calretinin as a marker for mature SGNs and their neurites. We focused on

the expression of these marker proteins in the nerve endings of ESC-derived neurons that were attached to HCs. Cochlear specimens obtained from P4 (before hearing onset), P12 (onset of hearing), and adult (10-week-old) mice were also stained for synapsin 1 and synaptophysin to correlate the expression patterns of these proteins with the development of mouse auditory function. Cochlear specimens obtained from adult mice were used as controls for ctbp2/ribeye or calretinin.

Whole mounts and cryostat sections were permeabilized and blocked with 10% normal goat serum (Chemicon, Teme-

cula, CA) in PBS-T (PBS containing 0.02% Triton X) for 30 min at RT, followed by overnight incubation at 4°C in the presence of primary antibodies diluted in an appropriate blocking solution. On the following day, sections were washed three times in PBS-T and incubated in diluted secondary antibody for 1 hr at RT. The primary antibodies used were rabbit antimyosin VIIa polyclonal antibody (1:250; Proteus BioSciences, Ramona, CA), mouse anti- β III tubulin monoclonal antibody (1:250; Covance Research Products, Berkeley, CA), mouse antisynapsin I monoclonal antibody (1:50; BD Biosciences Pharmingen, San Diego, CA), mouse antisynaptophysin monoclonal antibody (1:100; BD Biosciences Pharmingen), mouse anti-GAP43 monoclonal antibody (1:200; Chemicon), mouse anti-ctbp2/ribeye monoclonal antibody (1:50; BD Biosciences Pharmingen), and rabbit anti-calretinin polyclonal antibody (1:250; Chemicon). The secondary antibodies used were Alexa Fluor 546 goat anti-rabbit IgG and Alexa Fluor 633 goat anti-mouse IgG (1:500; Molecular Probes, Eugene, OR). Fluorescent images were obtained with a Leica TCS SP2 confocal laser scanning microscope (Leica Microsystems Inc., Wetzlar, Germany).

The ratio of synapsin 1, synaptophysin, or GAP43 expression in nerve endings of ESC-derived neurons adjacent to IHCs was quantitatively assessed by counting the number of nerve endings positive for each marker in three randomly selected cryostat sections from each specimen. The ratio was determined by dividing the number of positive nerve endings by the total number of nerve endings. Values are given as mean \pm SD.

Transmission Electron Microscopy

Cryostat sections adjacent to the section in which synaptophysin-positive nerve endings of ESC-derived neurons attached to IHCs were stained by immunohistochemistry using an anti-GFP rabbit polyclonal primary antibody (diluted 1:500 in PBS; Molecular Probes). GFP immunoreactivity was visualized using a Vectastain ABC kit and DAB substrate kit (Vector Laboratories, Burlingame, CA). Specimens were then postfixed with 1% OsO₄ in phosphate buffer for 90 min at 4°C. After dehydration in a graded ethanol series, specimens were embedded in Epon-812 and cut into 80-nm ultrathin sections. The sections were stained with lead citrate and viewed with a transmission electron microscope (TEM; H-7000, Hitachi, Tokyo, Japan).

RESULTS

Projection of neurites From ESC-Derived Neurons to HCs

Auditory epithelia explants were cocultured with ESC-derived neural progenitors labeled with EGFP for 7 days. After this, β III tubulin expression was detected in ESC-derived cells, as previously observed (Kim et al., 2005; Matsumoto et al., 2005), indicating that ESC-derived neural progenitors had differentiated into neurons. Seven days after plating, ESC-derived neurons formed colonies inside of a single row of IHCs and projected their neurites toward IHCs of the auditory epithel-




Fig. 2. Embryonic stem cell-derived neurons project neurites to auditory hair cells. A–D show xy projections and A'–D' show xz projections of cocultures with embryonic stem cell (ESC)-derived neurons and auditory epithelia on 7 days in vitro (DIV). A, B: ESC-derived neurons labeled with EGFP (green) and β III tubulin (blue) extend many processes toward IHCs (red). A', B': EGFP- and β III tubulin-expressing fibers (arrowhead in A, B) extend beneath the auditory epithelium and attach to the basal portion of IHCs. C, D: ESC-derived neurons project a few processes toward OHCs like cross-tunnel fibers (arrow in C, D). C', D': ESC-derived neurites attach to the basal portion of OHCs (arrow in C', D'). EGFP, enhanced green fluorescent protein; Myo, myosin VIIa; tubulin, β III tubulin; IHC, inner hair cell; OHC, outer hair cell. Scale bars = 20 μ m.

lia (Fig. 2A, B). EGFP-expressing fibers were shown to extend from ESC-derived neurons beneath the auditory epithelium and attach to the basal portion of IHCs (Fig. 2A', B'), where nerve fibers and nerve endings from type I SGNs also attach in vivo conditions. Some EGFP-labeled neurites were observed around OHCs (Fig. 2C, D). Three of twenty cocultured specimens contained ESC-derived neurites running through the auditory epithelium like the cross-tunnel fibers (Fig. 2C', D'). These findings demonstrate that ESC-derived neurons have the ability to project neurites to auditory HCs and that these neurites exhibit the same distribution as SGN-derived nerve fibers.




Fig. 3. Synapsin 1 and synaptophysin expression in auditory epithelia of postnatal day (P) 4, P12, and adult mice. **A:** In P4 mice, immunoreactivity for synapsin 1 is observed in the nerve fibers (arrows) and nerve endings adjacent to inner hair cells (arrowhead). **B,C:** In P12 and adult mice, the distribution of synapsin 1 immunoreactivity is limited to the nerve endings adjacent to inner hair cells (arrowhead). **D:** P4 mice exhibit synaptophysin immunoreactivity in the

nerve fibers (arrows) and nerve endings adjacent to inner hair cells (arrowhead) and in the cytoplasm of hair cells. **E,F:** In P12 and adult auditory epithelia, synaptophysin expression is accumulated in the nerve endings adjacent to inner hair cells (arrowhead). Cellular nuclei are stained with DAPI (blue). Asterisks indicate the location of hair cells. Synapto, synaptophysin. Scale bars = 20 μ m.

Expression of Synapsin 1, Synaptophysin, GAP43, Ctbp2/Ribeye, and Calretinin in ESC-Derived Neurons

To investigate the synapse formation of ESC-derived neurites with HCs, we examined the expression of the synaptic proteins synapsin 1, which is expressed in small synaptic vesicles containing neurotransmitters (De Camilli et al., 1983a,b; Huttner et al., 1983; Navone et al., 1984), and synaptophysin, which is an intrinsic membrane protein of small synaptic vesicles (Wiedemann and Franke, 1985; Navone et al., 1986; Sudhof et al., 1987) under normal conditions. In auditory epithelia of P4 mice, before the onset of hearing, synapsin 1 immunoreactivity was observed in the nerve fibers and nerve endings adjacent to HCs (Fig. 3A). In P12 mice, at the onset of auditory function, and in adult mice, the distribution of synapsin 1 immunoreactivity was limited to the nerve endings adjacent to HCs (Fig. 3B,C). P4 mice exhibited synaptophysin immunoreactivity in nerve fibers and nerve endings adjacent to IHCs and in the cytoplasm of HCs (Fig. 3D), whereas, in P12 and adult auditory epithelia, synaptophysin expression was observed in the nerve endings adjacent to HCs (Fig. 3E,F).

In cocultured specimens, the expression of synapsin 1 was identified in the nerve endings of ESC-derived neurons adjacent to the base of IHCs (Fig. 4), equivalent to the case in P12 mice. Synaptophysin immunoreactivity was found in the nerve endings of ESC-derived neurons adjacent to the base of IHCs and in the cytoplasm of IHCs (Fig. 5), equivalent to P4 mice. We quantified the ratio of the expression of synapsin 1 or synaptophysin in nerve endings of ESC-derived neurons adjacent to the base of IHCs as $90.5\% \pm 16.2\%$ and $95.2\% \pm 12.6\%$, respectively. These findings indicate active syn-

apse formation between ESC-derived neurons and IHCs in cocultured specimens. We also examined GAP43 immunoreactivity, which reflects neurite elongation and synaptic sprouting (Meiri et al., 1986; Skene et al., 1986; Skene, 1989). GAP43 expression was found in $81.0\% \pm 17.7\%$ of the nerve endings of ESC-derived neurons adjacent to the base of IHCs and the nerve fibers (Fig. 6).

We also performed immunohistochemistry for ctbp2/ribeye and calretinin to examine the types or maturation status of synaptic contacts between IHCs and ESC-derived neurons. We identified expression of calretinin in SGNs, their nerve fibers, and the HCs in adult mouse cochleae (Fig. 7). In cocultured specimens, the expression of calretinin was not observed in ESC-derived nerve fibers and endings, although HCs were positive (Fig. 7). The expression of ctbp2/ribeye was found in both the nucleus and the base of cytoplasm of IHCs in adult mouse cochleae (Fig. 7). In cocultured specimens, ctbp2/ribeye expression was also observed in the nuclei and the base of cytoplasm of IHCs, although few ctbp2/ribeye-positive dots were identified in IHC cytoplasm (Fig. 7).

Transmission Electron Microscopy

To investigate synaptic contacts between ESC-derived neurons and IHCs, we carried out TEM analysis of cryostat sections next to those containing the nerve endings of ESC-derived neurons expressing both EGFP and synaptophysin (Fig. 8A). EGFP expression was visualized by DAB staining, which was identified as electronically dense regions (Fig. 8B). High magnification revealed vesiculated nerve endings of ESC-derived neurons and their contacts with IHCs (Fig. 8C) as well as a parallel apposition of the two membranes with electronically dense material (Fig. 8D) showing the existence of




Fig. 4. Synapsin 1 expression in the nerve ending of embryonic stem cell-derived neurons. A–C: Low-magnification photographs of coculture with ESC-derived neurons and auditory epithelia on 7 days in vitro (DIV). A'–C': High magnification of the boxes shown in A–C, respectively. **A**: ESC-derived neurons labeled with EGFP (green) elongate their neurites toward IHCs (red). **A'**: ESC-derived nerve endings attach to the base of IHCs (arrow). **B, B'**: The expression of

synapsin 1 (blue) is accumulated in the nerve endings (arrow), not nerve fibers and soma of ESC-derived neurons. **C, C'**: Merged image. The expressions of EGFP (green) and synapsin 1 (blue) overlap in the nerve endings of ESC-derived neurons (arrow). EGFP, enhanced green fluorescent protein; Myo, myosin VIIa; IHCs, inner hair cells; OHCs, outer hair cells. Scale bars = 50 μ m in A–C; 10 μ m in A'–C'.

synaptic contacts. However, synapse ribbons, which are included in characteristics for afferent synaptic contacts between IHCs and SGN dendrites (Sobkowicz, 1992; Sobkowicz et al., 1997), were not observed in the base of IHCs attaching to ESC-derived nerve endings.

DISCUSSION

Our ultimate goal is to regenerate auditory HCs and SGNs and improve functional auditory systems after the degeneration of inner ears. Previous studies on human temporal bones have indicated that the loss of SGNs can cause SNHL (Schuknecht, 1993; Nelson and Hinojosa, 2001), and other studies have shown that cell transplantation is a realistic potential strategy for the replacement of SGNs (Naito et al., 2004; Tamura et al., 2004; Hu et al., 2005a,b; Okano et al., 2005; Corrales et al., 2006; Matsuoka et al., 2006).

In regeneration of the auditory afferent system, the formation of new synaptic contacts between IHCs and regenerated SGNs is crucial for its functionality. To investigate this, we have established an explant culture model in which SGNs are selectively eliminated by

using fine forceps and HCs are well maintained (Matsumoto et al., 2005). The procedure causes no damage to auditory epithelia and does not disrupt the arrangement of HC rows. By contrast, alternative elimination methods, such as the use of β -bungarotoxin that binds the potassium channels, have the potential to destroy HCs (Fuchs, 1996; Herkert et al., 2001; Shakhman et al., 2003; Martinez-Monedero et al., 2006). Our mechanical elimination method is preferable for in vitro experiments that examine the activity of engrafted cells for neurite outgrowth and the formation of new synaptic contacts with HCs.

The present findings demonstrate that ESC-derived neurons located at the corresponding position of SGNs used appropriate routes to extend their neurites, which were correctly terminated at the base of HCs. In the current study, ESC-derived neural progenitors were cocultured with explants of auditory epithelia just after denervation. Thus, the routes in which nerve fibers of SGNs had existed may not be scarred at the time point for starting coculture. In addition, previous studies have indicated that HCs have the potential to induce neurite outgrowth from neurons (Kawamoto et al., 2003;




Fig. 5. Synaptophysin expression in the nerve ending of embryonic stem cell-derived neurons. A–C: Low-magnification photographs of coculture with ESC-derived neurons and auditory epithelia on 7 days in vitro (DIV). A'–C': High magnification of the boxed shown in A–C, respectively. A: ESC-derived neurons labeled with EGFP (green) elongate their neurites toward IHCs (red). A': ESC-derived nerve endings attach to the base of IHCs (arrow). B, B': Synaptophysin

immunoreactivity (blue) is observed in the nerve endings of ESC-derived neurons adjacent to the base of IHCs (arrow) and in the cytoplasm of IHCs, not nerve fibers and soma of ESC-derived neurons. C, C': Merged image. The expressions of EGFP (green) and synaptophysin (blue) overlap in the nerve endings of ESC-derived neurons (arrow). EGFP, enhanced green fluorescent protein; Myo, myosin VIIa. Scale bars = 50 μ m in A–C; 10 μ m in A'–C'.

Kondo et al., 2005; Matsumoto et al., 2005; Martinez-Monedero et al., 2006). Therefore, ESC-derived neurons may elongate neurites through the route taken by host auditory nerves under the guidance of IHCs. However, the details of HC guidance cues are still unclear and will have to be elucidated prior to the development of cell therapy for functional restoration of the auditory system.

Immunohistochemical analyses in the present study revealed the active formation of synaptic contacts between ESC-derived neurons and IHCs. The distribution of synapsin 1 immunoreactivity was identical to that in the auditory epithelia of P12 mice, at the onset of hearing, and showed an accumulation in the nerve endings of ESC-derived neurons adjacent to IHCs. Moreover, the distribution of synaptophysin immunoreactivity in cocultured specimens was similar to that of P4 mice. In addition, the nerve endings and neurites of ESC-derived neurons exhibited GAP43 immunoreactivity, indicating active neurite outgrowth and synaptic sprout-

ing of ESC-derived neurons (Knipper et al., 1995). Immunohistochemistry for *ctbp2/ribeye* indicated the presence of synaptic ribbons in the basal pole of IHCs adjacent to ESC-derived nerve endings, similar to immature synaptic contacts between IHCs and afferent nerve fibers (Sobkowicz, 1992). Such immunohistochemical findings suggest active synapse formation between ESC-derived neurons and IHCs.

In the present study, TEM analysis revealed contacts between vesiculated nerve endings from ESC-derived neurons and IHC membranes, as well as synaptic densities in the membrane between IHCs and nerve endings of ESC-derived neurons. This represents the first evidence of synapse formation between ESC-derived neurons and IHCs. In normal cochleae of mature mice, afferent synapse contacts at the base of IHCs are characterized by the presence of synaptic ribbons and by a postsynaptic membrane density on the dendrite (Sobkowicz, 1992; Sobkowicz et al., 1997). However, in TEM analysis, no synaptic ribbons were found in the base of




Fig. 6. GAP43 expression in the nerve ending of embryonic stem cell-derived neurons. A-C: Low-magnification photographs of coculture with ESC-derived neurons and auditory epithelia on 7 days in vitro (DIV). A'-C': High magnification of the boxes shown in A-C, respectively. **A**: ESC-derived neurons labeled with EGFP (green) elongate their neurites toward IHCs (red). **A'**: ESC-derived nerve endings attach to the base of IHCs (arrows). **B, B'**: GAP43 immuno-

reactivity (blue) is found in the nerve endings of ESC-derived neurons adjacent to the base of IHCs (arrows) and the nerve fibers. **C, C'**: Merged image. The expressions of EGFP (green) and GAP43 (blue) overlap in the nerve endings of ESC-derived neurons (arrows) and the nerve fibers. EGFP, enhanced green fluorescent protein; Myo, myosin VIIa. Scale bars = 50 μm in A-C; 10 μm in A'-C'.




Fig. 7. Expression of *ctbp2/ribeye* and *calretinin* in adult mouse cochleae and cocultured specimens. **A-C**: In the adult mouse cochlea, *ctbp2/ribeye* expression (blue) is found in the nuclei and the basal pole of the IHC (arrows), and *calretinin* expression (red) is observed in hair cells and afferent nerve fibers. **D-F**: In cocultured

specimens, immunoreactivity for *ctbp2/ribeye* is found in the nucleus and in the basal pole of the IHC (arrow), and ESC-derived neurites expressing EGFP (green) are negative for *calretinin*. On the other hand, the IHC is positive for *calretinin*. Scale bars = 10 μm.




Fig. 8. Ultrastructure of contacts between inner hair cells and nerve endings of embryonic stem cell-derived neurons. **A:** Synaptophysin-positive nerve endings (blue) of ESC-derived neurons expressing EGFP (green) attach to the basal portion of the IHC (red; arrows). **B:** An EGFP-positive nerve ending visualized by DAB staining, which is identified as electronically dense regions, is found at the base of the IHC (arrows). **C:** A high magnification of the box shown in B demonstrates the nerve ending of ESC-derived neurons as a ves-

iculated nerve ending (arrows) and their contact with the membrane of the IHC (arrowheads). **D:** In a higher magnification of the box shown in C, this region exhibits a parallel apposition of the two membranes, with an electronically dense material between the nerve endings of ESC-derived neurons and the IHC (arrowheads). EGFP, enhanced green fluorescent protein; Myo, myosin VIIa; IHC, inner hair cell. Scale bars = 10 μm in A,B; 0.1 μm in C,D.

IHCs adjacent to ESC-derived nerve endings. Therefore, the demonstration that the ECS-derived neurons may be used for replacement of SGNs requires additional experiments.

The neurites of ESC-derived neurons adjacent to IHCs share some properties of presynaptic fibers, because synapsin 1 and synaptophysin are markers for small synaptic vesicles, which are usually located in the presynaptic compartments. In addition, the neurites of ESC-derived neurons were negative for calretinin, which is expressed in mature SGN and their neurites (Dechesne et al., 1994). However, previous studies have observed expression of synapsin 1 and synaptophysin in afferent growth cones (De Camilli et al., 1988). Immature afferent dendrites of SGNs are lack of calretinin expression

(Dechesne et al., 1994). Moreover, recent findings using cocultures of auditory epithelia explants and primary SGN cultures found that the processes from SGNs adjacent to HCs express markers for presynaptic fibers (Martinez-Monedero et al., 2006). In addition, coincident with synaptophysin and GAP-43 immunoreactivity in SGNs, outer spiral bundle fibers (OSB) exhibit immunoreactivity for synaptophysin on P2 but not P4 (Knipper et al., 1995), suggesting the localization of synaptophysin in the afferent growth cones of immature SGNs. Morphologically, immature efferent fibers are practically indistinguishable from those of afferent processes (Sobkowicz, 1992). Therefore, determining the type of synaptic contacts between ESC-derived neurons and IHCs will require further examinations.

CONCLUSIONS

The present findings demonstrate that ESC-derived neurons project neurites toward HCs in the auditory epithelium and have the ability to form new synaptic contacts with IHCs. In future experiments, we will investigate the maturation process of ESC-derived neurons after the formation of synaptic contacts with HCs and elucidate HC-derived guidance cues for neurite projection.

ACKNOWLEDGMENTS

The authors thank Marlies Knipper, Takayuki Okano, Yayoi S. Kikkawa, and Tetsuji Sekiya for critical reviews of this work; Hitoshi Niwa for providing mouse ESCs; Yoshiki Sasai for providing PA6 cells; and Makio Fujioka, Akemi Saito, Yoko Nishiyama, and Tomoyo Namura for their technical expertise.

REFERENCES

- Coleman B, Fallon JB, Pettingill LN, de Silva MG, Shepherd RK. 2007. Auditory hair cell explant co-cultures promote the differentiation of stem cells into bipolar neurons. *Exp Cell Res* 313:232–243.
- Corrales CE, Pan L, Li H, Liberman MC, Heller S, Edge AS. 2006. Engraftment and differentiation of embryonic stem cell-derived neural progenitor cells in the cochlear nerve trunk: growth of processes into the organ of Corti. *J Neurobiol* 66:1489–1500.
- De Camilli P, Cameron R, Greengard P. 1983a. Synapsin I (protein I), a nerve terminal-specific phosphoprotein. I. Its general distribution in synapses of the central and peripheral nervous system demonstrated by immunofluorescence in frozen and plastic sections. *J Cell Biol* 96:1337–1354.
- De Camilli P, Harris SM Jr, Huttner WB, Greengard P. 1983b. Synapsin I (protein I), a nerve terminal-specific phosphoprotein. II. Its specific association with synaptic vesicles demonstrated by immunocytochemistry in agarose-embedded synaptosomes. *J Cell Biol* 96:1355–1373.
- De Camilli P, Vitadello M, Canevini MP, Zaroni R, Jahn R, Gorio A. 1988. The synaptic vesicle proteins synapsin I and synaptophysin (protein P38) are concentrated both in efferent and afferent nerve endings of the skeletal muscle. *J Neurosci* 8:1625–1631.
- Dechesne CJ, Rabejac D, Desmadryl G. 1994. Development of calretinin immunoreactivity in the mouse inner ear. *J Comp Neurol* 346:517–529.
- Fuchs PA. 1996. Synaptic transmission at vertebrate hair cells. *Curr Opin Neurobiol* 6:514–519.
- Herkert M, Shakhman O, Schweins E, Becker CM. 2001. Beta-bungarotoxin is a potent inducer of apoptosis in cultured rat neurons by receptor-mediated internalization. *Eur J Neurosci* 14:821–828.
- Hooper M, Hardy K, Handyside A, Hunter S, Monk M. 1987. HPR T-deficient (Lesch-Nyhan) mouse embryos derived from germline colonization by cultured cells. *Nature* 326:292–295.
- Hu Z, Ulfendahl M, Olivius NP. 2005a. NGF stimulates extensive neurite outgrowth from implanted dorsal root ganglion neurons following transplantation into the adult rat inner ear. *Neurobiol Dis* 18:184–192.
- Hu Z, Wei D, Johansson CB, Holmstrom N, Duan M, Frisen J, Ulfendahl M. 2005b. Survival and neural differentiation of adult neural stem cells transplanted into the mature inner ear. *Exp Cell Res* 302:40–47.
- Huttner WB, Schiebler W, Greengard P, De Camilli P. 1983. Synapsin I (protein I), a nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly purified synaptic vesicle preparation. *J Cell Biol* 96:1374–1388.
- Kawamoto K, Ishimoto S, Minoda R, Brough DE, Raphael Y. 2003. Math1 gene transfer generates new cochlear hair cells in mature guinea pigs in vivo. *J Neurosci* 23:4395–4400.
- Kawasaki H, Mizuseki K, Nishikawa S, Kaneko S, Kuwana Y, Nakanishi S, Nishikawa SI, Sasai Y. 2000. Induction of midbrain dopaminergic neurons from ES cells by stromal cell-derived inducing activity. *Neuron* 28:31–40.
- Keithley EM, Ryan AF, Woolf NK. 1989. Spiral ganglion cell density in young and old gerbils. *Hear Res* 38:125–133.
- Kim TS, Nakagawa T, Kita T, Higashi T, Takebayashi S, Matsumoto M, Kojima K, Sakamoto T, Ito J. 2005. Neural connections between embryonic stem cell-derived neurons and vestibular hair cells in vitro. *Brain Res* 1057:127–133.
- Knipper M, Zimmermann U, Rohbock K, Kopschall I, Zenner HP. 1995. Synaptophysin and GAP-43 proteins in efferent fibers of the inner ear during postnatal development. *Brain Res Dev Brain Res* 89:73–86.
- Kondo T, Johnson SA, Yoder MC, Romand R, Hashino E. 2005. Sonic hedgehog and retinoic acid synergistically promote sensory fate specification from bone marrow-derived pluripotent stem cells. *Proc Natl Acad Sci U S A* 102:4789–4794.
- Lee JE, Nakagawa T, Kim TS, Iguchi F, Endo T, Dong Y, Yuki K, Naito Y, Lee SH, Ito J. 2003. A novel model for rapid induction of apoptosis in spiral ganglions of mice. *Laryngoscope* 113:949–999.
- Martinez-Monedero R, Corrales CE, Cuajungco MP, Heller S, Edge AS. 2006. Reinnervation of hair cells by auditory neurons after selective removal of spiral ganglion neurons. *J Neurobiol* 66:319–331.
- Matsumoto M, Nakagawa T, Higashi T, Kim TS, Kojima K, Kita T, Sakamoto T, Ito J. 2005. Innervation of stem cell-derived neurons into auditory epithelia of mice. *Neuroreport* 16:787–790.
- Matsuoka AJ, Kondo T, Miyamoto RT, Hashino E. 2006. In vivo and in vitro characterization of bone marrow-derived stem cells in the cochlea. *Laryngoscope* 116:1363–1367.
- Meiri KF, Pfenninger KH, Willard MB. 1986. Growth-associated protein, GAP-43, a polypeptide that is induced when neurons extend axons, is a component of growth cones and corresponds to pp46, a major polypeptide of a subcellular fraction enriched in growth cones. *Proc Natl Acad Sci U S A* 83:3537–3541.
- Nadol JB Jr, Young YS, Glynn RJ. 1989. Survival of spiral ganglion cells in profound sensorineural hearing loss: implications for cochlear implantation. *Ann Otol Rhinol Laryngol* 98:411–416.
- Naito Y, Nakamura T, Nakagawa T, Iguchi F, Endo T, Fujino K, Kim TS, Hiratsuka Y, Tamura T, Kanemaru S, Shimizu Y, Ito J. 2004. Transplantation of bone marrow stromal cells into the cochlea of chinchillas. *Neuroreport* 15:1–4.
- Navone F, Greengard P, De Camilli P. 1984. Synapsin I in nerve terminals: selective association with small synaptic vesicles. *Science* 226:1209–1211.
- Navone F, Jahn R, Di Gioia G, Stukenbrok H, Greengard P, De Camilli P. 1986. Protein p38: an integral membrane protein specific for small vesicles of neurons and neuroendocrine cells. *J Cell Biol* 103:2511–2527.
- Nelson EG, Hinojosa R. 2001. Aplasia of the cochlear nerve: a temporal bone study. *Otol Neurotol* 22:790–795.
- Okano T, Nakagawa T, Endo T, Kim TS, Kita T, Tamura T, Matsumoto M, Ohno T, Sakamoto T, Iguchi F, Ito J. 2005. Engraftment of embryonic stem cell-derived neurons into the cochlear modiolus. *Neuroreport* 16:1919–1922.
- Pujol R, Puel JL. 1999. Excitotoxicity, synaptic repair, and functional recovery in the mammalian cochlea: a review of recent findings. *Ann N Y Acad Sci* 884:249–254.
- Rubel EW, Fritsch B. 2002. Auditory system development: primary auditory neurons and their targets. *Annu Rev Neurosci* 25:51–101.
- Schuknecht HF. 1993. Pathology of the ear, 2nd ed. Boston: Lea & Febiger. 672 p.

- Shakhman O, Herkert M, Rose C, Humeny A, Becker CM. 2003. Induction by beta-bungarotoxin of apoptosis in cultured hippocampal neurons is mediated by Ca^{2+} -dependent formation of reactive oxygen species. *J Neurochem* 87:598–608.
- Skene JH. 1989. Axonal growth-associated proteins. *Annu Rev Neurosci* 12:127–156.
- Skene JH, Jacobson RD, Snipes GJ, McGuire CB, Norden JJ, Freeman JA. 1986. A protein induced during nerve growth (GAP-43) is a major component of growth-cone membranes. *Science* 233:783–786.
- Sobkowicz HM. 1992. The development of innervation in the organ of Corti. In: Romand R, editor. *Development of auditory and vestibular systems II*. Amsterdam: Elsevier. p 59–100.
- Sobkowicz HM, Slapnick SM, Nitecka LM, August BK. 1997. Compound synapses within the GABAergic innervation of the auditory inner hair cells in the adolescent mouse. *J Comp Neurol* 377:423–442.
- Sobkowicz HM, Slapnick SM, August BK. 2002. Differentiation of spinous synapses in the mouse organ of corti. *Synapse* 45:10–24.
- Spoendlin H. 1987. The afferent innervation of the cochlea. In: Naunton RF, Fernandez C, editors. *Evoked electrical activity in the auditory nervous system*. New York: Academic Press. p 21–41.
- Sudhof TC, Lottspeich F, Greengard P, Mehl E, Jahn R. 1987. A synaptic vesicle protein with a novel cytoplasmic domain and four transmembrane regions. *Science* 238:1142–1144.
- Tamura T, Nakagawa T, Iguchi F, Tateya I, Endo T, Kim TS, Dong Y, Kita T, Kojima K, Naito Y, Omori K, Ito J. 2004. Transplantation of neural stem cells into the modiolus of mouse cochleae injured by cisplatin. *Acta Otolaryngol Suppl* 551:65–68.
- Wiedenmann B, Franke WW. 1985. Identification and localization of synaptophysin, an integral membrane glycoprotein of Mr 38,000 characteristic of presynaptic vesicles. *Cell* 41:1017–1028.

Local Drug Delivery to Inner Ear for Treatment of Hearing Loss

Takayuki Nakagawa* and Juichi Ito

Department of Otolaryngology, Head and Neck Surgery, Graduate School of Medicine, Kyoto University, Kyoto, Japan

Abstract: Sensorineural hearing loss (SNHL) is one of the most common disabilities in our society. Experimentally, many candidates for use as therapeutic molecules have been discovered. However, a considerable obstacle to clinical application is the lack of an effective method for drug delivery to the cochlea. In order to overcome this obstacle, there needs to be development of a local cochlear drug delivery system. Advances in pharmacological technology have provided various drug delivery systems that use biomaterials, and which can be utilized for local drug delivery to the cochlea. Indeed, recent studies have demonstrated the potential of synthetic and natural biomaterials for local drug delivery to the cochlea, indicating that the clinical application of such local drug delivery systems could be used in the near future for therapeutic treatments. Recent progress in cell therapy research also offers a novel drug delivery method for the cochlea. In addition, transplantation of stem cells into the cochlea has been demonstrated to provide protective effects for the auditory function. Transplantation of genetically engineered cells has also resulted in the sustained delivery of aimed therapeutic molecules within the inner ear. Although problems involving clinical application still need to be resolved, these drug delivery systems for the inner ear may hold the future therapeutic options for treatment of SNHL.

Key Words: Drug delivery system, cochlea, biodegradable polymer, cell transplantation, gene therapy.

THERAPEUTIC TARGETS FOR TREATMENT OF HEARING LOSS

Sensorineural hearing loss (SNHL) is one of the most prevalent disabilities in our society. Sound stimuli are received by auditory hair cells (HCs) in the bony, snail-shaped cochlea, followed by transduction of the sound stimuli by the HCs to neural signals. Spiral ganglion neurons (SGNs), which are auditory primary neurons, are located in the central bony axis of the cochlea and responsible for transmitting auditory signals to the central auditory system. Excessive noise, ototoxic drugs, genetic disorders and aging all contribute to the causes of SNHL. Severe to profound SNHL affects 1 in 1000 newborns, and another 1 in 2000 children before they reach adulthood. About 60% of individuals older than 70 years will manifest SNHL. Previous studies on human temporal bones have demonstrated that the loss of HCs and/or SGNs is a major cause of SNHL [1]. Protecting HCs and SGNs from irreversible degeneration is therefore a primary objective due to the limited regeneration capacity of these cells. Acute SNHL sometimes responds to drug treatment; however, there are no therapeutic options for chronic SNHL except for hearing aids and cochlear implants, which are small devices that are surgically implanted into the cochlea in order to stimulate SGNs. However, the success of cochlear implants depends on the remaining SGNs and with their loss, this severely compromises the efficacy of this technique. HCs and SGNs are therefore the major targets for the treatment of SNHL.

WHY IS LOCAL DRUG DELIVERY REQUIRED FOR THE INNER EAR?

Based on the backgrounds described above, studies are being conducted with the hopes of providing an alternative

means of biological therapy. Thus far, research has identified a number of candidates for use as therapeutic molecules. Experimentally, protective effects of neurotrophins have been demonstrated [2,3], and inhibitors of apoptosis and glutamate antagonists have also been shown to have the ability to promote HC survival [4-6]. Recently, it has been found that local application of genes by virus vectors induces HC regeneration in the mammalian auditory epithelium [7,8], and additionally, by silencing the mutant gene *via* RNA interference, can restore hearing loss in the genetic mouse model [9]. These therapeutic strategies are attractive and promising for the restoration of SNHL. However, clinical application is still quite limited. The problem of how to deliver such therapeutic molecules to the inner ear has been a considerable obstacle in the development of treatments for SNHL. One of the reasons for the difficulty of drug delivery involves the limited blood flow to the cochlea [10]. In addition, the blood-inner ear barrier, which inhibits the transport of drugs from serum to the inner ear, represents a fundamental obstacle to the use of systemic applications [11]. The inner ear tissues are isolated from the surrounding organs by a bony construction, which allows for the topical introduction of drugs or genes. Based on these considerations, local application has generally been the preferred method for drug administration to the inner ear. The sustained delivery of therapeutic molecules is also critical for the efficient treatment of the cochlea, as bioactive molecules usually require a period of minutes or hours over which they produce their pharmacological actions. Consequently, a number of researchers are currently working to solve these problems and develop methods for the local direct application of these molecules into the cochlea [12].

STRATEGIES FOR LOCAL DRUG DELIVERY

The cochlea is connected to the tympanic cavity by the round window membrane (RWM). When substances are applied intratympanically, the assumption is that they will enter the scala tympani through the RWM and then be dis-

*Address correspondence to this author at the Department of Otolaryngology-Head and Neck Surgery, Graduate School of Medicine, Kyoto University, Kawaharacho 54, Shogoin, Sakyo-ku, 606-8507 Kyoto, Japan; Tel: +81-75-751-3346; Fax: +81-75-751-7225; E-mail: tnakagawa@ent.kuhp.kyoto-u.ac.jp

tributed throughout the cochlear fluids. The idea of using a topical application of medicine to the inner ear is not new, as local anesthetics and aminoglycosides were applied decades ago, with the compounds passing through the tympanic membrane into the tympanic cavity during the treatment of the inner ear disorders [13,14]. Intratympanic injections have been used for local application of aminoglycosides or steroids during therapy for Ménière's disease and sudden hearing loss. There are a number of clinical reports showing the efficacy of intratympanic injections of these drugs [see review in reference 12]. However, it is very difficult to predict the amounts of drugs that will actually reach the cochlear fluid space. Some reports have indicated that this method can lead to varying results during therapeutic treatment of Ménière's disease [15-17]. While intratympanic injection is a simple and easy method to perform, unfortunately, a controlled and sustained release of drugs cannot be achieved using this method. The pharmacokinetics of drug entry into cochlear fluids is crucial to determine the efficacy of the method for drug delivery into cochlear fluids [12]. Salt and Plontke have indicated importance of sustained delivery of drugs on the RWM by means of perilymph sampling from various regions of the cochlea [18] and computer simulation [19].

Implantable mini-pumps have also frequently been used for local drug delivery to the cochlea in animal experiments [20]. Several clinical reports have described the efficacy of local glucocorticoid application when using a semi-implantable mini-pump [21,22]. However, the use of an implantable mini-pump has not been widely adopted, given the need for surgical procedures similar to tympanoplasty that must be done in order to place the mini-pump. The use of a local viral gene transfer as a sustained treatment of the inner ear can provide sufficient protection from noise, drug toxicity and re-perfusion injury [23-28]. Today, adenoviral vectors or adeno-associated viral vectors are the most widely used for cochlear gene transfer, because of the high efficiency for the transfection, the availability of high titers, and the ease of production. However, their use can potentially initiate an immune response that results in the destruction of the recipient's cochlear cells.

The use of biomaterials for local drug delivery has recently gained attention as an alternative to the implantable mini-pumps or gene transfer using virus vectors. In general, biodegradable polymers containing therapeutic molecules are placed on the RWM, with the therapeutic molecules released into the cochlear fluids from the polymers in a controlled manner *via* the RWM [12,28].

DRUG DELIVERY VIA BIOMATERIALS

In the past decade, pharmaceutical technologists have paid increasing attention to controlled or sustained release technology using biomaterials for the delivery of drugs in order to avoid side effects and achieve sufficient drug levels in tissues. In an effort to develop a controlled-release system, a variety of methods using synthetic and natural materials have been undertaken. Recent publications have reported the use of a controlled-release system for local drug delivery to the inner ear. Two synthetic materials, siloxane-based polymers [29] and polylactic/glycolic acid (PLGA) polymers

[30], and several natural materials, which include hyaluronic acid [31] and gelatin [32-34], have been used for this purpose.

Siloxane-based polymers have been used for years in medical applications that involve contact with the human body. In the clinic, silicone-transdermal patches have been widely used. In this system, drug release is controlled by its diffusion through the silicone network [35]. The actual release rate is determined by the composition of the polymer. This system is particularly suitable for application of lipophilic and low-molecular weight molecules. Arnold *et al.* [29] have utilized this system for local application of beclomethasone into the cochlear fluids. When using this system, the silicone microimplant remains on the RWM, although it does not induce functional and histological damage in the cochlea. Therefore, repeated treatments require that there is extirpation of the material used during the procedure.

Encapsulating bioactive molecules in PLGA or polylactic acid (PLA) particles has been also used as a method of controlled-release application. Water-insoluble, low-molecular weight agents have been encapsulated in PLGA or PLA microparticles and nanoparticles [36,37]. PLGA and PLA are familiar substances to surgeons, as they are the materials that make up absorbable sutures. Tamura *et al.* [30] examined the potential of PLGA nanoparticles for drug delivery to the cochlea using guinea pigs. To evaluate the use of PLGA nanoparticles (140 to 180 nm in diameter) in the cochlea, rhodamine, which is a red fluorescent dye, was encapsulated and then following local application onto the RWM, its overall distribution was evaluated. PLGA nanoparticles containing rhodamine were observed in the cochlea, indicating that PLGA nanoparticles can penetrate through the RWM. Rhodamine is released from PLGA nanoparticles after penetration of the particles through the RWM. Compared to a silicone microimplant, PLGA nanoparticles have the advantage of being able to be repeatedly applied, as the PLGA is dissolved by hydrolysis. However, there is a limitation with regard to the variation of the drugs that can be applied, since the process of encapsulation in the PLGA particles requires that compounds must be dissolved in acetone. Therefore, this method is not suitable for the delivery of proteins or peptides.

GELATIN HYDROGEL

Gelatin is a commonly used natural polymer that is derived from collagen. In the clinic, gelatin polymers have been widely used as hemostats. Recently, gelatin-based controlled-release systems have been developed [38]. During the fabrication process, the isoelectric point of gelatin can be modified to yield either a negatively charged acidic gelatin or a positively charged basic gelatin. This allows for electrostatic interactions to take place between charged therapeutic molecules and gelatin of the opposite charge, leading to the formation of polyion complexes. The significance of such a system is that it provides the ability for application of water-soluble, comparatively high-molecular weight proteins and peptides. Additionally, this method is also capable of being used for the delivery of plasmid DNA [39]. In this system, therapeutic molecules are released by enzymatic degradation of gelatin polymers, for which the rates can be determined based on the crosslinking density of the gelatin polymers.

The potential use of the gelatin hydrogel system was initially investigated for cochlear delivery of brain-derived neurotrophic factor (BDNF) [32]. BDNF plays a crucial role in the development of the inner ears [38] and in the maintenance of the auditory function [41]. In addition, previous studies have demonstrated the effects of local BDNF application when using an osmotic mini pump [3] or adenovirus [26]. We measured BDNF concentrations in the cochlear fluid after placing a gelatin hydrogel that contained this agent onto the RWM [32]. The results revealed a sustained delivery of BDNF into the cochlear fluid *via* the hydrogel over a seven-day period. The functional and histological protection of the SGNs by BDNF that was applied through the gelatin hydrogel was then examined using a guinea pig model of SGN degeneration. The measurement of electrically evoked auditory-brainstem responses, which reflect SGN function, demonstrated that BDNF delivered *via* gelatin hydrogels was able to significantly reduce the threshold elevation [32]. Histological analysis demonstrated an increased survival of SGNs due to BDNF application through gelatin hydrogels. These findings indicate that gelatin hydrogel can be utilized for drug delivery to the cochlea.

Subsequently, we examined the efficacy of cochlear delivery of insulin-like growth factor-1 (IGF1) for the protection of auditory HCs against acoustic trauma [33]. IGF1 is a mitogenic peptide that plays essential roles in the regulation of growth and development in the inner ear [42]. In addition, previous studies on the inner ear have suggested the possibility of inner ear protection by IGF1 [43,44]. Moreover, recombinant human IGF-1 (rhIGF1) has already been approved for clinical use. Therefore, we selected rhIGF1 as a suitable trophic factor for local cochlear application using a gelatin hydrogel. Local rhIGF1 application through the gelatin hydrogel prior to noise exposure has been shown to efficiently protect the hearing from noise trauma. Additionally, histological analysis also revealed that local rhIGF-1 treatment ameliorated the loss of HCs [33].

Our ultimate goal is the clinical use of a local rhIGF1 application using gelatin hydrogel as a therapeutic option for the treatment of SNHL. Therefore, we examined whether post-traumatic application of rhIGF1 to the cochlea *via* gelatin hydrogels could attenuate noise-induced hearing loss. The results demonstrated that functional and histological efficacy of local rhIGF1 treatment on the attenuation of noise-induced hearing loss occurred in a dose-dependent manner [34]. We also measured IGF1 concentrations in the cochlear fluid, cerebrospinal fluid (CSF) and serum after placing a gelatin hydrogel containing rhIGF1 onto the RWM of guinea pigs. The results demonstrated that there was sustained delivery of rhIGF1 into the cochlear fluid, in addition to no alterations of the IGF1 levels in CSF and serum [34]. There were also no adverse effects due to local rhIGF1 treatment found in any of the experimental animals. These findings document both the effectiveness and the safety of local rhIGF1 treatment using gelatin hydrogels for noise-induced hearing loss.

CELL TRANSPLANTATION

Chronic SNHL is usually incurable because of the loss of HCs and SGNs, and which at the present time is irreversible.

Therefore, an alternative means of biological therapy, including cell therapy is required. Indeed, recent studies have indicated that cell therapy could be utilized to regenerate HCs [45] and SGNs [46]. In contrast, cell transplantation is an alternative that can be used as a method for drug delivery where the transplanted cells for this purpose have the ability to survive and generate therapeutic agents. Several stem cells have been reported to have the ability to secrete trophic factors [47-49]. Cell transplantation has been used as a means of delivering peptides or proteins into the central nervous system, demonstrating its viable use as a delivery vehicle for therapeutic molecules [50,51].

Iguchi *et al.* have reported on the ability of neural stem cell-derived cells being used for the production of BDNF and glial cell line-derived neurotrophic factor (GDNF) after engraftment into the cochlea [47]. In addition, transplantation of neural stem cells into the cochlea has the potential of being able to attenuate HC damages due to transient ischemia of the cochlea [48]. Bone marrow derived cells also have the potential for secreting trophic factors. Implantation of bone marrow stromal cells has been reported to contribute to functional recovery of the brain [52] and spinal cord [53] by means of producing trophic factors. Furthermore, previous studies have revealed the potential of bone marrow derived cells surviving in the cochlea [54,55]. Yoshida *et al.* have demonstrated a significant increase in the protein level of GDNF in cochlear specimens and the prevention of HC death due to transient cochlear ischemia by engraftment of hematopoietic stem cells [49]. These findings indicate that cell transplantation into the cochlea may be a novel strategy for treatment of SNHL by providing a means for local application of trophic factors within the cochlea.

Transplantation of cells that have been genetically manipulated *ex vivo* has been used as a means of delivering peptides or proteins into the central nervous system [56-58]. In comparison with the stem cell transplantation that has been described above, this strategy has an advantage in that aimed gene-encoded products are applicable. In addition, use of non-viral vectors for *ex vivo* gene transfer potentially could resolve the problem of viral vector toxicity in cochlear gene therapy. Therefore, we conducted an examination of the efficacy of cell-gene delivery in the application of therapeutic molecules into the cochlea [59]. NIH3T3 cells were chosen as a delivery vehicle for the gene. NIH3T3 cells are a well-established fibroblast cell line, thus, it is easy to optimize conditions for gene transfer and to select gene-expressing cells for use *in vitro*. In addition, such fibroblasts are available from various human sources, which may be advantageous for extending future clinical investigations. NIH3T3 cells were transfected with the BDNF gene using lipofection, with the cells expressing the BDNF gene being selected for use. We examined the potential for transplanting transfected NIH3T3 cells into the inner ear of the mouse. Immunohistochemistry and Western blotting demonstrated the survival of the grafted cells within the cochlea, and a BDNF-specific enzyme-linked immunosorbent assay revealed a significant increase in BDNF production in the inner ear following cell transplantations [59]. These findings indicate that cell-gene delivery with non-viral vectors may be applicable for the local, sustained delivery of therapeutic

molecules into the cochlea. Cell-gene delivery of therapeutic molecules into the inner ear is suitable for protection of inner ear cells against gradually progressive degeneration. Presbycusis, which is an age-related hearing loss, may also need to be included as one of the targets for cell-gene therapy. BDNF application *via* cell-gene delivery could be an effective strategy for survival promotion of SGNs in cases involving cochlear implants, which require the opening of the cochlea for the purpose of inserting an electrode.

CONCLUSIONS

The lack of effective methods for drug delivery to the cochlea has been a considerable obstacle with regard to developing novel therapeutic strategies for SNHL. However, recent findings in studies examining drug delivery systems using biomaterials and cell therapy demonstrate the efficacy of these strategies for cochlear drug delivery, which in the future may contribute to the establishment of novel therapeutic strategies for SNHL.

ACKNOWLEDGEMENTS

This study was supported by a Grant-in-Aid for Regenerative Medicine Realization, a Grant-in-Aid for Scientific Research and a Grant from the 21st Century COE program from the Ministry of Education, Science, Sports, Culture and Technology of Japan, and by a Grant-in-Aid for Researches on Sensory and Communicative Disorders from the Japanese Ministry of Health, Labour and Welfare.

REFERENCES

- [1] Schuknecht HF. Pathology of the ear. Cambridge, MA: Harvard University press; 1974.
- [2] Miller JM, Chi DH, O'Keefe LJ, Kruszka P, Raphael Y, Altschuler RA. Neurotrophins can enhance spiral ganglion cell survival after inner hair cell loss. *Int J Dev Neurosci* 1997; 15: 631-43.
- [3] Shinohara T, Bredberg G, Ulfendahl M, *et al.* Neurotrophic factor intervention restores auditory function in deafened animals. *Proc Natl Acad Sci USA* 2002; 99: 1657-60.
- [4] Nakagawa T, Kim TS, Murai N, *et al.* A novel technique for inducing local-inner ear damage. *Hear Res* 2003; 176: 122-7.
- [5] Cunningham LL, Cheng AG, Rubel EW. Caspase activation in hair cells of the mouse utricle exposed to neomycin. *J Neurosci* 2002; 22: 8532-40.
- [6] Duan ML, Ulfendahl M, Laurell G, *et al.* Protection and treatment of sensorineural hearing disorders caused by exogenous factors: experimental findings and potential clinical application. *Hear Res* 2002; 169: 169-78.
- [7] Kawamoto K, Ishimoto S, Minoda R, Brough DE, Raphael Y. Math1 gene transfer generates new cochlear hair cells in mature guinea pigs *in vivo*. *J Neurosci* 2003; 23: 4395-400.
- [8] Izumikawa M, Minoda R, Kawamoto K, *et al.* Auditory hair cell replacement and hearing improvement by *Atoh1* gene therapy in deaf mammals. *Nat Med* 2005; 11: 271-6.
- [9] Maeda Y, Fukushima K, Nishizaki K, Smith RJ. *In vitro* and *in vivo* suppression of GJB2 expression by RNA interference. *Hum Mol Genet* 2005; 14: 1641-1650.
- [10] Angelborg C, Hillerdal M, Hulterantz E, Larsen HC. The microsphere method for studies of inner ear blood flow. *ORL J Otorhinolaryngol Relat Spec* 1998; 50: 355-62.
- [11] Juhn SK, Rybak LP. Labyrinthine barriers and cochlear homeostasis. *Acta Otolaryngol* 1981; 91: 529-534.
- [12] Salt AN, Plontke S. Local inner-ear drug delivery and pharmacokinetics. *Drug Discov Today* 2005; 10: 1299-1306.
- [13] Emsner MS. Transstympanic injection of anesthetics for the treatment of Menière's Syndrome. *Arch Otorhinolaryngol* 1951; 43-52.
- [14] Schuknecht HF. Ablation therapy for the relief of Meniere's disease. *Laryngoscope* 1956; 66: 859-70.
- [15] Lange G, Maurer J, Mann W. Long-term results after interval therapy with intratympanic gentamicin for Meniere's disease. *Laryngoscope* 2004; 114: 102-5.
- [16] Thomsen J, Charabi S, Tos M. Preliminary results of a new delivery system for gentamicin to the inner ear in patients with Meniere's disease. *Eur Arch Otorhinolaryngol* 2000; 257: 362-5.
- [17] Schoendorf J, Neugebauer P, Michel O. Continuous intratympanic infusion of gentamicin *via* a microcatheter in Meniere's disease. *Otolaryngol Head Neck Surg* 2001; 124: 203-207.
- [18] Salt AN, Hale SA, Plontke SK. Perilymph sampling from the cochlear apex: a reliable method to obtain higher purity perilymph samples from scala tympani. *J Neurosci Methods* 2006; 153: 121-9.
- [19] Plontke SK, Salt AN. Simulation of application strategies for local drug delivery to the inner ear. *ORL J Otorhinolaryngol Relat Spec* 2006; 68: 386-92.
- [20] Takemura K, Komeda M, Yagi M, *et al.* Direct inner ear infusion of dexamethasone attenuates noise-induced trauma in guinea pig. *Hear Res* 2004; 196: 58-68.
- [21] Lefebvre PP, Staecker H. Steroid perfusion of the inner ear for sudden sensorineural hearing loss after failure of conventional therapy: a pilot study. *Acta Otolaryngol* 2002; 122: 698-702.
- [22] Plontke S, Lowenheim H, Preyer S, *et al.* Outcomes research analysis of continuous intratympanic glucocorticoid delivery in patients with acute severe to profound hearing loss: basis for planning randomized controlled trials. *Acta Otolaryngol* 2005; 125: 830-39.
- [23] Yagi M, Magal E, Sheng Z, Ang KA, Raphael Y. Hair cell protection from aminoglycoside ototoxicity by adenovirus-mediated overexpression of glial cell line-derived neurotrophic factor. *Hum Gene Ther* 1999; 10: 813-23.
- [24] Staecker H, Li D, O'Malley Jr BW, Van De Water TR. Gene expression in the mammalian cochlea: a study of multiple vector systems. *Acta Otolaryngol* 2001; 121: 157-63.
- [25] Luebke AE, Foster PK, Muller CD, Peel AL. Cochlear function and transgene expression in the guinea pig cochlea, using adenovirus- and adeno-associated virus-directed gene transfer. *Hum Gene Ther* 2001; 12: 773-81.
- [26] Nakaizumi T, Kawamoto K, Minoda R, Raphael Y. Adenovirus-Mediated expression of brain-derived neurotrophic factor protects SGNs from ototoxic damage. *Audiol Neurootol* 2004; 9: 135-143.
- [27] Hakuba N, Watabe K, Hyodo J, *et al.* Adenovirus-mediated overexpression of a gene prevents hearing loss and progressive inner hair cell loss after transient cochlear ischemia in gerbils. *Gene Ther* 2003; 10: 426-33.
- [28] Nakagawa T, Ito J. Drug delivery systems for the treatment of sensorineural hearing loss. *Acta Otolaryngol* 2007; Suppl 557: 30-5.
- [29] Arnold W, Senn P, Hennig M, *et al.* Novel slow- and fast-type drug release round-window microimplants for local drug application to the cochlea: An experimental study in guinea pigs. *Audiol Neurootol* 2005; 10: 53-63.
- [30] Tamura T, Kita T, Nakagawa T, *et al.* Drug delivery to the cochlea using PLGA nanoparticles. *Laryngoscope* 2005; 115: 2000-5.
- [31] Wang J, Ruel J, Ladrech S, *et al.* Inhibition of the c-Jun N-terminal kinase-mediated mitochondrial cell death pathway restores auditory function in sound-exposed animals. *Mol Pharmacol* 2007; 71: 654-66.
- [32] Endo T, Nakagawa T, Kita T, *et al.* A novel strategy for treatment of inner ears using a biodegradable gel. *Laryngoscope* 2005; 115: 2000-5.
- [33] Iwai K, Nakagawa T, Endo T, *et al.* Cochlear protection by local IGF-I application using biodegradable hydrogel. *Laryngoscope* 2006; 116: 526-33.
- [34] Lee KY, Nakagawa T, Okano T, *et al.* Novel therapy for hearing loss: Delivery of insulin-like growth factor-1 to the cochlea using gelatin hydrogel. *Otol Neurotol* 2007; 28: 976-81.
- [35] Colas A. Silicones in pharmaceutical applications. DowCorning Healthcare Industries. <http://www.dowcorning.com/content/publishedlit/51-993a-01.pdf>
- [36] Okada H, Yamamoto M, Heya Y, *et al.* Drug delivery using biodegradable microspheres. *J Control Release* 1994; 28: 121-9.
- [37] Niwa T, Takeuchi H, Hino T, Kunou N, Kawashima Y. Preparation of biodegradable nanospheres of water-soluble and insoluble drugs with D,L-lactide/glycolide copolymer by a novel spontaneous emulsification solvent diffusion method, and the drug release behavior. *J Control Release* 1993; 25: 89-98.