

Sato T, Kusaka S, Shimojo H, Fujikado T.	Vitreous Levels of Erythropoietin and Vascular Endothelial Growth Factor in Eyes with Retinopathy of Prematurity.	Ophthalmology	116	1599-603	2009
Sato T, Kusaka S, Hashida N, Saishin Y, Fujikado T, Tano Y.	Comprehensive Gene Expression Profile in Murine Oxygen-induced Retinopathy.	Br J Ophthalmol	93	96-103	2009
Maruko I, Iida T, Sugano Y, Ojima A, Ogasawara M, Spaide RF.	The subfoveal choroidal thickness following treatment of central serous chorioretinopathy.	Ophthalmology, in press.			
Saito M, Shiragami C, Shiraga F, kano M, Iida T.	Comparison of intravitreal triamcinolone acetonide with photodynamic therapy and intravitreal bevacizumab with photodynamic therapy for retinal angiomatous proliferation.	Am J Ophthalmol, in press.			
Maruko I, Iida T, Saito M, Nagayama D.	Combined cases of polypoidal choroidal vasculopathy and typical age-related macular degeneration.	Graefes Arch Clin Exp Ophthalmol, in press.			
Koizumi H, Iida T, Mori T, Furuta M, Yannuzzi LA.	Retinal arteriolar macroaneurysm and congenital retinal macrovessel.	Ophthalmic surgery, lasers & imaging	40	513-515	2009
Kaneko H, Iida T, Ishiko H, Ohguchi T, Ariga T, Tagawa Y, Aoki K, Ohno S, Suzutani T.	Analysis of the complete genome sequence of epidemic keratoconjunctivitis-related human adenovirus type 8, 19, 37 and a novel serotype.	J Gen Virol	90	1471-1476	2009
Sekiryu T, Iida T, Maruko I, Horiguchi M.	Clinical application of autofluorescence densitometry with a scanning laser ophthalmoscope.	Invest Ophthalmol Vis Sci	50	2994-3002	2009

Furuta A, Nakadomari S, Misaki M, Miyauchi S, Iida T.	Objective perimetry using functional magnetic resonance imaging in patients with visual field loss.	Experimental Neurology	217	401-406	2009
Maruko I, Iida T, Sekiryu T, Saito M.	Morphologic changes in the outer layer of the detached retina in rhegmatogenous retinal detachment and central serous chorioretinopathy.	Am J Ophthalmol	147	489-494	2009
Fujiwara T, Iida T, Saito K, Maruko I.	Arteriolar macroaneurysm on the optic disc associated with branch retinal vein occlusion.	Retinal Cases & Brief Reports	3	21-23	2009
Furuta M, Iida T, Kishi S.	Foveal thickness can predict visual outcome in patients with persistent central serous chorioretinopathy.	Ophthalmologica	223	28-31	2009.
小島 彰, 寺島寛隆, 宮坂英世, 室井 繁, 飯田知弘.	スズメバチによる角膜蜂刺症の1例	眼科, 印刷中.			
伊勢重之, 丸子一郎, 神田尚孝, 飯田知弘.	両眼性の脈絡膜新生血管を伴うVogt-小柳-原田病の1例	臨眼	63	1738-1741	2009
原田 学, 森 隆史, 橋本禎子, 八子恵子, 飯田知弘	輪部ヘルモイドに対する層状角膜移植術の予後.	眼科	51	1155-1161	2009
齋藤昌晃, 永山 大, 飯田知弘	ポリープ状脈絡膜血管症に対する光線力学的療法の長期成績	日眼会誌	113	792-799	2009
近藤剛史, 丸子一郎, 石龍鉄樹, 飯田知弘.	臨床報告 先天性トキソプラズマ症の再発性網脈絡膜炎の1例.	眼科	51	807-811	2009
丸子一郎	加齢黄斑変性: 最近の動向 1. 疾患頻度	眼科	51	973-980	2009
古田 実	眼窩疾患の取り扱い方 眼窩の非腫瘍性血管性病変	JOHNS	25	1147-1154	2009
森 隆史	涙道疾患の治療.	眼科ケア	11	97-100	2009

石龍鉄樹	特集OCT: 日常臨床での所見のみかた 1. 網膜硝子体界面病変 -Vitreoretinal interface disorders-	眼科	51	235 -244	2009
齋藤昌晃, 飯田知弘.	加齢黄斑変性とアンチエイジング.	Drug Delivery System	24	103-108	2009
森 隆史	電気生理学実験でみた両眼視機能の発達	神経眼科	26:	141 -144	2009
Masuyama K et al	Posturing time after macular hole surgery modified by optical coherence tomography images: a pilot study.	Am J Ophthalmol	147 (3)	481-8	2009
Arimura N et al	Intraocular expression and release of high-mobility group box 1 protein in retinal detachment.	Lab Invest	89(3)	278-89	2009
Arimura N et al	Vitreous mediators after intravitreal bevacizumab or triamcinolone acetonide in eyes with proliferative diabetic retinopathy.	Ophhtalmology	116(5)	921-6	2009
Shimura M et al	Drug reflux during posterior subtenon infusion of triamcinolone acetonide in diffuse diabetic macular edema not only brings insufficient reduction but also causes elevation of intraocular pressure.	Graefe Clin Exp Arch Ophthalmol	247 (7)	907-12	2009
Yamashita T et al	Sonothrombolysis for Intraocular Fibrin Formation in an Animal Model.	Ultrasound Med Biol	35(11)	1845-53	2009
Sakamoto T et al	Visualizing vitreous in vitrectomy by triamcinolone.	Graefes Arch Clin Exp Ophthalmol.	247(9)	1153-63	2009
中村大輔 吉永幸靖 岡田龍雄 石橋達朗 江内田寛	多波長分光画像による眼底酸素飽和度の計測	光アライアンス	3	6-9	2009

Nonobe NI, Kachi S, Kondo M, Takai Y, Takamoto K, Nakayama K, Hayakawa M, Terasaki H	Concentration of vascular endothelial growth factor in aqueous humor of eyes with advanced retinopathy of prematurity before and after intravitreal injection of bevacizumab	Retina	29	579-585	2009
Kurimoto Y, Kondo M, Ueno S, Sakai T, Machida S, Terasaki H	Asymmetry of focal macular photopic negative responses (PhNRs) in monkeys.	Experimental Eye Research	88	92-98	2009
Kazuyuki Hirooka, Masanori Mizote, Tetsuya Baba, Mai Takagishi, Fumio Shiraga	Risk factors for developing avascular filtering bleb after fornix based trabeculectomy with mitomycin C.	J Glaucoma	18	301-4	2009
Yusuke Oshima, Chiharu Shima, Taku Wakabayashi, Shunji Kusaka, Fumio Shiraga, Masahito Ohji, Yasuo Tano.	Microincision Vitrectomy Surgery and Intravitreal Bevacizumab as a Surgical Adjunct to Treat Diabetic Traction Retinal Detachment.	Ophthalmology	116	927-938	2009
Kazuyuki Hirooka, Hirokazu Takenaka, Tetsuya Baba, Mai Takagishi, Masanori Mizote, and Fumio Shiraga.	Effect of Trabeculectomy on Intraocular Pressure Fluctuation with Postural Change in Eyes with Open-angle Glaucoma	J Glaucoma	18	689-91	2009
Hiroyuki Nomoto , Fumio Shiraga, Noriyuki Kuno, Erika Kimura, Shinobu Fujii, Katsuhiko Shinomiya, Alex K. Nugent, Kazuyuki Hirooka, Tetsuya Baba.	Pharmacokinetics of Bevacizumab After Topical, Subconjunctival and Intravitreal Administration In Rabbits.	Invest Ophthalmol Vis Sci	50	4807-13	2009
Hongwei Yang , Kazuyuki Hirooka, Kouki Fukuda, Fumio Shiraga.	Neuroprotective Effects of Angiotensin II Type 1 Receptor Blocker in a Rat Model of Chronic Glaucoma.	Invest Ophthalmol Vis Sci	50	5800-4	2009

Chieko Shiragami, Fumio Shiraga, Hidetaka Yamaji, Kouki Fukuda, Mai Takagishi, Misako Morita, Takehiro Kishigami.	Unintentional Displacement of the Retina after Standard Vitrectomy for Rhegmatogenous Retinal Detachment. Ophthalmology	Ophthalmology	117	86-92	2010
Hikoya A, Sato M, Tsuzuki K, Koide YM, Asaoka R, Hotta Y.	Central corneal thickness in Japanese children.	Jpn J Ophthalmol	53(1)	7-11	2009
Motoko Maekawa, Yoshimi Iwayama, Kazuhiko Nakamura, Miho Sato( 他9名)	A novel missense mutation (Leu46Val) of PAX6 found in an autistic patient.	Newsciencel etters	462	267-271	2009
佐藤美保, 大平明彦, 大井長和	わからない症例・相 談したい症例 動揺 視を自覚する潜伏眼 振	神経眼科	26巻3号	321-324	2009
佐藤美保	斜視と眼精疲労	日本医事新報	4456	75-74	2009
佐藤美保	弱視研究の新たな波	視覚の科学	30巻2号	27-34	2009
根岸貴志, 佐藤美保	直筋への通糸と眼球 への縫着	眼科手術	22巻3号	370-372	2009
根岸貴志, 佐藤美保	結膜切開と直筋の同 定	眼科手術	22巻2号	210-211	2009
佐藤美保	幼小児の眼鏡	あたらしい眼 科	26巻6号	741-746	2009
西村香澄, 彦谷明子, 佐藤美保, 嘉鳥信 忠, 尾花明, 堀田喜 裕	外傷性下直筋断裂に 対する下斜筋短縮前 方移動術の効果	眼科臨床紀要	2巻3号	249-255	2009

# Vascular Abnormalities in Aggressive Posterior Retinopathy of Prematurity Detected by Fluorescein Angiography

Tadashi Yokoi, MD,<sup>1</sup> Miina Hiraoka, MD,<sup>1</sup> Mariko Miyamoto, MD, PhD,<sup>2</sup> Tae Yokoi, MD,<sup>1</sup> Yuri Kobayashi, MD,<sup>1</sup> Sachiko Nishina, MD, PhD,<sup>1</sup> Noriyuki Azuma, MD, PhD<sup>1</sup>

**Purpose:** To evaluate fluorescein angiography (FA) in eyes with aggressive posterior retinopathy of prematurity (AP-ROP).

**Design:** Retrospective, nonrandomized case series.

**Participants:** Three patients (6 eyes) with AP-ROP.

**Methods:** Three patients (6 eyes) diagnosed with AP-ROP during ROP screening between July 2007 and July 2008 were included in this study. Fundus photographs and FA were obtained before and after laser and surgical treatment using a wide-field digital pediatric imaging system.

**Main Outcome Measures:** Fluorescein angiography and fundus photographs.

**Results:** At the initial stage of AP-ROP, FA showed vascular abnormalities, including capillary nonperfusion throughout the vascularized retina, shunting in the vascularized retina, a circumferential demarcation line, and limited vessel development, which was difficult to identify only by ophthalmoscopy. After treatment, FA showed poorly developed retinal vessels, including 4 small major vessels without an arcade pattern, small macular vessels, an inhomogeneous capillary bed, and absence of a capillary-free zone in the fovea.

**Conclusions:** Capillary bed loss throughout the vascularized posterior retina is characteristic of AP-ROP and may exacerbate retinopathy.

**Financial Disclosure(s):** The authors have no proprietary or commercial interest in any of the materials discussed in this article. *Ophthalmology* 2009;116:1377-1382 © 2009 by the American Academy of Ophthalmology.

Aggressive posterior retinopathy of prematurity (AP-ROP) is a severe, atypical form of ROP characterized by posterior retinal location, rapidly progressive vascular changes, flat neovascularization, hemorrhages, and intraretinal shunting without ridge tissue. Often, AP-ROP results in retinal detachment despite adequate laser ablation without progressing through classic stages 1 to 5.<sup>1-6</sup> Although early recognition and early treatment seem crucial for improving the unfavorable outcome, the initial signs of AP-ROP, including a subtle demarcation line or extraretinal neovascularization before dilatation and tortuosity of the retinal vessels occur, and vascular loops and frizzy vessels that lie at the periphery of vascularized retina and resembles frizzy hair, especially in the superior nasal quadrant, are difficult to recognize ophthalmoscopically.<sup>7</sup>

Previous reports on fluorescein angiography (FA) in classical ROP showed leakage from the fibrovascular tissue, arteriovenous shunting at the junction of the vascularized and unvascularized retina, a neovascular tuft posterior to the ridge, a capillary-free zone behind the ridge, or a capillary-free zone along both veins and arteries.<sup>8-12</sup> However, little has been learned about the FA findings of AP-ROP. We investigated the characteristics of vascular abnormalities in eyes with AP-ROP at the initial stage and after a series of treatments.


## Patients and Methods

Three infants (6 eyes) with AP-ROP examined at our institute between July 2007 and July 2008 were included. The patient charts and medical records were reviewed retrospectively. All infants underwent an initial ROP screening at a postmenstrual age of 29 to 31 weeks. Fundus photographs were taken using RetCam (Massie Research Laboratories, Inc., Pleasanton, CA) when the initial signs of ROP were detected by ophthalmoscopy. We also performed FA by RetCam with a blue excitation light source and a yellow-green, barrier-filtered camera after administration of an intravenous bolus dose of 0.1 ml/kg fluorescein followed by a 3.0-ml isotonic saline flush. Fundus photographs and FA were evaluated after laser application or surgery. This study was approved by the ethics committee of the National Center for Child Health and Development.

## Results

### Case 1

A male infant (age, 22 weeks gestation) weighing 400 g was delivered vaginally with Apgar scores of 5 and 6 at 1 and 5 minutes, respectively. This patient had respiratory distress syndrome requiring mechanical ventilation and 1 dose of surfactant, a patent ductus arteriosus treated with 8 courses of ibuprofen and subsequent surgical closure, and chronic lung disease without home oxygen therapy.




**Figure 1.** Fundus photographs and fluorescein angiography (FA) of aggressive posterior retinopathy of prematurity in the left eye of patient 1. A, A fundus photograph obtained at the initial stage shows the posterior retinal vessels, a nasal demarcation line, and a hemorrhage at the junction between the vascularized and unvascularized retina. B, At the initial stage, FA shows vessels with 4-lobed topography, capillary nonperfusion around the vascularized retina, vessel proliferation at the junction, and circumferential shunting. C, A fundus photograph obtained after laser treatment shows stabilized vascular proliferation with foveal formation. D, After treatment, FA shows abnormal vessels, such as an inhomogeneous capillary bed, small major vessels without an arcade pattern, and insufficient macular vessels. Because the vitreous cavity was filled with fluorescein after administration, the findings in the left eye, which were clearer than those in the right eye, are representative of the 2 eyes.

Bilateral ROP screening at a postmenstrual age of 30 weeks identified immature development of the retinal vessels that stopped in zone I, a nasal demarcation line, vessel tortuosity around the vessel end, a hemorrhage at the junction, and slight shunting in the vascularized retina (Fig 1A). On FA, there was extension of the major vessels in 4 directions with 4-lobed topography (i.e., vessels develop more temporally than nasally) and capillary nonperfusion and shunting throughout the vascularized retina including the posterior pole, even though a relatively strong choroidal flush, probably owing to immature pigmentation of the retinal pigment epithelium, complicated the identification of capillary nonperfusion. At the junction of the vascularized and unvascularized retina, the branches of the retinal vessels had a brush-like appearance, with capillary nonperfusion and arteriovenous shunting that resulted in a circumferentially continuous demarcation line. Leakage from the neovascular tissue in the junction also was confirmed. In the posterior pole, a cilioretinal artery abnormally passed through the incipient fovea, and few macular vessels or branch vessels, including arterioles and venules from the major vessels, were observed (Fig 1B). The disease activity decreased successfully as the result of laser treatment with foveal formation (Fig 1C), and FA after laser treatment showed stabilized neovascularization, 4 small major vessels especially in the nasal vessels, decreased branch vessels, loss of the arcade pattern, an inhomogeneous capillary bed around the retina, insufficient macular vessels, absence of a capillary-free zone in the fovea, and disappearance of the preexisting vessels that had passed through the fovea (Fig 1D).

## Case 2

A female infant (age, 23 weeks gestation) weighing 480 g was delivered vaginally with Apgar scores of 4 and 5 at 1 and 5 minutes, respectively. The patient had respiratory distress syndrome that required mechanical ventilation and 1 dose of surfactant, a patent ductus arteriosus treated with 3 courses of ibuprofen, and chronic lung disease without home oxygen therapy.

Bilateral ROP screening at a postmenstrual age of 30 weeks identified immature development of retinal vessels that stopped in zone I, a few tortuous and dilated vessels, a hemorrhage at the junction of the vascularized and unvascularized retina, and a circumferential vessel at the junction (Fig 2A). On the same day, FA detected vessels extending more in the temporal than nasal quadrant, circumferential vessels, and shunting in the junction resulting in formation of a demarcation line, leakage from the neovascular tissue on the demarcation line, and capillary nonperfusion and shunting throughout the vascularized retina including the posterior pole. Macular vessels and branch vessels including arterioles and venules from major vessels were as hypoplastic as in case 1 (Fig 2B). Despite early laser treatment that was applied densely, fibrovascular proliferation and a regional retinal detachment (stage 4A) occurred in the coagulation scars at 40 weeks. Early vitrectomy with lensectomy was performed at 41 weeks, and the retina was reattached with foveal formation (Fig 2C). After surgery, FA identified an inhomogeneous capillary bed throughout the retina, 4 small and irregular major vessels, decreased branch



**Figure 2.** Fundus photographs and fluorescein angiography (FA) of aggressive posterior retinopathy of prematurity in the right eye of patient 2. **A,** A fundus photograph obtained at the initial stage shows posterior vessels, hemorrhage, and circumferential vessel in the inferotemporal junction. **B,** At the initial stage, FA shows limited vessel extension with circumferential vessels, shunting in the junction, subtle leakage from the neovascular tissue in the demarcation line, capillary nonperfusion around the vascularized retina, and hypoplastic macular vessels. **C,** Fundus photograph obtained after laser and surgery shows stabilized vascular proliferation with foveal formation and a localized tractional retinal detachment that developed in the laser scars. **D,** After laser and surgery, FA shows an inhomogeneous capillary bed throughout the retina, tiny and irregularly shaped major vessels, loss of the arcade pattern, insufficient macular vessels, and absence of a capillary-free zone in the fovea.

vessels, loss of the arcade pattern, insufficient macular vessels, and absence of a capillary-free zone in the fovea (Fig 2D).

### Case 3

A female infant (age, 27 weeks gestation) weighing 920 g was delivered by Cesarean section with Apgar scores of 1 and 4 at 1 and 5 minutes, respectively. The patient had respiratory distress syndrome requiring mechanical ventilation and 1 dose of surfactant, a patent ductus arteriosus treated with 2 courses of ibuprofen, and sepsis treated with antibiotics.


Bilateral ROP screening at a postmenstrual age of 34 weeks identified immature development of retinal vessels that stopped in zone I, a continual demarcation line, budding from the junction of the vascularized and unvascularized retina, flat neovascularization around the nasal junction, and a hemorrhage at the junction (Fig 3A). On the same day, FA showed vessels extended more in the temporal than nasal quadrant; obvious vessel tortuosity, capillary nonperfusion, and shunting in the entire vascularized retina; few macular vessels and branch vessels including arterioles and venules from the major vessels; and leakage from the flat nasal neovascular tissue and the neovascular tuft behind the junction (Fig 3B). Although the disease activity was stabilized successfully by laser treatment with foveal formation (Fig 3C), FA after a series of laser applications still showed insufficient vasculature, including an inhomogeneous capillary bed in the retina, 4 small major vessels, a decrease in branch vessels, loss of an arcade pattern, and few macular vessels or absence of a capillary-free zone in the

fovea; however, there was much better vessel formation than in the other cases (Fig 3D).

### Discussion

The major FA findings in the current case series of early-stage AP-ROP are: (1) vessels extending more temporally than nasally that sometimes show 4-lobed topography, (2) a continuous demarcation line composed of shunting and circumferential vessels along the junction of the vascularized and unvascularized retina, (3) shunting in the vascularized retina, (4) tortuous vessels without dilatation clearly detected on FA, (5) leakage from neovascular tissue, (6) few macular vessels and few branch vessels from the major vessels including arterioles and venules, and (7) inhomogeneous capillary bed loss throughout the entire vascularized retina, including the posterior pole. Interestingly, in the current case series of AP-ROP, capillary nonperfusion or arteriovenous shunting was present throughout the vascularized retina, which is in contrast to the capillary nonperfusion present only along the junction of the vascularized and unvascularized retina in classic ROP. Even after the ROP stabilized after a series of laser or surgical treatments, vascular abnormalities remained, including an inhomogeneous capillary bed throughout the vascularized retina, loss





**Figure 3.** Fundus photographs and fluorescein angiography (FA) of aggressive posterior retinopathy of prematurity in the right eye of patient 3. **A.** A fundus photograph obtained at the initial stage shows retinal vessels stopped in zone I, a continual demarcation line, budding from the junction, flat neovascularization in the nasal junction, and a hemorrhage at the junction. **B.** At the initial stage, FA shows vessel tortuosity and capillary nonperfusion around the vascularized retina, shunting in the vascularized retina, few vessels branching toward the fovea, and leakage from the flat nasal neovascular tissue and neovascular tuft behind the junction. **C.** A fundus photograph obtained after a series of laser treatments shows stabilized vascular proliferation with foveal formation. **D.** After treatment, FA shows an inhomogeneous capillary bed throughout the vascularized retina, loss of the arcade pattern, insufficient macular vessels, and deficiency of the capillary-free zone in the fovea.

of the arcade pattern, 4 small major vessels, decreased branch vessels, hypoplastic macular vessels, and insufficiency of the capillary-free zone in the fovea.

Recently, a retinal whole-mount preparation method has shown primary retinal vasculature in humans.<sup>13</sup> Retinal vasculature begins to develop from the optic nerve to form the inner vascular plexuses at about 14 weeks gestation as hyaloid vessels regress.<sup>13,14</sup> At 25 weeks' gestation, the inner plexus spreads toward the peripheral retina in a lopsided fashion showing the 4-lobed topography pattern.<sup>13-15</sup> Between 25 and 26 weeks' gestation, the outer vascular plexus begins to sprout from the inner plexus and elongates downward into the outer retina, where a second vascular network is established parallel to the first network.<sup>13,14,16</sup> By 32 weeks' gestation, the inner plexus reaches the peripheral retina, whereas the outer plexus does so after birth.<sup>13</sup> Because most infants with AP-ROP are born prematurely (gestational age  $\leq 25$  weeks),<sup>3,5,17</sup> the retinal vasculature has formed the inner plexus posterior to the middle retina, whereas the outer plexus has just started to develop. In our patients who were born at approximately 25 weeks' gestation, the inner vascular plexus may have been established at birth; however, FA showed a noncapillary area of arteriovenous shunting throughout the vascularized retina, probably because of loss of the preexisting inner vascular plexus rather than malformation of the plexus.

Most immature babies have a dramatic change in  $P_{aO_2}$  from 20 to 25 mmHg in utero<sup>18</sup> to about 60 to 100 mmHg after delivery.<sup>19</sup> High-level, long-term oxygen administration disturbs the development of the primary retinal vasculature. Hyperoxia suppresses production of vascular endothelial growth factor from the primary retinal cells and leads to subsequent vascular endothelial cell apoptosis<sup>20-24</sup> and decreased vascular density. In addition, the more immature the vessels, the greater the oxygen toxicity to the vessels.<sup>25,26</sup> Because patients with AP-ROP have immature vessels and often receive long-term, high oxygen therapy, the retinal capillary bed may be lost. Moreover, in a murine model of ROP, vaso-obliteration has been detected mainly in the posterior retina, possibly because the retina around the optic disc and arteries, through which oxygen-rich vessels initially pass, experiences hyperoxia-mediated vascular endothelial growth factor mRNA downregulation and subsequent vascular endothelial cell apoptosis, resulting in a surrounding capillary-free zone.<sup>22</sup> Because most patients with AP-ROP have vessel development posteriorly, the same mechanism may contribute to the severe capillary bed loss around the retina. Meanwhile, because not all patients with AP-ROP have vasculature localized in zone I and not all extremely premature infants develop AP-ROP, other factors, including insulin-like growth factor 1,<sup>27</sup> erythropoi-

etin,<sup>28</sup> vasculogenesis,<sup>15</sup> and genetic factors,<sup>29</sup> may be associated with the disease.

Laser has been applied only to avascular retinas with classical ROP,<sup>4</sup> whereas eyes with AP-ROP often develop retinal detachment despite early, dense laser treatment to the avascular retina.<sup>2-4</sup> In extremely low-birth-weight infants with ROP, Schulenburg and Tsanaktsidis<sup>30</sup> advocated laser application to the vascularized retina, demonstrating the areas of capillary nonperfusion posterior to the demarcation line. Because capillary beds were widely absent in the vascularized retina including the posterior pole, laser application even to the vascular retina may be needed to prevent progression of AP-ROP. Early vitreous surgery with lensectomy successfully stabilized AP-ROP; the vascularized retina was treated preoperatively with sufficient laser ablation 3 to 4 spots posterior to the junction of the vascularized and unvascularized retina.<sup>17</sup>

Interestingly, FA showed that the affected retinal vasculature, including abnormal formation of the major arteries and veins, decreased arterioles and venules, an inhomogeneous capillary bed throughout the retina, hypoplastic macular vessels, and loss of the avascular zone in the fovea and macular vessels are still affected, even after a series of treatments, although this was not well detected on ophthalmoscopy. Thus, once the developing vasculature is damaged, it may never recover. Joshi et al<sup>31</sup> examined 14 eyes with stage 4A ROP preoperatively and detected anatomic macular abnormalities, such as absence of a foveal avascular zone and foveal depression. Although the visual prognosis in AP-ROP improved with early vitreous surgery,<sup>17</sup> because the disease severity is much worse than classic ROP and the development of vasculature was affected in the current study, long-term follow-up of the visual acuity and further investigation of the retinal function are necessary.

## References

1. International Committee for the Classification of Retinopathy of Prematurity. The International Classification of Retinopathy of Prematurity revised. *Arch Ophthalmol* 2005;123:991-9.
2. Vander JF, Handa J, McNamara JA, et al. Early treatment of posterior retinopathy of prematurity: a controlled trial. *Ophthalmology* 1997;104:1731-6.
3. Hiraoka M, Watanabe T, Kawakami T, et al. Retinopathy of prematurity in extremely low birth weight infants: a Tokyo multicenter study [in Japanese]. *Nippon Ganka Gakkai Zasshi* 2004;108:600-5.
4. Early Treatment for Retinopathy of Prematurity Cooperative Group. Revised indications for the treatment of retinopathy of prematurity: results of Early Treatment for Retinopathy of Prematurity randomized trial. *Arch Ophthalmol* 2003;121:1684-96.
5. Vinekar A, Trese MT, Capone A Jr, Photographic Screening for Retinopathy of Prematurity (PHOTO-ROP) Cooperative Group. Evolution of retinal detachment in posterior retinopathy of prematurity: impact on treatment approach. *Am J Ophthalmol* 2008;145:548-55.
6. Kychenthal A, Dorta P, Katz X. Zone I retinopathy of prematurity: clinical characteristics and treatment outcomes. *Retina* 2006;26(suppl):S11-5.
7. Morizane H. Initial sign and clinical course of the most severe form of acute proliferative retrolental fibroplasias (type II) [in Japanese]. *Nippon Ganka Gakkai Zasshi* 1976;80:54-61.
8. Cantolino SJ, O'Grady GE, Herrera JA, et al. Ophthalmoscopic monitoring of oxygen therapy in premature infants: fluorescein angiography in acute retrolental fibroplasia. *Am J Ophthalmol* 1971;72:322-31.
9. Flynn JT, O'Grady GE, Herrera J, et al. Retrolental fibroplasias: I. Clinical observations. *Arch Ophthalmol* 1977;95:217-23.
10. Takagi I, Oka Y, Nishimura M. Regressive and progressive cases of retinopathy of prematurity: fluorescein angiographic observations [in Japanese]. *Nippon Ganka Gakkai Zasshi* 1977;81:950-7.
11. Flynn JT, Cassady J, Essner D, et al. Fluorescein angiography in retrolental fibroplasia: experience from 1969-1977. *Ophthalmology* 1979;86:1700-23.
12. Ng EY, Lanigan B, O'Keefe M. Fundus fluorescein angiography in the screening for and management of retinopathy of prematurity. *J Pediatr Ophthalmol Strabismus* 2006;43:85-90.
13. Hughes S, Yang H, Chan-Ling T. Vascularization of the human fetal retina: roles of vasculogenesis and angiogenesis. *Invest Ophthalmol Vis Sci* 2000;41:1217-28.
14. Chan-Ling T, McLeod DS, Hughes S, et al. Astrocyte-endothelial cell relationships during human retinal vascular development. *Invest Ophthalmol Vis Sci* 2004;45:2020-32.
15. Flynn JT, Chan-Ling T. Retinopathy of prematurity: two distinct mechanisms that underlie zone 1 and zone 2 disease. *Am J Ophthalmol* 2006;142:46-59.
16. Gariano RF, Iruela-Arispe ML, Hendrickson AE. Vascular development in primate retina: comparison of lamellar plexus formation in monkey and human. *Invest Ophthalmol Vis Sci* 1994;35:3442-55.
17. Azuma N, Ishikawa K, Hama Y, et al. Early vitreous surgery for aggressive posterior retinopathy of prematurity. *Am J Ophthalmol* 2006;142:636-43.
18. Fineman JR, Soifer SJ. The fetal and neonatal circulation. In: Chang AC, Hanley FL, Wernovsky G, Wessel DL, eds. *Pediatric Cardiac Intensive Care*. Baltimore, MD: Lippincott Williams & Wilkins; 1998:17-24.
19. Chen J, Smith LE. Retinopathy of prematurity. *Angiogenesis* 2007;10:133-40.
20. Alon T, Hemo I, Itin A, et al. Vascular endothelial growth factor act as a survival factor for newly formed retinal vessels and has implication for retinopathy of prematurity. *Nat Med* 1995;1:1024-8.
21. Pierce EA, Foley ED, Smith LE. Regulation of vascular endothelial growth factor by oxygen in a model of retinopathy of prematurity. *Arch Ophthalmol* 1996;114:1219-28.
22. Claxton S, Fruttiger M. Role of arteries in oxygen induced vaso-obliteration. *Exp Eye Res* 2003;77:305-11.
23. Gu X, El-Remmessy AB, Brooks SE, et al. Hyperoxia induces retinal vascular endothelial cell apoptosis through formation of peroxynitrite. *Am J Physiol Cell Physiol* 2003;285:C546-54.
24. Ashton N, Ward B, Serpell G. Effect of oxygen on developing retinal vessels with particular reference to the problem of retrolental fibroplasia. *Br J Ophthalmol* 1954;38:397-432.
25. Hughes S, Chan-Ling T. Characterization of smooth muscle cell and pericyte differentiation in the rat retina in vivo. *Invest Ophthalmol Vis Sci* 2004;45:2795-806.
26. Chan-Ling T, Page MP, Gardiner T, et al. Desmin ensheathment ratio as an indicator of vessel stability: evidence in normal development and in retinopathy of prematurity. *Am J Pathol* 2004;165:1301-13.
27. Smith LE. IGF-1 and retinopathy of prematurity in the preterm infant. *Biol Neonate* 2005;88:237-44.

28. Morita M, Ohneda O, Yamashita T, et al. HLF/HIF-2alpha is a key factor in retinopathy of prematurity in association with erythropoietin. *EMBO J* 2003;22:1134-46.
29. Bizzarro MJ, Hussain N, Jonsson B, et al. Genetic susceptibility to retinopathy of prematurity. *Pediatrics* 2006;118:1858-63.
30. Schulenburg WE, Tsanaktsidis G. Variations in the morphology of retinopathy of prematurity in extremely low birth-weight infants. *Br J Ophthalmol* 2004;88:1500-3.
31. Joshi MM, Trese MT, Capone A Jr. Optical coherence tomography findings in stage 4A retinopathy of prematurity: a theory for visual variability. *Ophthalmology* 2006;113:657-60.

## Footnotes and Financial Disclosures

---

Originally received: August 11, 2008.

Final revision: January 26, 2009.

Accepted: February 26, 2009.

Available online: May 30, 2009.

Manuscript no. 2008-963.

<sup>1</sup> Department of Ophthalmology, National Center for Child Health and Development, Tokyo, Japan.

<sup>2</sup> Department of Ophthalmology, Metropolitan Bokutoh Hospital, Tokyo, Japan.

Tadashi Yokoi and Miina Hiraoka contributed equally to this article as first authors.

Financial Disclosure(s):

The authors have no proprietary or commercial interest in any of the materials discussed in this article.

Correspondence:

Noriyuki Azuma, MD, PhD, Department of Ophthalmology, National Center for Child Health and Development, 2-10-1, Okura, Setagaya-ku, Tokyo 157-8535, Japan. E-mail: azuma-n@ncchd.go.jp.

# Evaluation of Scleral Buckling for Stage 4A Retinopathy of Prematurity by Fluorescein Angiography

TAE YOKOI, TADASHI YOKOI, YURI KOBAYASHI, MIINA HIRAOKA, SACHIKO NISHINA, AND NORIYUKI AZUMA

• **PURPOSE:** To determine the early efficacy of scleral buckling for active neovascularization by fundus fluorescein angiography (FA) in eyes with stage 4A retinopathy of prematurity.

• **DESIGN:** A retrospective, nonrandomized, observational case series.

• **METHODS:** Patients who underwent scleral buckling for stage 4A ROP at the National Center for Child Health and Development, Tokyo, Japan, from October 2007 through November 2008 were included. Preoperative and postoperative FA and fundus photographs obtained with a wide-field digital pediatric imaging system were reviewed. Three patients (5 eyes; gestational ages at birth, 23 to 25 weeks; birth weights, 574 to 811 g) with zone II stage 4A ROP who underwent postoperative FA, 2 weeks or less after scleral buckling (range, 7 to 12 days; postmenstrual ages at postoperative FA, 41 to 45 weeks) were evaluated. Patients who underwent postoperative FA 2 weeks or more after scleral buckling were excluded.

• **RESULTS:** Despite fluorescein leakage from fibrovascular tissue in all eyes before surgery, markedly decreased leakage occurred only between 7 to 12 days after surgery. The retinas were reattached completely in all eyes after surgery.

• **CONCLUSIONS:** Scleral buckling may prevent progression of retinal detachment in stage 4A ROP by reducing the tractional force and stabilizing the neovascular activity of the fibrovascular tissue. (*Am J Ophthalmol* 2009;148:544-550. © 2009 by Elsevier Inc. All rights reserved.)

**R**ETINOPATHY OF PREMATUREITY (ROP) SOMETIMES progresses to retinal detachment (RD) despite application of early dense photocoagulation. Surgery is considered at the next treatment. Scleral buckling<sup>1-7</sup> or lens-sparing vitrectomy<sup>8-14</sup> has been performed with good surgical outcomes in eyes with stage 4A ROP, with the exception of those with aggressive posterior ROP. Scleral buckling or lens-sparing vitrectomy cannot stop the progression of RDs in eyes with aggressive posterior ROP because fibrovascular tissue rapidly reaches the posterior

lens surface, extends toward the vitreous base, and generates higher contractile forces.<sup>15</sup> Scleral buckling has been performed in less advanced cases of stage 4A ROP, and the mechanism of action has been reported to be relief of the vitreoretinal traction.<sup>6,14,16,17</sup> However, it is unknown whether scleral buckling itself contributes to stabilization of the neovascular activity.

With an advanced digital camera that obtains panoramic images of the neonatal fundus, fundus fluorescein angiography (FA) was performed to assess the retinal vasculature of infants with many peripheral retinal diseases. FA enables easy visualization of the retinal vasculature and aids in early diagnosis and prompt management of patients with ROP.<sup>18</sup> FA findings in ROP have included arteriovenous shunting at the junction of the vascular and avascular retina, a neovascular tuft posterior to the ridge, and a capillary-free zone behind the ridge or capillary-free zone along both veins and arteries.<sup>19-21</sup> Of those seen on FA, the most important sign of activity is fluorescein dye leakage resulting from the neovascularization. We performed preoperative and postoperative FA in eyes with zone II stage 4A ROP with active neovascularization to assess the efficacy of scleral buckling for neovascular activity.


## METHODS

WE PERFORMED A RETROSPECTIVE REVIEW OF THE CASE notes of all surgeries for eyes with stage 4 ROP that had developed along classical stages 1 through 3 without any sign of aggressive posterior ROP, performed by one surgeon (N.A.) between October 2007 and November 2008 at the National Center for Child Health and Development, Tokyo, Japan. All patients previously had undergone bilateral peripheral photocoagulation at other clinics. However, the primary treatment did not stop progression of the fibrovascular tissue proliferation and RD, and the patients were referred to us for the next surgery. In the patients who required surgery, lens-sparing vitrectomy generally was performed as a first procedure. However, when fibrovascular tissue developed in the periphery or extended to the vitreous base, the lens was removed to resect the vitreous gel around the fibrovascular tissue and vitreous base. A few eyes underwent scleral buckling for lens preservation.

Accepted for publication May 22, 2009.

From the Department of Ophthalmology, National Center for Child Health and Development, Tokyo, Japan.

Inquiries to Noriyuki Azuma, Department of Ophthalmology, National Center for Child Health and Development, 2-10-1, Okura, Setagaya-ku, Tokyo 157-8535, Japan; e-mail: azuma-n@nchd.go.jp



**FIGURE 1.** Fundus photography and fluorescein angiography (FA) images from Case 1 carried out before and after surgery in both eyes with stage 4A retinopathy of prematurity. (Top left) Fundus photograph obtained before scleral buckling at postmenstrual age 42 weeks in the right eye showing expanded fibrovascular tissue at the temporal junction between the vascular and avascular retina. (Top right) FA image obtained before scleral buckling showing marked fluorescein leakage from the fibrovascular tissue at the junction. (Bottom left) Fundus photograph obtained after scleral buckling showing that the progression of the retinal detachment (RD) stopped after surgery. (Bottom right) FA image obtained after treatment showing markedly decreased fluorescein leakage from the fibrovascular tissue. The findings in the right eye are clearer than those in the left eye and are representative of the 2 eyes.


A scleral buckling procedure without simultaneous vitreous surgery was performed at a postmenstrual age of 41 to 45 weeks by placing a 2.0-mm-diameter silicone sponge on each eye (Mira Inc, Waltham, Massachusetts, USA). The sponge was secured with a 5-0 Dacron scleral suture (Alcon Laboratories, Fort Worth, Texas, USA) in each quadrant, and the ends of the sponge were sutured end to end with the same suture at the superior nasal region. The aqueous tap also was performed if necessary, but drainage of subretinal fluid was not.

Fluorescein angiography examination was performed before and after scleral buckling using the RetCam120 (Massie Research Laboratories Inc, Pleasanton, California, USA). The pupils were dilated using 0.5% phenylephrine and 0.5% tropicamide eye drops instilled 3 times at 10-minute intervals 1 hour before the procedure. A neonatal lid speculum and Scopisol Solution for Eyes (Senju Pharmaceutical Co, Ltd, Osaka, Japan) were used to maintain the cornea-camera interface. The RetCam120 obtained color fundus images and FA images using a blue excitation light source and the yellow filters in the system. FA was performed after administration of an intravenous bolus dose of 0.1 ml/kg 10% sodium fluorescein dye followed by an isotonic saline flush. Images of the posterior pole, peripheral retina, and fibrovascular tissue were re-

corded at 2-second intervals during all phases of the angiogram until 5 minutes from the injection of fluorescein dye. Dye leakage from the fibrovascular tissue was evaluated in the preoperative and postoperative images in which the recording time was approximately the same. To eliminate the natural course of cicatrization, eyes that underwent FA 2 weeks or less after scleral buckling were evaluated in this study. Five eyes of 3 consecutive infants (2 girls and 1 boy; gestational age range at birth, 23 to 25 weeks; birth weight range, 574 to 811 g) with stage 4A ROP were included. The patients were followed up for 4 to 18 months after surgery. The surgical outcomes at the latest follow-up examinations were determined by binocular ophthalmoscopy and photography.

To evaluate the preoperative and postoperative fluorescein leakage objectively, semi-automated computer-assisted image analyses were performed using ImageJ version 1.41 (Wayne Rasband; National Institutes of Health, Bethesda, Maryland, USA). The relative area of fluorescein leakage was outlined using the freehand tool, and the relative mean signal intensity of the area was quantified using ImageJ densitometry. The outlined area was fixed between preoperative and postoperative images. The illumination knob was fixed with maximum range throughout the FA examinations.





**FIGURE 2.** Fundus photography and FA images from Case 2 obtained before and after surgery in both eyes with stage 4A retinopathy of prematurity. (Top left) Fundus photograph obtained before scleral buckling in the left eye at postmenstrual age 41 weeks showing expanded fibrovascular tissue and that the retinal vessels stopped in zone II. (Top right) FA image obtained before scleral buckling was performed showing moderate fluorescein leakage from the nasal fibrovascular tissue. (Bottom left) Fundus photograph obtained after scleral buckling showing that the progression of the RD stopped after surgery. (Bottom right) FA image obtained after treatment showing markedly decreased fluorescein leakage from the fibrovascular tissue. The findings in the left eye are clearer than those in the right eye and are representative of the 2 eyes.

## RESULTS

• **CASE 1:** A 25-week-gestation female infant weighing 811 g was delivered vaginally with Apgar scores of 8 and 9 at 1 and 5 minutes, respectively. She also had respiratory distress syndrome requiring mechanical ventilation, patent ductus arteriosus treated with one course of ibuprofen, posthemorrhagic hydrocephalus treated with an implant of an Ommaya reservoir, and sepsis treated with antibiotics.


Bilateral ROP screening at the postmenstrual age of 32 weeks identified immature development of retinal vessels that stopped in zone II. During the next 2 weeks, the clinical condition deteriorated to stage 3 with plus disease bilaterally, and argon laser photocoagulation was applied to the avascular retina. The vascular activity was stabilized transiently, but it reactivated. The fibrovascular tissue grew circumferentially, and a regional tractional retinal detachment (RD) developed in both eyes 7 weeks after photocoagulation.

The patient was referred to us and scleral buckling was performed bilaterally at the postmenstrual age of 42 weeks. The fundus examination before scleral buckling showed expanded fibrovascular tissue nearly 4 clock hours in size at the temporal junction between the vascular and avascular retina, and no apparent skip lesions in photocoagulation

were seen in either eye (Figure 1). FA performed just before the intervention showed marked fluorescein leakage from the fibrovascular tissue. The progression of the RD stopped after surgery in both eyes. FA performed 7 days after surgery showed markedly decreased fluorescein leakage. Mean signal intensity of the outlined area (72,464 pixels) decreased from 69.7 to 40.6 in right eye after surgery. The retina was reattached, and the fovea was well formed at the correct retinal position at the last follow-up examination.

• **CASE 2:** A 24-week-gestation male infant weighing 656 g was delivered vaginally with Apgar scores of 5 and 7 at 1 and 5 minutes, respectively. He had respiratory distress syndrome requiring mechanical ventilation.

Bilateral ROP screening at the postmenstrual age of 30 weeks identified immature development of retinal vessels that stopped in zone II and a demarcation line at the junction of the vascular and avascular retina. During the next 3 weeks, the clinical situation deteriorated to stage 3 with plus disease bilaterally, and argon laser photocoagulation was applied to the avascular retina over the course of 4 weeks. The vascular activity stabilized transiently but later reactivated. The fibrovascular tissue grew circumferentially, and a tractional RD developed gradually after the



**FIGURE 3.** Fundus photography and FA images from Case 3 obtained before and after surgery in the left eye with stage 4A retinopathy of prematurity. (Top left) Fundus photograph obtained before scleral buckling in the left eye at postmenstrual age 45 weeks showing that the retinal vessels stopped in zone II and showing expanded fibrovascular tissue at the temporal junction between the vascular and avascular retina. Skip lesions are present. (Top right) FA image obtained before scleral buckling showing marked fluorescein leakage from the temporal fibrovascular tissue. (Bottom left) Fundus photograph obtained after scleral buckling showing that the progression of the RD stopped after surgery. (Bottom right) FA image obtained after treatment showing markedly decreased fluorescein leakage from the fibrovascular tissue.

last session of photocoagulation at the postmenstrual age of 37 weeks.

The patient was referred to us and scleral buckling was performed bilaterally at the postmenstrual age of 41 weeks. The fundus examination before scleral buckling showed expanded fibrovascular tissue 5 clock hours in size at the junction of the vascular and avascular retina on the temporal inferior side of the right eye and for nearly 5 clock hours nasally in the left eye (Figure 2). No apparent skip lesions in photocoagulation were seen in either eye. FA performed just before the intervention showed fluorescein leakage from the fibrovascular tissue. Progression of the RD stopped in both eyes after surgery, and FA performed 12 days after surgery showed markedly decreased fluorescein leakage. Because there seemed to be some mask effect by blood after scleral buckling in the right eye, the mean signal intensity of fluorescein leakage was measured except the area covered with blood. The mean signal intensity of the outlined area (16,595 pixels) decreased 79.5 to 28.2. The retina reattached, and the fovea was well formed at the correct retinal position at the last follow-up examination.

• **CASE 3:** A 23-week-gestation female infant weighing 574 g was delivered by cesarean section with Apgar scores

of 5 and 7 at 1 and 5 minutes, respectively. The patient had respiratory distress syndrome requiring mechanical ventilation, patent ductus arteriosus treated with three courses of ibuprofen, subsequent surgical closure, and chronic lung disease without home oxygen therapy.

Bilateral ROP screening at the postmenstrual age of 28 weeks identified immature development of the retinal vessels that stopped in zone II. During the next 10 weeks, the clinical condition deteriorated to stage 3 with plus disease bilaterally, and argon laser photocoagulation was applied during a number of sessions to the avascular retina. The vascular activity stabilized transiently but later reactivated. The fibrovascular tissue grew circumferentially in both eyes after the last photocoagulation session at the postmenstrual age of 40 weeks. A tractional and serous RD developed in the right eye, and a regional tractional RD developed in the left eye.

Because of the rapid progression of bilateral RD, the patient was referred to us, and scleral buckling to both eyes was performed immediately at the postmenstrual age of 45 weeks. Although some skip lesions were present bilaterally, additional laser photocoagulation was not performed because of the risk of retinal tear after operation. The fundus examination before scleral buckling showed expanded fibrovascular tissue nearly 5 clock hours in size at the

**TABLE.** Characteristics of Eyes Undergoing Fluorescein Angiography before and after Scleral Buckling for Stage 4A Retinopathy of Prematurity

Patient	Gender	GA		Eye	Zone	Stage	Extent of FT (clock hours)	PMA of Ablation (wks)	PMA at SB (wks)	Preoperative Dye Leakage from FT	PMA at FA (wks)	Postoperative Dye Leakage from FT	Follow-up (mos)	Retinal Attachment	Foveal Formation (final)	SCF
1	F	25	811	Right	II	4A	4	34	42	Severe	43	Decrease <sup>a</sup>	7	Y	Y	Y
				Left	II	4A	4	34	42	Severe	43	None	7	Y	Y	Y
2	M	24	656	Right	II	4A	5	33 to 38	41	Moderate	44	Decrease <sup>a</sup>	17	Y	Y	Y
				Left	II	4A	5	33 to 38	41	Moderate	44	Decrease <sup>a</sup>	17	Y	Y	Y
3	F	23	574	Right	II	4A	5	38 to 40	45	Severe	47	Decrease	2	Y	Y	Y

BW = birth weight; F = female; FA = fluorescein angiography; FT = fibrovascular tissue; GA = gestational age; M = male; mos = months; PMA = postmenstrual age; POD = postoperative day; SB = scleral buckling; SCF = steady central fixation; wks = weeks; Y = yes.  
<sup>a</sup>More than 40% decrease of mean signal intensity of dye leakage.

junction of the vascular and avascular retina temporally in both eyes (Figure 3). A serous RD in the right eye was diagnosed based on the presence of exudates in the posterior pole and a bullous RD involving the macula with no retinal tear. FA performed just before the intervention detected marked fluorescein leakage from the fibrovascular tissue temporally in both eyes and diffuse fluorescein leakage from the retinal vessels in the right eye. Although serous RD in the right eye was exacerbated after scleral buckling, the progression of the tractional RD in the left eye stopped after surgery. FA performed 8 days after surgery showed marked cessation of fluorescein leakage in the left eye. Mean signal intensity of the outlined area (51,194 pixels) decreased from 84.3 to 49.7 in the left eye after surgery. Retinal reattachment was achieved and the fovea was well formed at the correct retinal position in left eye at the last follow-up examination. Results are summarized in the Table.

## DISCUSSION

SCLERAL BUCKLING ORIGINALLY WAS THOUGHT TO ADDRESS RDs by offsetting the vitreoretinal traction as the buckle indented the globe. The procedure allows the retinal pigment epithelium (RPE) to absorb the subretinal fluid and to support the fixed surface area of the developing retina while the eyeball grows rapidly in infants. Scleral buckling also may prevent death of the immature retinal cells by maintaining contact between the retina and the RPE.<sup>3,6,14,16,17</sup> One study reported that scleral buckling contributes to a notable change in neovascularization, which was identified by funduscopy.<sup>3</sup> In the current series, we found that leakage from the fibrovascular tissue markedly decreased only 7 to 12 days after scleral buckling. This course obviously was different from the natural course of fibrovascular tissue regression, which usually requires 1 month or more, suggesting that scleral buckling itself stabilizes the neovascular activity.

Neovascularization of ROP is induced by several growth factors,<sup>22-25</sup> of which vascular endothelial growth factor (VEGF) is the most important in the development of pathologic neovascularization, which is released from the vascular endothelial cells in hypoxia. A recent in vitro study reported that retinal vascular endothelial cells and the RPE, when stretched, release VEGF.<sup>26-28</sup> Integrin is the major receptor that connects vascular endothelial cells and the extracellular matrix and plays a key role in development of pathologic neovascularization.<sup>29-31</sup> Inhibition of integrin reduced VEGF in a murine model of ROP.<sup>32</sup> Because integrin is the mechanotransduction receptor that transforms mechanical stress into chemical signaling,<sup>33</sup> surgically offsetting the mechanical stress on the retinal vascular endothelial cells may reduce the release VEGF by inhibiting the integrin-mediated mechanotransduction. Vitrectomy may wash out VEGF directly from the vitreous cavity and may remove the vitreous gels that may contain the source of VEGF.<sup>34</sup> In contrast, scleral buckling does not have such effects. Thus, relieving the mechanical stress may decrease VEGF expression in the vascular endothelial cells, which may have contributed to decreased neovascular activity in our series.

Vascular endothelial growth factor expression is regulated primarily by hypoxia.<sup>35</sup> Hypoxia in the retinal outer layers caused by RD increases VEGF expression in the subretinal fluid.<sup>36</sup> In the eyes with stage 4 ROP, elevated VEGF levels also are detected in the subretinal fluid.<sup>37</sup> Retinal reattachment around the scleral buckling region may improve the oxygen supply from the choroid to the retina, contributing to the reduction of VEGF and preventing exacerbation of the neovascular activity.

In ROP and other proliferative retinopathies, the active stage characterized by neovascular formation is followed by the cicatricial stage with formation of contractile connective tissue.<sup>38</sup> A recent report demonstrated that the connective tissue growth factor, which is established as a profibrotic factor in retina,<sup>39,40</sup> in a critical balance with VEGF, drives the angiofibrotic switch and subsequent fibrosis in eyes with



proliferative diabetic retinopathies.<sup>41</sup> Compared with the natural course, pathologic neovascular inactivation in our series was achieved rapidly with subsequent cicatricial phase, which is not explained only by a decrease in the release of growth factors and suggests that scleral buckling also may act as the modulator of angiofibrotic switch.

Good surgical results have been reported after scleral buckling<sup>1-7</sup>; however, it also has some disadvantages, including high myopia resulting from changes in the corneal or lens refractive power or the need for another surgery to divide or remove the buckle to reduce the intrusion and promote ocular growth.<sup>38,42</sup> Recently, lens-sparing vitrectomy has become the surgery of choice for

stage 4 ROP, although application is limited to eyes with posterior proliferation. Thus, scleral buckling that allows lens preservation still may be chosen to treat eyes with peripheral proliferation in stage 4 ROP. Although the purpose of the surgery has been primarily to reduce traction,<sup>6,14,16,17</sup> the current small series showed that scleral buckling also can reduce the neovascular activity of peripheral proliferation in progressive stage 4A ROP. Thus, scleral buckling may be considered a treatment option to reduce the neovascular activity in eyes with peripheral proliferation where lens-sparing vitrectomy is difficult to perform, although further prospective studies are needed.

THE AUTHORS INDICATE NO FINANCIAL SUPPORT OR FINANCIAL CONFLICT OF INTEREST. INVOLVED IN DESIGN AND CONDUCT OF STUDY (T.Y., T.Y., N.A.); COLLECTION AND MANAGEMENT OF DATA (Tae.Y., N.A.); PROVISION OF PATIENTS AND RESOURCES (Tae.Y., Tad.Y., Y.K., M.H., S.N., N.A.); ANALYSIS AND INTERPRETATION OF THE DATA (Tae.Y., Tad.Y., Y.K., M.H., S.N., N.A.); PREPARATION OF MANUSCRIPT (Tae.Y., N.A.); REVIEW AND APPROVAL OF THE MANUSCRIPT (Tae.Y., Tad.Y., Y.K., M.H., S.N., N.A.). THIS STUDY WAS SUPPORTED BY GRANTS FOR SENSORY DISORDERS FROM THE MINISTRY OF HEALTH, LABOR AND WELFARE, TOKYO, JAPAN. ALL ASPECTS OF THIS STUDY WERE APPROVED BY THE INSTITUTIONAL ETHICS COMMITTEE.

## REFERENCES

1. Trese MT. Scleral buckling for retinopathy of prematurity. *Ophthalmology* 1994;101:23-26.
2. Ricci B, Santo A, Ricci F, et al. Scleral buckling surgery in stage 4 retinopathy of prematurity. *Graefes Arch Clin Exp Ophthalmol* 1996;234:S38-S41.
3. Hinz BJ, de Juan E Jr, Repka MX. Scleral buckling surgery for active stage 4A retinopathy of prematurity. *Ophthalmology* 1998;105:1827-1830.
4. Chuang YC, Yang CM. Scleral buckling for stage 4 retinopathy of prematurity. *Ophthalmic Surg Lasers* 2000;31:374-379.
5. Beyrau K, Danis R. Outcomes of primary scleral buckling for stage 4 retinopathy of prematurity. *Can J Ophthalmol* 2003;38:267-271.
6. Ertzbischoff LM. A systematic review of anatomical and visual function outcomes in preterm infants after scleral buckle and vitrectomy for retinal detachment. *Adv Neonatal Care* 2004;4:10-19.
7. Ratanasukon M, Visaetsilpanonta S, Tengtrisorn S, et al. Outcomes of scleral buckling for stage 4 retinopathy of prematurity in Thai children. *J Med Assoc Thai* 2006;89:1659-1664.
8. Capone A, Trese MT. Lens-sparing vitreous surgery for tractional stage 4A retinopathy of prematurity retinal detachments. *Ophthalmology* 2001;108:2068-2070.
9. Hartnett ME. Features associated with surgical outcome in patients with stages 4 and 5 retinopathy of prematurity. *Retina* 2003;23:322-329.
10. Prenner JL, Capone A, Trese MT. Visual outcome after lens-sparing vitrectomy for stage 4A retinopathy of prematurity. *Ophthalmology* 2004;111:2271-2273.
11. Hubbard III GB, Cherwick DH, Burian G. Lens-sparing vitrectomy for stage 4 retinopathy of prematurity. *Ophthalmology* 2004;111:2274-2277.
12. Hartnett ME, Maguluri S, Thompson HW, et al. Comparison of retinal outcomes after scleral buckle or lens-sparing vitrectomy for stage 4 retinopathy of prematurity. *Retina* 2004;24:753-757.
13. Yu YS, Kim SJ, Kim SY, et al. Lens-sparing vitrectomy for stage 4 and stage 5 retinopathy of prematurity. *Korean J Ophthalmol* 2006;20:113-117.
14. Sears JE, Sonnie C. Anatomic success of lens-sparing vitrectomy with and without scleral buckle for stage 4 retinopathy of prematurity. *Am J Ophthalmol* 2007;143:810-813.
15. Azuma N, Ishikawa K, Hama Y, et al. Early vitreous surgery for aggressive posterior retinopathy of prematurity. *Am J Ophthalmol* 2006;142:636-643.
16. McPherson A, Hittner HM. Scleral buckling in 2 1/2 to 11-month-old premature infants with retinal detachment associated with acute retrolental fibroplasia. *Mod Probl Ophthalmol* 1979;20:431-434.
17. Mintz-Hittner HA, Kretzer FL. The rationale for cryotherapy with a prophylactic scleral buckle for zone I threshold retinopathy of prematurity. *Doc Ophthalmol* 1990;74:263-268.
18. Azad R, Chandra P, Khan MA, et al. Role of intravenous fluorescein angiography in early detection and regression of retinopathy of prematurity. *J AAPOS* 2008;45:36-39.
19. Cantolino SJ, O'Grady GE, Herrera JA, et al. Ophthalmoscopic monitoring of oxygen therapy in premature infants. Fluorescein angiography in acute retrolental fibroplasia. *Am J Ophthalmol* 1971;72:322-331.
20. Flynn JT, O'Grady GE, Herrera J, et al. Retrolental fibroplasia. I. Clinical observations. *Arch Ophthalmol* 1977;95:217-223.
21. Flynn JT, Cassady J, Essner D, et al. Fluorescein angiography in retrolental fibroplasia: experience from 1969-1977. *Ophthalmology* 1979;86:1700-1723.
22. Pierce EA, Foley ED, Smith LE. Regulation of vascular endothelial growth factor by oxygen in a model of retinopathy of prematurity. *Arch Ophthalmol* 1996;114:1219-1228.
23. Lashkari K, Hirose T, Yazdany J, et al. Vascular endothelial growth factor and hepatocyte growth factor levels are differ-

- entially elevated in patients with advanced retinopathy of prematurity. *Am J Pathol* 2000;156:1337–1344.
24. Morita M, Ohneda O, Yamashita T, et al. HLF/HIF-2alpha is a key factor in retinopathy of prematurity in association with erythropoietin. *EMBO J* 2003;22:1134–1146.
  25. Hellstrom A, Engstrom E, Hard AL, et al. Postnatal serum insulin-like growth factor I deficiency is associated with retinopathy of prematurity and other complications of premature birth. *Pediatrics* 2003;112:1016–1020.
  26. Seko Y, Fujikura H, Pang J, et al. Induction of vascular endothelial growth factor after application of mechanical stress to retinal pigment epithelium of the rat in vitro. *Invest Ophthalmol Vis Sci* 1999;40:3287–3291.
  27. Suzuma I, Hata Y, Clermont A, et al. Cyclic stretch and hypertension induce retinal expression of vascular endothelial growth factor and vascular endothelial growth factor receptor-2: potential mechanisms for exacerbation of diabetic retinopathy by hypertension. *Diabetes* 2001;50:444–454.
  28. Suzuma I, Suzuma K, Ueki K, et al. Stretch-induced retinal vascular endothelial growth factor expression is mediated by phosphatidylinositol 3-kinase and protein kinase C (PKC)-zeta but not by stretch-induced ERK1/2, Akt, Ras, or classical/novel PKC pathways. *J Biol Chem* 2002;277:1047–1057.
  29. Luna J, Tobe T, Mousa SA, et al. Antagonists of integrin alpha vs beta 3 inhibit retinal neovascularization in a murine model. *Lab Invest* 1996;75:563–573.
  30. Friedlander M, Theesfeld CL, Sugita M, et al. Involvement of integrins alpha vs beta 3 and alpha vs beta 5 in ocular neovascular diseases. *Proc Natl Acad Sci U S A* 1996;93:9764–9769.
  31. Lahdenranta J, Sidman RL, Pasqualini R, et al. Treatment of hypoxia-induced retinopathy with targeted proapoptotic peptidomimetic in a mouse model of disease. *FASEB J* 2007;21:3272–3278.
  32. Wilkinson-Berka JL, Jones D, Taylor G, et al. SB-267268, a nonpeptidic antagonist of alpha(vs)beta3 and alpha(vs)beta5 integrins, reduces angiogenesis and VEGF expression in a mouse model of retinopathy of prematurity. *Invest Ophthalmol Vis Sci* 2006;47:1600–1605.
  33. Heerkens EH, Izzard AS, Heagerty AM, et al. Integrins, vascular remodeling, and hypertension. *Hypertension* 2007;49:1–4.
  34. Hanneken A, Maher PA, Baird A. High affinity immunoreactive FGF receptors in the extracellular matrix of vascular endothelial cells—implications for the modulation of FGF-2. *J Cell Biol* 1995;128:1221–1228.
  35. Stein I, Itin A, Einat P, et al. Translation vascular endothelial growth factor mRNA by internal ribosome entry: implications for translation under hypoxia. *Mol Cell Biol* 1998;18:3112–3119.
  36. Su CY, Chen MT, Wu WS, et al. Concentration of vascular endothelial growth factor in the subretinal fluid of retinal detachment. *J Ocul Pharmacol Ther* 2000;16:463–469.
  37. Lashkari K, Hirose T, Yazdany J, et al. Vascular endothelial growth factor and hepatocyte growth factor levels are differentially elevated in patients with advanced retinopathy of prematurity. *Am J Pathol* 2000;156:1337–1344.
  38. Chow DR, Ferrone PJ, Trese MT. Refractive changes associated with scleral buckling and division in retinopathy of prematurity. *Arch Ophthalmol* 1998;116:1446–1448.
  39. Kuiper EJ, De Smet MD, van Meurs JC, et al. Association of connective tissue growth factor with fibrosis in vitreoretinal disorders in the human eye. *Arch Ophthalmol* 2006;124:1457–1462.
  40. He S, Chen Y, Khankan R, et al. Connective tissue growth factor as a mediator of intraocular fibrosis. *Invest Ophthalmol Vis Sci* 2008;49:4078–4088.
  41. Kuiper EJ, Van Nieuwenhoven FA, De Smet MD, et al. The angio-fibrotic switch of VEGF and CTGF in proliferative diabetic retinopathy. *PLoS ONE* 2008;16:3:e2675.
  42. Choi MY, Young SY. Efficacy of removal of buckle after scleral buckling surgery for retinopathy of prematurity. *J AAPOS* 2000;4:362–365.

# Effect of Early Vitreous Surgery for Aggressive Posterior Retinopathy of Prematurity Detected by Fundus Fluorescein Angiography

Sachiko Nishina, MD, PhD, Tadashi Yokoi, MD, Tae Yokoi, MD, Yuri Kobayashi, MD, Miina Hiraoka, MD, Noriyuki Azuma, MD, PhD

**Objective:** To assess the effect of early vitrectomy for aggressive posterior retinopathy of prematurity (APROP) using fundus fluorescein angiography.

**Design:** Retrospective, observational case series.

**Participants:** Eleven eyes of 7 patients with APROP that underwent early vitreous surgery.

**Methods:** All eyes underwent vitrectomy with lensectomy that removed the vitreous gel around the fibrovascular proliferative tissue, but not the proliferative tissue when fibrovascular proliferation and retinal detachment occurred despite retinal photocoagulation. Fundus fluorescein angiography was performed before and after the early vitreous surgery.

**Main Outcome Measures:** Dye leakage from the fibrovascular tissue, dilation and tortuosity of the retinal vasculature, and shunt vessels were evaluated by fundus fluorescein angiography. The status of the retinal reattachment was assessed postoperatively.

**Results:** Nine eyes had severe dye leakage from the fibrovascular tissue and 2 eyes had moderate leakage seen by preoperative fluorescein angiography. Severe dilation and tortuosity of the retinal vessels were detected in 10 eyes and shunt vessels in 7 eyes. Six to 12 days after successful surgery, the retina reattached and dilation and tortuosity of the retinal vessels decreased substantially. Dye leakage diminished markedly in all eyes, resolved completely in 7 eyes, and was still apparent slightly in 4. At the final examination, fibrovascular proliferation and retinal detachment did not progress in any eyes; however, 2 eyes had a dragged or folded retina. Follow-up ranged from 6 to 19 months (mean, 9.2).

**Conclusions:** Early vitrectomy that removes vitreous gel from around the proliferative tissue promptly reduces vascular activity and may limit progression of retinal detachment in APROP.

**Financial Disclosure(s):** The authors have no proprietary or commercial interest in any of the materials discussed in this article. *Ophthalmology* 2009;116:2442-2447 © 2009 by the American Academy of Ophthalmology.



The revised International Classification of Retinopathy of Prematurity (ROP)<sup>1</sup> described aggressive posterior retinopathy of prematurity (APROP) and recognized it as an unusual form of ROP that rapidly progresses to a closed funnel of tractional retinal detachment within 1 or 2 weeks despite application of early, dense photocoagulation. The features and pathogenesis of this severe form of ROP have been investigated as type II ROP since 1974 in Japan.<sup>2,3</sup> Identification of the initial signs of APROP and immediate application of dense, extensive laser photocoagulation to the nonvascularized retina and adjacent vascularized retina including the area of the shunting vessels are essential for treating APROP.<sup>2-5</sup> Laser ablation performed early, densely, and repeatedly may transiently stabilize APROP; however, fibrovascular proliferation sometimes recurs in the area of the photocoagulation scars, and the tractional retinal detachment rapidly progresses.<sup>4-6</sup> Recently, we have re-

ported that early vitrectomy with lensectomy for APROP and early photocoagulation prevented progression of retinal detachment.<sup>5</sup> Early vitreous surgery for APROP also was evaluated in another report.<sup>6</sup> The aim of the surgery was not to remove the fibrovascular proliferative tissue, but to remove the surrounding vitreous gel along which the fibrovascular tissue grows. However, the mechanism by which the surgery stabilizes the retinopathy has not been clarified fully.

Fundus fluorescein angiography for ROP, first performed by Flynn et al,<sup>7</sup> is now available with a digital fundus camera, the RetCam 120 (Massie Research Laboratories, Inc., Pleasanton, CA), that can view a wide angle of the fundus including the periphery. With this camera, bedside fluorescein angiography can be performed easily to detect vascular areas of nonperfusion and early neovascularization of active retinopathy.<sup>8,9</sup>

We conducted the current study to evaluate the preoperative status and the efficacy of early vitreous surgery for APROP using fundus fluorescein angiography with the Retcam 120.

## Patients and Methods

This study was approved by the institutional ethics committee; the parents of the patients provided informed consent before the infants were enrolled. Eleven eyes of 7 patients with APROP were studied. All eyes had undergone previous laser ablation at other clinics when the initial signs of APROP<sup>1-3</sup> were identified, and laser was applied repeatedly and densely to the nonvascularized and adjacent vascularized retina. However, when laser treatment could not stop the progression of fibrovascular proliferation and tractional retinal detachment, the patients were referred immediately to our clinic at the National Center for Child Health and Development, Tokyo, Japan.

All eyes underwent early vitrectomy with lensectomy as a second treatment performed by 1 surgeon (NA) in our clinic between December 2006 and March 2008. Three-port vitrectomy was performed using the Accurus 25-Gauge Surgical System (Alcon, Fort Worth, TX) and a small contact lens was designed for premature eyes. The details of the surgery have been described previously<sup>5</sup> and are shown in online Video Clips 1 through 3 (available at <http://aaojournal.org>). We did not remove the fibrovascular tissue to avoid intraoperative bleeding, but we did remove the surrounding vitreous gel. Fluid-air exchange and endophotocoagulation were unnecessary, because no iatrogenic breaks occurred during the surgery.

Fundus fluorescein angiography was performed within 2 days before the surgery and 6 to 12 days after the surgery. The pupils were dilated using 0.5% phenylephrine and 0.5% tropicamide eye drops instilled 3 times at 10-minute intervals 1 hour before the examination. A neonatal lid speculum and hydroxyethyl cellulose (Scopisol 15; Senju Pharmaceutical Co., Ltd., Osaka, Japan) were used to maintain the cornea-camera interface. The Retcam 120 was used to obtain color fundus photographs and perform fluorescein angiography using a blue excitation light source and the yellow filters in the system. The vital signs were monitored by a neonatologist during the procedure. Fundus fluorescein angiography was performed after administration of an intravenous dose of 0.1 ml/kg of 10% sodium fluorescein dye followed by a saline flush to obtain a bolus dose. Images of the posterior pole, peripheral retina, and fibrovascular tissue were recorded at 2-second intervals during all phases of the angiogram until 5 minutes. Dye leakage from the fibrovascular tissue, dilation and tortuosity of the vasculature, and shunt vessels were evaluated.

The patients were followed for 6 to 19 months (mean, 9.2) postoperatively. Aphakic eyes were corrected with spectacles or contact lenses designed for premature babies about 2 weeks after surgery. The final anatomic outcomes were determined by binocular ophthalmoscopy and photography.

## Results

The characteristics of eyes that underwent early vitreous surgery for APROP and the results of fundus fluorescein angiography before and after surgery are shown in Table 1. The patients included 1 boy and 6 girls. The gestational ages at birth ranged from 23 to 28 weeks (mean, 25), and the birth weights ranged from 389 to 1194 g (mean, 734). The ROP extended to all 4 quadrants and underwent laser ablation within zone I in 9 eyes and to

posterior zone II in 2 eyes. The extent of the fibrovascular tissue ranged from 6 to 10 clock hours and all eyes had a tractional retinal detachment; the fovea was not involved (stage 4A) in 10 eyes and was involved (stage 4B) in 1 eye. The tip of the fibrovascular tissue was not attached to the vitreous base in any eyes. The postmenstrual ages at the time of vitreous surgery ranged from 35 to 41 weeks (mean, 38). Intraoperative complications did not develop in any eyes.

Preoperative fundus fluorescein angiography showed severe dye leakage (the extent of dye leakage was 100%) from the fibrovascular tissue in 9 eyes and moderate leakage (the extent of leakage was >70%) in 2 eyes. Severe dilation and tortuosity of the vasculature at the posterior pole were seen clearly in 10 eyes and the shunt vessels remained in 7 eyes. Six to 12 days after successful operations, the dye leakage decreased markedly in all eyes, that is, it resolved in 7 eyes (Figs 1 and 2) and was still apparent in 4 eyes (Fig 3). The extent of the fibrovascular tissue with residual dye leakage in the 4 eyes was <50% of the preoperative findings. The dilation and tortuosity of vasculature also decreased in all eyes (Figs 1 to 3). Fluorescein did not cause any adverse effects.

At the final examination, there was no progression of fibrovascular proliferation or retinal detachment in any eyes. Two eyes had a dragged or folded retina (Fig 3).

## Discussion

We perform vitrectomy with lensectomy in eyes with APROP as early as possible when fibrovascular tissue extends from the posterior retina to the posterior lens surface and causes a regional retinal detachment (stage 4A) despite sufficient laser ablation. Retinopathy of prematurity rapidly progresses to stage 5 within 1 to 2 weeks, when the growing fibrovascular tissue attaches to the posterior lens surface or the retina and ciliary body beneath the vitreous base. Thus, a wide-field vitrectomy with lensectomy to remove the vitreous gel between the fibrous tissue and vitreous base is needed to prevent progression of APROP.<sup>5</sup> If the tip of the fibrovascular tissue has attached to the vitreous base, the tractional retinal detachment rapidly progresses to involve the fovea. In this case, cutting the fibrovascular tissue and removing the firm adhesion to the vitreous base are difficult without iatrogenic retinal breaks, resulting in retinal folds or a dragged retina as a result of the strong contraction of the residual fibrovascular tissue and no foveal formation.<sup>5</sup>

In the current series, all but 1 eye had stage 4A disease in which the fibrovascular tissue increased to 6 to 10 clock hours without attachment to the vitreous base. The eye with stage 4B also had 6 clock hours of fibrovascular tissue without attachment to the vitreous base. We successfully treated these eyes with vitrectomy and lensectomy to obtain retinal reattachment. Preoperative fundus fluorescein angiography showed extreme vascular activity based on extensive dye leakage from the fibrovascular tissue, dilation and tortuosity of the retinal vasculature at the posterior pole, and atypical shunt vessels. However, only 6 to 12 days after surgery, fluorescein angiography showed markedly decreased vascular activity and stabilized APROP. The dilation and tortuosity of the retinal vessels decreased, and the dye leakage from the fibrovascular tissue decreased substantially. These findings strongly support the effectiveness of early vitrectomy, not only for preventing fibrovascular pro-