

- Seroepidemiological study in a Puumala virus outbreak area in South-East Germany. *Med Microbiol Immunol* 2009. 198:83-91.
25. Kariwa H, Tkachenko EA, Morozov VG, Seto T, Tanikawa Y, Kolominov SI, Belov SN, Nakamura I, Hashimoto N, Balakiev AE, Dzagurnova TK, Daud NH, Miyashita D, Medvedkina OA, Nakauchi M, Ishizuka M, Yoshii K, Yoshimatsu K, Arikawa J, and Takashima I. Epidemiological study of hantavirus infection in the Samara Region of European Russia. *J Vet Med Sci* 2009. 71:1569-1578.
 26. Chandy S, Yoshimatsu K, Boorugu HK, Chrispal A, Thomas K, Peedicyil A, Abraham P, Arikawa J, and Sridharan G. Acute febrile illness caused by hantavirus: serological and molecular evidence from India. *Trans R Soc Trop Med Hyg* 2009. 103:407-412.
 27. Chandy S, Okumura M, Yoshimatsu K, Ulrich RG, John GT, Abraham P, Arikawa J, and Sridharan G. Hantavirus species in India: A retrospective study. *Indian J Med Res* 2009. 27:348-350.
 28. Lim, C.K., Nishibori, T., Watanabe, K., Ito, M., Kotaki, A., Tanaka, K., Kurane, I., Takasaki, T. Chikungunya virus isolated from a returnee to Japan from Sri Lanka: Isolation of two sub-strains with different characteristics. *Am. J. Trop. Med. Hyg.*, 2009. 81(5):865-868.
 29. Lim, C.K., Kurane, I., and Takasaki, T. (2010) Re-emergence of chikungunya virus, pp. 1-22. In Maeda, A. (ed), *Animal Viruses*. Transworld Research Network., Kerala, India.
 30. Aoyama, I., Uno, K., Yumisashi, T., Takasaki, T., Lim, C.K., Kurane, I., Kase, T., Takahashi, K. A Case of Chikungunya Fever Imported from India to Japan, Follow-Up of Specific IgM and IgG Antibodies over a 6-Month Period. *Jpn. J. Infect. Dis.*, 2010. 63(1):65-66.
 31. 林 昌宏. チクングニヤウイルス. *臨床と微生物*, 36(3): 211-216, 2009
 32. Saijo, M.: Emerging and re-emerging infection threats to society. *J. Disaster Res.* 4:291-297, 2009
 33. Morimoto, K., Saijo, M.: Imported rabies cases and preparedness for rabies in Japan. *J. Disaster Res.* 4:346-357, 2009
 34. Matsuyama S, and Taguchi F. (2009) Two step conformational changes mediated by receptor binding and proteolysis of a coronavirus envelope glycoprotein. *J. Virol.* 83: 11133-11141
 35. Fukushima A, Fukuda N, Lai Y, Ueno T, Moriyama M, Taguchi F, Iguchi A, Shimizu K and Kuroda K. (2009) Development of a chimeric DNA-RNA hammerhead ribozyme targeting SARS virus. *Intervirology* 52: 99-92
 36. Terahara K, Yoshida M, Taguchi F, Igarashi O, Nochi T, Gotoh Y, Yamamoto T, Tsunetsugu-Yokota Y, Beauchemin N, and Kiyono H. (2009) Expression of newly identified secretory CEACAM1a isoforms in the intestinal epithelium. *Biochem. Biophys. Res. Commun.* 383: 340-346
 37. Ishii K, Hasegawa H, Nagata N, Ami Y, Fukushi S, Taguchi F, and Tsunetsugu- Yokota Y. (2009) Neutralizing antibody against severe acute respiratory syndrome (SARS)-coronavirus spike is highly effective for the protection of mice in the murine SARS model. *Microbiol. Immunol.* 53:75-82.
 38. Shirato K, and Taguchi F. (2009) Indirect induction of mast cell degranulation and migration

- by respiratory syncytial virus infection. *Virology* 386: 88-93
39. Kawase M, Shirato K, Matsuyama S and Taguchi F. (2009) Protease-mediated entry via endosome of human coronavirus 229E. *J. Virol.* 83: 712-721.
 40. Nakamura S, Yang CS, Sakon N, Ueda M, Tougan T, Yamashita A, Goto N, Takahashi K, Yasunaga T, Ikuta K, Mizutani T, Okamoto Y, Tagami M, Morita R, Maeda N, Kawai J, Hayashizaki Y, Nagai Y, Horii T, Iida T, Nakaya T. Direct metagenomic detection of viral pathogens in nasal and fecal specimens using an unbiased high-throughput sequencing approach. *PLoS ONE.* 2009. 4:e4219.
 41. Shumpei Watanabe, Naoya Ueda, Koichiro Iha, Joseph S. Masangkay, Hikaru Fujii, Phillip Alviola, Tetsuya Mizutani, Ken Maeda, Daisuke Yamane, Azab Walid, Kentaro Kato, Shigeru Kyuwa, Yukinobu Tohya, Yasuhiro Yoshikawa, Hiroomi Akashi. Detection of a new bat gammaherpesvirus in the Philippines. *Virus Genes.* 2009. 39. 90-93.
 42. Tomomitsu Satho, Hamady Dieng, Tetsuya Mizutani, Yuki Eshita, Takeshi Miyata, Parimal Talukder, Nobuhiro Kashige, Abu Hassan Ahmad and Fumio Miake. Fluorescence can be used to trace the fate of exogenous micro-organisms inside the alimentary tract of mosquitoes. *Journal of Parasitology and Vector Biology.* 1. 013-018. 2009

