

5. Betts, J. C., P. T. Lukey, L. C. Robb, R. A. McAdam, and K. Duncan. 2002. Evaluation of a nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and protein expression profiling. *Mol. Microbiol.* **43**: 717–731.
6. Bloom, B. R. 2002. Tuberculosis—the global view. *N. Engl. J. Med.* **346**: 1434–1435.
7. Brennan, P. J., and H. Nikaido. 1995. The envelope of mycobacteria. *Annu. Rev. Biochem.* **64**:29–63.
8. Chen, C. C., H. Y. Wu, M. Y. Chou, C. H. Huang, and A. Majumder. 2005. LeuO protein delimits the transcriptionally active and repressive domains on the bacterial chromosome. *J. Biol. Chem.* **280**:15111–15421.
9. Cole, S. T., R. Brosch, J. Parkhill, T. Garnier, C. Churcher, D. Harris, S. V. Gordon, K. Eiglmeier, S. Gas, C. E. Barry III, F. Tekaiia, K. Badcock, D. Basham, D. Brown, T. Chillingworth, R. Connor, R. Davies, K. Devlin, T. Feltwell, S. Gentles, N. Hamlin, S. Holroyd, T. Hornsby, K. Jagels, A. Krogh, J. McLean, S. Moule, L. Murphy, K. Oliver, J. Osborne, M. A. Quail, M. A. Rajandream, J. Rogers, S. Rutter, K. Seeger, J. Skelton, R. Squares, S. Squares, J. E. Sulston, K. Taylor, S. Whitehead, and B. G. Barrell. 1998. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* **393**:537–544.
10. Cole, S. T., K. Eiglmeier, J. Parkhill, K. D. James, N. R. Thomson, P. R. Wheeler, N. Honore, T. Garnier, C. Churcher, D. Harris, K. Mungall, D. Basham, D. Brown, T. Chillingworth, R. Connor, R. M. Davies, K. Devlin, S. Duthoy, T. Feltwell, A. Fraser, N. Hamlin, S. Holroyd, T. Hornsby, K. Jagels, C. Lacroix, J. Maclean, S. Moule, L. Murphy, K. Oliver, M. A. Quail, M. A. Rajandream, K. M. Rutherford, S. Rutter, K. Seeger, S. Simon, M. Simmonds, J. Skelton, R. Squares, S. Squares, K. Stevens, K. Taylor, S. Whitehead, J. R. Woodward, and B. G. Barrell. 2001. Massive gene decay in the leprosy bacillus. *Nature* **409**:1007–1011.
11. Costanzi, C., and J. R. Pehrson. 1998. Histone macroH2A1 is concentrated in the inactive X chromosome of female mammals. *Nature* **393**:599–601.
12. De Voss, J. J., K. Rutter, B. G. Schroeder, H. Su, Y. Zhu, and C. E. Barry III. 2000. The salicylate-derived mycobactin siderophores of *Mycobacterium tuberculosis* are essential for growth in macrophages. *Proc. Natl. Acad. Sci. USA* **97**:1252–1257.
13. Fleischmann, R. D., D. Alland, J. A. Eisen, L. Carpenter, O. White, J. Peterson, R. DeBoy, R. Dodson, M. Gwinn, D. Haft, E. Hickey, J. F. Kolonay, W. C. Nelson, L. A. Umayam, M. Ermolaeva, S. L. Salzberg, A. Delcher, T. Utterback, J. Weidman, H. Khouri, J. Gill, A. Mikula, W. Bishai, W. R. Jacobs, Jr., J. C. Venter, and C. M. Fraser. 2002. Whole-genome comparison of *Mycobacterium tuberculosis* clinical and laboratory strains. *J. Bacteriol.* **184**:5479–5490.
14. Fujita, Y., T. Naka, T. Doi, and I. Yano. 2005. Direct molecular mass determination of trehalose monomycolate from 11 species of mycobacteria by MALDI-TOF mass spectrometry. *Microbiology* **151**:1443–1452.
15. Harlow, E., and D. Lane. 1988. *Antibodies: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
16. Jackson, M., C. Reynaud, M. A. Laneelle, C. Guilhot, C. Laurent-Winter, D. Ensergueix, B. Gicquel, and M. Daffe. 1999. Inactivation of the antigen 85C gene profoundly affects the mycolate content and alters the permeability of the *Mycobacterium tuberculosis* cell envelope. *Mol. Microbiol.* **31**:1573–1587.
17. Jacobs, W. R., Jr., M. Tuckman, and B. R. Bloom. 1987. Introduction of foreign DNA into mycobacteria using a shuttle phasmid. *Nature* **327**:532–535.
18. Karakousis, P. C., T. Yoshimatsu, G. Lamichhane, S. C. Woolwine, E. L. Nuermberger, J. Grosset, and W. R. Bishai. 2004. Dormancy phenotype displayed by extracellular *Mycobacterium tuberculosis* within artificial granulomas in mice. *J. Exp. Med.* **200**:647–657.
19. Lee, B. H., B. Murugasu-Oei, and T. Dick. 1998. Upregulation of a histone-like protein in dormant *Mycobacterium smegmatis*. *Mol. Gen. Genet.* **260**: 475–479.
20. Liu, J., C. E. Barry III, G. S. Besra, and H. Nikaido. 1996. Mycolic acid structure determines the fluidity of the mycobacterial cell wall. *J. Biol. Chem.* **271**:29545–29551.
21. Lucchini, S., G. Rowley, M. D. Goldberg, D. Hurd, M. Harrison, and J. C. Hinton. 2006. H-NS mediates the silencing of laterally acquired genes in bacteria. *PLoS Pathog.* **2**:e81.
22. Manabe, Y. C., and W. R. Bishai. 2000. Latent *Mycobacterium tuberculosis*—persistence, patience, and winning by waiting. *Nat. Med.* **6**:1327–1329.
23. Matsumoto, S., M. Furugen, H. Yukitake, and T. Yamada. 2000. The gene encoding mycobacterial DNA-binding protein 1 (MDP1) transformed rapidly growing bacteria to slowly growing bacteria. *FEMS Microbiol. Lett.* **182**:297–301.
24. Matsumoto, S., M. Matsumoto, K. Umemori, Y. Ozeki, M. Furugen, T. Tatsuo, Y. Hirayama, S. Yamamoto, T. Yamada, and K. Kobayashi. 2005. DNA augments antigenicity of mycobacterial DNA-binding protein 1 and confers protection against *Mycobacterium tuberculosis* infection in mice. *J. Immunol.* **175**:441–449.
25. Matsumoto, S., M. Tamaki, H. Yukitake, T. Matsuo, M. Naito, H. Teraoka, and T. Yamada. 1996. A stable *Escherichia coli*-mycobacteria shuttle vector 'pSO246' in *Mycobacterium bovis* BCG. *FEMS Microbiol. Lett.* **135**:237–243.
26. Matsumoto, S., H. Yukitake, M. Furugen, T. Matsuo, T. Mineta, and T. Yamada. 1999. Identification of a novel DNA-binding protein from *Mycobacterium bovis* bacillus Calmette-Guerin. *Microbiol. Immunol.* **43**:1027–1036.
27. Mikusova, K., M. Mikus, G. S. Besra, I. Hancock, and P. J. Brennan. 1996. Biosynthesis of the linkage region of the mycobacterial cell wall. *J. Biol. Chem.* **271**:7820–7828.
28. Naito, M., N. Ohara, S. Matsumoto, and T. Yamada. 1998. Immunological characterization of alpha antigen of *Mycobacterium kansasii*: B-cell epitope mapping. *Scand. J. Immunol.* **48**:73–78.
29. Naito, M., N. Ohara, S. Matsumoto, and T. Yamada. 1998. The novel fibronectin-binding motif and key residues of mycobacteria. *J. Biol. Chem.* **273**:2905–2909.
30. Park, H. D., K. M. Guinn, M. I. Harrell, R. Liao, M. I. Voskuil, M. Tompa, G. K. Schoolnik, and D. R. Sherman. 2003. Rv3133c/dosR is a transcription factor that mediates the hypoxic response of *Mycobacterium tuberculosis*. *Mol. Microbiol.* **48**:833–843.
31. Pethe, K., P. Bifani, H. Drobecq, C. Sergheraert, A. S. Debric, C. Locht, and F. D. Menozzi. 2002. Mycobacterial heparin-binding hemagglutinin and laminin-binding protein share antigenic methyllysines that confer resistance to proteolysis. *Proc. Natl. Acad. Sci. USA* **99**:10759–10764.
32. Randall, L. L., and S. J. Hardy. 1995. High selectivity with low specificity: how SecB has solved the paradox of chaperone binding. *Trends Biochem. Sci.* **20**:65–69.
33. Ronning, D. R., T. Klabunde, G. S. Besra, V. D. Vissa, J. T. Belisle, and J. C. Sacchettini. 2000. Crystal structure of the secreted form of antigen 85C reveals potential targets for mycobacterial drugs and vaccines. *Nat. Struct. Biol.* **7**:141–146.
34. Sassetti, C. M., D. H. Boyd, and E. J. Rubin. 2003. Genes required for mycobacterial growth defined by high density mutagenesis. *Mol. Microbiol.* **48**:77–84.
35. Schnappinger, D., S. Ehrt, M. I. Voskuil, Y. Liu, J. A. Mangan, I. M. Monahan, G. Dolganov, B. Efron, P. D. Butcher, C. Nathan, and G. K. Schoolnik. 2003. Transcriptional adaptation of *Mycobacterium tuberculosis* within macrophages: insights into the phagosomal environment. *J. Exp. Med.* **198**:693–704.
36. Shimoji, Y., V. Ng, K. Matsumura, V. A. Fischetti, and A. Rambukkana. 1999. A 21-kDa surface protein of *Mycobacterium leprae* binds peripheral nerve laminin-2 and mediates Schwann cell invasion. *Proc. Natl. Acad. Sci. USA* **96**:9857–9862.
37. Shires, K., and L. Steyn. 2001. The cold-shock stress response in *Mycobacterium smegmatis* induces the expression of a histone-like protein. *Mol. Microbiol.* **39**:994–1009.
38. Soares de Lima, C., L. Zulianello, M. A. Marques, H. Kim, M. I. Portugal, S. L. Antunes, F. D. Menozzi, T. H. Ottenhoff, P. J. Brennan, and M. C. Pessolani. 2005. Mapping the laminin-binding and adhesive domain of the cell surface-associated Hlp/LBP protein from *Mycobacterium leprae*. *Microbes Infect.* **7**:1097–1109.
39. Stover, C. K., V. F. de la Cruz, T. R. Fuerst, J. E. Burlein, L. A. Benson, L. T. Bennett, G. P. Bansal, J. F. Young, M. H. Lee, G. F. Hatfull, et al. 1991. New use of BCG for recombinant vaccines. *Nature* **351**:456–460.
40. Takayama, K., C. Wang, and G. S. Besra. 2005. Pathway to synthesis and processing of mycolic acids in *Mycobacterium tuberculosis*. *Clin. Microbiol. Rev.* **18**:81–101.
41. Tropis, M., X. Meniche, A. Wolf, H. Gebhardt, S. Strelkov, M. Chami, D. Schomburg, R. Kramer, S. Murbach, and M. Daffe. 2005. The crucial role of trehalose and structurally related oligosaccharides in the biosynthesis and transfer of mycolic acids in Corynebacteriaceae. *J. Biol. Chem.* **280**:26573–26585.
42. Voskuil, M. I., D. Schnappinger, K. C. Visconti, M. I. Harrell, G. M. Dolganov, D. R. Sherman, and G. K. Schoolnik. 2003. Inhibition of respiration by nitric oxide induces a *Mycobacterium tuberculosis* dormancy program. *J. Exp. Med.* **198**:705–713.
43. Wayne, L. G., and C. D. Sohaskey. 2001. Nonreplicating persistence of *Mycobacterium tuberculosis*. *Annu. Rev. Microbiol.* **55**:139–163.
44. Zhou, J., and Z. Xu. 2005. The structural view of bacterial translocation-specific chaperone SecB: implications for function. *Mol. Microbiol.* **58**:349–357.

Original article

## Contribution of GM-CSF on the enhancement of the T cell-stimulating activity of macrophages

Masahiko Makino\*, Yumi Maeda, Yasuo Fukutomi, Tetsu Mukai

Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan

Received 9 November 2005; accepted 10 October 2006

Available online 11 December 2006

### Abstract

*Mycobacterium leprae* is an intracellular parasitic organism that multiplies in macrophages (MØ). It inhibits the fusion of mycobacterial phagosome with lysosome and induces interleukin (IL)-10 production from macrophages. However, macrophages are heterogenous in various aspects. We examined macrophages that differentiated from monocytes using either recombinant (r) granulocyte-MØ colony-stimulating factor (GM-CSF) (these MØ are named as GM-MØ) or rMØ colony-stimulating factor (M-CSF) (cells named as M-MØ) in terms of the T cell-stimulating activity. Although both macrophages phagocytosed the mycobacteria equally, GM-MØ infected with *M. leprae* and subsequently treated with IFN- $\gamma$ - and CD40 ligand (L) stimulated T cells to produce interferon-gamma (IFN- $\gamma$ ), but M-MØ lacked the ability to stimulate T cells. While M-MØ mounted a massive IL-10 production, GM-MØ did not produce the cytokine on infection with *M. leprae*. *M. leprae*-infected, IFN- $\gamma$ - and CD40L-treated GM-MØ expressed a higher level of HLA-DR and CD86 Ags than those of M-MØ, and expressed one of the dominant antigenic molecules of *M. leprae*, Major Membrane Protein-II on their surface. These results indicate that GM-CSF, but not M-CSF, contributes to the up-regulation of the T cell-stimulating activity of *M. leprae*-infected macrophages.

© 2006 Elsevier Masson SAS. All rights reserved.

**Keywords:** Macrophage; *M. leprae*; GM-CSF; IFN- $\gamma$

### 1. Introduction

*Mycobacterium leprae* (*M. leprae*), a causative agent of human leprosy, is a representative parasitic pathogen that induces skin lesions and chronic progressive peripheral nerve injury, leading to systemic deformity [1,2]. Leprosy represents a clinical spectrum, in which clinical manifestations are associated with different levels of immune responses to *M. leprae*

infection [3]. One representative type is a tuberculoid leprosy, in which patients exhibit innate and adaptive immunities to *M. leprae* and manifest a localized form of the disease with granuloma formation in infected tissues [4–6]. For the activation of an adaptive immunity, dendritic cells (DC) derived from inflammatory monocytes, play a central role [7,8]; and, in *in vitro* experiments, both CD4<sup>+</sup> and CD8<sup>+</sup> T cells are activated by DC infected with *M. leprae*, and these DC expressed Major Membrane Protein-II (MMP-II) as a dominant antigenic molecule [9,10]. Another representative manifestation is lepromatous leprosy, in which patients show reduced levels of host defense associated immunities and manifest a disseminated form of the disease with a broad spread of foamy MØ, in which an abundance of bacilli are usually involved [11,12]. *M. leprae* resides in the phagosome in MØ and replicates there without being digested by lysosomal enzymes [13]. Furthermore, *M. leprae* stimulates MØ to produce IL-10 [5,6] and suppresses the DC-mediated Ag-specific

**Abbreviations:** Ag, antigen; APC, Ag-presenting cells; BCG, *Mycobacterium bovis* BCG; DC, dendritic cells; GFP-BCG, BCG expressing GFP; GM-CSF, granulocyte-macrophage colony-stimulating factor; IFN- $\gamma$ , interferon-gamma; IL, interleukin; L, ligand; mAb, monoclonal antibody; MØ, macrophages; M-CSF, MØ colony-stimulating factor; *M. leprae*, *Mycobacterium leprae*; MMP-II, Major Membrane Protein-II; PBMC, peripheral blood mononuclear cells; r, recombinant.

\* Corresponding author. Tel.: +81 42 391 8059; fax: +81 42 391 8212.

E-mail address: mmaki@nih.go.jp (M. Makino).

adaptive immunity [14,15]. These observations may indicate that the induction of intracellular processing of *M. leprae* and that of expression of molecules, such as MMP-II could lead to the activation of IFN- $\gamma$  producing type 1 CD4<sup>+</sup> T cells. Another important element that should be considered for the full activation of T cells is the suppression of IL-10 production from *M. leprae*-infected M $\phi$ .

So far, a variety of methods and tools, including cytokines, have been used for the differentiation of M $\phi$  from human peripheral monocytes *in vitro* [16–18]. One representative M $\phi$  can be differentiated by using M-CSF, termed M-M $\phi$ , and another by using GM-CSF, termed GM-M $\phi$ . Both M $\phi$  represent different functions on infection with mycobacteria. However, much remains not fully understood with regard to *M. leprae* infection and the T cell-stimulating activity of these M $\phi$ .

In this report, we analyzed the characteristics of *M. leprae*-infected GM-M $\phi$  and M-M $\phi$ , and tried to develop immunological methods to enhance the M $\phi$ -mediated host defense activities against the bacteria.

## 2. Materials and methods

### 2.1. Preparation of cells and bacteria

Peripheral blood was obtained from healthy PPD-positive individuals under informed consent. We are aware that PPD-negative individuals would help to provide more information for our study; however, in Japan, most healthy individuals are PPD-positive, because *Mycobacterium bovis* BCG vaccination is compulsory for children (0–4 years old). Moreover, PPD-negative individuals in the Japanese population are those who do not respond to BCG vaccination, and therefore, it is likely that they suffer from some immune insufficiency. Therefore, these individuals cannot be used as controls for our experiments. Peripheral blood mononuclear cells (PBMC) were isolated using Ficoll-Paque Plus (Pharmacia, Uppsala, Sweden) and cryopreserved in liquid nitrogen until use, as previously described [19]. For preparation of peripheral monocytes, CD3<sup>+</sup> T cells were removed from either freshly isolated heparinized blood, or cryopreserved PBMC using immunomagnetic beads coated with anti-CD3 monoclonal antibody (mAb) (Dynabeads 450, Dynal, Oslo, Norway). The CD3<sup>-</sup> PBMC fraction was plated on collagen-coated plates and the non-plastic adherent cells were removed by extensive washing. The remaining adherent cells were used as monocytes [19]. M $\phi$  were differentiated by culturing monocytes in the presence of 20% fetal calf serum and either rM-CSF (R and D Systems, Abingdon, UK) (M-M $\phi$ ) or rGM-CSF (Pepro Tech EC LTD, London, UK) (GM-M $\phi$ ) [20]. Both GM-M $\phi$  and M-M $\phi$  were pulsed with *M. leprae*, treated with an optimal dose of IFN- $\gamma$  on day 3 of culture, further treated with CD40L on day 4, and were used as a stimulator of T cells on day 5 [21]. *M. leprae* (Thai-53) was isolated from the footpads of BALB/c-*nulnu* mice [22]. The isolated bacteria were counted by Shepard's method [22]. Killed *M. leprae* was prepared by heating the bacteria at 60 °C for 18 h. BCG (Pasteur strain) was cultured *in vitro* using Middlebrook 7H9 broth

supplemented with 0.05% Tween 80 and albumin-dextrose-catalase. BCG expressing GFP was constructed as follows. The GFP sequence was amplified from pEGFP-1 vector (CLONTECH, Palo Alto, CA), and cloned into pMV261 [23]. Transformants were selected on a 7H10 plate containing 25  $\mu$ g/ml kanamycin. The phagocytosis of BCG by GM-M $\phi$  and M-M $\phi$  after culture was determined using FACScalibur (Becton Dickinson Immunocytometry System, San Jose, CA). The multiplicity of infection (MOI) was determined based on the assumption that M $\phi$  were equally susceptible to infection with *M. leprae* [24].

### 2.2. Analysis of cell surface antigen (Ag)

The expression of cell surface Ag on M $\phi$  was analyzed using FACScalibur. Dead cells were eliminated from the analysis by staining with propidium iodide (Sigma Chemical Co., St. Louis, MO) and  $1 \times 10^4$  live cells were analyzed. For analysis of cell surface Ag, the following mAbs were used: FITC-conjugated mAb against HLA-ABC (G46-2.6, PharMingen, San Diego, CA), HLA-DR (L243, PharMingen), CD14 (M5E2, BD Biosciences, San Jose, CA), TLR2 (TL2.3, Serotec, Oxford, UK), TLR4 (HTA125, Santa Cruz Biotech, Santa Cruz, CA), CD209 (DCN46, PharMingen), CD86 (FUN-1, PharMingen), and CD40 (5C3, PharMingen).

The expression of MMP-II, which is one of the dominant antigenic molecules of *M. leprae* [9] on *M. leprae*-infected M $\phi$  was determined using the mAb (IgM, kappa) against MMP-II, followed by FITC-conjugated anti-mouse Igs Ab (Tago-immunologicals, Camarillo, CA).

### 2.3. APC function of *M. leprae*-infected M $\phi$

The ability of *M. leprae*-infected M $\phi$  to stimulate T cells was assessed using an autologous M $\phi$ -T cell co-culture as previously described [24,25]. Freshly thawed PBMC were depleted of CD56<sup>+</sup>, MHC class II<sup>+</sup> and CD8<sup>+</sup> cells by using magnetic beads coated with mAb to CD56, MHC class II and CD8 Ags (Dynabeads 450; Dynal) [25]. The purity of CD4<sup>+</sup> T cells was more than 98% as assessed by FACS analyses. The purified responder cells ( $1 \times 10^5$  per well) were plated in 96-well round-bottom tissue culture plates and M $\phi$  were added to give the indicated M $\phi$ :CD4<sup>+</sup> T cell ratio. Supernatants of M $\phi$ -T cell co-cultures were collected on day 4 and the concentration of cytokines was determined.

### 2.4. Cytokine production

Levels of the following cytokines were measured; IFN- $\gamma$  produced by CD4<sup>+</sup> T cells, IL-10, IL-1 $\beta$ , TNF $\alpha$  and IL-12p40 produced by M $\phi$  stimulated for 24 h with *M. leprae*. The concentrations of these cytokines were quantified using the enzyme assay kits, Opt EIA Human ELISA Set (BD PharMingen International).

### 2.5. Statistical analysis

Student's *t*-test was applied to determine the statistical differences.

## 3. Results

### 3.1. Characteristics of MØ differentiated from monocytes using GM-CSF

MØ were differentiated from monocytes using either GM-CSF (GM-MØ) or M-CSF (M-MØ). To characterize these two types of MØ, surface markers expressed on GM-MØ and M-MØ were analyzed using the monocytes obtained from the same donor by flow cytometry (Fig. 1). MHC class I (HLA-ABC) and class II (HLA-DR) Ags were similarly expressed on GM-MØ and M-MØ, but the expression of CD14 Ag was significantly reduced in GM-MØ. While the expression level of TLR2, CD209, CD40 and TLR4 Ags was not different between GM-MØ and M-MØ, the expression of CD86 was significantly higher on GM-MØ than M-MØ. Then, we examined the phagocytic capacity of GM-MØ and M-MØ by using BCG expressing GFP (GFP-BCG), since *M. leprae* cannot be cultured *in vitro* or express GFP. The percentage of MØ expressing GFP after co-culture of MØ with GFP-BCG was similar

between GM-MØ and M-MØ (Fig. 2). These results indicate that GM-MØ and M-MØ differed in the expression of some surface markers, but they similarly phagocytosed the mycobacteria.

### 3.2. Effect of *M. leprae* infection to GM-MØ on the T cell-stimulating activity

Since *M. leprae* is an intracellular parasitic bacterium and is hardly digested with lysosomal enzyme in MØ unless MØ are activated [26], we analyzed the T cell-stimulating activity of *M. leprae*-infected GM-MØ and M-MØ (Table 1). When M-MØ were infected with up to MOI 80 of *M. leprae* and treated with IFN- $\gamma$  and CD40L, they did not stimulate CD4<sup>+</sup> T cells to secrete a significant dose of IFN- $\gamma$ . In contrast to M-MØ, when *M. leprae*-infected, IFN- $\gamma$ - and CD40L-treated GM-MØ were used as Ag-presenting cells (APC), T cells produced significant levels of IFN- $\gamma$  in a manner dependent on the dose of *M. leprae*. Since GM-MØ express CD40, and are activated by IFN- $\gamma$ , we examined the effect of treatment with IFN- $\gamma$  and CD40L on the T cell-stimulating activity of *M. leprae*-infected GM-MØ (Table 2). While IFN- $\gamma$  production from CD4<sup>+</sup> T cells was not significantly induced by GM-MØ untreated or treated with either IFN- $\gamma$  or CD40L, the cytokine production was significantly enhanced by the treatment of GM-MØ with




Fig. 1. Phenotype of GM-MØ and M-MØ differentiated from monocytes. Plastic adherent monocytes were differentiated into MØ by 3 days culture with either rGM-CSF or rM-CSF. Dashed lines, isotype-matched control IgG; solid lines, mAb staining. The number represents the difference in mean fluorescence intensity between dashed and solid lines. Representatives of three independent experiments are shown.




Fig. 2. Phagocytic activity of GM-MØ and M-MØ. Plastic adherent monocytes were differentiated into MØ by 3 days culture with either rGM-CSF or rM-CSF. For analysis of the phagocytic activity of MØ, GM-MØ and M-MØ were pulsed with BCG expressing GFP and assessed on day 4 of culture. Dashed lines, unpulsed cells; solid lines, GFP-BCG pulsed cells. The number represents the difference in mean fluorescence intensity between the dashed and solid lines. The number in parenthesis indicates the percent GFP-positive cell number. Representatives of three independent experiments are shown.

both IFN- $\gamma$  and CD40L. Then, we compared the T cell-stimulating activity of live and heat-killed *M. leprae* (Table 3). Both forms of *M. leprae* stimulated CD4<sup>+</sup> T cells when pulsed to GM-MØ, but the heat-killed *M. leprae* more efficiently induced T cell activation than live bacteria. When we examined the effect of heat-killed *M. leprae* on M-MØ, they did not stimulate CD4<sup>+</sup> T cells significantly, even when IFN- $\gamma$  and CD40L were administered (data not shown). Also note that, when GM-MØ and monocyte-derived DC were compared in terms of their T cell-stimulating activity, GM-MØ were less efficient in this respect (data not shown).

### 3.3. Factors associated with the enhancement of the T cell-stimulating activity of GM-MØ

Various factors may be responsible for enhancing the T cell-stimulating activity of APC. When we examined the expression of APC associated molecules on *M. leprae*-infected MØ (Fig. 3), the expression of HLA-DR and CD86 on GM-MØ was higher than on M-MØ, although there was no

significant difference in the expression of HLA-ABC between GM-MØ and M-MØ. The cytokines produced from APC should also be considered to be another important factor that should be monitored and MØ produce a variety of cytokines, including IL-10, IL-1 $\beta$ , TNF $\alpha$  and IL-12 [6,11,27]. IL-10 was efficiently produced from M-MØ by stimulation with *M. leprae*, but it was hardly produced from GM-MØ (Fig. 4a). When macrophages were differentiated by using both GM-CSF and M-CSF, the function of GM-CSF was dominant and, the production of IL-10 was suppressed (Fig. 4a). Similarly to the production of IL-10, IL-1 $\beta$  was more efficiently produced from M-MØ than GM-MØ (Fig. 4b). In contrast, TNF $\alpha$ , which is important for granuloma formation, was more efficiently produced from GM-MØ (Fig. 4c). However, there was no significant difference in the production of IL-12p40 between GM-MØ and M-MØ (Fig. 4d). Finally, we assessed whether *M. leprae*-infected GM-MØ expressed dominant antigenic molecules of *M. leprae* on the surface (Fig. 5). To this end, we examined the expression of MMP-II on GM-MØ and M-MØ. No apparent expression of

Table 1  
T cell-stimulating activity of *M. leprae*-infected GM-MØ and M-MØ<sup>a</sup>

| Stimulator of CD4 <sup>+</sup> T cells | <i>M. leprae</i> infection of macrophages (MOI) | IFN- $\gamma$ (pg/ml) production by CD4 <sup>+</sup> T cells after stimulation with macrophages at ratio (T:MØ) |                               |                               |
|--|---|---|-------------------------------|-------------------------------|
|  |   | 2:1   | 4:1                           | 8:1                           |
| GM-MØ                                  | 0   | 0.6 $\pm$ 0.2 <sup>a,†</sup>  | 0.5 $\pm$ 0.1 <sup>†,‡</sup>  | 1.4 $\pm$ 0.2 <sup>¶,  </sup> |
|  | 40  | 38.1 $\pm$ 3.8 <sup>*</sup>   | 34.2 $\pm$ 2.3 <sup>‡</sup>   | 23.4 $\pm$ 3.8 <sup>¶</sup>   |
|  | 80  | 230.7 $\pm$ 21.4 <sup>†</sup>   | 120.5 $\pm$ 16.9 <sup>‡</sup> | 74.7 $\pm$ 6.8 <sup>  </sup>  |
| M-MØ                                   | 0   | 0.9 $\pm$ 0.1   | 3.1 $\pm$ 1.2                 | 13.9 $\pm$ 2.2                |
|  | 40  | 0.9 $\pm$ 0.1   | 2.6 $\pm$ 1.3                 | 12.2 $\pm$ 3.1                |
|  | 80  | 11.8 $\pm$ 0.3  | 17.5 $\pm$ 2.1                | 12.2 $\pm$ 2.9                |

<sup>\*</sup> $p < 0.005$ ; <sup>†</sup> $p < 0.005$ ; <sup>‡</sup> $p < 0.01$ ; <sup>§</sup> $p < 0.01$ ; <sup>¶</sup> $p < 0.01$ ; <sup>||</sup> $p < 0.005$ .

<sup>a</sup> CD4<sup>+</sup> T cells ( $1 \times 10^5$ /well) were stimulated for 4 days with autologous GM-MØ or M-MØ at the indicated dose of macrophage. GM-MØ or M-MØ were pulsed with *M. leprae* and IFN- $\gamma$  (100 IU/ml) on day 3, treated with CD40L (1  $\mu$ g/ml) on day 4, and were used as APC on day 5. Representatives of three separate experiments are shown. Assays were done in triplicate, and results are expressed as mean  $\pm$  SD. Groups with identical symbols were compared using Student's *t*-test.

Table 2  
Contribution of IFN- $\gamma$  and CD40L on T cell-stimulating activity of GM-M $\phi$ <sup>a</sup>

| <i>M. leprae</i> infection of GM-M $\phi$ (MOI: 80) | Treatment of <i>M. leprae</i> -infected GM-M $\phi$ with |                        | IFN- $\gamma$ (pg/ml) production by CD4 <sup>+</sup> T cells after stimulation with GM-M $\phi$ at ratio (T:M $\phi$ ) |                                |                                 |
|---|--|------------------------|--|--------------------------------|---------------------------------|
|   | IFN- $\gamma$ (100 IU/ml)                                | CD40L (1.0 $\mu$ g/ml) | 2:1  | 4:1                            | 8:1                             |
| –   | +  | +                      | 2.3 $\pm$ 0.3  | 0.1 $\pm$ 0.2                  | 0.8 $\pm$ 0.5                   |
| +   | –  | –                      | 4.0 $\pm$ 1.1*   | 5.5 $\pm$ 1.9 <sup>h</sup>     | 6.0 $\pm$ 2.1 <sup>h</sup>      |
| +   | –  | +                      | 21.4 $\pm$ 3.1 <sup>†</sup>  | 22.7 $\pm$ 4.0 <sup>¶</sup>    | 14.8 $\pm$ 2.2 <sup>**</sup>    |
| +   | +  | –                      | 20.3 $\pm$ 1.7 <sup>†</sup>  | 15.9 $\pm$ 1.3 <sup>  </sup>   | 10.7 $\pm$ 2.3 <sup>††</sup>    |
| +   | +  | +                      | 226.1 $\pm$ 20.9 <sup>*††</sup>  | 107.8 $\pm$ 13.7 <sup>h*</sup> | 94.8 $\pm$ 9.7 <sup>h**††</sup> |

\* $p < 0.005$ ; <sup>†</sup> $p < 0.005$ ; <sup>††</sup> $p < 0.005$ ; <sup>‡</sup> $p < 0.005$ ; <sup>¶</sup> $p < 0.005$ ; <sup>||</sup> $p < 0.01$ ; <sup>h</sup> $p < 0.005$ ; <sup>\*\*</sup> $p < 0.005$ ; <sup>†††</sup> $p < 0.005$ .

<sup>a</sup> CD4<sup>+</sup> T cells ( $1 \times 10^5$ /well) were stimulated for 4 days with autologous GM-M $\phi$  at the indicated dose of macrophage. GM-M $\phi$  were pulsed with *M. leprae* and IFN- $\gamma$  (100 IU/ml) on day 3, treated with CD40L (1  $\mu$ g/ml) on day 4, and were used as APC on day 5. Representatives of three separate experiments are shown. Assays were done in triplicate, and the results are expressed as the mean  $\pm$  SD. Groups with identical symbols were compared using Student's *t*-test.

MMP-II was observed on M-M $\phi$ , but, on GM-M $\phi$ , significant expression of MMP-II was induced. The expression was dependent on the dose of *M. leprae* (Fig. 5). However, the MMP-II expression on *M. leprae*-infected GM-M $\phi$  required both IFN- $\gamma$  and CD40L, and apparent expression was not induced by sole treatment of macrophages with either IFN- $\gamma$  or CD40L (data not shown).

#### 4. Discussion

In order to avoid the intracellular multiplication and intercellular spread of *M. leprae*, the activation of adaptive immunity, especially of IFN- $\gamma$ -producing type 1 T cells, plays an important role [5,6]. In fact, paucibacillary (tuberculoid) leprosy patients activate CD4<sup>+</sup> T cells through DC, although the bacteria cannot be eliminated completely [8,28]. The *M. leprae*-infected DC digest the bacteria and express dominant antigenic molecules for the efficient IFN- $\gamma$  production from T cells [9]. In contrast, multibacillary (lepromatous) leprosy patients retain a large number of *M. leprae* in their M $\phi$ , and concordantly induce reduced levels or completely lack the ability to effectively stimulate T cells [11,12]. Since tissue resident M $\phi$  are heterogenous with regard to functional aspects [17,29], we assessed two different types of M $\phi$ : GM-M $\phi$  and M-M $\phi$ , and found that GM-M $\phi$ , but not M-M $\phi$ , stimulated T cells. GM-M $\phi$  were generated from monocytes using cytokine GM-CSF whilst M-M $\phi$  were produced using M-CSF.

Although there were some differences in the expression levels of MHC class I, II, CD14 and CD209 Ags on GM-M $\phi$  and M-M $\phi$ , both forms were equally susceptible to mycobacteria as far as phagocytosis of BCG-GFP was examined. However, there was a striking difference between *M. leprae*-infected GM-M $\phi$  and M-M $\phi$  in the expression of antigenic molecules; only GM-M $\phi$  expressed MMP-II, which is one of the dominant antigenic molecules capable of stimulating T cells in *M. leprae*-infected individuals. The induction of MMP-II expression on GM-M $\phi$  requires not only GM-CSF, but also the co-stimulation of M $\phi$  with IFN- $\gamma$  and CD40L. In case of *M. leprae*-infected DC, the phagosomal bacteria could be processed by lysosomal enzymes, and MMP-II expression was observed on DC [9]. The MMP-II expression observed on GM-M $\phi$  may indicate that at least some intracellular *M. leprae* were processed. However, the processing of *M. leprae* by GM-M $\phi$  still seemed partial, since the heat-killed *M. leprae* induced T cell activation more vigorously than live bacteria, and *M. leprae*-infected DC stimulated T cells more efficiently than GM-M $\phi$ , although other factors, such as an induction of IL-12, cannot be ruled out completely. The cell wall architecture including surface-exposed molecules, of heat-killed mycobacteria is globally altered [29,30], resulting in the exudation of some soluble antigenic molecules which may be feasibly digested in macrophages (unpublished observation). Therefore, T cells are more efficiently activated by heat-killed bacteria than by live bacteria.

Table 3  
Comparison of T cell-stimulating activity of live and heat-inactivated *M. leprae*<sup>a</sup>

| <i>M. leprae</i> pulsed on GM-M $\phi$ (MOI) | IFN- $\gamma$ (pg/ml) production by CD4 <sup>+</sup> T cells after stimulation with GM-M $\phi$ at ratio (T:M $\phi$ ) |                               |                                |
|--|--|-------------------------------|--------------------------------|
|  | 2:1  | 4:1                           | 8:1                            |
| None   | 2.3 $\pm$ 1.1  | 2.1 $\pm$ 1.2                 | 2.4 $\pm$ 0.9                  |
| HK (40)                                      | 406.5 $\pm$ 49.3*  | 157.3 $\pm$ 20.1 <sup>†</sup> | 75.4 $\pm$ 6.8 <sup>¶</sup>    |
| HK (80)                                      | 399.8 $\pm$ 33.2 <sup>†</sup>  | 187.7 $\pm$ 17.8 <sup>h</sup> | 106.9 $\pm$ 11.2 <sup>  </sup> |
| Live (40)                                    | 101.5 $\pm$ 8.8*   | 30.2 $\pm$ 4.6 <sup>†</sup>   | 3.2 $\pm$ 1.9 <sup>¶</sup>     |
| Live (80)                                    | 152.0 $\pm$ 12.7 <sup>†</sup>  | 82.9 $\pm$ 7.4 <sup>h</sup>   | 32.7 $\pm$ 2.8 <sup>  </sup>   |

\* $p < 0.01$ ; <sup>†</sup> $p < 0.005$ ; <sup>††</sup> $p < 0.005$ ; <sup>‡</sup> $p < 0.005$ ; <sup>¶</sup> $p < 0.005$ ; <sup>||</sup> $p < 0.005$ .

<sup>a</sup> CD4<sup>+</sup> T cells ( $1 \times 10^5$ /well) were stimulated for 4 days with autologous GM-M $\phi$  at the indicated dose of macrophage. GM-M $\phi$  were pulsed with either heat-killed (HK) or live *M. leprae* and IFN- $\gamma$  (100 IU/ml) on day 3, treated with CD40L (1  $\mu$ g/ml) on day 4, and were used as APC on day 5. Representatives of three separate experiments are shown. Assays were done in triplicate, and the results are expressed as the mean  $\pm$  SD. Groups with identical symbols were compared using Student's *t*-test.

**Macrophage  
(*M. leprae*)**




Fig. 3. Phenotype of *M. leprae*-infected GM-MØ and M-MØ. GM-MØ and M-MØ differentiated from monocytes by 3 days culture with rGM-CSF or rM-CSF were infected with *M. leprae*, treated with IFN- $\gamma$  (100 IU/ml) on day 3, and further treated with CD40L (1  $\mu$ g/ml) on day 4 of culture. On day 5, the phenotype of GM-MØ and M-MØ was analyzed. Dashed lines, isotype-matched control IgG; solid lines, mAb staining. The number represents the difference in mean fluorescence intensity between dashed and solid lines. Representatives of three independent experiments are shown.

Ottenhoff et al. have also reported that GM-CSF up-regulates the T cell-stimulating activity of MØ, but not M-CSF, and mycobacteria-infected GM-MØ promoted the type 1 cell-mediated immunity against pathogens [31]. Our

observations are in line with their data and provide ways to enhance the cell-mediated immunity, especially in cases progressing towards lepromatous leprosy. To facilitate the T cell activation and MMP-II expression, it was required to use




Fig. 4. Cytokine production from GM-MØ and M-MØ. MØ were differentiated by 3 days culture with rGM-CSF, rM-CSF or rGM-CSF + rM-CSF, and were stimulated with *M. leprae* for 24 h. The cytokines: (a) IL-10; (b) IL-1 $\beta$ ; (c) TNF $\alpha$ ; and (d) IL-12p40 were measured by ELISA. Representatives of three independent experiments are shown.




Fig. 5. Expression of MMP-II on the surface of GM-MØ and M-MØ. GM-MØ and M-MØ were differentiated from monocytes by 3 days culture with either rGM-CSF or rM-CSF, respectively. These macrophages were infected with an indicated dose of *M. leprae* and treated with IFN- $\gamma$  (100 IU/ml) on day 3 of monocyte culture, further treated with CD40L (1  $\mu$ g/ml) on day 4, and were analyzed for MMP-II expression on day 5. Dashed lines, control IgM; solid lines, MMP-II mAb staining. The number represents the difference in mean fluorescence intensity between dashed and solid lines. Representatives of three independent experiments are shown.

both IFN- $\gamma$  and CD40L, in addition to the lineage-determining cytokine GM-CSF. IFN- $\gamma$  and CD40L are probably required to compensate for the lower antigenic characteristics of *M. leprae* [24]. Studies using other mycobacteria, such as *M. bovis* BCG, may provide further useful information. Although IFN- $\gamma$  is known to activate MØ for bacterial digestion and to induce IL-12p35 gene transcription [26,31], it remains to be determined if CD40L treatment on GM-MØ furthers the intracellular processing of phagosomal bacteria or whether conditioning of GM-MØ through CD40-CD40L interaction, such as in DC [32], is required for T cell stimulation. In addition to MMP-II expression, there were some differences in the phenotypic features of *M. leprae*-infected GM-MØ and M-MØ. A higher level of HLA-DR and CD86 Ags was expressed on the GM-MØ infected with *M. leprae* and co-stimulated, than on similarly treated M-MØ. The mechanism, leading to the enhanced Ag expression, especially of Ag processing by IFN- $\gamma$ , has not been clearly demonstrated and remains to be elucidated, but GM-CSF, IFN- $\gamma$  and CD40L seem to at least partially co-ordinate and induce the higher expression of HLA-DR and CD86. These phenotypic differences between *M. leprae*-infected GM-MØ and M-MØ again contribute to the differences in T cell stimulatory activity.

Another peculiar difference observed between the two types of MØ, GM-MØ and M-MØ, was the cytokines that they induced. IL-10 suppresses DC-dependent as well as DC-independent T cell activation [14,15,33], and creates a situation in which *M. leprae* can feasibly parasitize in the cells. While M-MØ secreted a large amount of IL-10, GM-MØ completely lacked in the production of cytokine upon stimulation with *M. leprae* (Fig. 4) or lipopolysaccharide (data not shown). Furthermore, the presence of GM-CSF diminished the IL-10 production from M-MØ by *M. leprae*. Thus, treatment of monocytes with GM-CSF can wipe off the favorable conditions for *M. leprae* survival. On the other hand, GM-MØ produced a higher level of TNF $\alpha$  than M-MØ. TNF $\alpha$  plays an

important role in the granuloma formation, and TNF $\alpha$  is an important mediator of host defense activity in MØ, in mycobacterial lesions [34,35]. The treatment of monocyte with GM-CSF would be beneficial for MØ-mediated host defense in this respect. These observations were consistent with the previous findings that IL-10-deficient mice display increased antimycobacterial immunity with concordant higher levels of TNF $\alpha$  and a lower bacterial burden [36]. Our previous studies show that T cells from lepromatous leprosy can mount a significant production of IFN- $\gamma$  by appropriate stimulation [10], therefore, the present studies may provide useful information for the development of immunotherapeutic tools, such as endogenous or exogenous treatment of macrophages with GM-CSF, and thus prevent the dissemination of *M. leprae*.

In this study, we analyzed the two types of MØ with regard to T cell-stimulating activity, and found that GM-CSF and co-stimulators enhance the host defense activity of *M. leprae*-infected MØ.

#### Acknowledgments

We acknowledge the contribution of Ms. N. Makino in the preparation of this manuscript. We also thank Ms. Y. Harada for technical support, and the Japanese Red Cross Society for kindly providing PBMC from healthy donors.

This work was supported in part by a Grant-in-Aid for Research on Emerging and Re-emerging Infectious Diseases and by a Grant-in-Aid for Research on HIV/AIDS from the Ministry of Health, Labour and Welfare of Japan.

#### References

- [1] G.L. Stoner, Importance of the neural predilection of *Mycobacterium leprae* in leprosy, *Lancet* 2 (1979) 994–996.



- [2] C.K. Job. Nerve damage in leprosy, *Int. J. Lepr. Other Mycobact. Dis.* 57 (1989) 532–539.
- [3] S.N.C. Wemambu, J.L. Turk, M.F.R. Waters, R.J.W. Rees. *Erythema nodosum leprosum*: a clinical manifestation of the Arthus phenomenon, *Lancet* 2 (1969) 933–935.
- [4] S. Verghese, D.G. Healey, J. Curtis, J.L. Turk. Accessory cell function of dendritic cells from lymph nodes containing *Mycobacterium leprae* induced granulomas, *Int. Arch. Allergy Appl. Immunol.* 87 (1988) 392–399.
- [5] P. Salgame, J.S. Abrams, C. Clayberger, H. Goldstein, J. Convit, R.L. Modlin, B.R. Bloom. Differing lymphokine profiles of functional subsets of human CD4 and CD8 T cell clones, *Science* 254 (1991) 279–282.
- [6] M. Yamamura, K. Uyemura, R.J. Deans, K. Weinberg, T.H. Rea, B.R. Bloom, R.L. Modlin. Defining protective responses to pathogens: cytokine profiles in leprosy lesions, *Science* 254 (1991) 277–279.
- [7] P.S. Yamauchi, J.R. Bleharski, K. Uyemura, J. Kim, P.A. Sieling, A. Miller, H. Brightbill, K. Schlienger, T.H. Rea, R.L. Modlin. A role for CD40–CD40 ligand interactions in the generation of type 1 cytokine responses in human leprosy, *J. Immunol.* 165 (2000) 1506–1512.
- [8] S.R. Krutzik, B. Tan, H. Li, M.T. Ochoa, P.T. Liu, S.E. Sharfstein, T.G. Graeber, P.A. Sieling, Y.-J. Liu, T.H. Rea, B.R. Bloom, R.L. Modlin. TLR activation triggers the rapid differentiation of monocytes into macrophages and dendritic cells, *Nat. Med.* 11 (2005) 653–660.
- [9] Y. Maeda, T. Mukai, J. Spencer, M. Makino. Identification of an immunomodulating agent from *Mycobacterium leprae*, *Infect. Immun.* 73 (2005) 2744–2750.
- [10] M. Makino, Y. Maeda, N. Ishii. Immunostimulatory activity of major membrane protein-II from *Mycobacterium leprae*, *Cell. Immunol.* 233 (2005) 53–60.
- [11] D.S. Ridley, W.H. Jopling. Classification of leprosy according to immunity. A five-group system, *Int. J. Lepr. Other Mycobact. Dis.* 34 (1966) 255–273.
- [12] D.A. Hagge, N.A. Ray, J.L. Krahenbuhl, L.B. Adams. An *in vitro* model for the lepromatous leprosy granuloma: fate of *Mycobacterium leprae* from target macrophages after interaction with normal and activated effector macrophages, *J. Immunol.* 172 (2004) 7771–7779.
- [13] C. Frehel, N. Rastogi. *Mycobacterium leprae* surface components intervene in the early phagosome–lysosome fusion inhibition event, *Infect. Immun.* 55 (1987) 2916–2921.
- [14] H. Jonuleit, E. Schmitt, K. Steinbrink, A.H. Enk. Dendritic cells as a tool to induce anergic and regulatory T cells, *Trends Immunol.* 22 (2001) 394–400.
- [15] A. Granelli-Piperno, A. Golebiowska, C. Trumpfheller, F.P. Siegal, R.M. Steinman. HIV-1-infected monocyte-derived dendritic cells do not undergo maturation but can elicit IL-10 production and T cell regulation, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 7669–7674.
- [16] K. Nakata, K.S. Akagawa, M. Fukayama, Y. Hayashi, M. Kadokura, T. Tokunaga. Granulocyte-macrophage colony-stimulating factor promotes the proliferation of human alveolar macrophages *in vitro*, *J. Immunol.* 147 (1991) 1266–1272.
- [17] K.S. Akagawa. Functional heterogeneity of colony-stimulating factor-induced human monocyte-derived macrophages, *Int. J. Hematol.* 76 (2002) 27–34.
- [18] M.U. Martin, H. Wesche. Summary and comparison of the signaling mechanisms of the Toll/interleukin-1 receptor family, *Biochim. Biophys. Acta* 1592 (2002) 265–280.
- [19] M. Makino, M. Baba. A cryopreservation method of human peripheral blood mononuclear cells for efficient production of dendritic cells, *Scand. J. Immunol.* 45 (1997) 618–622.
- [20] G.J. Randolph, K. Inaba, D.F. Robbani, R.M. Steinman, W.A. Muller. Differentiation of phagocytic monocytes into lymph node dendritic cells *in vivo*, *Immunity* 11 (1999) 753–761.
- [21] M. Makino, S. Shimokubo, S. Wakamatsu, S. Izumo, M. Baba. The role of human T-lymphotropic virus type 1 (HTLV-1)-infected dendritic cells in the development of HTLV-1-associated myelopathy/tropical spastic paraparesis, *J. Virol.* 73 (1999) 4575–4581.
- [22] R.D. McDermott-Lancaster, T. Ito, K. Kohsaka, C.C. Guelpa-Lauras, J.H. Grosset. Multiplication of *Mycobacterium leprae* in the nude mouse, and some applications of nude mice to experimental leprosy, *Int. J. Lepr. Other Mycobact. Dis.* 55 (1987) 889–895.
- [23] C.K. Stover, V.F. de la Cruz, T.R. Fuerst, J.E. Burlein, L.A. Benson, L.T. Bennett, G.P. Bansal, J.F. Young, M.H. Lee, G.F. Hatfull, S.B. Snapper, R.G. Barletta, W.R. Jacobs Jr., B.R. Bloom. New use of BCG for recombinant vaccines, *Nature* 351 (1991) 456–460.
- [24] K. Hashimoto, Y. Maeda, H. Kimura, K. Suzuki, A. Masuda, M. Matsuoka, M. Makino. *Mycobacterium leprae* infection in monocyte-derived dendritic cells and its influence on antigen presenting function, *Infect. Immun.* 70 (2002) 5167–5176.
- [25] S. Wakamatsu, M. Makino, C. Tei, M. Baba. Monocyte-driven activation-induced apoptotic cell death of human T-lymphotropic virus type 1-infected T cells, *J. Immunol.* 163 (1999) 3914–3919.
- [26] H. Kimura, Y. Maeda, F. Takeshita, L.E. Takaoka, M. Matsuoka, M. Makino. Upregulation of T-cell-stimulating activity of mycobacteria-infected macrophages, *Scand. J. Immunol.* 60 (2004) 278–286.
- [27] M.M. Stefani, C.M. Martelli, T.P. Gillis, J.L. Krahenbuhl, Brazilian Leprosy Study Group. *In situ* type 1 cytokine gene expression and mechanisms associated with early leprosy progression, *J. Infect. Dis.* 188 (2003) 1024–1031.
- [28] P.A. Sieling, D. Jullien, M. Dahlem, T.F. Tedder, T.H. Rea, R.L. Modlin, S.A. Porcelli. CD1 expression by dendritic cells in human leprosy lesions: correlation with effective host immunity, *J. Immunol.* 162 (1999) 1851–1858.
- [29] P.B. Kang, A.K. Azad, J.B. Torrelles, T.M. Kaufman, A. Beharka, E. Tibesar, L.E. Desjardin, L.S. Schlesinger. The human macrophage mannose receptor directs *Mycobacterium tuberculosis* lipoarabinomannan-mediated phagosome biogenesis, *J. Exp. Med.* 202 (2005) 987–999.
- [30] I. Vergne, J. Chua, H.H. Lee, M. Lucas, J. Belisle, V. Deretic. Mechanism of phagolysosome biogenesis block by viable *Mycobacterium tuberculosis*, *Proc. Natl. Acad. Sci. U.S.A.* 102 (2005) 4033–4038.
- [31] F.A. Verreck, T. de Boer, D.M. Langenberg, M.A. Hoeve, M. Kramer, E. Vaisberg, R. Kastelein, A. Kolk, R. de Waal-Malefyt, T.H. Ottenhoff. Human IL-23-producing type 1 macrophages promote but IL-10-producing type 2 macrophages subvert immunity to (myco)bacteria, *Proc. Natl. Acad. Sci. U.S.A.* 101 (2004) 4560–4565.
- [32] C. Caux, C. Massacrier, B. Vanbervliet, B. Dubois, C. Van Kooten, I. Durand, J. Banchereau. Activation of human dendritic cells through CD40 cross-linking, *J. Exp. Med.* 180 (1994) 1263–1272.
- [33] K. Mochida-Nishimura, K.S. Akagawa, E.A. Rich. Interleukin-10 contributes development of macrophage suppressor activities by macrophage colony-stimulating factor, but not by granulocyte-macrophage colony-stimulating factor, *Cell. Immunol.* 214 (2001) 81–88.
- [34] J.H. Davis, H. Clay, J.L. Lewis, N. Ghorri, P. Herbomel, L. Ramakrishnan. Real-time visualization of mycobacterium–macrophage interactions leading to initiation of granuloma formation in zebrafish embryos, *Immunity* 17 (2002) 693–702.
- [35] E. Giacomini, E. Iona, L. Ferroni, M. Miettinen, L. Fattorini, G. Orefici, I. Julkunen, E.M. Coccia. Infection of human macrophages and dendritic cells with *Mycobacterium tuberculosis* induces a differential cytokine gene expression that modulates T cell response, *J. Immunol.* 166 (2001) 7033–7041.
- [36] P.J. Murray, R.A. Young. Increased antimycobacterial immunity in interleukin-10-deficient mice, *Infect. Immun.* 67 (1999) 3087–3095.



# Evaluation of major membrane protein-II as a tool for serodiagnosis of leprosy

Yumi Maeda<sup>1</sup>, Tetsu Mukai<sup>1</sup>, Masanori Kai<sup>1</sup>, Yasuo Fukutomi<sup>1</sup>, Hiroko Nomaguchi<sup>1</sup>, Chiyoji Abe<sup>3</sup>, Kazuo Kobayashi<sup>4</sup>, Seigo Kitada<sup>5</sup>, Ryoji Maekura<sup>5</sup>, Ikuya Yano<sup>6</sup>, Norihisa Ishii<sup>2</sup>, Toru Mori<sup>1,2</sup> & Masahiko Makino<sup>1</sup>

<sup>1</sup>Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan; <sup>2</sup>Department of Bioregulation, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan; <sup>3</sup>Research Institute of Tuberculosis, Tokyo, Japan; <sup>4</sup>Department of Host Defense, Graduate School of Medicine, Osaka City University, Osaka, Japan; <sup>5</sup>Department of Internal Medicine, Toneyama National Hospital, Osaka, Japan; and <sup>6</sup>Japan BCG Central Laboratory, Kiyose, Tokyo, Japan

**Correspondence:** Yumi Maeda, Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases (NIID), 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan. Tel.: +81 42 391 8211; fax: +81 42 391 8212; e-mail: yumi@nih.go.jp

Received 24 January 2007; revised 3 April 2007; accepted 10 April 2007.  
First published online 22 May 2007.

DOI: 10.1111/j.1574-6968.2007.00754.x

Editor: Roger Buxton

## Keywords

serology; diagnosis; infection; mycobacteria; leprosy.

## Introduction

Leprosy represents a broad-spectrum disease caused by *Mycobacterium leprae*, with lepromatous leprosy at one pole and tuberculoid leprosy at the other end of the pole, depending on the clinical manifestation, which is an ultimate effect of the immunity of the host (Ridley & Jopling, 1996). In all forms of the disease, *M. leprae* induces skin lesions and a chronic progressive peripheral nerve injury, to a lesser or greater extent, which leads to systemic deformity (Stoner, 1979; Job, 1989). Therefore, early detection of *M. leprae* infection is the key to avoiding deformities. The diagnosis of leprosy is based on microscopic detection of acid-fast bacteria (AFB) in skin smears or biopsies, along with clinical and histopathological evaluation. Acid-fast staining requires at least a thousand organisms per gram of tissue for reliable detection (Shepard & McRae, 1968), resulting in an extremely low sensitivity, especially for the tuberculoid form of the disease, where AFB are rare or

## Abstract

As serodiagnosis is the easiest way of diagnosing a disease, the utility of *Mycobacterium leprae*-derived major membrane protein-II (MMP-II), one of the immuno-dominant antigens, in the serodiagnosis of leprosy was examined. The percent positivity by an enzyme-linked immunosorbent assay for anti-MMP-II antibody was 82.4% for multi-bacillary leprosy, and the specificity of the test was 90.1%. For pauci-bacillary leprosy where cell-mediated immunity predominates, 39.0% showed positive results. These percentage values were significantly higher than these values obtained for existing phenolic glycolipid-I based methods, suggesting that MMP-II antibody detection would facilitate the diagnosis of leprosy.

absent. Recently, real-time PCR-based methods have been developed (Martinez *et al.*, 2006), but the sensitivity of the test for clinical specimens is still problematic. In developing countries, where leprosy is endemic, diagnosis still relies on clinical observations and easy inexpensive tests. Serodiagnosis is the most easy and tangible way of diagnosing a disease. For leprosy, the only antigen currently used is phenolic glycolipid-I (PGL-I), which is supposed to be *M. leprae* specific. Since the discovery of PGL-I in 1981 by Hunter and Brennan, considerable progress has been made in the development of serological tools (Hunter & Brennan, 1981). In this process, simple user-friendly assays such as Serodia leprae<sup>®</sup>, a simple lateral flow test, and dipstick assays, based on PGL-I antigen, have been developed to detect leprosy patients in leprosy-endemic areas (Izumi *et al.*, 1990; Buhrer-Sekula *et al.*, 2003). However, these tests seem to be insufficient to detect both multi-bacillary and pauci-bacillary patients, as well as for early diagnosis, and have not been used as widely as would be expected, in the field situations. Therefore, there is a need to look for more sensitive

antigens. To date, various antigens of *M. leprae* have been studied (Hunter *et al.*, 1990), but due to a lack of either specificity or sensitivity, their use has been limited. Major membrane protein-II (MMP-II, #ML2038c, gene name *bfrA*, also known as bacterioferritin) had been identified previously from the cell membrane fraction of *M. leprae* as an antigenic molecule capable of activating both antigen-presenting cells and T cells (Pessolani *et al.*, 1994; Maeda *et al.*, 2005). These findings prompted examination of the role of MMP-II in the humoral responses of patients. Here, MMP-II was expressed and purified in *Escherichia coli* and the use of MMP-II as an antigen for the serodiagnosis of leprosy was evaluated.

## Materials and methods

### Study population

Sera were obtained with informed consent from healthy individuals, leprosy patients, and tuberculosis patients from Japan. The samples were frozen at  $-30\text{ }^{\circ}\text{C}$  before use. The population studied included multi-bacillary ( $n=74$ ) and pauci-bacillary ( $n=77$ ) leprosy patients, either treated or untreated, from the National Sanatorium Oshimaseishoen, and tuberculosis patients ( $n=55$ ) from Fukujiji Hospital. Classification of leprosy was performed according to WHO recommendations. The home page is available at <http://www.who.int/lep/classification/en/index.html>. Individuals who have not been vaccinated with *Mycobacterium bovis* bacillus Calmette–Guerin (BCG) are unavailable in Japan, due to compulsory vaccination at least once in childhood. Therefore, sera from BCG-vaccinated healthy volunteers ( $n=81$ ) residing in Japan were used as negative controls in the enzyme-linked immunosorbent assay (ELISA) to determine the cut-off value for positivity. The age and sex of normal individuals may not be fully matched with those of patients because some of the details of leprosy patients as well as normal individuals are unknown. This study was approved by the ethics committee of the National Institute of Infectious Diseases, Tokyo.

### Expression and purification of protein

The MMP-II gene (ML2038c, *bfrA*) was expressed in *E. coli* as a fusion construct using a pMAL-c2X expression vector (New England BioLabs). The protein was affinity purified to almost homogeneity using an amylose column (data not shown), and used as the antigen for the detection of anti-MMP-II antibody levels in the leprosy patients. The synthetic bovine serum albumin (BSA)-conjugated trisaccharide-phenyl propionate (NTP-BSA) for the detection of PGL-I antibodies was kindly provided by Dr T. Fujiwara, Institute for Natural Sciences, Nara University. The procedure for synthesis of the antigen is described elsewhere (Fujiwara *et al.*, 1984).

### Assay method for the detection of antibodies

The ELISA for the detection of anti-MMP-II immunoglobulin G (IgG) antibodies or anti-PGL-I IgM antibodies was performed as described previously, with modifications (Izumi *et al.*, 1990). Ninety-six well plates (Immunosorb, Nunc) were coated overnight, with MMP-II at a concentration of  $2\text{ }\mu\text{g mL}^{-1}$  in 0.1 M carbonate buffer (pH 9.5). After blocking with 2% skim milk, the plates were washed with phosphate-buffered saline containing 0.1% Tween 20 (PBST), and human sera (normal or patient's sera) diluted 100-fold were added and incubated at  $37\text{ }^{\circ}\text{C}$  for 2 h. After washing with PBST, biotinylated anti-human IgG (Vector Laboratories) was added at a concentration of  $0.5\text{ }\mu\text{g mL}^{-1}$  and incubated for 1 h. The plates were incubated with reagents from a Vectastain ABC Kit (Vector Laboratories) for 30 min. After further washing with PBST, a substrate solution consisting of  $0.2\text{ mg mL}^{-1}$  of 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) and 0.02%  $\text{H}_2\text{O}_2$  in 0.1 M citrate buffer was added until a blue color developed and the OD was measured at 405 nm using a spectrophotometer. Plate-to-plate variations in OD readings were controlled using a common standard serum with an OD value of 0.350. For detecting anti-PGL-I antibody, NTP-BSA was coated at a concentration of  $0.5\text{ }\mu\text{g mL}^{-1}$ , and the same procedure used to detect anti-MMP-II antibody was followed, except that the secondary antibody used was biotinylated anti-human IgM at a concentration of  $1\text{ }\mu\text{g mL}^{-1}$ . The volume of all solutions used in the 96-well plate was  $100\text{ }\mu\text{L well}^{-1}$ .

### Statistical analyses

The data were analyzed using MEDCALC software (MedCalc, Belgium). A receiver operator characteristics (ROC) curve was drawn to calculate the cut-off levels. The McNemar test was applied (MMP-II vs. PGL-I test) to determine the *P* value. The *P* value of  $<0.05$  was considered to be statistically significant. The  $\kappa$  value was calculated to determine the agreement between the two tests.

## Results and discussion

MMP-II (*bfrA*) has been previously identified as one of the components of *M. leprae* capable of stimulating  $\text{CD4}^+$  and  $\text{CD8}^+$  T cells (Maeda *et al.*, 2005). In multi-bacillary leprosy, *M. leprae* is widely disseminated and abundant antibody production is observed due to polyclonal B cell activation, while in pauci-bacillary leprosy, the bacilli are usually localized in skin lesions and type 1 T cells are predominantly activated, and so the level of antibodies to *M. leprae* antigens is usually low. However, preliminary experiments showed that the pooled pauci-bacillary leprosy sera reacted to MMP-II on polyvinylidene difluoride (PVDF) membranes.

**Table 1.** Positivity rates of MMP-II and PGL-I tests in various groups of subjects by an enzyme-linked immunosorbent assay

|                         | MMP-II |          |      |           | PGL-I  |          |      |           | McNemar test<br>(MMP-II vs. PGL) | Rate difference<br>(95% CI) | Inter-rater<br>agreement<br>( $\kappa$ ) |
|-------------------------|--------|----------|------|-----------|--------|----------|------|-----------|----------------------------------|-----------------------------|--|
|                         | Tested | Positive | %    | 95% CI    | Tested | Positive | %    | 95% CI    |                                  |                             |  |
| Multi-bacillary leprosy | 74     | 61       | 82.4 | 71.8–90.3 | 74     | 51       | 68.9 | 57.1–79.2 | $P=0.063$                        | 13.5% (–0.7 to 24.2)        | 0.14                                     |
| Pauci-bacillary leprosy | 77     | 30       | 39   | 28.8–50.1 | 77     | 15       | 19.5 | 12.2–29.7 | $P=0.007$                        | 19.5% (5.4 to 28.9)         | 0.189                                    |
| Tuberculosis            | 55     | 9        | 16.4 | 8.9–28.3  | ND     | ND       | ND   | ND        | –                                | –                           | –  |
| Healthy subjects        | 81     | 8        | 9.9  | 5.1–18.3  | 81     | 8        | 9.9  | 5.1–18.3  | –                                | –                           | –  |

ND, not detected.

Therefore, a systematic study was conducted, measuring the anti-MMP-II antibody IgG levels in the leprosy patients' sera. The study population consisted of multi-bacillary leprosy, pauci-bacillary leprosy, tuberculosis patients, and normal healthy BCG-vaccinated volunteers from Japan. Some of the leprosy patients were already under treatment, so that all patients were not active leprosy patients. The cut-off value of OD 0.130 was defined by an ROC curve analysis (MEDCALC software) using the OD titers from 81 normal individuals and 74 multi-bacillary leprosy patients. Using this cut-off value, it was observed that 61 (82.4%, 95% CI; 71.8–90.3) out of 74 multi-bacillary patients had positive results, and 30 (39.0%, 95% CI; 28.8–50.1) out of 77 pauci-bacillary patients had positive titers (Table 1). The only serological test for leprosy that is currently available is the detection of antibodies to PGL-I of *M. leprae*. The haptenic trisaccharide of PGL-I is known to be *M. leprae* specific, and this trisaccharide unit could be chemically synthesized (Fujiwara *et al.*, 1987). Several reports show the performance of PGL-I for serodiagnosis (Agis *et al.*, 1988; Cho *et al.*, 1991). When the anti-PGL-I IgM antibody levels were examined in the same Japanese leprosy patients by ELISA, it was found that only 68.9% (95% CI; 57.1–79.2) of multi-bacillary patients ( $n=74$ ) and 19.5% (95% CI; 12.2–29.7) of pauci-bacillary patients ( $n=77$ ) showed positive values (Table 1). These percentages were far lower than expected, which may be due to the influence of chemotherapy. However, the percent positivity for anti-MMP-II antibodies was significantly higher than that for anti-PGL-I antibodies of the same sera ( $P=0.0008$ , McNemar test,  $n=152$ ) when both multi-bacillary and pauci-bacillary leprosy were considered together. When multi-bacillary sera were taken separately, the significance of the MMP-II test was statistically marginal compared with that of the PGL-I test ( $P=0.06$ ), but there was a significant difference ( $P=0.007$ ) between the two tests for pauci-bacillary leprosy (Table 1). The agreement between these two tests was low ( $\kappa$  value for multi-bacillary leprosy: 0.140,  $\kappa$  value for pauci-bacillary leprosy: 0.189), so that when positive titers for either or both PGL-I and MMP-II were considered, the percent positivity increased to 91.9% in multi-bacillary patients and 48.7% in pauci-bacillary patients. The specificity of both the MMP-II and PGL-I tests was 90.1%.

When the anti-MMP-II IgG levels were measured in the normal individuals (BCG-vaccinated), a low level of seropositivity was found, 9.9% (95% CI; 5.1–18.3) ( $n=81$ ). As the amino acid homology between MMP-II protein from *M. leprae* and its homologue in *M. bovis* BCG is 90.6%, it was expected that a higher percentage of normal individuals would be positive. But this result was to the contrary and advantageous to the assay system. Also, tuberculosis patients showed a low positive result of 16.4% (95% CI; 9.9–28.3) ( $n=55$ ). As, the genes encoding MMP-II were conserved between *M. leprae* and *M. tuberculosis*, the low positivity in tuberculosis was unexpected. The exact reason for the low positivity in tuberculosis patients is not clear, but might be due to slight conformational differences in the antibody recognition site on MMP-II between *M. leprae* and *M. tuberculosis*. More intensive studies need to be carried out using active tuberculosis patients and native MMP-II protein derived from *M. tuberculosis* to fully realize the significance of MMP-II homologue in the detection of tuberculosis.

There is only one report showing that the sera from leprosy patients had a higher IgG titer to MMP-II, regardless of the clinical type of leprosy (Deshpande *et al.*, 1995). However, their study was carried out with a limited number of individuals ( $n=10$  in each group), and so, here, a more systematic study has been conducted for evaluating the expediency of measuring anti-MMP-II antibody levels. In addition, when monitoring of patients on multidrug treatment was conducted ( $n=4$ ) for a period of 2 years, there was a definite decline in the MMP-II antibody levels after chemotherapy (data not shown). Further intensive studies have to be carried out with a larger cohort of patients to fully realize the significance of MMP-II in the monitoring of chemotherapy.

This report showed that MMP-II could contribute to the sero-detection of multi-bacillary as well as pauci-bacillary leprosy patients. Further study will be pursued to evaluate its efficacy for serodiagnosis in developing countries and for the development of user-friendly tools.

### Acknowledgement

This work was supported by grants from Health Sciences Research Grants-Research on Emerging and Re-emerging

Infectious Diseases, from the Ministry of Health, Labour and Welfare, Tokyo, Japan.

## References

- Agis F, Schlich P, Cartel JL, Guidi C & Bach MA (1988) Use of anti-*M. leprae* phenolic glycolipid-I antibody detection for early diagnosis and prognosis of leprosy. *Int J Lepr Other Mycobact Dis* **56**: 527–535.
- Buhrer-Sekula S, Smits HL, Gussenhoven GC, Leeuwen van J, Amador S, Fujiwara T, Klatser PR & Oskam L (2003) Simple and fast lateral flow test for classification of leprosy patients and identification of contacts with high risk of developing leprosy. *J Clin Microbiol* **41**: 1991–1995.
- Cho SN, Cellona RV, Fajardo TT Jr, Abalos RM, dela Cruz EC, Walsh GP, Kim JD & Brennan PJ (1991) Detection of phenolic glycolipid-I antigen and antibody in sera from new and relapsed lepromatous patients treated with various drug regimens. *Int J Lepr Other Mycobact Dis* **59**: 25–31.
- Deshpande RG, Khan MB, Bhat DA & Navalkar RG (1995) Immunoaffinity chromatographic isolation of a high molecular weight seroreactive protein from *Mycobacterium leprae* cell sonicate. *FEMS Immunol Med Microbiol* **11**: 163–169.
- Fujiwara T, Hunter SW, Cho SN, Aspinall GO & Brennan PJ (1984) Chemical synthesis and serology of disaccharides and trisaccharides of phenolic glycolipid antigens from the leprosy bacillus and preparation of a disaccharide protein conjugate for serodiagnosis of leprosy. *Infect Immun* **43**: 245–252.
- Fujiwara T, Aspinall GO, Hunter SW & Brennan PJ (1987) Chemical synthesis of the trisaccharide unit of the species-specific phenolic glycolipid from *Mycobacterium leprae*. *Carbohydr Res* **163**: 41–52.
- Hunter SW & Brennan PJ (1981) A novel phenolic glycolipid from *Mycobacterium leprae* possibly involved in immunogenicity and pathogenicity. *J Bacteriol* **147**: 728–735.
- Hunter SW, Rivoire B, Mehra V, Bloom BR & Brennan PJ (1990) The major native proteins of the leprosy bacillus. *J Biol Chem* **265**: 14065–14068.
- Izumi S, Fujiwara T, Ikeda M, Nishimura Y, Sugiyama K & Kawatsu K (1990) Novel gelatin particle agglutination test for serodiagnosis of leprosy in the field. *J Clin Microbiol* **28**: 525–529.
- Job CK (1989) Nerve damage in leprosy. *Int J Lepr Other Mycobact Dis* **57**: 532–539.
- Maeda Y, Mukai T, Spencer J & Makino M (2005) Identification of an immunomodulating agent from *Mycobacterium leprae*. *Infect Immun* **73**: 2744–2750.
- Martinez AN, Britto CFPC, Nery JAC, Sampaio EP, Jardim MR, Sarno EN & Moraes MO (2006) Evaluation of real-time and conventional PCR targeting complex 85 genes for detection of *Mycobacterium leprae* DNA in skin biopsy samples from patients diagnosed with leprosy. *J Clin Microbiol* **44**: 3154–3159.
- Pessolani MC, Smith DR, Rivoire B, McCormick J, Hefta SA, Cole ST & Brennan PJ (1994) Purification, characterization, gene sequence, and significance of a bacterioferritin from *Mycobacterium leprae*. *J Exp Med* **180**: 319–327.
- Ridley DS & Jopling WH (1996) Classification of leprosy according to immunity. A five-group system. *Int J Lepr Other Mycobact Dis* **34**: 255–273.
- Shepard CC & McRae DH (1968) A method for counting acid-fast bacteria. *Int J Lepr Other Mycobact Dis* **36**: 78–82.
- Stoner GL (1979) Importance of the neural predilection of *Mycobacterium leprae* in leprosy. *Lancet* **2**: 994–996.

# Identification of trehalose dimycolate (cord factor) in *Mycobacterium leprae*

Masanori Kai<sup>a,\*</sup>, Yukiko Fujita<sup>b</sup>, Yumi Maeda<sup>a</sup>, Noboru Nakata<sup>a</sup>, Shinzo Izumi<sup>c</sup>, Ikuya Yano<sup>b</sup>,  
Masahiko Makino<sup>a</sup>

<sup>a</sup> Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, Higashimurayama-shi, Tokyo 189-0002, Japan

<sup>b</sup> BCG Laboratory, Kiyose, Tokyo 204-0022, Japan

<sup>c</sup> Leprosy Study Group, Tropical Disease Center, Airlangga University Kampus C UNAIR, Jl. Mulyorejo 60115, Surabaya, Indonesia

Received 15 May 2007; revised 11 June 2007; accepted 13 June 2007

Available online 21 June 2007

Edited by Sandro Sonnino

**Abstract** Glycolipids of *Mycobacterium leprae* obtained from armadillo tissue nodules infected with the bacteria were analyzed. Mass spectrometric analysis of the glycolipids indicated the presence of trehalose 6,6'-dimycolate (TDM) together with trehalose 6-monomycolate (TMM) and phenolic glycolipid-I (PGL-I). The analysis showed that *M. leprae*-derived TDM and TMM possessed both  $\alpha$ - and keto-mycolates centering at C78 in the former and at C81 or 83 in the latter subclasses, respectively. For the first time, MALDI-TOF mass analyses showed the presence of TDM in *M. leprae*.

© 2007 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

**Keywords:** Trehalose 6,6'-dimycolate; Trehalose 6-monomycolate; Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; Thin-layer chromatography; *Mycobacterium leprae*

## 1. Introduction

Mycolic acids and mycolyl glycolipids are unique and ubiquitous components of mycobacterial cell envelopes. Among such components, trehalose 6,6'-dimycolate (TDM) was first isolated as cord factor from highly virulent *Mycobacterium tuberculosis* showing cord-like growth on the surface culture in liquid media [1–3]. TDM of *M. tuberculosis* was recognized as one of the virulence factors capable of inhibiting fusion of phagosome with lysosome in infected macrophage [4]. However, on the other hand, the TDM was considered to be associated with host defence against the mycobacterial infection since it induced immune responses such as type 1 T cell activation and the formation of granuloma in the mycobacteria-infected lesion [5–7]. Wang et al. have reported that a high proportion of patients infected with *Mycobacterium leprae* possess IgG antibody against TDM of unknown origin as well as trehalose 6-monomycolate (TMM), a biosynthetic precursor of TDM [8]. These observa-

tions suggest the possible existence of both TDM and TMM in *M. leprae*. Previously, TDM has been isolated from almost all species of culturable mycobacteria [9,10], and also TMM was isolated from *M. leprae*, however, the search for TDM in *M. leprae* has been unsuccessful [11]. The possible reason for not being able to identify TDM, may be due to (1) inadequate supply of *M. leprae*, (2) negligible amount of the product, (3) technically inefficient to identify TDM. Recent development of newer techniques such as MALDI-TOF mass spectrometry has enabled us to identify even several pg amounts of products. Therefore, the present study was designed as an attempt to directly detect TDM in *M. leprae*, by use of newer technologies. In the process, higher amounts of *M. leprae* phenolic glycolipid-I (PGL-I) was obtained which was analyzed by MALDI-TOF mass spectrometry.

## 2. Materials and methods

### 2.1. Sources for extraction of glycolipids

*M. tuberculosis* Aoyama B (ATCC 31726) and *Mycobacterium bovis* BCG Connaught (ATCC 35745) were grown at 37 °C on Sauton's medium for four weeks as surface pellicles until early stationary phase. Cultivated mycobacterial strains were used for extraction of glycolipids. Because *M. leprae* cannot be cultivated in any artificial media, armadillo tissue nodules infected with *M. leprae* (Thai 53 strain) were used for the extraction of glycolipids.

### 2.2. Extraction of glycolipids and mycolic acid methyl esters

Glycolipids were extracted according to the methods described previously [12]. In brief, bacterial culture or tissues infected with *M. leprae* [13] were autoclaved at 121 °C for 15 min and collected by centrifugation. Lipids were extracted from homogenized tissue with 20 volumes of chloroform/methanol (2:1, v/v) three times with vigorous grinding. The two phases were separated in a funnel, the lower organic phase was collected, and the solvent was evaporated from the organic phase. The total lipids were separated by solvent fractionation and tetrahydrofuran-soluble fraction was further separated by thin-layer chromatography (TLC) on silica gel plates (Uniplate; Analtech Inc. Newark, DE) with the solvent system of chloroform/methanol/water (90:10:1, by vol.) or chloroform/methanol/acetone/acetic acid (90:10:6:1, by vol.). Glycolipid spots were visualized with a 9 M H<sub>2</sub>SO<sub>4</sub> spray followed by charring at 200 °C for 10 min or with iodine vapor for preparative purposes.

To determine the subclass composition of the mycolic acids in each mycobacterial TMM and TDM, mycolic acid methyl esters were prepared by alkaline hydrolysis of glycolipids. The glycolipids were hydrolyzed with 1.25 M NaOH in 90% methanol at 70 °C for 1 h and the resultant mycolic acids were then extracted with *n*-hexane after acidification with HCl, followed by methylation with benzene/methanol/H<sub>2</sub>SO<sub>4</sub> (10:20:1, by vol.) [14]. Mycolic acid methyl esters from each

\*Corresponding author. Fax: +81 42 391 8807.

E-mail address: mkai@nih.go.jp (M. Kai).

**Abbreviations:** TDM, trehalose 6,6'-dimycolate; TMM, trehalose 6-monomycolate; *M. leprae*, *Mycobacterium leprae*; *M. tuberculosis*, *Mycobacterium tuberculosis*; *M. bovis*, *Mycobacterium bovis*; TLC, thin-layer chromatography; MALDI-TOF mass, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry

glycolipid were fully separated into subclasses by TLC with the solvent system of benzene in a draft chamber under reduced pressure.

### 2.3. Mass spectrometry analysis

Analysis by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF mass) was carried out on a Voyager DE-STR (Applied Biosystems, Tokyo, Japan) with pulsed UV light (337 nm) from an N<sub>2</sub> laser, essentially according to the method reported previously [14]. TDM and TMM were analyzed in the reflectron mode by the instrument operated at 20 kV in the positive ion mode. The 2,5-dihydroxybenzoic acid (2,5-DHB) matrix was used at a concentration of 10 mg/ml in chloroform/methanol (2:1 v/v). Typically, 5 µl of TDM or TMM samples (5 µg) in chloroform/methanol (2:1 v/v) solution and 5 µl of the matrix solution were mixed, and 1.5 µl of the mixture was applied on a sample plate. An external mass spectrum calibration was performed using calibration mixture 2 of the Sequazyme Peptide Mass Standards kit (Applied Biosystems), including known peptide standards in a mass range from 1290 to 5700 Da. The molecular mass of mycobacterial TMM, TDM, and PGL-I was determined based on the quasimolecular mass ions [M+Na]<sup>+</sup> by the reflectron mode. In general, nominal number of atomic mass is used for calculation of number of molecular mass. However, there is a slight difference between nominal mass number and accurate mass number read from spectrometry [15]. For instance, when both of the two molecules of mycocerosates (R<sub>1</sub> and R<sub>2</sub>) in PGL-I are C32, the nominal mass number [M+Na]<sup>+</sup> of PGL-I (C<sub>124</sub>H<sub>232</sub>O<sub>19</sub>) is [(C × 124) + (H × 232) + (O × 19) + (Na × 1)] = 2047 (2024 + 23), but the accurate number is 2050.15 [(12.0107 × 124) + (1.00794 × 232) + (15.9994 × 19) + 22.9898]. The nominal mass numbers are given in the text.

## 3. Results

### 3.1. TLC analysis of mycolic acids methyl ester

To determine the subclass composition of the mycolic acids in each mycobacterial TMM and/or TDM, mycolic acid methyl esters were analyzed by TLC. The TLC analysis indicated that fatty acid methyl esters had two spots corresponding to either α- and keto-mycolic acid (Fig. 1A). The same two spots pattern was observed for *M. bovis* BCG Connaught (BCG-C) and *M. leprae* while three spots were detected for *M. tuberculosis* corresponding to α-, methoxy-, and keto-mycolic acid [16,17]. The bottom spot in ML lane could be cholesterol from armadillo's tissues (data not shown).

### 3.2. MALDI-TOF-MS analysis of TMM

We separated the final solvent extracts of *M. leprae* into four fractions (M1–M4). Fig. 1B shows a thin-layer chromatogram of the TMM and TDM from *M. tuberculosis* and the solvent fractionated glycolipids from *M. leprae*. *M. leprae* exhibited bands, which were faint, but significantly reddish glycolipid-like, migrating close to bands of TDM or TMM of *M. tuberculosis*. The major bands of M2 and M4 migrated close to TDM and TMM positions of *M. tuberculosis*, respectively. Therefore, we tried to carefully analyze the bands which may correspond to TDM and TMM by mass spectrometry.

Major band in M4 in Fig. 1B was analyzed using BCG-C as a reference. The mass spectra of TMM from BCG-C showed a biphasic distribution of pseudomolecular ions, [M+Na]<sup>+</sup> (Fig. 2A). In the higher mass ranges of BCG-C TMM, dominant ions were detected at *m/z* 1555, 1583, 1597, 1611 and 1625 due to [M+Na]<sup>+</sup> of C82, C84, C85, C86 and C87 keto mycolyl TMM, and the major mass ions in the lower mass ranges were detected at *m/z* 1455, 1483, 1511 and 1539 due to [M+Na]<sup>+</sup> of α-mycolyl TMM centering at C78, respectively (Fig. 2A) [18]. On the other hand, *M. leprae* derived TMM showed in lower mass ranges at *m/z* 1427, 1455, 1483, 1511




Fig. 1. Thin-layer chromatograms (TLC) of solvent extracts from tetrahydrofuran soluble fraction and mycolic acid methyl ester subclasses of *M. tuberculosis*, *M. bovis* BCG Connaught, and *M. leprae*. (A) TLC of mycolic acid methyl ester subclasses with the solvent system of benzene/methanol/H<sub>2</sub>SO<sub>4</sub> (10:20:1, by vol.). Mycolic acid methyl ester subclasses of *M. tuberculosis* (TB); α-, methoxy- and keto-mycolic acid methyl esters are shown with the mycolic acid methyl ester mixture of glycolipids derived from armadillo tissue nodules infected with *M. leprae* (ML), and those of *M. bovis* BCG Connaught (BCG-C). (B) TLC of solvent extracts from *M. leprae* (ML) and *M. tuberculosis* (TB) with the solvent system of chloroform/methanol/acetone/acetic acid (90:10:6:1, by vol.). Trehalose monomycolate (TMM) and trehalose di-mycolate (TDM) bands of TB were identified previously and used as references in this TLC. Fractions 1–4 separated from the final extracts of *M. leprae* and designated M1–4.

and 1539 due to [M+Na]<sup>+</sup> of α-mycolyl TMM centering at C78 same to that of BCG-C (Fig. 2B). In the higher mass ranges, dominant ions were shifted lower and the major ions were detected at *m/z* 1541, 1569 and 1597 (Fig. 2B), indicating the major keto-mycolyl TMM consisted of C81, C83 or C85 mycolate, respectively. The molecular species of TMM from *M. leprae* and that from BCG-C are summarized in Table 1. These results indicate that *M. leprae* possess trehalose 6-monomycolate, with C78 α- and C83 keto-mycolates, as the major molecular species.

### 3.3. MALDI-TOF-MS analysis of TDM

TDM from BCG-C showed a diverse distribution of mass ions according to the combination of di α-, α- and keto-, and di keto-mycolic acid subclasses and each molecular species, that leads to a multiphasic distribution of mass ions due to the dominant combination of α-α, α-keto and keto-keto dimycolyl TDM (Fig. 2C). Given the small sample size, the identification of *M. leprae* TDM was achieved primarily on the thin-layer chromatographic behavior and MALDI-TOF mass analysis, in comparison with the analytical results from BCG-C possessing the same mycolic acid subclasses. Fig. 2D shows the positive MALDI-TOF mass spectra of *M. leprae* TDM (Fig. 1B, M2). In contrast to TDM from BCG-C, TDM from *M. leprae* showed a distinctive mass ion distribution shifted to lower mass ranges due to the major combinations of α-α dimycolyl TDM and α-keto dimycolyl TDM with a small shoulder due to keto-keto dimycolyl TDM. In *M. leprae* TDM, major




Fig. 2. MALDI-TOF-MS spectra of TMM and TDM. (A) TMM of *M. bovis* BCG Connaught, (B) TMM of *M. leprae* Thai 53, (C) TDM of *M. bovis* BCG Connaught, (D) TDM of *M. leprae* Thai 53.



Table 1  
MALDI-TOF mass spectrometry data of the individual types of TMM from *M. leprae* (Thai 53) and *M. bovis* BCG (Connaught)

| Species (strain)                | Mycolic acid type <sup>a</sup> | Total carbon number of TMM mycolate |             |      |             |      |             |             |             |             |             |             |             |             |      |      |      |      |    |  |
|---------------------------------|--------------------------------|-------------------------------------|-------------|------|-------------|------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|------|------|------|------|----|--|
|                                 |                                | 74                                  | 75          | 76   | 77          | 78   | 79          | 80          | 81          | 82          | 83          | 84          | 85          | 86          | 87   | 88   | 89   | 90   | 91 |  |
| <i>M. leprae</i> (Thai 53)      | I                              | 1427                                | <b>1455</b> |      | <b>1483</b> | 1497 | <b>1511</b> | 1525        | 1539        | 1553        |             |             |             |             |      |      |      |      |    |  |
|                                 | II                             |                                     | 1471        |      | 1499        |      | 1527        | <b>1541</b> | 1555        | <b>1569</b> | 1583        | 1597        |             |             |      |      |      |      |    |  |
| <i>M. bovis</i> BCG (Connaught) | I                              |                                     | <b>1455</b> | 1469 | <b>1483</b> |      | <b>1511</b> |             | 1539        |             |             |             |             |             |      |      |      |      |    |  |
|                                 | II                             |                                     |             |      |             |      |             |             | <b>1555</b> | 1569        | <b>1583</b> | <b>1597</b> | <b>1611</b> | <b>1625</b> | 1639 | 1653 | 1667 | 1681 |    |  |

<sup>a</sup>I;  $\alpha$ -dicyclopropanoic or dienolic, II; keto-monocyclopropanoic or monoenoic. Major homologues are shown in bold.

mass ions due to molecular species possessing C76, C78, C80  $\alpha$ - $\alpha$  dimycolic acids were detected at  $m/z$  2573, 2601 and 2629, and those possessing  $\alpha$ -keto dimycolic acids at  $m/z$  2659, 2687 and 2715, and those possessing keto-keto dimycolic acids at  $m/z$  2731, 2745, 2773 and 2801, respectively. The deduced molecular species of TDM from *M. leprae* and that from BCG-C are summarized in Table 2.

### 3.4. MALDI-TOF-MS analysis of PGL-I

The MALDI-TOF mass spectra of M2 in Fig. 1B showed the existence of TDM as described above, however, the major cluster ions in mass spectra of M2 were observed in lower mass ranges from around  $m/z$  1990 to 2100. So, we analyzed the spectra in more detail. The result indicates that the cluster ions are derived from phenolic glycolipid-I (PGL-I) (Fig. 3A). The nominal mass number of the PGL-I, which is assumed to have two molecules of C32 mycocerosates, was  $m/z$  2047 (Fig. 3A and B). Thus, the deduced combination of mycocerosic acids with different carbon numbers in PGL-I are shown in Fig. 3A and the general structure of PGL-I is shown in Fig. 3B.

## 4. Discussion

In the present study, we have directly detected TDM and TMM from armadillo tissues infected with *M. leprae*. The presence of TMM in *M. leprae* possessing C74-82  $\alpha$ -mycolic acids has been reported previously [8,18], however, the existence of keto-mycolate in *M. leprae* TMM was not clear. We identified keto-mycolate clearly in *M. leprae* TDM and TMM, and the chain length of keto-mycolate in *M. leprae* was found to be shorter than those from slow-growing culturable mycobacteria such as *M. tuberculosis* and *M. bovis* BCG [19].

Previously, no detectable TDM was identified by the analysis of the lipids obtained from *M. leprae* infected armadillo [9,20]. However, in our hands, we observed a meager but significant spot on TLC from *M. leprae* extract which migrated to the position of TDM of *M. tuberculosis* (Aoyama B strain). When this spot was analyzed by mass spectrometry in reference to TDM from *M. bovis* BCG Connaught (BCG-C), TDM having  $\alpha$ - $\alpha$  dimycolates were observed in the lower mass ranges than  $m/z$  2673 as seen in TDM from BCG-C (Fig. 2C). Therefore, both *M. leprae* and *M. bovis*

Table 2  
Most probable combination of mycolic acids constructing TDM from *M. leprae* (Thai 53) and *M. bovis* BCG (Connaught)

| Mass no. of TDM ( $m/z$ ) | <i>M. leprae</i> (Thai 53)  | <i>M. bovis</i> BCG (Connaught)                   |
|---------------------------|---|---|
| 2573                      | $\alpha$ 76: $\alpha$ 78  |   |
| 2601                      | $\alpha$ 76: $\alpha$ 80, $\alpha$ 78: $\alpha$ 78                                  |   |
| 2617                      | $\alpha$ 74:k82, $\alpha$ 76:k80, $\alpha$ 78:k78, $\alpha$ 80:k76                  |   |
| 2629                      | $\alpha$ 78: $\alpha$ 80  |   |
| 2645                      | $\alpha$ 74:k84, $\alpha$ 76:k82, $\alpha$ 78:k80, $\alpha$ 80:k78, $\alpha$ 82:k76 |   |
| 2659                      | $\alpha$ 76:k83, $\alpha$ 78:k81  |   |
| 2673                      | $\alpha$ 76:k84, $\alpha$ 78:k82, $\alpha$ 79:k81, $\alpha$ 80:k80, $\alpha$ 82:k78 | $\alpha$ 76:k84, $\alpha$ 78:k82                  |
| 2687                      | $\alpha$ 78:k83, $\alpha$ 80:k81  |   |
| 2701                      | $\alpha$ 78:k84, $\alpha$ 79:k83, $\alpha$ 80:k82, $\alpha$ 81:k81, $\alpha$ 82:k80 | $\alpha$ 76:k86, $\alpha$ 78:k84, $\alpha$ 80:k82 |
| 2715                      | $\alpha$ 80:k83   | $\alpha$ 76:k87, $\alpha$ 78:k85                  |
| 2729                      |   | $\alpha$ 78:k86, $\alpha$ 80:k84                  |
| 2731                      | k80:k83, k81:k82  |   |
| 2743                      |   | $\alpha$ 78:k87, $\alpha$ 80:k85                  |
| 2745                      | k81:k83   | k82:k82   |
| 2757                      |   | $\alpha$ 80:k86                                   |
| 2773                      | k83:k83   | k82:k84   |
| 2787                      |   | k82:k85   |
| 2801                      | k83:k85, k84:k84  | k82:k86, k84:k84                                  |
| 2815                      |   | k82:k87, k84:k85                                  |
| 2829                      |   | k84:k86, k85:k85                                  |
| 2843                      |   | k84:k87, k85:k86                                  |
| 2857                      |   | k85:k87, k86:k86                                  |
| 2871                      |   | k86:k87   |
| 2885                      |   | k87:k87   |

$\alpha$ ,  $\alpha$ -dicyclopropanoic or dienolic; k, keto-monocyclopropanoic or monoenoic mycolic acid. Molecular ions of TDM with intensities  $\geq 30\%$  of the highest intensity observed are listed.




Fig. 3. MALDI-TOF mass spectra of *M. leprae* PGL-I and the deduced chemical structure. (A) The mass spectra of the major cluster ions ranging from around  $m/z$  1997 (C30:C30) to 2103 (C34:C34) due to PGL-I separated by solvent fractionation, followed by TLC. C30–C34 shows carbon numbers of two sets of mycocerosate in PGL-I. (B) The chemical structure of *M. leprae* PGL-I. R indicates mycocerosate.

BCG-C can synthesize  $\alpha$ -mycolic acids with similar sizes (C74 to C82 centering at C78), however, the subclass composition of keto-mycolate differed. *M. bovis* BCG-C synthesizes multi sub-classes of ketomycolic acids such as *cis*-monoenoic, *cis*-monocyclopropanoic, *cis*-monoenoic *cis*-monocyclopropanoic, *trans*-monoenoic and *trans*-monocyclopropanoic mycolates [21], while *M. leprae* synthesizes only *cis*-monoenoic ketomycolate centering at C81 or C83 (Tables 1 and 2) [22]. Recently, the immunological role of mycolic acid has become a point of focus. The *trans*-, but not *cis*-cyclopropanation of mycolic acids in TDM induce massive inflammation in the lesion infected with *M. tuberculosis* and, in this respects, the *trans*-cyclopropane structure may be contributing to virulence [23,24]. Furthermore, recently, it was reported that the  $\Delta kasB$  (a  $\beta$ -keto acyl-ACP synthase gene) mutant strain of *M. tuberculosis* synthesized shorter chain mycolate than the wild ones, and resulted in the loss of keto-mycolic acid *trans*-cyclopropanation. The *kasB* deletion induced the ability of the mutant strain to persist in infected mice for up to 600 days without causing disease [25]. Thus, the lack of *trans*-cyclopropanoic keto-mycolic acid in *M. leprae* may

contribute to the low virulence in contrary to substantial quantities found in most of the slow growing pathogenic mycobacteria.

In the search for TDM, we also identified a major cluster ions in M2 fraction. Detailed analyses showed that the ions were derived from phenolic glycolipid (PGL-I), which is a major species specific glycolipid of *M. leprae*. Previously, combined GLC-mass spectroscopic analysis of the trisaccharide of PGL-I had been achieved, but TOF-mass spectrometric data have not been reported [26]. We analyzed the M1 fraction by MALDI TOF mass spectrometry and detected some clusters. Mass spectra of major clusters look like monoglycosyl phthiocerol diester and diglycosyl phthiocerol diester (data not shown), which differ from PGL-I in lacking the terminal sugars [27]. The co-existence of both PGL-I and TDM in armadillo tissue nodules infected with *M. leprae* indicates that these lipids were originated actually from *M. leprae*. Thus, this paper identifies TDM structure of *M. leprae* for the first time and shows that *M. leprae* TDM contains only  $\alpha$ -mycolic acids and shorter chain *cis*-monoenoic keto-mycolic acids, exclusively.

**Acknowledgments:** We thank Dr. M. Matsuoka, and Dr. M. Gidoh, for providing armadillo tissues infected with *M. leprae*. This work was supported in part by a Health Science Research Grant for Research on Emerging and Re-emerging Infectious Diseases, Ministry of Health, Labour and Welfare, Japan.

## References

- [1] Minnikin, D. E. (1982) Presented at the Part I. Physiology of the mycobacteria. In *The Biology of the Mycobacteria*.
- [2] Rastogi, N., Legrand, E. and Sola, C. (2001) The mycobacteria: an introduction to nomenclature and pathogenesis. *Rev. Sci. Tech.* 20, 21–54.
- [3] Ryll, R., Kumazawa, Y. and Yano, I. (2001) Immunological properties of trehalose dimycolate (cord factor) and other mycolic acid-containing glycolipids – a review. *Microbiol. Immunol.* 45, 801–811.
- [4] Fujiwara, N. (1997) Distribution of antigenic glycolipids among *Mycobacterium tuberculosis* strains and their contribution to virulence. *Kekkaku* 72, 193–205.
- [5] Coelho, E.A., Tavares, C.A., de Melo Lima, K., Silva, C.L., Rodrigues Jr., J.M. and Fernandes, A.P. (2006) *Mycobacterium* hsp65 DNA entrapped into TDM-loaded PLGA microspheres induces protection in mice against *Leishmania* (*Leishmania*) *major* infection. *Parasitol. Res.* 98, 568–575.
- [6] Oiso, R., Fujiwara, N., Yamagami, H., Maeda, S., Matsumoto, S., Nakamura, S., Oshitani, N., Matsumoto, T., Arakawa, T. and Kobayashi, K. (2005) Mycobacterial trehalose 6,6'-dimycolate preferentially induces type 1 helper T cell responses through signal transducer and activator of transcription 4 protein. *Microb. Pathog.* 39, 35–43.
- [7] Hunter, R.L., Olsen, M.R., Jagannath, C. and Actor, J.K. (2006) Multiple roles of cord factor in the pathogenesis of primary, secondary, and cavitary tuberculosis, including a revised description of the pathology of secondary disease. *Ann. Clin. Lab. Sci.* 36, 371–386.
- [8] Dhariwal, K.R., Yang, Y.-M., Fales, H.M. and Goren, M.B. (1987) Detection of trehalose monomycolate in *Mycobacterium leprae* grown in armadillo tissues. *J. Gen. Microbiol.* 133, 201–209.
- [9] Prome, J.C., Lacave, C.A., Ahibo-Coffy and Savagnac, A. (1976) Separation of structural studies of the molecular species of monomycolates and dimycolates of alpha-D-trehalose present in *Mycobacterium phlei*. *Eur. J. Biochem.* 63, 543–552.
- [10] Wang, L., Izumi, S., He, H., Fujiwara, N., Saita, N., Yano, I., Kobayashi, K. and Tatsumi, N. (1999) Serodiagnosis of Hansen's disease/leprosy by enzyme-linked immunosorbent assay using cord factor (trehalose 6,6'-dimycolate) as an antigen. *Nihon Hansenbyo Gakkai Zasshi* 68, 165–174.
- [11] Goren, M.B., Brokl, O. and Roller, P. (1979) Cord factor (trehalose-6,6'-dimycolate) of in vivo-derived *Mycobacterium lepraemurium*. *Biochim. Biophys. Acta* 574, 70–78.
- [12] Fujita, Y., Ogata, H. and Yano, I. (2005) Clinical evaluation of serodiagnosis of active tuberculosis by multiple-antigen ELISA using lipids from *Mycobacterium bovis* BCG Tokyo 172. *Clin. Chem. Lab. Med.* 43, 1253–1262.
- [13] Sasaki, N., Kawatsu, K., Tsutsumi, S., Gidoh, M., Nakagawa, H., Kashiwabara, Y., Matsuki, G. and Endo, H. (1997) Pathological investigation of armadillos infected with *Mycobacterium leprae*. *Nihon Hansenbyo Gakkai Zasshi* 66, 227–235.
- [14] Fujita, Y., Naka, T., Doi, T. and Yano, I. (2005) Direct molecular mass determination of trehalose monomycolate from 11 species of mycobacteria by MALDI-TOF mass spectrometry. *Microbiology* 151, 1443–1452.
- [15] Fujita, Y., Naka, T., McNeil, M.R. and Yano, I. (2005) Intact molecular characterization of cord factor (trehalose 6,6'-dimycolate) from nine species of mycobacteria by MALDI-TOF mass spectrometry. *Microbiology* 151, 3403–3416.
- [16] Minnikin, D.E. and Polgar, N. (1967) Synthesis of methyl 3,12,15-trimethyldocosanoate and 3,15-dimethyldocosanoate. *J. Chem. Soc.* 7, 575–577.
- [17] Yuan, Y., Crane, D.C., Musser, J.M., Sreevatsan, S. and Barry 3rd., C.E. (1997) MMAS-1, the branch point between *cis*- and *trans*-cyclopropane-containing oxygenated mycolates in *Mycobacterium tuberculosis*. *J. Biol. Chem.* 272, 10041–10049.
- [18] Draper, P., Dobson, G., Minnikin, D.E. and Minnikin, S.M. (1982) The mycolic acids of *Mycobacterium leprae* harvested from experimentally infected nine-banded armadillos. *Ann. Microbiol.* 133, 39–47.
- [19] Takayama, K., Wang, C. and Besra, G.S. (2005) Pathway to synthesis and processing of mycolic acids in *Mycobacterium tuberculosis*. *Clin. Microbiol. Rev.* 18, 81–101.
- [20] Minnikin, D.E., Dobson, G., Goodfellow, M., Draper, P. and Magnusson, M. (1985) Quantitative comparison of the mycolic and fatty acid compositions of *Mycobacterium leprae* and *Mycobacterium goodii*. *J. Gen. Microbiol.* 131, 2013–2021.
- [21] Watanabe, M., Aoyagi, Y., Mitome, H., Fujita, T., Naoki, H., Ridell, M. and Minnikin, D.E. (2002) Location of functional groups in mycobacterial meromycolate chains; the recognition of new structural principles in mycolic acids. *Microbiology* 148, 1881–1902.
- [22] Watanabe, M., Aoyagi, Y., Ridell, M. and Minnikin, D.E. (2001) Separation and characterization of individual mycolic acids in representative mycobacteria. *Microbiology* 147, 1825–1837.
- [23] Glickman, M.S., Cahill, S.M. and Jacobs Jr., W.R. (2001) The *Mycobacterium tuberculosis* *cmaA2* gene encodes a mycolic acid *trans*-cyclopropane synthetase. *J. Biol. Chem.* 276, 2228–2233.
- [24] Yuan, Y., Zhu, Y., Crane, D.D. and Barry 3rd., C.E. (1998) The effect of oxygenated mycolic acid composition on cell wall function and macrophage growth in *Mycobacterium tuberculosis*. *Mol. Microbiol.* 29, 1449–1458.
- [25] Bhatt, A., Fujiwara, N., Bhatt, K., Gurcha, S.S., Kremer, L., Chen, B., Chan, J., Porcelli, S.A., Kobayashi, K., Besra, G.S. and Jacobs Jr., W.R. (2007) Deletion of *kasB* in *Mycobacterium tuberculosis* causes loss of acid-fastness and subclinical latent tuberculosis in immunocompetent mice. *Proc. Natl. Acad. Sci. USA* 104, 5157–5162.
- [26] Hunter, S.W. and Brennan, P.J. (1981) A novel phenolic glycolipid from *Mycobacterium leprae* possibly involved in immunogenicity and pathogenicity. *J. Bacteriol.* 147, 728–735.
- [27] Daffe, M. and Laneelle, M.A. (1989) Diglycosyl phenol phthiocerol diester of *Mycobacterium leprae*. *Biochim. Biophys. Acta* 1002, 333–337.

## Characterization of the Fucosylation Pathway in the Biosynthesis of Glycopeptidolipids from *Mycobacterium avium* Complex<sup>∇</sup>

Yuji Miyamoto,<sup>1</sup> Tetsu Mukai,<sup>1\*</sup> Yumi Maeda,<sup>1</sup> Noboru Nakata,<sup>1</sup> Masanori Kai,<sup>1</sup>  
Takashi Naka,<sup>2</sup> Ikuya Yano,<sup>2</sup> and Masahiko Makino<sup>1</sup>

Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan,<sup>1</sup> and Japan BCG Central Laboratory, 3-1-5 Matsuyama, Kiyose, Tokyo 204-0022, Japan<sup>2</sup>

Received 3 March 2007/Accepted 2 May 2007

The cell envelopes of several species of nontuberculous mycobacteria, including the *Mycobacterium avium* complex, contain glycopeptidolipids (GPLs) as major glycolipid components. GPLs are highly antigenic surface molecules, and their variant oligosaccharides define each serotype of the *M. avium* complex. In the oligosaccharide portion of GPLs, the fucose residue is one of the major sugar moieties, but its biosynthesis remains unclear. To elucidate it, we focused on the 5.0-kb chromosomal region of the *M. avium* complex that includes five genes, two of which showed high levels of similarity to the genes involved in fucose synthesis. For the characterization of this region by deletion and expression analyses, we constructed a recombinant *Mycobacterium smegmatis* strain that possesses the *rfA* gene of the *M. avium* complex to produce serovar 1 GPL. The results revealed that the 5.0-kb chromosomal region is responsible for the addition of the fucose residue to serovar 1 GPL and that the three genes *mdhA*, *merA*, and *gtfD* are indispensable for the fucosylation. Functional characterization revealed that the *gtfD* gene encodes a glycosyltransferase that transfers a fucose residue via 1→3 linkage to a rhamnose residue of serovar 1 GPL. The other two genes, *mdhA* and *merA*, contributed to the formation of the fucose residue and were predicted to encode the enzymes responsible for the synthesis of fucose from mannose based on their deduced amino acid sequences. These results indicate that the fucosylation pathway in GPL biosynthesis is controlled by a combination of the *mdhA*, *merA*, and *gtfD* genes. Our findings may contribute to the clarification of the complex glycosylation pathways involved in forming the oligosaccharide portion of GPLs from the *M. avium* complex, which are structurally distinct.

Mycobacteria have a unique multilayer cell envelope composed of peptidoglycan, arabinogalactan, mycolic acids, and an outer layer that contains abundant species-specific glycolipids. It is thought that these cell wall characteristics allow mycobacteria to survive in host cells (8, 11). Among the glycolipids present in the outer layer of the cell envelope, glycopeptidolipids (GPLs) are recognized as highly antigenic surface molecules in several species of nontuberculous mycobacteria, including the *Mycobacterium avium* complex (29). The common structure in which the fatty acyl-tetrapeptide core is glycosylated with both 6-deoxy-talose (6-d-Tal) and *O*-methyl-rhamnose (*O*-Me-Rha), termed core GPLs, is present in all types of GPLs (2, 4, 12). The core GPLs are further glycosylated with a Rha residue or various oligosaccharides linked to the 6-d-Tal residue. These glycosylation events give rise to structural variations, especially in the GPLs from the *M. avium* complex, that are responsible for the creation of serotypes, and they are also known as serovar-specific GPLs (ssGPLs). The ssGPLs define approximately 30 serotypes, some of which are found predominantly in isolates from patients. For example, the serotypes 1, 4, 6, and 8 are frequently isolated from AIDS patients coinfecting with the *M. avium* complex (17, 30). In addition, ssGPLs are associated with host responses to infection and func-

tion as agonists of cell surface receptors such as Toll-like receptor 2 (28).

The biosynthesis of GPLs, including methylation, acetylation, and peptide synthesis, has been investigated mainly with *M. smegmatis*, but it has not been fully elucidated (6, 15, 21, 23). We have recently identified the glycosyltransferase genes involved in the formation of core GPLs in *M. smegmatis* and *M. avium* (19), but the glycosylation pathways of ssGPLs have not been clarified, except for that of serovar 1 GPL (14). The presence of a Rha residue linked to 6-d-Tal, as observed with serovar 1 GPL, is structurally required for the formation of all types of ssGPLs. The Rha residue is further extended by the subsequent addition of one of the three different sugars, Rha, fucose (Fuc), or glucose, to form the structural group of ssGPLs (9). In these groups, the Fuc-containing ssGPLs, such as serovar 2, 3, 4, and 9 GPLs, shown in Table 1, comprise the major group in the ssGPLs of the *M. avium* complex, and serovar 2 GPL is the basic structure for these ssGPLs (9). The biosynthetic pathway for the Fuc-containing ssGPLs is still unclear. In the *M. avium* complex, a 22- to 25-kb chromosomal region with the ability to synthesize the serovar 2 GPL has been identified and cloned from two strains (4, 5). However, the key genes responsible for the formation of the Fuc residue of serovar 2 GPL have not been identified in this chromosomal region. In this study, we have focused on a 5.0-kb segment that includes the genes predicted to be associated with Fuc synthesis and have characterized their functions in the biosynthesis of GPLs in the *M. avium* complex.

\* Corresponding author. Mailing address: Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan. Phone: 81-42-391-8211. Fax: 81-42-394-9092. E-mail: tmukai@nih.go.jp.

<sup>∇</sup> Published ahead of print on 25 May 2007.