

FIGURE 2. A, Up-regulated expression of APC-associated molecules on DC by infection with BCG-70M. PBMCs were obtained from one donor. Monocyte-derived immature DC were infected with either BCG-261H or BCG-70M at a MOI of 0.5 and cultured for another 2 days in the presence of rGM-CSF and rIL-4. The DC from day 5 were gated and analyzed. Dotted lines, isotype-matched control IgG; solid lines, the indicated test mAb. Representative results of three separate experiments are shown. The number in the top right-hand corner of each panel represents the mean \pm SD for three independent experiments of the difference in mean fluorescence intensity between the control IgG and the test mAb. Titers were statistically compared using Student's *t* test. *, $p < 0.01$; †, $p < 0.05$; ‡, $p < 0.01$; §, $p < 0.05$. B, Cytokine production from DC stimulated with BCG. PBMCs were obtained from one donor. Monocyte-derived DC from 4 days of culture in the presence of rGM-CSF and rIL-4 were stimulated with the indicated dose of either BCG-261H or BCG-70M for 24 h. The concentration of the indicated cytokine was determined by the ELISA method. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test.

though faintly, expressed MMP-II on their surface, and the MMP-II expression was inhibited by the pretreatment of DC with chloroquine, an inhibitor of phagosomal acidification, before BCG-70M infection (data not shown). Moreover, when we examined the influence of BCG-70M infection in DC in terms of the production of proinflammatory cytokines, BCG-70M was superior to BCG-261H in the production of IL-12p70, TNF- α , and IL-1 β (Fig. 2B). These results indicate again that BCG-70M more efficiently activated DC than did the parental BCG.

T cell activation by BCG-70M

Enhanced activation of DC may be closely associated with the efficient activation of both CD4⁺ and CD8⁺ T cell subsets. Because BCG-infected mitomycin C-treated DC were confirmed not to produce IFN- γ (data not shown), the T cell-activating ability of




FIGURE 3. A, IFN- γ production from CD4⁺ T cells by stimulation with BCG. PBMCs were obtained from one donor. Monocyte-derived DC were infected with either BCG-261H or BCG-70M at the indicated MOI, and used as a stimulator of memory-type or naive CD4⁺ T cells in a 4-day culture. Responder CD4⁺ T cells (1×10^5) were stimulated with the indicated dose of BCG-infected DC. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test. B, Inhibition of naive CD4⁺ T cell activation by the treatment of BCG-70M-infected DC with mAb or the treatment of immature DC with chloroquine. PBMCs were obtained from one donor. Monocyte-derived DC were infected with BCG-70M at a MOI of 0.5, and subsequently treated with 10 μ g/ml mAb to HLA-DR, CD86 Ags, or normal murine IgG. Immature DC were treated with 50 μ M chloroquine for 2 h and subsequently infected with BCG-70M at a MOI of 0.5. These DC were used as the stimulator of naive CD4⁺ T cells (1×10^5 /well) at T:DC = 20:1. IFN- γ produced from T cells was measured. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test.

BCG-70M was assessed by using DC as APCs. Memory-type and naive CD4⁺ T cells were purified from healthy BCG-vaccinated individuals, and were stimulated with DC infected with rBCG (Fig. 3A). As expected, memory T cells responded more vigorously to DC infected with smaller dose of BCG than naive CD4⁺ T cells. Although both BCG-261H and BCG-70M stimulated memory and naive CD4⁺ T cells, BCG-70M induced a significantly higher level of IFN- γ production in both types of CD4⁺ T cells than BCG-261H. Note that high levels of IFN- γ could be produced from naive CD4⁺ T cells by BCG-70M. In addition, various MOIs and T:DC ratios were assessed, and a similar difference between BCG-261H and BCG-70M was observed (data not shown). To address the mechanism leading to the activation of naive CD4⁺ T cells, BCG-70M-infected DC were treated with mAbs to HLA-DR and CD86 molecules and subsequently used as a stimulator of the T cells. The IFN- γ production from naive CD4⁺ T cells was significantly inhibited by the surface treatment of BCG-70M-infected DC with the mAb to HLA-DR or CD86 (Fig. 3B), and similarly, IL-2 production from naive CD4⁺ T cells




FIGURE 4. A, IFN- γ production from CD8⁺ T cells by stimulation with BCG. PBMCs were obtained from one donor. Monocyte-derived DC were infected with either BCG-261H or BCG-70M at the indicated MOI, and used as a stimulator of memory-type or naive CD8⁺ T cells in a 4-day culture. Responder CD8⁺ T cells (1×10^5) were stimulated with the indicated dose of BCG-infected DC. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test. B, Inhibition of naive CD8⁺ T cell activation by the treatment of BCG-70M-infected DC with mAb. PBMCs were obtained from one donor. Monocyte-derived DC were infected with BCG-70M at a MOI of 1.0, and subsequently treated with 10 μ g/ml mAb to HLA-ABC, CD86, or MMP-II (M270-13). The DC were used as the stimulator of naive CD8⁺ T cells (1×10^5 /well) at T:DC = 10:1. IFN- γ produced from the T cells was measured by the ELISA method. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test. C, Effect of treatment of immature DC with reagents on the activation of naive CD8⁺ T cells. PBMCs were obtained from one donor. Monocyte-derived immature DC were treated with the indicated dose of either chloroquine, brefeldin A, or lactacystin, and subsequently infected with BCG-70M at a MOI of 1.0. These DC were used as the stimulator of naive CD8⁺ T cells (1×10^5 /well) at T:DC = 10:1. IFN- γ produced from the T cells was measured. A representative of three separate experiments is shown. Assays were performed in triplicate, and the results are expressed as the mean \pm SD. Titers were statistically compared using Student's *t* test.




FIGURE 5. Influence of naive CD4⁺ T cells on the activation of naive CD8⁺ T cells. PBMCs were obtained from one donor. Monocyte-derived DC were infected with BCG (MOI 0.5) and cocultured with unseparated naive T cells (T:DC = 10:1) for 7 days. The stimulated CD8⁺ T cells were gated and analyzed for perforin production and for expression of CD62L Ag. Numbers indicate either percentage of perforin-positive CD8⁺ T cells or CD62L^{low} CD8⁺ T cells among CD8⁺ T cell population. A representative of three separate experiments is shown.

was inhibited (data not shown). Furthermore, pretreatment of immature DC with chloroquine before infection with BCG-70M significantly inhibited the IFN- γ production from naive CD4⁺ T cells (Fig. 3B). These results indicated that the secreted HSP70-MMP-II protein or BCG-70M itself may be processed in the DC, and some of the antigenic peptides were used for the stimulation of autologous Ag-specific naive CD4⁺ T cells. Similarly, BCG-70M-infected DC stimulated memory CD8⁺ T cells more efficiently than BCG-261H-infected DC, although a higher dose of BCG-70M was necessary to induce a similar level of IFN- γ production from CD8⁺ T cells than the dose of BCG-70M required to produce the cytokine from memory CD4⁺ T cells. As reported, BCG-261H did not activate naive CD8⁺ T cells efficiently (15); however, BCG-70M-infected DC induced a significant level of IFN- γ production from naive CD8⁺ T cells (Fig. 4A). Using a higher dose of BCG-70M (MOI 1.0) and a larger number of BCG-70M-infected DC (T:DC = 10:1), a high concentration (\sim 200 pg/ml) of IFN- γ could be produced from naive CD8⁺ T cells. In addition, various MOIs and T:DC ratios were assessed, and a similar difference between BCG-261H and BCG-70M was observed (data not shown). To clarify the mechanism leading to the activation of naive CD8⁺ T cells by BCG-70M, BCG-70M-infected DC were treated with mAbs. Again, the activation of naive CD8⁺ T cells by BCG-70M-infected DC was significantly inhibited by the treatment of the DC with the mAb to HLA-ABC or CD86. However, surface treatment of the DC with the mAb to MMP-II significantly, but only partially, inhibited the T cell activation (Fig. 4B). These results may indicate that BCG-70M-infected DC cross-primed naive CD8⁺ T cells in an Ag-specific manner. To elucidate the mechanisms leading to the cross-presentation by BCG-70M-infected DC, we treated immature DC with various reagents in advance of the BCG-70M infection (Fig. 4C). On the pretreatment of DC with chloroquine, IFN- γ production from naive CD8⁺ T cells was significantly inhibited, indicating that protein derived from BCG-70M was degraded in presumably the phagolysosome. Furthermore, on the pretreatment of DC with brefeldin A, an inhibitor of antegrade Golgi transportation, and lactacystin, an inhibitor of proteosomal protein degradation, IFN- γ production from naive CD8⁺ T cells was inhibited significantly in a manner dependent on the concentration of the reagents. Because BCG-70M activated both naive CD4⁺ T cells and naive CD8⁺ T cells, we stimulated naive CD8⁺ T cells with BCG-70M in the presence of the CD4⁺ T cells (Fig. 5). The expression level of CD62L on some CD8⁺ T cells was




FIGURE 6. A, Production of memory-type T cells in C57BL/6 mice by infection with BCG. Five-week-old C57BL/6 mice were infected with 1×10^3 CFU/mouse of either BCG-261H or BCG-70M s.c. Four weeks after the inoculation, splenocytes (2×10^5 cells/well) were stimulated with the indicated dose of either MMP-II-, HSP70-, or BCG-derived cytosolic protein for 4 days, and IFN- γ produced in the cell supernatant was measured. Assays were performed in triplicate for each mouse, and the results of three mice per group are shown as the mean \pm SD. Representative results of four separate experiments are shown. Titers were statistically compared using Student's *t* test. B, Intracellular IFN- γ production from CD4⁺ T cells and CD8⁺ T cells in C57BL/6 mice s.c. inoculated with BCG by secondary stimulation. Groups of 5-wk-old C57BL/6 mice were infected with 1×10^3 CFU/mouse BCG-261H or BCG-70M intradermally. Four weeks after the inoculation, splenocytes (2×10^5 cells/well) were stimulated with 1.0 μ g/ml rMMP-II for 3 days. The CD4⁺ T cells and CD8⁺ T cells were gated separately and analyzed for intracellular production of IFN- γ . The number in the top right-hand corner of each panel represents the mean \pm SD for three mice in the percentage of IFN- γ -producing cells among the gated T cell population. Representative results of four separate experiments are shown. Titers were statistically compared using Student's *t* test. *, $p < 0.05$; †, $p < 0.01$.

significantly reduced by stimulation with BCG-70M-infected DC, and a significant amount of intracellular perforin was produced in naive CD8⁺ T cells by the stimulation. These changes were more efficiently induced by BCG-70M-infected DC than by BCG-261H-infected DC (Fig. 5). The CD62L^{low}CD8⁺ T cells and perforin-producing CD8⁺ T cells were not produced when naive CD8⁺ T cells were stimulated in the absence of naive CD4⁺ T cells (data not shown).

Memory T cell production by BCG-70M in vivo

Another important aspect to be studied is the production of Ag-specific memory T cells in vivo. C57BL/6 mice were infected with




FIGURE 7. Inhibition of *M. leprae* multiplication by s.c. vaccination with BCG-70M. Five-week-old C57BL/6 mice (8 mice per group) were vaccinated with 1×10^2 or 1×10^3 CFU/mouse either BCG-261H or BCG-70M s.c., and were challenged with 5×10^3 bacilli/mouse *M. leprae* in footpad 4 wk after the vaccination. The number of *M. leprae* recovered from the footpad at 32 wk after the challenge was enumerated by Shepard's methods (29). Representative results of two separate experiments are shown. Titers were statistically compared using Student's *t* test.

1000 CFU/mouse rBCG for 4 wk, and their splenocytes were secondarily stimulated in vitro with rMMP-II protein, or recall Ags, like BCC (Fig. 6A). When a lower dose of MMP-II was used as a stimulator, only T cells obtained from BCG-70M-infected mice responded to the stimulator. Because BCG-Tokyo, the parental strain of the rBCGs, has the gene encoding MMP-II, a higher dose of *M. leprae*-derived MMP-II induced IFN- γ production from both T cells obtained from BCG-261H- and BCG-70M-inoculated mice; however, T cells from BCG-70M-infected mice more efficiently produced the cytokine than those from BCG-261H-infected mice. Also, T cells from BCG-70M-inoculated mice produced significantly higher level of IFN- γ than T cells from mice inoculated with BCG-261H on stimulation with HSP70 in vitro. The efficient generation of memory T cells responding to mycobacteria-derived Ags in mice infected with BCG-70M was confirmed, because only T cells from mice infected with BCG-70M significantly responded to BCC (Fig. 6A). To clarify the T cell subsets responsible for the IFN- γ production on secondary MMP-II stimulation, T cells producing the cytokine intracellularly were analyzed (Fig. 6B). Both CD4⁺ T cells and CD8⁺ T cells derived from not only non-BCG-inoculated mice, but also BCG-infected mice, produced intracellular IFN- γ by MMP-II stimulation. Both noninoculated and PBS-inoculated mice showed the similar responses (data not shown). However, significantly larger populations of both CD4⁺ T cells (~1.3%) and CD8⁺ T cells (~0.5%) obtained from BCG-70M-infected mice produced the cytokine. There were no CD4⁺ T cells or CD8⁺ T cells that were positively stained with the isotype control IgG (data not shown).

Effect of BCG-70M vaccination on the multiplication of *M. leprae* in vivo

C57BL/6 mice vaccinated with either BCG-261H or BCG-70M (1×10^2 or 1×10^3 CFU/mouse) for 4 wk were challenged with 5×10^3 of *M. leprae* in footpad. Thirty-two weeks later, footpad was removed and *M. leprae* recovered from the footpad was enumerated (Fig. 7). A total of 2×10^5 *M. leprae* was recovered from mice inoculated with PBS and challenged with *M. leprae*. Both mice inoculated with BCG-261H or BCG-70M inhibited the multiplication of *M. leprae* in the manner dependent on the dose of rBCG vaccinated; however, BCG-70M vaccination was significantly more efficient than BCG-261H vaccination in inhibiting the

multiplication, and only 2×10^4 *M. leprae* was recovered from mice vaccinated with 1×10^3 CFU/mouse BCG-70M.

Discussion

M. leprae is a representative mycobacterium among slow-growing prokaryotes, which needs 12–14 days for one division and 10–70 years for the manifestation of the disease, depending on the dose of bacilli entering the hosts. Host defense against *M. leprae* is chiefly conducted by adaptive immunity involving both CD4⁺ and CD8⁺ T cell subsets (3–5). Although CD4⁺ T cells usually act at the initial phase of infection, CD8⁺ T cells inhibit the multiplication of *M. leprae* in the chronic phase or when it is reactivated from a dormant state (31). Therefore, the vaccine should have an ability to competently activate not only CD4⁺ T cells, but also CD8⁺ T cells to produce memory T cells. To date, BCG is used as sole, but safe, vaccine against leprosy; however, nowadays, its efficacy is considered not so convincing (13). We have made several attempts to improve the potency of BCG, especially its immunostimulatory activities. We chiefly focused on achieving the fusion of BCG-infected phagosomes with lysosomes, without which the full and polyclonal activation of Ag-specific T cells cannot be expected. One approach we tried was the production of an ure C-deficient rBCG that successfully produces acidic phagosomes and facilitates their fusion with lysosomes (15, 28, 32, 33). Actually, the rBCG efficiently colocalized with lysosomes and effectively stimulated CD4⁺ T cells when DC were targeted (28). However, it did not activate naive CD8⁺ T cells. Then, we produced a second rBCG that secretes MMP-II (BCG-SM) in the phagosome (14). BCG-SM was useful in the activation of not only naive CD4⁺ T cells, but also naive CD8⁺ T cells to some extent (14). The T cell activation presumably occurs as a consequence of the translocation of the antigenic determinants of the secreted MMP-II to the cell surface, although the precise mechanism has not been clarified. Therefore, the intracellular secretion of immunodominant Ag by BCG is thought necessary to enhance the T cell-stimulating activity of BCG. However, BCG-SM vaccinations only partially inhibited the multiplication of *M. leprae* in the footpads of mice (our unpublished observation). These observations indicate the need for another rBCG capable of activating both naive CD4⁺ and CD8⁺ T cells more strongly.

Because the strong activation of naive CD8⁺ T cells by mycobacteria required the cross-presentation of mycobacteria-derived Ags to CD8⁺ T cells, in this study, we used BCG-derived HSP70 as a mediator facilitating the cross-presentation by DC, because HSPs of both mammalian host cells and bacterial origin are reported to have chaperone activity (34) and can effectively prime a cytolytic response (35). The usefulness of HSP65 as a convincing stimulator of CD8⁺ T cells was revealed in animal studies to develop vaccine against *M. tuberculosis* (36). A DNA vaccine containing the *hsp65* gene activated naive CD8⁺ T cells, and consequently inhibited the development of tuberculosis, although precisely how was not clarified. Our newly constructed rBCG (BCG-70M) secreted a fusion protein comprising HSP70 and MMP-II in vitro in the absence of any exogenous secretion signal. The secretion of HSP70-MMP-II fusion was confirmed by Western blotting analyses (Fig. 1A) and by the surface expression of MMP-II on DC (data not shown). The exact mechanism leading to the secretion of the fusion protein from BCG-70M is not known, but the secretion could be due to the inherent characteristics of HSP70 to be secreted (16, 37). Although we tried to enhance the secreting activity of BCG-70M by additionally inserting *M. tuberculosis*-derived secretion signal to BCG-70M, the secretion efficacy was rather inhibited and the construct stimulated naive T cells less efficiently than BCG-70M (data not shown).

BCG-70M secreted a 90-kDa fusion protein composed of HSP70 and MMP-II (14). The activation of naive CD8⁺ T cells by BCG-70M was only partially inhibited by the treatment of DC with the mAb to MMP-II. Therefore, it can be speculated that BCG-70M activated CD8⁺ T cells polyclonally by using various epitopes, originating from MMP-II, HSP70, or other Ags of BCG, because T cells from BCG-70M-infected mice vigorously responded to MMP-II, HSP70, and BCG-derived cytosolic protein (Fig. 6A). Therefore, HSP70 may alter the clonality of responding CD8⁺ T cells, and the production of such polyclonal Ag-specific CD8⁺ T cells might be beneficial for the broad coverage of a heterogeneous MHC population.

BCG-70M induced higher level of cytokine production, including IL-12p70, TNF- α , and IL-1 β , than the vector control BCG. The enhanced cytokine production by BCG-70M could be due to the intracellular secretion of HSP70 as a part of the fusion protein by the BCG. MMP-II can ligate TLR2 (6), but HSP70 is also known to bind TLRs (38); thus, the secreted HSP70-MMP-II protein seems to activate DC strongly. The contribution of TLR2 to cytokine production was confirmed by the inhibition of IL-12p70 production by antagonistic Ab to TLR2. The cytokines released from DC by BCG-70M stimulation could facilitate skewing of the direction of T cell activation to type 1 and induce the efficient and strong production of IFN- γ from naive CD8⁺ T cells.

HSPs play a varied role in enhancing the ability of APCs to stimulate T cells (39). For the activation of Ag-specific CD4⁺ and CD8⁺ T cells, peptides should be loaded onto the corresponding MHC pathways. For the loading of BCG-derived Ags on these pathways, the proteins secreted from phagocytosed BCG should be transported to functional lysosomes. In the phagolysosome, some portions of HSP70-MMP-II fusion protein could be degraded, but the rest would be sequestered into the cytosol, where they are degraded and used for cross-priming CD8⁺ T cells. BCG-70M-infected DC expressed derivatives of MMP-II and the other proteins on their surface, and they activated both naive CD4⁺ T cells and naive CD8⁺ T cells. However, both MMP-II expression on DC and the activation of the T cells by DC were inhibited by the pretreatment of DC with chloroquine (24, 40). These results indicate that secreted HSP70-MMP-II fusion protein was efficiently processed in lysosomes and its derivatives are used for the activation of both subsets of naive T cells. When naive CD8⁺ T cells were stimulated by BCG-70M in the presence of naive CD4⁺ T cells, CD62L^{low}CD8⁺ T cells and perforin-producing CD8⁺ T cells were efficiently produced. The activation of naive T cells was confirmed by the production of memory-type T cells by BCG-70M infection to unprimed mice, because both CD4⁺ and CD8⁺ T cell subsets from BCG-70M-infected mice responded to the restimulation with MMP-II in vitro. Furthermore, BCG-70M significantly and more efficiently inhibited the multiplication of *M. leprae*, which were challenged in footpad of mice, than BCG-261H.

There are two pathways of cross-presentation, as follows: cytosolic (ER-Golgi-dependent) and vacuolar pathways (20). It is known that HSP can enhance both pathways (20). In the present study, IFN- γ production from naive CD8⁺ T cells was largely blocked by the treatment of DC with brefeldin A, an inhibitor of antegrade Golgi transportation and of TAP-dependent transportation, and also with lactacystin, a proteasomal protein degradation blocker (20, 40). Therefore, it can be presumed that the fusion protein was sequestered into the cytosol from the lysosome, degraded in the proteasome, and used for loading on MHC class I molecules through the TAP-dependent pathway. Furthermore, it has been reported that proteins that are intracellularly secreted are usually processed by a cytosolic (ER-Golgi-dependent) pathway, and DC prefer this pathway for cross-priming CD8⁺ T cells with

protein Ag (20). Our present observations seem to fit well with these previous findings. Therefore, we concluded that BCG-70M activates naive CD8⁺ T cells through the ER-Golgi-dependent cytosolic cross-presentation pathway. However, *M. tuberculosis*-derived HSP70 activated T cells through a post-Golgi, proteasomal-independent mechanism, and both brefeldin A and lactacystin may inhibit vacuolar pathway in some cases (20). Also, the vacuolar pathway is used more frequently by macrophages (20). Therefore, the possibility that BCG-70M may also use the post-Golgi pathway in vivo cannot be ruled out.

Taken together, in this study, we constructed a rBCG that secretes HSP70-MMP-II fusion protein, which effectively activates not only DC, but also naive T cells. Therefore, the combination of HSP70 and MMP-II may be useful for stimulating both subsets of naive T cells.

Acknowledgments

We acknowledge the contribution of N. Makino in the preparation of the manuscript. We also thank Y. Harada for her technical support, and the Japanese Red Cross Society for kindly providing PBMCs from healthy donors.

Disclosures

The authors have no financial conflict of interest.

References

- Hagge, D. A., N. A. Ray, J. L. Krahenbuhl, and L. B. Adams. 2004. An in vitro model for the lepromatous leprosy granuloma: fate of *Mycobacterium leprae* from target macrophages after interaction with normal and activated effector macrophages. *J. Immunol.* 172: 7771–7779.
- Ridley, D. S., and W. H. Jopling. 1966. Classification of leprosy according to immunity: a five-group system. *Int. J. Lepr. Other Mycobact. Dis.* 34: 255–273.
- Modlin, R. L., J. Melancon-Kaplan, S. M. M. Young, C. Piemez, H. Kino, J. Convit, T. H. Rea, and B. R. Bloom. 1988. Learning from lesions: patterns of tissue inflammations in leprosy. *Proc. Natl. Acad. Sci. USA* 85: 1213–1217.
- Ramasesh, N., L. B. Adams, S. G. Franzblau, and J. L. Krahenbuhl. 1991. Effects of activated macrophages on *Mycobacterium leprae*. *Infect. Immun.* 59: 2864–2869.
- Hashimoto, K., Y. Maeda, H. Kimura, K. Suzuki, A. Masuda, M. Matsuoka, and M. Makino. 2002. Infection of *M. leprae* to monocyte derived dendritic cells and its influence on antigen presenting function. *Infect. Immun.* 70: 5167–5176.
- Maeda, Y., T. Mukai, J. Spencer, and M. Makino. 2005. Identification of immunomodulating agent from *Mycobacterium leprae*. *Infect. Immun.* 73: 2744–2750.
- Makino, M., Y. Maeda, and N. Ishii. 2005. Immunostimulatory activity of major membrane protein-II from *Mycobacterium leprae*. *Cell. Immunol.* 233: 53–60.
- Sieling, P. A., D. Jullien, M. Dahlem, T. F. Tedder, T. H. Rea, R. L. Modlin, and S. A. Porcelli. 1999. CD1 expression by dendritic cells in human leprosy lesions: correlation with effective host immunity. *J. Immunol.* 162: 1851–1858.
- Seiling, P. A., M. T. Ochoa, D. Jullien, D. S. Leslie, S. Sabet, J. P. Rosat, A. E. Burdick, T. H. Rea, M. B. Brenner, S. A. Porcelli, and R. A. Modlin. 2000. Evidence for human CD4⁺ T cells in the CD1-restricted repertoire: derivation of mycobacteria reactive T cells from leprosy lesions. *J. Immunol.* 164: 4790–4796.
- Lombardi, C., E. S. Pedrazzani, J. C. Pedrazzani, P. F. Filho, and F. Zicker. 1996. Protective efficacy of BCG against leprosy in San Paulo. *Bull. Pan Am. Health Organ.* 30: 24–30.
- Ponnighaus, J. M., P. E. Fine, J. A. Sterne, R. J. Wilson, E. Msosa, P. J. Gruer, P. A. Jenkins, S. B. Lucas, N. G. Liomba, and L. Bliss. 1992. Efficacy of BCG vaccine against leprosy and tuberculosis in northern Malawi. *Lancet* 14: 636–639.
- Sharma, P., R. Mukherjee, G. P. Talwar, K. G. Sarathchandra, R. Walia, S. K. Parida, R. M. Pandey, R. Rani, H. Kar, A. Mukherjee, et al. 2005. Immunoprophylactic effects of the anti-leprosy *Mw* vaccine in household contacts of leprosy patients: clinical field trails with a follow up of 8–10 years. *Lepr. Rev.* 76: 127–143.
- Setia, M. S., C. Steinmaus, C. H. Ho, and G. W. Rutherford. 2006. The role of BCG in prevention of leprosy: a meta-analysis. *Lancet Infect. Dis.* 6: 162–170.
- Makino, M., Y. Maeda, and K. Inagaki. 2006. Immunostimulatory activity of recombinant *Mycobacterium bovis* BCG that secretes major membrane protein II of *Mycobacterium leprae*. *Infect. Immun.* 74: 6264–6271.
- Grode, L., P. Seiler, S. Baumann, J. Hess, V. Brinkmann, A. N. Eddine, P. Mann, C. Goosmann, S. Bandermann, D. Smith, et al. 2005. Increased vaccine efficacy against tuberculosis of recombinant *Mycobacterium bovis* bacille Calmette-Guérin mutants that secrete listeriolysin. *J. Clin. Invest.* 115: 2472–2479.
- Horwitz, M. A., B. W. Lee, B. J. Dillon, and G. Harth. 1995. Protective immunity against tuberculosis induced by vaccination with major extracellular proteins of *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* 92: 1530–1534.
- Binder, R. J., and P. K. Srivastava. 2005. Peptides chaperoned by heat-shock proteins are necessary and sufficient source of antigen in the cross-priming of CD8⁺ T cells. *Nat. Immunol.* 6: 593–599.
- Flechtner, J. B., K. P. Cohane, S. Mehta, P. Slusarewicz, A. K. Leonard, B. H. Barber, D. L. Levey, and S. Andjelic. 2006. High-affinity interactions between peptides and heat shock protein 70 augment CD8⁺ T lymphocyte immune responses. *J. Immunol.* 177: 1017–1027.
- Tobian, A. A. R., D. H. Canaday, W. H. Boom, and C. V. Harding. 2004. Bacterial heat shock proteins promote CD91-dependent class I MHC cross-presentation of chaperoned peptide to CD8⁺ T cells by cytosolic mechanisms in dendritic cells versus vacuolar mechanisms in macrophages. *J. Immunol.* 172: 5277–5286.
- Tobian, A. A. R., C. V. Harding, and D. H. Canaday. 2005. *Mycobacterium tuberculosis* heat shock fusion protein enhances class I MHC cross-processing and -presentation by B lymphocytes. *J. Immunol.* 174: 5209–5214.
- Makino, M., and M. Baba. 1997. A cryopreservation method of human peripheral blood mononuclear cells for efficient production of dendritic cells. *Scand. J. Immunol.* 45: 618–622.
- Wakamatsu, S., M. Makino, C. Tei, and M. Baba. 1999. Monocyte-driven activation-induced apoptotic cell death of human T-lymphotropic virus type I-infected T cells. *J. Immunol.* 163: 3914–3919.
- Makino, M., S. Shimokubo, S. Wakamatsu, S. Izumo, and M. Baba. 1999. The role of human T-lymphotropic virus type I (HTLV-I)-infected dendritic cells in the development of HTLV-I-associated myelopathy/tropical spastic paraparesis. *J. Virol.* 73: 4575–4581.
- Maeda, Y., M. Gidoh, N. Ishii, C. Mukai, and M. Makino. 2003. Assessment of cell mediated immunogenicity of *Mycobacterium leprae*-derived antigens. *Cell. Immunol.* 222: 69–77.
- Bardarov, S., S. Bardarov, Jr., M. S. Pavelka, Jr., V. Sambandamurthy, M. Larsen, J. Tufariello, J. Chan, G. Hatfull, and W. R. Jacobs, Jr. 2002. Specialized transduction: an efficient method for generating marked and unmarked targeted gene disruptions in *Mycobacterium tuberculosis*, *M. bovis* BCG and *M. smegmatis*. *Microbiology* 148: 3007–3017.
- Snapper, S. B., L. Iugosi, A. Jekkel, R. E. Melton, T. Kieser, B. R. Bloom, and W. R. Jacobs, Jr. 1988. Lysogeny and transformation in mycobacteria: stable expression of foreign genes. *Proc. Natl. Acad. Sci. USA* 85: 6987–6991.
- Sambrook, J., and D. W. Russell. 2001. *Molecular Cloning: A Laboratory Manual*, 3rd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Mukai, T., Y. Maeda, T. Tamura, Y. Miyamoto, and M. Makino. 2008. CD4⁺ T-cell activation by antigen-presenting cells infected with urease-deficient recombinant *Mycobacterium bovis* bacillus Calmette-Guérin. *FEMS Immunol. Med. Microbiol.* 53: 96–106.
- Shepard, C. C., and D. H. McRae. 1968. A method for counting acid-fast bacteria. *Int. J. Lepr. Other Mycobact. Dis.* 36: 78–82.
- Wang, Y., T. Whittall, E. McGowan, J. Younson, C. Kelly, L. A. Bergmeier, M. Singh, and T. Lehner. 2005. Identification of stimulating and inhibitory epitopes within the heat shock protein 70 molecule that modulate cytokine production and maturation of dendritic cells. *J. Immunol.* 174: 3306–3316.
- Kaufmann, S. H. E., and A. J. McMichael. 2005. Annulling a dangerous liaison: vaccination strategies against AIDS and tuberculosis. *Nat. Med.* 11: S33–S44.
- Clements, D. I., B. Y. Lee, and M. A. Horwitz. 1995. Purification, characterization, and genetic-analysis of *Mycobacterium tuberculosis* urease, a potentially critical determinant of host-pathogen interaction. *J. Bacteriol.* 177: 5644–5652.
- Reyrat, J. M., F. X. Berthet, and B. Gicquel. 1995. The urease locus of *Mycobacterium tuberculosis* and its utilization for the demonstration of allelic exchange in *Mycobacterium bovis* bacillus Calmette-Guérin. *Proc. Natl. Acad. Sci. USA* 92: 8768–8772.
- Flynn, G. C., T. G. Chappell, and J. E. Rothman. 1989. Peptide binding and release by proteins implicated as catalysts of protein assembly. *Science* 245: 385–390.
- Moroi, Y., M. Mayhew, J. Trcka, M. H. Hoe, Y. Takechi, F. U. Hartl, J. E. Rothman, and A. N. Houghton. 2000. Induction of cellular immunity by immunization with novel hybrid peptides complexed to heat shock protein 70. *Proc. Natl. Acad. Sci. USA* 97: 3485–3490.
- Yoshida, S., T. Tanaka, Y. Kita, S. Kuwayama, N. Kanamura, Y. Muraki, S. Hashimoto, Y. Inoue, M. Sakatani, B. Kobayashi, et al. 2006. DNA vaccine using hemagglutinating virus of Japan-liposome encapsulating combination encoding mycobacterial heat shock protein 65 and interleukin-12 confers protection against *Mycobacterium tuberculosis* by T cell activation. *Vaccine* 24: 1191–1204.
- Belisle, J. T., V. D. Vissa, T. Sievert, K. Takayama, P. J. Brennan, and G. S. Besra. 1997. Role of the major antigen of *Mycobacterium tuberculosis* in cell wall biogenesis. *Science* 276: 1420–1422.
- Asea, A., M. Rehli, E. Kabingu, J. A. Boch, O. Bare, P. E. Auron, M. A. Stevenson, and S. K. Calderwood. 2002. Novel signal transduction pathway utilized by extracellular HSP70: role of Toll-like receptor (TLR) 2 and TLR4. *J. Biol. Chem.* 277: 15028–15304.
- Wang, Y., C. G. Kelly, M. Singh, E. G. McGowan, A. S. Carrara, L. A. Bergmeier, and T. Lehner. 2002. Stimulation of Th1-polarizing cytokines, C-C chemokines, maturation of dendritic cells, and adjuvant function by the peptide binding fragment of heat shock protein 70. *J. Immunol.* 169: 2422–2429.
- Schiavo, R., D. Baatar, P. Olkhaund, F. E. Indig, N. Restifo, D. Taub, and A. Biragyn. 2006. Chemokine receptor targeting efficiently directs antigens to MHC class I pathways and elicits antigen-specific CD8⁺ T-cell responses. *Blood* 107: 4597–4605.

Temperature dependency for survival of *Mycobacterium leprae* in macrophages

Yasuo FUKUTOMI * , Yumi MAEDA, Masanori MATSUOKA,
and Masahiko MAKINO

Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan

[Received: 4 Aug. 2008 / Accepted: 15 Oct. 2008]

Key words : human, macrophage, mouse, *Mycobacterium leprae*, survival

Hansen's disease is caused by an infection with an intracellular pathogen, *Mycobacterium leprae*, which mainly inhabits macrophages and Schwann cells. However, little is known about the survival or growth mechanisms of the bacilli in mouse and human macrophages. In the present study, by using radiorespirometry analysis for the evaluation of the viability of *M.leprae*, we observed that *in vitro* incubation of *M.leprae*-infected macrophages at 35°C was more growth permissive than at 37°C, and supplementation with the immunosuppressive cytokine IL-10 supported the survival of the bacilli in the macrophages for 3 weeks, whereas viability of the bacilli was gradually lost if cultured without IL-10. In human macrophages, *M.leprae* retained its viability when cultured at 35°C for at least 4 weeks without IL-10. However, the viability of *M.leprae* was almost lost within 2 weeks if cultured at 37°C. These data suggest that temperature is a crucial factor for the survival of *M.leprae* in host cells.

Introduction

Hansen's disease is caused by an infection with *Mycobacterium leprae*. *M.leprae* is an intracellular pathogen, mainly residing in macrophages and Schwann cells. In patients,

M.leprae is predominantly observed in the skin, nasal mucosa and peripheral nerves, particularly the more superficial ones. This clinical observation suggests that the optimal temperature of *M.leprae* for survival in human cells is less than 37°C¹⁾. In animal models, *M.leprae* multiplies in the mouse footpad where the temperature is lower than the core temperature, and the optimal temperature for the growth of *M.leprae* is reported to be in the range of several degrees above and below 30°C²⁾.

From another aspect, the growth of *M.leprae* seems to be largely affected by the host immune

*Corresponding author :

Department of Microbiology, Leprosy Research Center,
National Institute of Infectious Diseases

4-2-1, Aoba-cho, Higashimurayama-shi, Tokyo 189-0002, Japan

TEL : 81-42-391-8211 FAX: 81-42-394-9092

E-mail : fukutomi@nih.go.jp

response. Hansen's disease is characterized by a broad spectrum of the host immune response, such as lepromatous type (towards the increased load of bacteria) and tuberculoid type (towards the decreased bacterial load). In lepromatous type leprosy, Th-2 cytokines (IL-4, IL-5 and IL-10) are predominantly expressed in local lesions. In contrast, in tuberculoid type, Th-1 cytokines (IFN- γ , IL-2) are predominantly expressed³⁾. Among cytokines, IFN- γ has been demonstrated to play a central role in activating macrophages to kill intracellular pathogens including *M.leprae*, whereas IL-10 is reported to inhibit the microbicidal activity of macrophages, resulting in the survival of the intracellular pathogen⁴⁾. However, little is known about the survival and growth mechanisms of *M.leprae* in human macrophages since the viability of these uncultivable bacilli cannot be easily measured by in vitro study.

Previously we reported that metabolically active *M.leprae* were maintained in monolayer cultures of mouse peritoneal macrophages and supplemental IL-10 bolstered *M.leprae* metabolism in the macrophages for as long as 8 weeks. In the cell culture system temperature is extremely important and 31-33°C incubation temperature is more growth permissive than 37°C⁵⁾. In the present study, we observed that incubation of mouse macrophages at 35°C was also more permissive than at 37°C and supplemental IL-10, but not TGF- β , supported the metabolic activity of *M.leprae* in the macrophages for several weeks. Moreover, *M.leprae* from infected human macrophages cultured in vitro sustained metabolic activity for at least 4 weeks if cultured at 35°C but not at 37°C. Collectively, these data demonstrate that temperature is one of the crucial factors for *M.leprae* survival in human host cells.

Materials and Methods

***M.leprae* inoculum:** The Thai-53 strain of *M.leprae*⁶⁾ was maintained in continuous passage in athymic *nu/nu* mice (Clea Co, Tokyo, Japan) by inoculation of bacilli into both hind foot pads. Experiments with mice were performed in compliance with the guidelines of the Experimental Animal Committee of the National Institute of Infectious Diseases. At approximately one year post inoculation, the foot pads were processed to recover *M.leprae* by Nakamura's method with a slight modification⁷⁾. Briefly, tissue was minced and homogenized with Hanks' balanced salt solution (HBSS) containing 0.05% Tween 80. The homogenate was centrifuged at 150×g for 10 min and supernatant of the sample homogenate was treated with 0.05% trypsin at 37°C for 60min. The suspension was centrifuged at 4,000×g for 20min and sediment was re-suspended in HBSS followed by treatment with 1% sodium hydroxide at 37°C for 15min. The treated material was washed and re-suspended in HBSS at the desired bacillary concentration. Bacillary number in each foot pad was enumerated individually according to standard techniques⁸⁾.

Cytokines: Murine recombinant IL-10 was obtained from Genzyme Corp. TGF- β was obtained from Kurashiki Bouseki (Kurashiki, Japan). Both cytokines were stored at -80°C until use.

Mouse macrophage culture: Mouse peritoneal resident cells (approximately 50% macrophages) were harvested from retired ICR mice and suspended as previously described⁹⁾ at a concentration of 2×10⁶/ml in RPMI 1640 (Gibco BRL, Invitrogen Corp., Carlsbad, CA) supplemented with 15% fetal bovine serum (FBS, HyClone Laboratories, Logan UT), 25 mM N-2-

hydroxyethylpiperazine -N'- 2-ethanesulfonic acid (HEPES), 2 mM glutamine and 100µg/ml ampicillin (Sigma Chemical Co., St. Louis, MO). One half ml was seeded into 24 well tissue culture plates (Corning) containing 13 mm LUX coverslips (Nunc Thermanox coverslips, NalgeNunc, Thermo Scientific, Rochester, NY). After 20 hr adherence of the cells, macrophage monolayers were obtained after washing non-adherent cells from the coverslip with Hanks Balanced Salt Solution (HBSS, Sigma) leaving approximately 1×10^6 macrophages adhered per coverslip.

Human macrophage culture: Human peripheral blood was obtained under informed consent from healthy individuals. Peripheral blood mononuclear cells (PBMC) were isolated using Ficoll-Paque Plus (GE Healthcare Life Sciences, Buckinghamshire, HP7 9NA, UK) gradient centrifugation¹⁰. The cells were suspended in AIM-V medium (Gibco BRL, Invitrogen Corp., Carlsbad, CA) and 1×10^6 PBMC were cultured in a well of a 24-well tissue culture plate (Falcon, Becton Dickinson Labware, Becton Dickinson and Company, Franklin Lakes) containing 13 mm LUX coverslips at 37°C in a 5%-CO₂ incubator for adherence of monocytes. After 1 hr incubation, the coverslips were washed with HBSS to remove non-adherent cells. The monocytes on the coverslips were cultured in a new 24-well plate containing RPMI1640 medium (Sigma) supplemented with 20% FBS (Whittaker Co., Walkersville, MD), 25mM HEPES, 2mM L-glutamine and 100µg/ml ampicillin in the presence of 10 ng/ml of human M-CSF (R&D Systems, Minneapolis, MN) or 40 ng/ml of GM-CSF (R&D Systems). After 7 days, the M-CSF-conditioned macrophages (M-macrophages) and the GM-CSF-conditioned macrophages (GM-macrophages) were used for infection with *M.leprae*.

Infection of macrophages with *M.leprae*:

Purified mouse macrophage monolayers were infected with fresh *M.leprae* suspended in 0.5 ml medium per well. After 4 hr incubation for mouse macrophages and 20 hr incubation for human macrophages, non-phagocytosed bacilli were removed by washing and the cultures were incubated in 1.0 ml media supplemented with the appropriate cytokine in 5% CO₂ at the appropriate experimental temperatures⁹. Media were changed and cytokines were replenished at 5 days interval.

Radiorespirometry: The macrophages were lysed with 0.1 N NaOH to release the phagocytosed *M.leprae*, and the viability of the bacilli was determined by evaluating the oxidation of ¹⁴C-palmitic acid to ¹⁴CO₂ by radiorespirometry as described previously¹¹. Total isotope release was usually analyzed after one week of incubation at 31°C⁹.

Staining of *M.leprae*-infected macrophages: Coverslips of *M.leprae*-infected adherent macrophages were prefixed with absolute methanol and acid-fast stained. The specimens were observed under Nikon Optiphot light microscopy.

Results

Viability of *M.leprae* in mouse macrophages cultured *in vitro*: Mouse peritoneal resident macrophages (1×10^6 cells per well) were incubated with freshly harvested *M.leprae* (multiplicity of infection (MOI), 5:1 or 10:1) for 4 hr to allow phagocytosis. Non-phagocytosed bacilli were washed off and the culture of the macrophages continued for up to 14 days. Viability (metabolic activity) of *M.leprae* in macrophages was assessed by radiorespirometry. As shown in Fig. 1, the viability of the bacilli was gradually decreased in macrophages cultured at 35°C. In contrast, the viability was significantly lost, if the macrophages were cultured at 37°C. Next, the mouse peritoneal

resident macrophages were incubated with 3 doses of *M.leprae* (MOI, 1:1, 4:1 and 10:1) for 4 hr to allow phagocytosis, and the culture continued for longer periods up to 21 days. Viability of *M.leprae* in macrophages was assessed at 7 day intervals. As

shown in Fig. 2, in each dose of *M.leprae* infection, decrease in viability was significant after 21 days.

Effects of cytokines on viability of *M.leprae* in mouse macrophages: Supplementation of IL-10 to the infected macrophage culture was




Fig.1. Viability of *M.leprae* in mouse macrophages cultured *in vitro*. Mouse peritoneal resident macrophages were incubated with 5×10^6 or 1×10^7 per well of *M.leprae* (MOI, 5:1 or 10:1), for 4 hr at 37°C to allow phagocytosis. Non-phagocytosed bacilli were washed off and the culture of the macrophages continued up to 14 days at 35°C or 37°C. The cells were lysed to obtain *M.leprae* and metabolism of the bacilli was assessed by radiorespirometry.




Fig.2. Viability of *M.leprae* in mouse macrophages cultured *in vitro*. Mouse peritoneal resident macrophages were incubated with 3 doses, 1×10^6 , 4×10^6 and 1×10^7 per well of *M.leprae* (MOI, 1:1, 4:1 and 10:1) at 35°C for 4 hr to allow phagocytosis, and the culture continued at 35°C for longer periods up to 21 days. The cells were lysed to obtain *M.leprae* and metabolism of the bacilli was assessed by radiorespirometry.

clearly associated with sustained viability of intracellular *M.leprae* cultured at 35°C (Fig.3). In the presence of 3 U/ml of IL-10, *M.leprae* maintained their viability, whereas viability was

steadily lost without IL-10. We also examined the effect of TGF-β, another suppressive cytokine for macrophage activation, on the viability of the bacilli. To the contrary, supplementation of TGF-β




Fig.3. Effect of IL-10 on *M.leprae* survival in mouse macrophages. Mouse peritoneal resident macrophages were incubated with 1×10^7 per well of *M.leprae* (MOI, 10:1) at 35°C for 4 hr to allow phagocytosis, and the culture continued at 35°C for 9, 18 and 27 days. The cells were lysed to obtain *M.leprae* and metabolism of the bacilli was assessed by radiorespirometry.




Fig.4. Viability of *M.leprae* in human macrophages cultured *in vitro*. Human M- or GM-macrophages were incubated with *M.leprae* (MOI, 50:1) for 20 hr either at 35°C or 37°C for infection and incubated at the same temperatures for indicated periods. By observation of the acid fast-stained cells under light microscopy, no difference was recognized in the number of *M.leprae* phagocytosed by macrophage cultured between at 35°C and at 37°C. So the viability at day 0 is considered equal. After 7 days and 13 days incubation period, the cells were lysed to obtain *M.leprae* and metabolism of the bacilli was assessed by radiorespirometry (dpm: disintegrations per minute).

significantly decreased the viability of *M. leprae*, when incubated for longer than 28 days post infection (Table 1).

Viability of *M. leprae* in human macrophages cultured *in vitro*: Human macrophages were obtained by culturing monocytes in the presence

of either M-CSF or GM-CSF for 7 days. These macrophages (1×10^5 cells per well) were incubated with *M. leprae* (MOI, 50:1) for 20 hr either at 35°C or 37°C for infection and incubated again at the same temperatures. By observation of the acid fast-stained cells under light microscopy,




Fig.5. Viability of *M. leprae* in human macrophages cultured *in vitro*. Human M- or GM-macrophages were infected with *M. leprae* (MOI, 50:1) for 20 hr at 35°C and incubated again at 35°C for indicated periods. The cells were lysed to obtain *M. leprae* and metabolism of the bacilli was assessed by radiorespirometry. The results at day 7, 14, 21 and 28 are expressed as percentages of *M. leprae* metabolic activity at time 0. Radiorespirometry data obtained from *M. leprae* in M-macrophages at time 0 was $5,932 \pm 399$ and those in GM-macrophages was $3,084 \pm 78$. *P values calculated in comparison to day 0 viability.

Table 1. Effect of TGF- β on survival of *M. leprae* in mouse macrophages cultured at 35°C^a

Experiment 1				
Days of incubation period	At time 0	7	14	28
Medium only	$5,222 \pm 936^b$	$2,774 \pm 295$	$3,086 \pm 425$	$2,828 \pm 1,815$
+TGF- β		$2,919 \pm 535$	$3,119 \pm 1,339$	$1,973 \pm 126$
Experiment 2				
Days of incubation period	At time 0	14	28	49
Medium only	$26,791 \pm 1,428$	$19,103 \pm 621$	$7,420 \pm 2,986$	$5,713 \pm 1,144$
+TGF- β		$14,306 \pm 2,240$	$3,728 \pm 410$	$1,594 \pm 317$

^aMouse peritoneal resident macrophages were incubated with *M. leprae* (MOI, 1:10) for 4 hr to allow phagocytosis, and the culture continued for indicated periods. The cells were lysed to obtain *M. leprae* and metabolism of the bacilli was assessed by radiorespirometry.

^bRadiorespirometry data, cpm.

Dose of TGF- β , 500pg/ml.

N.D., not determined.

no difference was recognized in the number of *M.leprae* phagocytosed by macrophage cultured at 35°C and at 37°C (data not shown). Viability of *M.leprae* was assessed after 7 and 13 days. The results clearly showed that the viability of *M.leprae* incubated at 35°C was maintained, whereas the viability was lost if cultured at 37°C (Fig. 4). Next, *M.leprae*-infected human M- and GM-macrophages were cultured for prolonged periods at 35°C. Viability was sustained well for 4 weeks in human macrophages, especially in M-macrophages (Fig. 5).

Discussion

In vivo M.leprae is able to enter and survive in a wide variety of tissues and cell types¹²⁾. The preferred host cell for the leprosy bacillus appears to be the macrophages and a number of unsuccessful attempts have been made to grow *M.leprae* in macrophages *in vitro*. For example, Sharp and Banerjee¹³⁾ employed macrophages from conventional mice and rats, *nu/nu* mice or *nu/nu* rats and armadillos. The *M.leprae* inocula were derived from 3 sources (human leproma, *nu/nu* mouse footpad and frozen infected armadillo tissue). Incubation temperature was varied from 31°C to 35°C and *M.leprae*-infected cells were maintained for up to 200 days. Fieldsteel and McIntosh¹⁴⁾ employed a range of rat, mouse and human tissue. The conclusion of these reports is that no significant multiplication of *M.leprae* occurred in any of the cells or tissues.

Previously, we reported that metabolically active *M.leprae* could be maintained in monolayer cultures of mouse peritoneal macrophages and that supplemental IL-10 bolstered *M.leprae* metabolism in the macrophages for as long as 8 weeks. In the cell culture system, temperature is an extremely important factor for growth and 31-

33°C incubation temperature is more permissive than 37°C⁵⁾. In the present study, we further observed that incubation of mouse macrophages infected with *M.leprae* at 35°C was also more growth permissive than at 37°C. We chose 35°C as the incubation temperature, and not 31°C, because the maintenance of the integrity of the macrophage monolayer was better at 35°C than at 31-33°C. Moreover, the monolayer of *M.leprae*-infected human macrophages at 31-33°C could not be maintained for longer than one week. We observed that maintenance of the monolayer was good at 35°C, and *M.leprae* at 35°C was also more growth permissive than those at 37°C in human macrophages (Fig. 4 and 5). Our starting inoculum of *M.leprae* was freshly obtained for each experiment from infected *nu/nu* mice. We also were able to rapidly quantify the metabolic activity of *M.leprae* using the radiorespirometry technique adapted by Franzblau¹¹⁾. This assay is accurate and highly sensitive with the results available in a short duration of 1 wk (compared to 6-12 months when titrated in mouse footpads). Radiorespirometry data correlates well with other *in vitro* systems¹¹⁾ but, more importantly, the data correlated well with "viability" as observed in the mouse footpad system¹²⁾.

Various clinical evidence suggests that *M.leprae* prefer a growth temperature of less than 37°C¹⁾. In animal models, *M.leprae* multiplies in the mouse foot pad where the temperature is lower than the body temperature²⁾. In addition, *Dasypus novemcinctus*, the nine-banded armadillo has a core temperature of ~33°C, which renders it permissive as a host for the leprosy bacillus¹³⁾. Mononuclear phagocytes in virtually every organ of the natural or experimentally infected armadillo become heavily parasitized with propagating *M.leprae*¹⁴⁾. Whether intracellular or extracellular, *M.leprae* clearly prefers temperatures cooler than

normal human body temperature ¹²⁾, and 37°C appeared to be highly detrimental to *M.leprae* viability. The *in vitro* results obtained in the present study confirmed the preference of lower temperature (35°C) by *M.leprae* residing in human macrophages.

In this study, supplemental IL-10, but not TGF- β supported the metabolic activity of *M.leprae* in mouse macrophages for several weeks, similar to the results obtained previously ⁵⁾. In choosing TGF- β and IL-10 as the cytokines that might bolster the intracellular survival of *M.leprae*, we were attempting to down-regulate any innate ability of the normal macrophages to cope with the organism. TGF- β is produced by activated macrophages and other inflammatory cells and has a broad array of modulatory functions on the immune response. TGF- β has been shown to interfere with macrophage antimicrobial mechanisms including the generation of reactive oxygen intermediates ¹⁵⁾ and reactive nitrogen intermediates ¹⁶⁾, and has been shown to enhance the intracellular growth of *M.tuberculosis* in human monocytes ¹⁷⁾. However, in the present studies with mouse macrophages, exogenous TGF- β had no detectable effect on sustaining intracellular *M.leprae* viability, and in fact decreased the viability (Table 1). In contrast, supplementing media with IL-10 clearly affected the long term viability of *M.leprae* in mouse macrophages (Fig. 3). IL-10 is produced by T cells, B cells and macrophages ^{18, 19)}. IL-10 has been shown to be a potent down-regulator of cell-mediated immunity to intracellular pathogens ²⁰⁾. *In vivo*, endogenous IL-10 dampened the cell-mediated immune response to avirulent mycobacterial infection ⁴⁾ and appeared to lead to loss of control of *M.tuberculosis* infection with widespread dissemination ²¹⁾. IL-10 functions in part at the level of the macrophage by attenuating iNOS mRNA expression, iNOS activity

and, by inference, NO production ²²⁾. In human macrophages, however, the viability of *M.leprae* was maintained for 4 weeks in the absence of IL-10 (Fig. 5), suggesting that human cells seem to be better hosts than mouse cells for *M.leprae* survival. Viability of *M.leprae* in M-macrophages seems to be maintained for a longer period (up to one month) than that in GM-macrophages (Fig. 5). One of the reasons for this may be due to the production of IL-10 by M-macrophages ²³⁾, although the mechanism by which IL-10 contributes to the maintenance and growth of *M.leprae* is unclear.

In conclusion, the present study showed that the metabolism, and presumably the viability, of *M.leprae* could be sustained under culture conditions at 35°C, which is also a moderate temperature necessary to maintain the integrity of macrophages.

Acknowledgments

The study was supported partly by a Health Science Research Grant of Emerging and Re-emerging Infectious Diseases, from the Ministry of Health, Labour and Welfare of Japan. We are also grateful to the Japanese Red Cross Society for kindly providing PBMCs from healthy donors.

References

- 1) Brand PW: Temperature variation and leprosy deformity. *Int J Lepr* 27:1-7, 1959.
- 2) Shepard CC: Temperature Optimum of *Mycobacterium leprae* in Mice. *J Bacteriol* 90:1271-1275, 1965.
- 3) Yamamura M, Uyemura KD, Aeans RJ, Weinberg K, Rea TH, Bloom BR, Modlin RL: Defining protective response to pathogens: cytokine profiles in leprosy lesions. *Science* 254:277-282, 1991.

- 4) Sharma S, Bose M: Role of cytokines in immune response to pulmonary tuberculosis. *Asian Pac J Allergy Immunol* 19: 213-219, 2001.
- 5) Fukutomi Y, Matsuoka M, Minagawa F, Toratani S, McCormick G, Krahenbuhl J: Subversion of macrophage anti-microbial function bolsters intracellular survival of *M.leprae*. *Int J Lepr* 72:16-26, 2004.
- 6) Kohsaka K, Mori T, Ito T: Lepromatoid lesion developed in nude mouse inoculated with *Mycobacterium leprae*. *Lepro* 45:177-187, 1976.
- 7) Nakamura M: Elimination of contaminants in a homogenate of nude-mouse footpad experimentally infected with *Mycobacterium leprae*. *Jpn J Lepr* 64:47-50, 1994.
- 8) Shepard CC, McRae DH: A method for counting acid-fast bacteria. *Int J Lepr* 36:78-82, 1968.
- 9) Adams LB, Franzblau S, Taintor R, Hibbs J Jr, Krahenbuhl JL: L- arginine-dependent macrophage effector functions inhibit metabolic activity of *Mycobacterium leprae*. *J Immunol* 147: 1642- 1646, 1991.
- 10) Maeda Y, Mukai T, Spencer J, Makino M: Identification of an immunomodulating agent from *Mycobacterium leprae*. *Infect Immun* 73:2744-2750, 2005.
- 11) Franzblau SG: Oxidation of palmitic acid by *Mycobacterium leprae* in an axenic medium. *J Clin Microbiol* 26:18-2624, 1988.
- 12) Truman RW, Krahenbuhl JL: Viable *Mycobacterium leprae* as a research reagent. *Int J Lepr* 69: 1-12, 2001.
- 13) Kirchheimer W F, Storrs EH: Attempts to establish the armadillo (*Dasypus novemcinctus*) as a model for the study of leprosy. *Int J Lepr* 39:693-702, 1971.
- 14) Fieldsteel AH, McIntosh AH: Attempts to cultivate and determine the maximum period of viability of *M.leprae* in tissue culture. *Int J Lepr* 40: 271-277, 1972.
- 15) Tsunawaki S, Sporn M, Ding A, Nathan C: Deactivation of macrophage by transforming growth factor-beta. *Nature* 334:260-262, 1988.
- 16) Ding A, Nathan CF, Graycar J, Derynck R, Stuehr DJ, Srimal S: Macrophage deactivating factor and transforming growth factors-beta 1, - beta 2 and -beta 3 inhibit induction of macrophage nitrogen oxide synthesis by IFN-gamma. *J Immunol* 145:940-944, 1990.
- 17) Hirsch CS, Yoneda T, Averill L, Ellner JJ, Toosi Z: Enhancement of intracellular growth of *Mycobacterium tuberculosis* in human monocytes by transforming growth factor-1. *J Inf Dis* 170:1229-1237, 1994.
- 18) Fiorentino DF, Zlotnik A, Viera P, Mosmann TR, Howard M, Moore KW, O'Garra A: IL-10 acts on the antigen presenting cell to inhibit cytokine production by Th1 cells. *J Immunol* 146:3444-3451, 1991.
- 19) O'Garra A, Chang R, Go N, Hastings R, Houghton G, Howard M: Ly-1 B (B-1) cells are the main source of B cell - derived IL-10. *Eur J Immunol* 22:711-717, 1992.
- 20) Redpath S, Ghazal P, Gascoigne NR: Hijacking and exploitation of IL-10 by intracellular pathogens. *Trends Microbiol* 9:86-92, 2001.
- 21) Dugas N, Palacios-Calender M, Dugas B, Riveros-Moreno V, Delfraissy J, Kolb J, Moncada S: Regulation by endogenous IL-10 of the expression of nitric oxide synthase induced by ligation of CD23 in human macrophage. *Cytokine* 10:680-689, 1998.
- 22) Huang C, Stevens B, Nielsen E, Slovin P, Fang X, Nelson D, Skimming J: Interleukin-10 inhibition of nitric oxide biosynthesis involves suppression of CAT-2 transcription. *Nitric Oxide* 6:79-84, 2002.
- 23) Makino M, Maeda Y, Fukutomi Y, Mukai T: Contribution of GM-CSF on the enhancement of the T cell-stimulating activity of macrophages. *Microbes Infect* 91:70-77, 2007.

GM-CSF-mediated T-cell activation by macrophages infected with recombinant BCG that secretes major membrane protein-II of *Mycobacterium leprae*

Masahiko Makino, Yumi Maeda, Masanori Kai, Toshiki Tamura & Tetsu Mukai

Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan

Correspondence: Masahiko Makino, Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan. Tel.: +81 42 3918059; fax: +81 42 3918212; e-mail: mmaki@nih.go.jp

Received 14 July 2008; revised 18 October 2008; accepted 23 October 2008.
First published online 8 December 2008.

DOI:10.1111/j.1574-695X.2008.00495.x

Editor: Patrick Brennan

Keywords

macrophage; *Mycobacterium bovis* Bacillus Calmette–Guerin (BCG); *Mycobacterium leprae*; GM-CSF; T-cell activation; interleukin-10.

Introduction

Leprosy is a chronic infectious disease induced by parasitic infection with *Mycobacterium leprae* (Stoner, 1979). Despite the marked reduction in the number of both registered leprosy cases and new cases, a significant number of new cases (254 525 for the year 2007) are still detected each year (World Health Organization, 2008). The emergence of multidrug-resistant *M. leprae* (Kai *et al.*, 2004), although still few in number, and the complexity of leprosy reactions are distressing (Moschella, 2004). These observations indicate the urgent need to develop an efficacious vaccine against leprosy. *Mycobacterium bovis* Bacillus Calmette–Guerin (BCG) has been known to provide partial protection against the development of leprosy (Ponnighaus *et al.*, 1992). However, meta-analyses conducted by Setia *et al.* (2006) demonstrated an overall protective effect of only 26% against leprosy. There seem to be several reasons why BCG is not as effective as previously predicted. One of them may be

Abstract

The potential of *Mycobacterium bovis* Bacillus Calmette–Guerin (BCG) needs to be augmented to efficiently activate CD4⁺ T cells through macrophages. *Mycobacterium leprae*-derived recombinant major membrane protein (MMP)-II induced GM-CSF production from macrophages. A recombinant BCG-SM that secretes MMP-II more efficiently produced GM-CSF and activated interferon (IFN)- γ -producing CD4⁺ T cells than did vector control BCG when infected with macrophages. The T-cell activation by BCG-SM was dependent on the GM-CSF production by macrophages. Interleukin (IL)-10 production by macrophages stimulated with *M. leprae* was inhibited in a GM-CSF-dependent manner when the precursor monocytes were infected with BCG-SM. BCG inducing GM-CSF production was effective in macrophage-mediated T-cell activation partially through IL-10 inhibition.

that the human immune cells most susceptible to BCG infection are macrophages (Grode *et al.*, 2005). On entry into macrophages, mycobacteria inhibit phagosome–lysosome fusion, which results in a less efficient stimulation of interferon (IFN)- γ -producing type 1 CD4⁺ T cells (Ridley & Jopling, 1966; Frehel & Rastogi, 1987). Further, BCG as well as pathogenic mycobacteria can induce the production of an abundant amount of interleukin (IL)-10 from macrophages (Yamamura *et al.*, 1991), which inhibits activation of CD4⁺ T cells (Jonuleit *et al.*, 2001; Granelli-Piperno *et al.*, 2004). Moreover, it has been demonstrated in a murine study that BCG primarily infects macrophages *in vivo*, and the active proliferation of T cells *in vivo* needs the enrolment of dendritic cells (DC). Further, DC are known to be the most professional antigen-presenting cells (APC) in terms of T-cell activation. Thus, the transfer to DC of antigens produced by the processing of intracellular BCG or of proteins secreted from the mycobacteria in macrophages, seems to be important (Winau *et al.*, 2006).

We previously identified major membrane protein (MMP)-II (gene name, *bfrA* or ML2038), which is originally identified as bacterioferritin (Pessolani *et al.*, 1994) and localized in the cell membrane, as one of the dominant antigen of *M. leprae* (Maeda *et al.*, 2005; Makino *et al.*, 2005). Recombinant (r) MMP-II-pulsed DC activate naïve CD4⁺ T cells to produce IFN- γ in an antigen-specific manner, and also stimulate T cells from not only paucibacillary leprosy, a representative clinical leprosy at one pole of the clinical spectrum, but also multibacillary leprosy, a representative leprosy at the opposite pole (Makino *et al.*, 2005). The activation of type 1 CD4⁺ T cells is closely associated with the inhibition of the spread of *M. leprae* *in vivo* as observed in paucibacillary leprosy (Sieling *et al.*, 1999). In this respect, it was interesting to find that T cells from some paucibacillary leprosy patients seemed to be primed with MMP-II antigen *in vivo* (Makino *et al.*, 2005). Therefore, MMP-II was considered to be an immunodominant antigen of *M. leprae*. We constructed an rBCG strain (BCG-SM) that secretes MMP-II of *M. leprae* (Makino *et al.*, 2006). BCG-SM-infected DC stimulated quite efficiently both human naïve CD4⁺ T cells and naïve CD8⁺ T cells *in vitro*, and MMP-II-specific memory T cells were produced in mice inoculated with BCG-SM (Makino *et al.*, 2006).

Macrophages are heterogeneous in various aspects (Randolph *et al.*, 1999), and their differentiation is largely influenced by the cytokine milieu (Nakata *et al.*, 1991; Akagawa, 2002). Previously, we analysed the characteristics of two distinct macrophage subsets: rGM-CSF-mediated macrophages (GM-M ϕ) and rM-CSF-mediated macrophages (M-M ϕ) (Makino *et al.*, 2007). Both macrophages were equally susceptible to mycobacterial infection *in vitro*, but M-M ϕ infected with *M. leprae* did not activate CD4⁺ T cells even after activation using both CD40 ligand and exogenous IFN- γ . Likewise, *Mycobacterium tuberculosis*-infected M-M ϕ failed to stimulate T cells (Verreck *et al.*, 2004). Further, a large amount of IL-10 was produced from M-M ϕ on stimulation with mycobacteria. Therefore, the fact that mycobacteria are highly susceptible to phagocytosis by M-M ϕ and poorly stimulate T cells through M-M ϕ , may be closely associated with the affinity of mycobacteria to macrophages, the induction of a latent infection and, in some cases, the development of disease. Likewise, M-M ϕ is one of the major target immune cells of BCG infection. However, to control the subsequently invading pathogenic mycobacteria, such as *M. leprae*, by producing memory T cells, modified BCG including the newly developed recombinant BCG-SM is required to be able to fully stimulate T cells even if M-M ϕ are the initial target host cells.

In this report, we examined the T-cell-stimulating ability of BCG-SM-infected M-M ϕ , and further assessed the influence of BCG-SM on the IL-10-producing activity of M ϕ upon a challenge with *M. leprae*.

Materials and methods

Preparation of cells and bacteria

Peripheral blood was obtained from healthy purified protein derivative (PPD)-positive individuals with informed consent. In Japan, most healthy individuals are PPD-positive due to a compulsory BCG vaccination for children (0–4 years old). Peripheral blood mononuclear cells (PBMC) were isolated using Ficoll-Paque Plus (Pharmacia, Uppsala, Sweden) and cryopreserved in liquid nitrogen until use as previously described (Makino & Baba, 1997). For the preparation of peripheral monocytes, CD3⁺ T cells were removed from freshly isolated heparinized blood or from cryopreserved PBMC using immunomagnetic beads coated with anti-CD3 monoclonal antibody (mAb) (Dynabeads 450, Dynal, Oslo, Norway). The CD3⁻ PBMC fraction was plated on collagen-coated plates and nonadherent cells were removed by extensive washing. The remaining adherent cells were used as monocytes (Makino & Baba, 1997). Macrophages were generated by culturing monocytes in the presence of 20% foetal calf serum and either rM-CSF (R and D Systems, Abingdon, UK) (M-M ϕ) or rGM-CSF (PeproTech EC Ltd, London, UK) (GM-M ϕ) (Makino *et al.*, 2007). Both GM-M ϕ and M-M ϕ were pulsed with rBCGs on day 3 or 5 of culture, and were used as a stimulator of T cells on day 5 or 7 (Makino *et al.*, 2007). A recombinant BCG that secretes *M. leprae*-derived MMP-II was constructed as described previously (Makino *et al.*, 2006). In brief, a shuttle vector, pMV-261, having a kanamycin resistance gene and origins of replication for *Escherichia coli* and mycobacteria was used to construct pMV-SM (Secreting MMP-II) having the MMP-II cDNA fragment. The BCG substrain Pasteur was cultured *in vitro* using Middlebrook 7H9 broth (BD Biosciences Pharmingen, San Jose, CA) supplemented with 0.05% Tween 80 and 10% albumin–dextrose–catalase (BD). Expression vectors were introduced into BCG by electroporation (Snapper *et al.*, 1988). Transformants were selected on Middlebrook 7H10 agar (BD) plates supplemented with 10% OADC (BD) and 25 $\mu\text{g mL}^{-1}$ kanamycin. Mycobacteria were subsequently grown in Middlebrook 7H9 broth containing 25 $\mu\text{g mL}^{-1}$ of kanamycin. BCG containing pMV-SM as an extrachromosomal plasmid is referred to as BCG-SM, and BCG containing pMV-261 is referred to as BCG-pMV (vector control BCG). *Mycobacterium leprae* (Thai-53) was isolated from the footpads of BALB/*c-nu/nu* mice (McDermott-Lancaster *et al.*, 1987). The isolated bacteria were counted using Shepard's method (McDermott-Lancaster *et al.*, 1987). The multiplicity of infection (MOI) was determined based on the assumption that macrophages were equally susceptible to infection with BCG or *M. leprae* (Hashimoto *et al.*, 2002). A recombinant MMP-II protein was produced as reported previously (Maeda *et al.*, 2005). Briefly,

the MMP-II gene (ML2038) was inserted into the expression plasmid pET28 (Novagen, Madison, WI) and transformed into *E. coli* strain ER2566 (New England BioLabs, Ipswich, MA). The expressed protein was eluted using Whole Gel Eluter (Bio-Rad Laboratories, Hercules, CA). As a control for *M. leprae* antigen, we have purified hsp18 (ML1795) in *E. coli* using the PET expression system. The cytosolic fraction of the parental BCG was obtained as described previously (Maeda *et al.*, 2003).

Antigen-presenting function of rBCG-infected macrophages

The ability of rBCG-infected macrophages to stimulate T cells was assessed using an autologous macrophage-T cell coculture system as previously described (Wakamatsu *et al.*, 1999; Hashimoto *et al.*, 2002). The responder CD4⁺ T cells were purified from freshly thawed PBMC using a CD4-negative isolation kit (Dynabeads 450) (Wakamatsu *et al.*, 1999). The purity of CD4⁺ T cells was > 95% as assessed by FACS. The purified responder cells (1×10^5 per well) were plated in 96-well round-bottom tissue culture plates and macrophages were added to give the indicated macrophage:CD4⁺ T-cell ratio. Supernatants of macrophage-T cell cocultures were collected on day 4. To identify molecules restricting T-cell activation, the following purified mAbs were used: anti-HLA-DR Ab (L243) and anti-CD86 Ab [IT2.2, Becton Dickinson (BD), San Jose, CA]. The concentration of IFN- γ produced by CD4⁺ T cells was quantified using an enzyme assay kit, OptEIA Human enzyme-linked immunosorbent assay (ELISA) Set (BD). In some cases, M-M ϕ were pulsed with BCG-SM in the presence of $10 \mu\text{g mL}^{-1}$ of either normal rat IgG or neutralizing mAb to GM-CSF (rat IgG2a) (BD).

Production of IL-10 and GM-CSF by macrophages

The ability of M-M ϕ to produce IL-10 on stimulation with *M. leprae* was assessed. The monocytes were pretreated with the indicated dose of rBCG and subsequently made to differentiate into M-M ϕ by culturing for 5 days in the presence of rBCG and M-CSF. These macrophages were stimulated with *M. leprae* at the indicated MOI for 24 h. In some cases, monocytes were infected with BCG-SM in the presence of $10 \mu\text{g mL}^{-1}$ of neutralizing mAb to GM-CSF. Also, the ability of M-M ϕ to produce GM-CSF on stimulation with rBCG for 24 h was assessed. The concentration of these cytokines was quantified using OptEIA Human ELISA Set (BD).

Statistical analysis

Student's *t*-test was applied to determine statistical differences.

Results

Effect of rBCG-infected macrophages on T-cell-stimulating activity

We analysed the T-cell-stimulating activity of rBCG-infected GM-M ϕ and M-M ϕ (Fig. 1). GM-M ϕ infected with either BCG-SM or BCG-pMV significantly stimulated CD4⁺




Fig. 1. Production of IFN- γ by CD4⁺ T cells. (a) GM-M ϕ , differentiated by 3 days of culture with rGM-CSF from monocytes, were infected with BCG-pMV (vector control BCG) or BCG-SM (rBCG that secretes MMP-II) at the indicated MOI, and cultured for another 2 days. These GM-M ϕ were used as a stimulator of CD4⁺ T cells (1×10^5 cells per well) at a T cell:GM-M ϕ ratio of 40:1 in a 4-day culture. (b) M-M ϕ , differentiated by 5 days of culture with rM-CSF from monocytes, were infected with BCG-pMV or BCG-SM at the indicated MOI, and cultured for another 2 days. M-M ϕ were then used as a stimulator of CD4⁺ T cells (1×10^5 cells per well) at a T cell:M-M ϕ ratio of 20:1 in a 4-day culture. A representative experiment based on three separate experiments conducted using three separate PPD-positive individuals is shown. Assays were performed in triplicate and the results are expressed as the mean \pm SD. Titres were statistically compared using Student's *t*-test.

T cells. However, a larger amount of IFN- γ was produced by the T cells when GM-M ϕ were infected with BCG-SM (Fig. 1a), although BCG vaccination did not prime for MMP-II-specific T-cell response (not shown). We then analysed the T-cell-stimulating activity of BCG-infected M-M ϕ (Fig. 1b). Again, M-M ϕ infected with BCG-SM induced a higher amount of IFN- γ production by T cells than did BCG-pMV-infected M-M ϕ , although the IFN- γ production was less efficient than that induced by rBCG-infected GM-M ϕ even though higher doses of BCG-infected M-M ϕ were used as a stimulator.

Factors associated with the induction of the T-cell-stimulating activity of M-M ϕ

To define the factors associated with the CD4⁺ T-cell activation by BCG-SM-infected M-M ϕ , we phenotypically analysed M-M ϕ infected with either BCG-pMV or BCG-SM. There was no significant difference between BCG-pMV-infected M-M ϕ and BCG-SM-infected M-M ϕ in the expression of HLA-DR, CD86 or CD40 molecules (not shown). The cytokines produced by M-M ϕ stimulated with rBCGs, including GM-CSF and IL-23, were examined. Both rBCGs induced GM-CSF production, but BCG-SM did so more efficiently than BCG-pMV (Fig. 2). However, IL-23 was not produced by M-M ϕ on stimulation with either BCG-pMV or BCG-SM. We also assessed whether rMMP-II protein can induce GM-CSF production in macrophages. Whereas *M. leprae*-derived cytosolic protein (not shown), other mycobacterial proteins such as BCG-derived cytosolic protein (5–10 $\mu\text{g mL}^{-1}$), control recombinant *M. leprae* antigen (hsp18), and lipopolysaccharide (amount present with rMMP-II protein) did not stimulate M ϕ , MMP-II induced GM-CSF production in a concentration-dependent manner (Fig. 3). rMMP-II also efficiently induced the production of other cytokines including tumour necrosis factor (TNF) α and IL-12p40 from M ϕ (not shown).

We examined the influence of surface antigens on M-M ϕ . The T-cell-stimulating activity of BCG-SM-infected M-M ϕ was significantly inhibited when the infected M-M ϕ were pretreated with the mAb to HLA-DR or CD86 antigens, whereas the control IgG did not affect IFN- γ production by T cells (Fig. 4a). However, IFN- γ production was partially inhibited when BCG-SM-infected M-M ϕ were treated with the mAb to MMP-II (not shown). Next, we examined the effect on T-cell activation of GM-CSF produced by M-M ϕ stimulated with BCG-SM (Fig. 4b). When M-M ϕ were infected with BCG-SM in the presence of the neutralizing mAb to GM-CSF, IFN- γ production by CD4⁺ T cells was significantly inhibited. The T-cell-stimulating activity of BCG-SM-infected M-M ϕ was not affected by normal rat IgG.




Fig. 2. Production of GM-CSF by M-M ϕ . M-M ϕ differentiated by 5 days of culture with rM-CSF from monocytes, were stimulated with BCG-pMV or BCG-SM for 24 h at the indicated MOI. A representative experiment based on three separate experiments conducted using three separate PPD-positive individuals is shown. Assays were performed in triplicate and the results are expressed as the mean \pm SD. Titres were statistically compared using Student's *t*-test.




Fig. 3. GM-CSF production by M-M ϕ . M-M ϕ obtained after 5 days of culture with rM-CSF were stimulated for 24 h with rMMP-II, the BCG-derived cytosolic fraction or *Mycobacterium leprae*-derived hsp18 antigen [dose of antigen in (a) 5 $\mu\text{g mL}^{-1}$ and that in (b) 10 $\mu\text{g mL}^{-1}$]. Lipopolysaccharide, assumed to be present with rMMP-II protein (660 ng mg^{-1} MMP-II protein), was used as a negative control (lipopolysaccharide in (a) 3.3 ng mL^{-1} and that in (b) 6.6 ng mL^{-1}). A representative experiment based on three separate experiments conducted using three separate PPD-positive individuals is shown. Assays were performed in triplicate and the results are expressed as the mean \pm SD. Titres were statistically compared using Student's *t*-test.

Effect of infection of monocytes with BCG on IL-10 production by M-M ϕ

Macrophages are one of the cells most sensitive to *M. leprae* infection and M-M ϕ produce abundant IL-10 when infected with the bacteria (Makino et al., 2007). As precursor




Fig. 4. (a) Inhibition of IFN- γ production by CD4⁺ T cells by pretreatment of BCG-SM-infected M-M ϕ with mAb to HLA-DR or CD86. M-M ϕ differentiated from monocytes using rM-CSF were infected with BCG-SM at an MOI of 0.25 on day 5 of culture and cultured for another 2 days. The BCG-SM-infected M-M ϕ were treated with mAb to HLA-DR or CD86, or isotype-matched control IgG (10 $\mu\text{g mL}^{-1}$), and used as a stimulator of CD4⁺ T cells, at a T cell : M-M ϕ ratio of 20 : 1 and cultured for another 4 days. The optimal concentration of mAb was determined in advance. Non-pretreated BCG-SM-infected M-M ϕ induced the production of 220.8 pg mL^{-1} of IFN- γ by CD4⁺ T cells. This titre was taken as 0% inhibition. (b) Inhibition of IFN- γ production by CD4⁺ T cells by neutralizing GM-CSF produced from BCG-SM-infected M-M ϕ . M-M ϕ , differentiated from monocytes by culturing for 5 days with rM-CSF, were infected with BCG-SM (MOI 0.25) in the presence of neutralizing mAb to GM-CSF or isotype-matched control IgG (10 $\mu\text{g mL}^{-1}$). These M-M ϕ were used as a stimulator of CD4⁺ T cells as in (a). The optimal concentration of mAb was determined in advance. M-M ϕ infected with BCG-SM in the absence of any Ab induced the production of 168.3 pg mL^{-1} of IFN- γ by CD4⁺ T cells. This titre was taken as 0% inhibition. A representative experiment based on three separate experiments conducted using three separate PPD-positive individuals is shown. Assays were performed in triplicate and the results are expressed as the mean \pm SD. Titres were statistically compared using Student's *t*-test.




Fig. 5. (a) Production of IL-10 by M-M ϕ . Monocytes were infected with the indicated dose of BCG-pMV or BCG-SM, and subsequently differentiated into M-M ϕ by culturing for 5 days in the presence of M-CSF and rBCGs. These rBCG-preinfected M-M ϕ were stimulated with *Mycobacterium leprae* at the indicated MOI for 24 h. (b) Recovery of IL-10 production by M-M ϕ . Monocytes were infected with BCG-SM (MOI 0.063) in the presence of neutralizing GM-CSF mAb or isotype-matched control IgG and were subsequently differentiated into M-M ϕ by culturing for 5 days. These BCG-SM-preinfected M-M ϕ were stimulated with *M. leprae* at the indicated MOI for 24 h. The optimal concentration of mAb was determined in advance. A representative experiment based on three separate experiments conducted using three separate PPD-positive individuals is shown. Assays were performed in triplicate and the results are expressed as the mean \pm SD. Titres were statistically compared using Student's *t*-test.

monocytes also produced GM-CSF on stimulation with BCG in a BCG-SM-predominant fashion (not shown), we examined the effect of infection with rBCGs in monocytes on IL-10 production by M-M ϕ challenged with *M. leprae* (Fig. 5). M-M ϕ differentiated from monocytes untreated with any bacteria produced > 400 pg mL^{-1} of IL-10 on

stimulation with *M. leprae*; however, the production of cytokine by M-M ϕ pretreated with rBCGs was significantly inhibited (Fig. 5a). The inhibition was more significant when BCG-SM was used as a stimulator of monocytes, and IL-10 production by M-M ϕ was almost completely inhibited. The inhibition was dependent on the dose of BCGs

used for pretreatment. In addition, pretreatment of monocytes with BCG-SM inhibited the IL-10 production induced even by lipopolysaccharide (not shown).

Furthermore, M-M ϕ differentiated from monocytes infected with BCG-SM in the presence of normal IgG did not produce IL-10 on stimulation with *M. leprae* (Fig. 5b). However, a significant level of IL-10 was produced when monocytes were infected with BCG-SM in the presence of the neutralizing mAb to GM-CSF. These results indicate that endogenously produced GM-CSF can inhibit IL-10 production.

Discussion

The host defence against intracellular parasitic pathogens such as *M. leprae* is closely associated with the activation of IFN- γ -producing type 1 T cells (Hashimoto *et al.*, 2002). In fact, in patients with paucibacillary leprosy, the activation of CD4⁺ T cells results in inhibition of the intracellular multiplication and intercellular spread of *M. leprae* (Sieling *et al.*, 1999). The T-cell activation largely depends on the extent of the activation of APC, in which DC play an extremely important role, as DC are the most powerful T-cell stimulators among the APC (Hashimoto *et al.*, 2002). However, if T cells are not efficiently activated due to poor participation of DC, *M. leprae* may be predominantly retained in macrophages. In fact, multibacillary leprosy patients retain numerous *M. leprae* in their macrophages which, in some cases, allow the multiplication and intercellular spread of the bacteria (Ridley & Jopling, 1966).

The tissue resident macrophages, represented by GM-M ϕ and M-M ϕ , are heterogeneous in terms of function (Nakata *et al.*, 1991; Randolph *et al.*, 1999; Akagawa, 2002), despite being similarly susceptible to mycobacterial infection (Makino *et al.*, 2007). GM-M ϕ infected with *M. tuberculosis* or *M. leprae* significantly stimulated CD4⁺ T cells, whereas M-M ϕ failed to stimulate CD4⁺ T cells (Verreck *et al.*, 2004; Makino *et al.*, 2007). In this study, we found that, similar to those pathogenic mycobacteria, vector control BCG (BCG-pMV)-infected GM-M ϕ significantly stimulated CD4⁺ T cells, whereas the BCG-pMV-infected M-M ϕ were less efficient in stimulating these cells. These results indicate the possibility that parental BCG may long reside in M-M ϕ and stimulate T cells inadequately, like the *M. leprae*-infected resident macrophages in multibacillary leprosy. In contrast to BCG-pMV, rBCG that secretes MMP-II (BCG-SM) has the ability to enlist not only GM-M ϕ , but also M-M ϕ , for T-cell activation. Further, the production of IFN- γ by CD4⁺ T cells stimulated with BCG-SM-infected M-M ϕ was significantly inhibited by pretreatment of the M-M ϕ with the mAb to HLA-DR or CD86 antigens. In addition, the pretreatment of M-M ϕ infected with both BCG-SM and BCG-pMV effectively inhibited CD4⁺ T-cell activation (not

shown). Therefore, the BCG-SM-infected M-M ϕ seemed to stimulate CD4⁺ T cells in an antigen-specific manner. Furthermore, there was a striking difference between BCG-pMV and BCG-SM in the induction of GM-CSF production. Not only from M-M ϕ , but also from GM-M ϕ , BCG-SM more efficiently induced GM-CSF production than BCG-pMV, and, further, rMMP-II protein, though less efficient, induced significant GM-CSF production. Previously, we reported that rMMP-II is highly immunogenic and induces production of various cytokines, including IL-12 and TNF- α , from APCs such as macrophages and DC (Maeda *et al.*, 2005). These findings indicate that the enhanced production of GM-CSF on stimulation by BCG-SM was at least partially associated with the secretion of MMP-II from BCG-SM.

As the activation of T cells by BCG-SM-infected M-M ϕ was largely inhibited when endogenously produced GM-CSF was neutralized by the mAb to GM-CSF, the endogenously produced GM-CSF may be closely associated with the enhanced T-cell activation by BCG-SM. Although we could not identify the most relevant antigen for T-cell activation, GM-CSF may change the activation status of macrophages or may at least partially transform the BCG-SM-infected M-M ϕ to GM-M ϕ (Makino *et al.*, 2007). Therefore, BCG-SM seems to be a unique rBCG capable of producing GM-CSF and utilizing M-M ϕ for T-cell stimulation.

Another important characteristic of mycobacteria which contributes to the inhibition of T-cell activation is the abundant production of IL-10 by M-M ϕ (Jonuleit *et al.*, 2001; Mochida-Nishimura *et al.*, 2001; Granelli-Piperno *et al.*, 2004). The major purpose of a vaccination is the production of memory T cells which can rapidly respond to subsequently invading pathogenic mycobacteria. However, IL-10 inhibits the re-activation of memory T cells *in vitro*. We found that the ability of BCG-SM to induce production of GM-CSF is useful to inhibit IL-10 production. Monocytes were quite sensitive in the production of GM-CSF, and both BCG-pMV and BCG-SM induced cytokine production by monocytes, although BCG-SM predominated at lower doses (not shown). Thus, even BCG-pMV inhibited IL-10 production at higher doses. However, macrophages differentiated from monocytes which were infected with a small dose of BCG-SM completely inhibited IL-10 production upon subsequent challenge with *M. leprae*, and the inhibitory activity was at least partially cancelled out by the neutralization of endogenously produced GM-CSF. Further, heat-killed BCG-SM, which does not secrete MMP-II (Makino *et al.*, 2006), did not inhibit IL-10 production. These observations indicate that macrophages treated with GM-CSF endogenously diminished the ability to produce IL-10 upon stimulation with *M. leprae*. Previously, we observed that addition of GM-CSF exogenously blocked the ability to produce IL-10 (Makino *et al.*, 2007), which agrees with the