

## ANTI-HCV NEUTRALIZING ANTIBODIES IN SERA OF HCV-INFECTED PATIENTS

Patients chronically infected with HCV-1b, who were treated with pegylated interferon  $\alpha$ -2b (1.5  $\mu$ g per kilogram body weight, once weekly, subcutaneously) and ribavirin (600-800 mg daily, per os), were described previously (8, 9). Sera were collected and stored at  $-80^{\circ}\text{C}$  until ready for use. The sera were inactivated at  $56^{\circ}\text{C}$  for 30 min before being used for the virus neutralization test, as describe below.

### HCV focus reduction neutralization assay

An HCV focus reduction neutralization assay was performed, as described elsewhere (Sasayama et al., in preparation). Briefly, one-tenth volume of serum obtained from uninfected healthy human, which had been inactivated at  $56^{\circ}\text{C}$  for 30 min, was mixed with HCV solutions at  $37^{\circ}\text{C}$  for 1 h to avoid nonspecific inhibition. Serial 3-fold dilutions of each serum sample were mixed with pre-treated HCV solution containing  $10^4$  FFU. After incubation at  $37^{\circ}\text{C}$  for 1 h, the mixtures were inoculated to naïve Huh-7.5 cells ( $2 \times 10^5$  cells per well in 24-well plates) and incubated in a 5%  $\text{CO}_2$  incubator. After 3 h of virus adsorption, the inocula were removed and fresh complete DMEM were added to the cells. At 24 h postinfection, cells were washed with PBS, fixed with 100% methanol, blocked with 5% goat serum in PBS and subjected to immunofluorescence analysis using mouse monoclonal antibody against HCV core antigen (clone 2H9; a kind gift from Dr. T. Wakita, Department of Virology II, National Institute of Infectious Diseases, Tokyo, Japan) (15) and Alexa Fluor 488-conjugated goat anti-mouse IgG (H+L) (Invitrogen). The immunostained cells were washed with PBS, counterstained with Hoechst 33342 solution (Invitrogen) at room temperature for 5 min, mounted on glass slides, and observed under a fluorescence microscope (BZ-9000; Keyence). The number of HCV-infected foci in each well was counted by a software BZ-H1C (Keyence). The dilution that neutralized 50% of the initial virus infectivity was calculated by curvilinear regression analysis. Each neutralization titer was determined as the logarithmic value of the reciprocal antibody dilution that reduced 50% of viral foci to the total number. Titers were expressed as logarithmic values and means  $\pm$  standard deviation (SD) were calculated.

### Statistical analysis.

The statistical significance of comparisons between the two groups of patients was determined using Student's *t*-test. In all tests, a *p*-value lower than 0.05 was considered statistically significant.

## RESULTS

### Virological response of the patients underwent treatment with PEG-IFN/RBV

In this study, 65 patients infected with HCV-1b were enrolled. During and after PEG-IFN/RBV therapy, their clinical responses to the treatment were evaluated (Table 1.). Of all the 65 patients, 34 (52%) patients achieved early virological response (EVR) by week 12, with the remaining 31 patients (48%) being Non-EVR. Twenty-seven patients (42%) achieved sustained virological response (SVR), with the remaining 38 patients (58%) being Non-SVR, which is divided into 2 categories, complete non-response (CNR) and relapse as described previously (14). CNR was observed with 16 (25%) of the 65 patients and relapse was observed with 22 patients (34%).

Table 1. Proportion of various virological responses of patients treated with PEG-IFN/RBV.

Virological response	Proportion
EVR	52% (34 / 65)*
Non-EVR	48% (31 / 65)
SVR	42% (27 / 65)
Non-SVR	58% (38 / 65)
CNR	25% (16 / 65)
Relapse	34% (22 / 65)

\* Number of patients/Number of total.

### Pre-treatment NAb titers in the serum and PEG-IFN/RBV treatment outcome

To study the possible impact of NAb on PEG-IFN/RBV treatment outcome of patients infected with HCV-1b, we measured NAb titers in patients sera using the J6/JFH1 strain of HCV. The mean dilution of antibodies required for 50% neutralization (NAb<sub>50</sub> titer) of HCV J6/JFH1 in patients who achieved an EVR (2.28±0.36) was significantly higher than that of non-EVR (1.97±0.38) ( $p < 0.01$ ) (Fig. 1). Also, the NAb<sub>50</sub> titers in patients who achieved an SVR (2.27±0.40) was significantly higher than that of non-SVR (2.04±0.37) ( $p < 0.05$ ) or CNR (1.91±0.43) ( $p < 0.05$ ) (Fig. 1). These data suggest that NAb<sub>50</sub> titers in the pre-treatment sera are associated with treatment outcome.





Fig. 1. NAb<sub>50</sub> titers in the sera of HCV-infected patients. NAb<sub>50</sub> titers in the pre-treatment sera of HCV-1b-infected patients were compared between RVR and Non-RVR (A), EVR and Non-EVR (B), SVR and Non-SVR (C), SVR and CNR (C), and SVR and Relapsers (C). \*,  $P < 0.05$ ; †, 0.01.

**Fate and change ratios of NAb titers before and after disappearance of HCV RNA in the serum of patients treated with PEG-IFN/RBV**

In order to investigate as to whether or not the NAb titers decrease after disappearance of HCV RNA in the serum of patients treated with PEG-IFN/RBV, we measured NAb titers in

patients' sera collected at both pre-treatment (before disappearance) and after disappearance of HCV RNA in the serum. Rather unexpectedly, NAb titers did not decrease when measured even one year after disappearance of HCV RNA. On the other hand, when change ratios of NAb titers before and after disappearance of HCV RNA were compared between patients with different treatment outcomes, we noticed that the change ratio of NAb<sub>50</sub> titers of patients who achieved an EVR was significantly lower than that of Non-EVR (Fig. 2).

Fig. 2. ratios before and after



Change of NAb and

## ANTI-HCV NEUTRALIZING ANTIBODIES IN SERA OF HCV-INFECTED PATIENTS

disappearance of HCV RNA in the serum. Change ratios (post-treatment/pre-treatment) in the NAb<sub>50</sub> titers before and after disappearance of HCV RNA in the sera of patients who were treated with PEG-IFN/RBV were compared between RVR and Non-RVR (A), EVR and Non-EVR (B), and ETR and Non-ETR (C). \*,  $P < 0.05$ .

### DISCUSSION

We previously reported that anti-NS5A antibodies were more frequently detected in sera of patients who achieved EVR compared to Non-EVR (8). In this study, we demonstrated that NAb<sub>50</sub> titers in the pre-treatment patients' sera were associated with the good responses (EVR and SVR) to PEG-IFN/RBV combination therapy (Fig. 1). Consistent with our observations, it was reported that NAb titers to HCV-LPs were higher in patients who achieved an SVR with IFN/RBV therapy than in relapsers and non-responders (2). The better humoral responses, such as NAb and anti-NS5A antibodies, might be associated with better cell-mediated immune responses, which involve CD4<sup>+</sup> and CD8<sup>+</sup> T cells. It is well known that peripheral and intrahepatic CD8<sup>+</sup> T cell responses, which play an important role in the control of and recovery from HCV infection, are also important in determining an SVR in response to PEG-IFN/RBV treatment (6). Thus our present results imply the possibility that individuals who can maintain harmonized good immune responses are able to achieve good responses to PEG-IFN/RBV therapy, such as EVR and SVR.

We also found that NAb<sub>50</sub> titers did not decrease significantly when measured even one year after disappearance of HCV RNA in the serum (Fig. 2).

Another important finding in this study is that the NAb in patients infected with HCV-1b significantly cross-reacts to HCV-2a; we observed that average NAb titers of HCV-2a-infected patients were ca. 3 times higher than those of HCV-1b-infected patients when measured with the same experimental system using the J6/JFH1 strain of HCV-2a ( $439 \pm 2.72$  vs.  $139 \pm 2.48$ ;  $P < 0.0001$ ). This information would be helpful when considering immunological prophylaxis against HCV infection, either active or passive immunizations using vaccines and NAb.

In conclusion, NAb<sub>50</sub> titers were significantly higher in sera of patients who achieved EVR and SVR than those of Non-EVR and Non-SVR, respectively. Also, NAb<sub>50</sub> titers declined only slightly during the course of one year after disappearance of HCV RNA in the sera.

### ACKNOWLEDGEMENTS

The authors are grateful to Dr C. M. Rice (Center for the Study of Hepatitis C, the Rockefeller University, New York, NY, USA) for providing pFL-J6/JFH1 and Huh-7.5 cells. This work was supported in part by grants-in-aid for Scientific Research from the Ministry of Health, Labour and Welfare, Japan. This study was also carried out as part of the Program for Promoting Science and Technology Diplomacy, International Joint Research Collaborating with Official Development Assistance (ODA) supported by Japan Science and Technology Agency (JST) and Japan International Cooperation Agency (JICA), and part of the Program of Founding Research Centers for Emerging and Reemerging Infectious Diseases, Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. This study was also part of the Global Center of Excellence (COE) Program at Kobe University Graduate School of Medicine.

### REFERENCES

1. **Bartosch, B., Bukh, J., Meunier, J. C., Granier, C., Engle, R. E., Blackwelder, W. C., Emerson, S. U., Cosset, F. L., and Purcell, R. H.** 2003. In vitro assay for neutralizing antibody to hepatitis C virus: evidence for broadly conserved neutralization epitopes. *Proc Natl Acad Sci U S A.* **100**: 14199-204.
2. **Baumert, T. F., Wellnitz, S., Aono, S., Sato, J., Herion, D., Tilman, G. J., Pape, G. R., Lau, J. Y., Hoofnagle, J. H., Blum, H. E., and Liang, T. J.** 2000. Antibodies against hepatitis C virus-like particles and viral clearance in acute and chronic hepatitis C. *Hepatology* **32**: 610-7.
3. **Blight, K. J., MaKeating, J. A., and Rice, C. M.** 2002. Highly permissive cell lines for subgenomic and genomic hepatitis C virus RNA replication. *J. Virol.* **76**: 13001-13014.
4. **Bowen, D. G., and Walker, C. M.** 2005. Adaptive immune responses in acute and chronic hepatitis C virus infection. *Nature* **436**: 946-52.
5. **Bungyoku, Y., Shoji, I., Makine, T., Adachi, T., Hayashida, K., Nagano-Fujii, M., Ide, YH., Deng, L., and Hotta, H.** 2009. Efficient production of infectious hepatitis C virus with adaptive mutations in cultured hepatoma cells. *J Gen Virol.* **90**: 1681-91.
6. **Caetano, J., Martinho, A., Paiva, A., Pais, B., Valente, C., and Luxo, C.** 2008. Differences in hepatitis C virus (HCV)-specific CD8 T-cell phenotype during pegylated alpha interferon and ribavirin treatment are related to response to antiviral therapy in patients chronically infected with HCV. *J Virol.* **82**: 7567-77.
7. **Cramp, M. E., Rossol, S., Chokshi, S., Carucci, P., Williams, R., and Naoumov, N. V.** 2000. Hepatitis

- C virus-specific T-cell reactivity during interferon and ribavirin treatment in chronic hepatitis C. *Gastroenterology*. **118**: 346-55.
8. El-Shamy, A., Sasayama, M., Nagano-Fujii, M., Sasase, N., Imoto, S., Kim, S. R., and Hotta, H. 2007. Prediction of efficient virological response to pegylated interferon/ribavirin combination therapy by NS5A sequences of hepatitis C virus and anti-NS5A antibodies in pre-treatment sera. *Microbiol Immunol*. **51**: 471-82.
  9. El-Shamy, A., Nagano-Fujii, M., Sasase, N., Imoto, S., Kim, S. R., and Hotta, H. 2008. Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *Hepatology*. **48**: 38-47.
  10. Lavillette, D., Morice, Y., Germanidis, G., Donot, P., Soulier, A., Pagkalos, E., Sakellariou, G., Intrator, L., Bartosch, B., Pawlotsky, JM., and Cosset, FL. 2005. Human serum facilitates hepatitis C virus infection, and neutralizing responses inversely correlate with viral replication kinetics at the acute phase of hepatitis C virus infection. *J Virol*. **79**: 6023-34.
  11. Lindenbach, B. D., Evans, M. J., Syder, A. J., Wölk, B., Tellinghuisen, T. L., Liu, C. C., Maruyama, T., Hynes, R. O., Burton, D. R., McKeating, J. A., and Rice, C. M. 2005. Complete replication of hepatitis C virus in cell culture. *Science*. **309**: 623-626.
  12. Moradpour, D., Francois P. F., and Rice, C. M. 2007. Replication of hepatitis C virus. *Nat. Rev. Microbiol*. **5**: 453-63.
  13. Youn, J. W., Park, S. H., Lavillette, D., Cosset, F. L., Yang, S. H., Lee, C. G., Jin, H. T., Kim, C. M., Shata, M. T., Lee, D. H., Pfahler, W., Prince, A. M., and Sung, Y. C. 2005. Sustained E2 antibody response correlates with reduced peak viremia after hepatitis C virus infection in the chimpanzee. *Hepatology*. **42**: 1429-36.
  14. Takaki, A., Wiese, M., Maertens, G., Depla, E., Seifert, U., Liebetrau, A., Miller, J. L., Manns, M. P., and Rehermann, B. 2000. Cellular immune responses persist and humoral responses decrease two decades after recovery from a single-source outbreak of hepatitis C. *Nat Med*. **6**: 578-82.
  15. Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Kräusslich, H. G., Mizokami, M., Bartenschlager, R., and Liang, T. J. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med*. **2005 11**: 791-6.
  16. Youn, J. W., Park, S. H., Lavillette, D., Cosset, F. L., Yang, S. H., Lee, C. G., Jin, H. T., Kim, C. M., Shata, M. T., Lee, D. H., Pfahler, W., Prince, A. M., and Sung, Y. C. 2005. Sustained E2 antibody response correlates with reduced peak viremia after hepatitis C virus infection in the chimpanzee. *Hepatology*. **42**: 1429-36.



