

Polymorphisms of DNA damage response genes in radiation-related and sporadic papillary thyroid carcinoma

Natallia M Akulevich¹, Vladimir A Saenko², Tatiana I Rogounovitch¹,
Valentina M Drozd³, Eugeny F Lushnikov⁴, Victor K Ivanov⁴,
Norisato Mitsutake¹, Ryo Kominami⁵ and Shunichi Yamashita^{1,2}

Departments of ¹Molecular Medicine and ²International Health and Radiation Research, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8523, Japan

³Department of Thyroid Disease Research, Belarusian Medical Academy for Postgraduate Education, Minsk 220013, Republic of Belarus

⁴Medical Radiological Research Center, Obninsk 249036, Russian Federation

⁵Department of Molecular Genetics, Graduate School of Medical and Dental Sciences, Niigata University, Niigata 951-8122, Japan

(Correspondence should be addressed to V A Saenko; Email: saenko@net.nagasaki-u.ac.jp)

Abstract

Papillary thyroid carcinoma (PTC) etiologically occurs as a radiation-induced or sporadic malignancy. Genetic factors contributing to the susceptibility to either form remain unknown. In this retrospective case-control study, we evaluated possible associations between single-nucleotide polymorphisms (SNPs) in the candidate DNA damage response genes (*ATM*, *XRCC1*, *TP53*, *XRCC3*, *MTF1*) and risk of radiation-induced and sporadic PTC. A total of 255 PTC cases (123 Chernobyl radiation-induced and 132 sporadic, all in Caucasians) and 596 healthy controls (198 residents of Chernobyl areas and 398 subjects without history of radiation exposure, all Caucasians) were genotyped. The risk of PTC and SNPs interactions with radiation exposure were assessed by logistic regressions. The *ATM*G5557A and *XRCC1* Arg399Gln polymorphisms, regardless of radiation exposure, associated with a decreased risk of PTC according to the multiplicative and dominant models of inheritance (odds ratio (OR)=0.69, 95% confidence interval (CI) 0.45–0.86 and OR=0.70, 95% CI 0.59–0.93 respectively). The *ATM* IVS22-77 T>C and *TP53* Arg72Pro SNPs interacted with radiation ($P=0.04$ and $P=0.01$ respectively). *ATM* IVS22-77 associated with the increased risk of sporadic PTC (OR=1.84, 95% CI 1.10–3.24) whereas *TP53* Arg72Pro correlated with the higher risk of radiogenic PTC (OR=1.80, 95% CI 1.06–2.36). In the analyses of *ATM/TP53* (rs1801516/rs664677/rs609429/rs1042522) combinations, the GG/TC/CG/GC genotype strongly associated with radiation-induced PTC (OR=2.10, 95% CI 1.17–3.78). The GG/CC/GG/GG genotype displayed a significantly increased risk for sporadic PTC (OR=3.32, 95% CI 1.57–6.99). The results indicate that polymorphisms of DNA damage response genes may be potential risk modifiers of ionizing radiation-induced or sporadic PTCs.

Endocrine-Related Cancer (2009) 16 491–503

Introduction

Thyroid cancer accounts for more than 90% of all endocrine malignancies. The incidence of thyroid cancer in the world is increasing during the past three decades, mainly due to the papillary thyroid carcinoma (PTC) which is the predominant type of malignant thyroid tumors (Davies & Welch 2006).

Most thyroid cancer patients do not have the history of radiation exposure, yet ionizing radiation (IR) is a recognized etiological factor of the disease. An

increased risk of thyroid cancer has been documented after external irradiation (Ron *et al.* 1995) and after environmental exposure to ¹³¹I, such as after the Chernobyl fallouts in Belarus, Ukraine, and Russia (Bennett *et al.* 2006).

Although radiation thyroid doses in Chernobyl PTC cases are generally greater than in controls in epidemiological studies (Cardis *et al.* 2005, Jacob *et al.* 2006, Likhtarev *et al.* 2006), thus confirming radiation to be a risk factor for thyroid cancer, those in

controls are non-zero. Furthermore, there were some 14 million residents in the contaminated territories at the time of exposure (Bennett *et al.* 2006). Conceivably, at least some of them might have accumulated thyroid doses comparable with doses in diseased individuals. However, thyroid cancer developed only in a small fraction of irradiated population.

Among the variety of DNA damage types induced by radiation, double-strand DNA breaks are considered to be the most significant for chromosomal aberrations, mutagenesis, genetic instability, and carcinogenesis (Khanna & Jackson 2001). PTC is one of the rare human cancers of epithelial origin in whose oncogenesis gene rearrangements play a noticeable role. Several variants of rearrangements are described in PTC, with RET/PTC occurring most frequently (Nikiforov *et al.* 1997, Rabes *et al.* 2000).

While in the exposed individuals DNA damage could be attributed to IR, the origination of genetic alterations in sporadic cancers remains obscure. Nevertheless, the spectrum of oncogenic changes in radiation-related and sporadic PTCs is largely common. Such similarities imply the resemblance of molecular reactions on DNA damage in exposed and non-exposed thyrocytes. These reactions involve first of all DNA damage response factors, including DNA repair and checkpoint complexes.

The vast majority of Chernobyl thyroid malignancies were PTCs which displayed wide variations in clinical course, from highly aggressive tumors developing after the shorter latency to more indolent carcinomas with the longer latent period (Williams 2006). The randomness and multiplicity of forms of genetic alterations caused by IR can only partly explain these differences in the individual reactions on exposure as well as why cancer develops only in some of the exposed individuals.

It is attractive to hypothesize that inherited variability in the genes directly or indirectly involved in the maintenance of genome stability in response to environmental carcinogens such as IR or chemicals that may play a role in susceptibility for radiation-related or sporadic PTC or may be a marker of it. In this work, we tested the relation of genetic variants of some of such genes, namely *ATM*, *TP53*, *XRCC1*, *XRCC3*, and *MTF1* to PTC of different etiology.

The ataxia-telangiectasia mutated (*ATM*) gene plays a key role in the sensing and repair of DNA double-strand breaks. Activation of the ATM protein kinase by IR results in the subsequent initiation of several molecular pathways of DNA damage repair (Shiloh 2003). One of the ATM targets is the p53 pathway.

Overexpression of *TP53* arrests the cell cycle and affects DNA repair and apoptosis.

The *ATM* and *TP53* genes play a significant role especially in the tumors that are induced by IR. A number of single-nucleotide polymorphisms (SNPs) in the *ATM* and *TP53* genes studied in populations of different ethnicities have been reported to associate with the risk of different radiogenic tumors (Hu *et al.* 2002, Angele *et al.* 2003, Thorstenson *et al.* 2003, Malmer *et al.* 2007). By contrast, studies of post-Chernobyl pediatric thyroid cancers demonstrated a low mutation and polymorphism rate in the *TP53* gene (Nikiforov *et al.* 1996, Hillebrandt *et al.* 1997). It, however, should be mentioned that after exposure to radiation p53 facilitates DNA repair in normal thyrocytes *in vitro* (Yang *et al.* 1997).

The base excision repair (BER) and homologous recombination repair (HRR) pathways are particularly important for genomic integrity restoration (Hoeijmakers 2001). The product of the X-ray repair cross complementing 1 (*XRCC1*) gene acts as a scaffold and a modulator of different enzymes involved in BER. The *XRCC1* Arg399Gln and Arg280His variants have been extensively investigated for their function and association with cancer risk; however, the results remain contradictory rather than conclusive (Hu *et al.* 2005). The *XRCC3* gene is a member of the *Rad51* DNA-repair gene family. Its product is a factor of the HRR. The *XRCC3* Thr241Met polymorphism has been controversially associated with different human malignancies (Han *et al.* 2006). Sturgis *et al.* (2005) reported 241Met allele association with the risk of differentiated thyroid cancer.

The metal-responsive transcription factor 1 (*MTF1*) gene has been implicated in tumor initiation and progression to malignant growth. MTF1 protein interacts with metallothioneins that are able to suppress cellular stresses generated by IR and other agents (Tamura *et al.* 2005). Polymorphism in murine *Mtf1* gene has been found to associate with the susceptibility to experimental γ -ray-induced thymic lymphomas. This observation points at possible involvement of human *MTF1* polymorphisms in the modulation of radiation-induced malignancies (Tamura *et al.* 2005).

To date no polymorphisms of the *ATM*, *XRCC1*, and *MTF1* genes have been studied neither in human sporadic or radiation-induced PTCs. Data on the *TP53* and *XRCC3* polymorphisms associations are quite limited (Hillebrandt *et al.* 1997, Boltze *et al.* 2002, Granja *et al.* 2004, Sturgis *et al.* 2005, Rogounovitch *et al.* 2006). Therefore, in this study, we addressed the relation of SNPs in aforementioned DNA damage response genes to the risk of PTCs of different etiology.

Materials and methods

Study population

A total of 255 histologically verified PTC cases and 596 healthy controls, all Caucasians, were included in the study. Among the patients, 123 individuals with PTC (24 males and 99 females) lived in the areas of the Russian Federation (38 patients) and Belarus (85 patients) contaminated with radionuclides from Chernobyl fallouts. At the time of the Chernobyl accident, these subjects were younger than 18 years old (mean age at exposure \pm s.d., 9.8 ± 5.1 years old; 1–18 years old, range). The mean age at diagnosis was 24.4 ± 4.9 years old, range 19–37 years old (IR-induced PTCs). Information about individual radiation thyroid doses was available for PTC cases from Russia as reconstructed in previous studies (Davis *et al.* 2004, Stepanenko *et al.* 2004). The doses varied from 43 to 2640 mGy. Radiation thyroid doses for PTC patients and controls from Belarus evaluated in dosimetric investigations at the places of residence ranged 21–1500 mGy (Bouville *et al.* 2007). Among the controls, 198 individuals (65 males and 133 females, mean age at sampling 22.2 ± 3.2 years old; 19–35 years old, range) were residents of the Chernobyl areas (60 from the Russian Federation and 138 from Belarus). The averaged thyroid radiation dose in the exposed control subjects from Russia is 41 mGy (Bouville *et al.* 2007). All exposed control individuals were aged <18 years at the time of the accident (mean age at exposure 1.8 ± 3.2 years old; 1–16 years old, range) (IR-exposed controls). IR-exposed controls and patients with IR-induced PTCs not were individually matched; however, they were residents of the same settlements. This, given the uncertainty with individual radiation thyroid doses, was supposed to partly reduce exposure bias. Age of IR-exposed control subjects was set to be ± 3 years of that of IR-induced PTC individuals.

One hundred and thirty-two PTC cases (21 males and 111 females, mean age at diagnosis 47.8 ± 11.4 years old; 19–76 years old, range) were adults without history of radiation exposure (sporadic PTCs). The remaining 398 control participants (180 males and 218 females, mean age at sampling 45.0 ± 10.3 years old; 16–65 years old, range) had no previous history of radiation exposure (non-exposed controls); their age was also set to be ± 3 years of that of patients with sporadic PTC. Both sporadic PTCs and non-exposed controls originated from the European part of Russia not contaminated by the Chernobyl fallouts.

Thyroid tissues and/or blood samples were collected from patients during surgery or further follow-up. Blood samples and information from the controls were

obtained during a routine health examination or complex screening for thyroid diseases.

Written informed consent was obtained from all participants. Protocols of the present study were approved by the Committee for Ethical Issues of Human Genome Analysis of Nagasaki University.

SNP selection

The candidate SNPs (Table 1) were selected based on their reported functional role (if available), associations with radiosensitivity or (thyroid) cancer risk. Accordingly, we did not search for tag SNPs or account for the genetic variability in the regions of SNP location. All SNPs are listed in a public database, dbSNP (<http://www.ncbi.nlm.nih.gov/SNP/>), with validated status in ethnically diverse populations. To ensure sufficient power for calculations, only SNPs with minor allele frequency (MAF) of >1% were included.

SNP genotyping

DNA was extracted from normal thyroid tissues using proteinase K/phenol–chloroform method or from the whole blood lymphocytes with Puregene DNA Purification Kit (Gentra Systems, Inc., Minneapolis, PA, USA). All specimens were genotyped using various techniques (Table 1). Primers and probes (Table 2) were designed with Primer Express Version 1.0 (Applied Biosystems, Foster City, CA, USA) software.

Briefly, 25 μ l PCR mixtures generally contained 50 ng DNA, 1.5 mM MgCl₂, 200 μ M each dNTP, optimized concentrations of corresponding primers and 0.625 U AmpliTaq Gold (Applied Biosystems). All restriction endonucleases for PCR/RFLP were from New England BioLabs (Ipswich, MA, USA). TaqMan allelic discrimination assay for *TP53* variants was done essentially as described previously (Rogounovitch *et al.* 2006). Melting curve T_m -shift assay for *MTF1* genotyping was designed according to the described technology (Wang *et al.* 2005) and done in a Thermal Cycler Dice Real Time System TP800 (TaKaRa, Ohtsu, Japan). Technical details are available from the authors upon request.

For every SNP, some 20–30 randomly chosen DNA samples, unless otherwise specified, were also analyzed by direct sequencing with a Big Dye Terminator sequencing kit v 3.1 (Applied Biosystems) in an ABI Prism 3100 Genetic Analyzer (Applied Biosystems). A complete concordance between different techniques was observed.

Table 1 Single-nucleotide polymorphism (SNP), genotyping methods and possible functional role

| SNP nucleotide/amino acid change | Database ID | Genotyping method | Chromosome/exon or intron | MAF (%) in different populations (NCBI dbSNP) | SNP effects, minor allele versus wild-type (reference) |
|----------------------------------|-------------|--|---------------------------|--|--|
| ATM G5557A Asp1853Asn | rs1801516 | PCR/RFLP ^a (AflII) GG (187, 30) GA (217, 187, 30) AA (217) | 11/exon 39 | European: 7–22 Asian: 0–2 Global: 5 | Alters an exonic splicing enhancer, modulates correct splicing of exon 39 (Thorstenson et al. 2003) Decreases ATM expression level and capacity of DNA damage recognition (Heikkinen et al. 2005) |
| ATM IVS22–77 T>C T60136C | rs664677 | PCR/RFLP (RsaI) TT (299) TC (299, 265, 34) CC (265, 34) | 11/intron 22 | European: 34–50 Asian: 44–70 Global: 35–36 | No reports |
| ATM IVS48+238 C>G C113450G | rs609429 | PCR/RFLP (KpnI) CC (172, 35) CG (207, 172, 35) GG (207) | 11/intron 48 | European: 60 Asian: 37 Global: 53 | Generates a week additional donor splice site and decreases gene expression (Angele et al. 2003) |
| XRCC1 G25211A Arg280His | rs25489 | PCR/RFLP (RsaI) GG (155, 123) GA (278, 155, 123) AA (278) | 19/exon 9 | European: 3–10 Asian: 0 Global: 7 | Compromises DNA repair (reviewed by Hu et al. 2005) |
| XRCC1 G25897A Arg399Gln | rs25487 | PCR/RFLP (MspI) GG (327, 107) GA (434, 327, 107) AA (434) | 19/exon 10 | European: 30–46 Asian: 27 Global: 23–26 | Affects IR-induced mitotic delay and hypersensitivity to IR (Hu et al. 2002) Compromises single-strand DNA breaks repair (controversially) (reviewed by Taylor et al. 2002) |
| TP53 G640C Arg72Pro | rs1042522 | TaqMan | 17/exon 4 | European: 23–27 Asian: 40–51 Global: 35 | Lower efficiency in apoptosis induction; higher level of G1 arrest (Pim & Banks 2004) |
| XRCC3 C18067T Thr241Met | rs861539 | TaqMan | 14/exon 7 | European: 41–45 Asian: 6–14 Global: 22 | Decreased DNA repair capacity (reviewed by Han et al. 2006) |
| MTF1 T2193A | rs11488567 | Melting curve T_m -shift | 1/intron 1 | Unknown | No reports |
| MTF1 G20433A | rs3912368 | Melting curve T_m -shift | 1/intron 5 | European: 25–37 Asian: 21 | No reports |

^aRestriction enzymes, genotypes and corresponding restriction fragments sizes (bp) are indicated for the SNPs analyzed by PCR/RFLP.

Table 2 Primers and probes for genotyping

| SNP | Primer/probe sequences (5'–3') ^a | Primer/probe concentration (μM) | Annealing temperature (°C) |
|-------------------|---|---------------------------------|----------------------------|
| ATM G5557A | F: CCATACTTGATTCATGATATTTTACcttAA R: TTCATCTTAAATCCATCTTTCTC | 0.2 0.2 | 57 |
| ATM IVS22–77 T>C | F: AGTTTAGCACAGAAAGACATATTGGAAGTAACgTA R: CGGGAAAAGAAGCTGGTTAAATATGAAA | 0.2 0.2 | 57 |
| ATM IVS48+238 C>G | F: CTCATTTCTGGTTATAAAATGAGAAGgTAC R: TTAACACTTGTGTCAGGGACTATCTTAAGGAC | 0.2 0.2 | 57 |
| XRCC1 G25211A | F: GTCTGAGGGAGGAGGGTCTG R: TTCTGGAAGCCACTCAGCAC | 0.2 0.2 | 59 |
| XRCC1 G25897A | F: CCACCAGCTGTGCCTTTG R: CCGGGACTCACTTTGAATGA | 0.2 0.2 | 55 |
| TP53 G640C | F: CGTCCAAGCAATGGATGATT R: CCGGTGTAGGAGCTGCTGG w/t allele probe (FAM): CTCCCCGCGTGGCCCC Variant allele probe (VIC): CTCCCCCGTGGCCCC | 0.8 0.8 0.4 0.4 | 61 |
| XRCC3 C18067T | F: AGGGCCAGGCATCTGCA R: CTTCCGCATCCTGGCTAA w/t allele probe (FAM): TCACGCAGCGTGGCCCCCAG Variant allele probe (VIC): TCACGCAGCATGGCCCCCAG | 0.8 0.8 0.5 0.5 | 61 |
| MTF1 T2193A | F1: GCGGGCAGGGCGGCTTAACTTTAAACCATCAAGTCATTTTAgA F2: GCGGGCTTAACTTTAAACCATCAAGTCATTTTAAAT R: ACGCCCAGTCGGCATTGCT | 0.2 0.2 0.2 | 58 |
| MTF1 G20433A | F1: GCGGGCAGGGCGGCTAATTATGCTCACCTGAATATATACAGGG F2: GCGGGCCTAATTATGCTCACCTGAATATATACAGGA R: GAGACCTGTAGAGCTAGTGGATATACAGAGATAT | 0.075 0.2 0.2 | 63 |

^aThe bases shown in lowercase are mismatches introduced to generate restriction endonuclease sites (PCR/RFLP) or to optimize allelic specificity (T_m -shift). The underlined 5' portions of primer sequences correspond to GC tails in the T_m -shift method.

Raw genotyping outputs were interpreted by at least two independent investigators. Missing results due to genotyping procedure failures accounted for <1% for any SNP tested.

Statistical analysis

Genotype frequencies in each group were determined by univariate analysis and evaluated for departure from Hardy-Weinberg equilibrium by the χ^2 test. SNP associations with PTC were assessed by multivariate logistic regression analysis for codominant, multiplicative, dominant, and recessive models to avoid assumptions regarding the mode of inheritance (see notes below Table 4). All analyses were adjusted for gender (male or female, nominal), age (years, continuous), and IR-exposure (yes or no, nominal). Besides all the parameters above, the full model included disease status (yes or no, nominal) and, depending on the mode of inheritance, genotype for each SNP (nominal variable in the codominant, dominant, and recessive models and ordinal in the multiplicative model).

Power calculations were done with the PS software (<http://biostat.mc.vanderbilt.edu/twiki/bin/view/Main/PowerSampleSize>). With given sample size, the study

had a power of 54–99% to detect an OR of 2.0 at the significance level of 5% with MAF ranging 4–45%.

Interaction between SNPs, cancer and radiation exposure were hypothesized *a priori* and evaluated by multivariate analysis with corresponding adjustments. Separate calculations of OR were done in irradiated and non-exposed case-control groups when *P* value for an interaction term did not exceed 0.05.

Statistical analysis was done using SPSS for Windows version 17.0 (SPSS, Inc., Chicago, IL, USA).

Results

The distribution of genotypes and MAF for each SNP in the four study groups is shown in Table 3. The observed distributions in the control groups were not statistically different from those expected from Hardy-Weinberg equilibrium for all SNP except for *ATM* G5557A and *ATM* IVS22-77 T>C in the non-exposed controls. Since such deviation might point at possible genotyping error (Hosking *et al.* 2004), we reanalyzed 96 non-exposed controls for these SNPs by direct sequencing. There were no inconsistencies between PCR/RFLP and sequencing results (data not shown) ruling out technical flaw. Furthermore, allelic frequencies determined in our study are in a good agreement

Table 3 Distribution of genotypes and minor allele frequencies by study groups

| SNP, genotype | IR-induced PTC <i>n</i> (%) | IR-exposed controls <i>n</i> (%) | Sporadic PTC <i>n</i> (%) | Non-exposed controls <i>n</i> (%) |
|------------------------------------|--------------------------------|-------------------------------------|------------------------------|--------------------------------------|
| ATM G5557A | <i>n</i> =122 | <i>n</i> =198 | <i>n</i> =132 | <i>n</i> =398 |
| GG | 95 (77.9) | 138 (69.7) | 105 (79.5) | 293 (73.6) |
| GA | 25 (20.5) | 53 (26.8) | 24 (18.2) | 90 (22.6) |
| AA | 2 (1.6) | 7 (3.5) | 3 (2.3) | 15 (3.8) |
| <i>P</i> | 0.24 | | 0.36 | |
| A, % | 11.9 | 16.9 | 11.4 | 15.1 |
| ATM IVS22 – 77 T>C | <i>n</i> =123 | <i>n</i> =195 | <i>n</i> =132 | <i>n</i> =398 |
| TT | 35 (28.4) | 62 (31.8) | 45 (34.1) | 135 (33.9) |
| TC | 76 (61.8) | 102 (52.3) | 61 (46.2) | 216 (54.3) |
| CC | 12 (9.8) | 31 (15.9) | 26 (19.7) | 47 (11.8) |
| <i>P</i> | 0.17 | | 0.06 | |
| C, % | 40.6 | 42.0 | 42.8 | 38.9 |
| ATM IVS48 + 238 C>G | <i>n</i> =122 | <i>n</i> =196 | <i>n</i> =132 | <i>n</i> =398 |
| CC | 37 (30.3) | 68 (34.7) | 41 (31.1) | 131 (32.9) |
| CG | 69 (56.6) | 97 (49.5) | 61 (46.2) | 201 (50.5) |
| GG | 16 (13.1) | 31 (15.8) | 30 (22.7) | 66 (16.6) |
| <i>P</i> | 0.47 | | 0.28 | |
| G, % | 41.4 | 40.3 | 45.8 | 41.8 |
| XRCC1 Arg280His^a | <i>n</i> =123 | <i>n</i> =195 | <i>n</i> =132 | <i>n</i> =398 |
| GG | 113 (91.9) | 176 (90.3) | 117 (88.6) | 366 (92.0) |
| GA | 10 (8.1) | 19 (9.7) | 15 (11.4) | 32 (8.0) |
| <i>P</i> | 0.63 | | 0.24 | |
| A, % | 4.1 | 4.9 | 5.7 | 4.0 |
| XRCC1 Arg399Gln | <i>n</i> =123 | <i>n</i> =197 | <i>n</i> =132 | <i>n</i> =398 |
| GG | 55 (44.7) | 75 (38.1) | 65 (49.2) | 158 (39.7) |
| GA | 50 (40.7) | 100 (50.7) | 53 (40.2) | 193 (48.5) |
| AA | 18 (14.6) | 22 (11.2) | 14 (10.6) | 47 (11.8) |
| <i>P</i> | 0.20 | | 0.15 | |
| A, % | 35.1 | 36.5 | 30.7 | 36.1 |
| TP53 Arg72Pro | <i>n</i> =122 | <i>n</i> =197 | <i>n</i> =129 | <i>n</i> =395 |
| GG | 53 (43.4) | 115 (58.4) | 69 (53.5) | 196 (49.6) |
| GC | 57 (46.7) | 73 (37.0) | 49 (38.0) | 161 (40.8) |
| CC | 12 (9.9) | 9 (4.6) | 11 (8.5) | 38 (9.6) |
| <i>P</i> | 0.02 | | 0.74 | |
| C, % | 33.2 | 23.1 | 27.5 | 30.0 |
| XRCC3 Thr241Met | <i>n</i> =120 | <i>n</i> =198 | <i>n</i> =132 | <i>n</i> =398 |
| CC | 53 (44.2) | 82 (41.4) | 55 (41.7) | 161 (40.5) |
| CT | 51 (42.5) | 89 (45.0) | 65 (49.2) | 192 (48.2) |
| TT | 16 (13.3) | 27 (13.6) | 12 (9.1) | 45 (11.3) |
| <i>P</i> | 0.89 | | 0.78 | |
| T, % | 34.6 | 36.1 | 33.7 | 35.4 |
| MTF1 T2193A | <i>n</i> =122 | <i>n</i> =198 | <i>n</i> =131 | <i>n</i> =397 |
| TT | 45 (36.9) | 82 (41.4) | 44 (33.6) | 133 (33.5) |
| TA | 64 (52.5) | 91 (46.0) | 67 (51.1) | 188 (47.4) |
| AA | 13 (10.6) | 25 (12.1) | 20 (15.3) | 76 (19.1) |
| <i>P</i> | 0.52 | | 0.57 | |
| A, % | 36.8 | 35.6 | 40.8 | 42.8 |
| MTF1 G20433A | <i>n</i> =123 | <i>n</i> =198 | <i>n</i> =132 | <i>n</i> =398 |
| GG | 62 (50.4) | 100 (50.5) | 66 (50.0) | 192 (48.2) |
| GA | 53 (43.1) | 88 (44.4) | 56 (42.4) | 151 (38.0) |
| AA | 8 (6.5) | 10 (5.1) | 10 (7.6) | 55 (13.8) |
| <i>P</i> | 0.85 | | 0.16 | |
| A, % | 28.0 | 27.3 | 28.8 | 32.8 |

NOTE. Total numbers of samples in each group vary slightly due to genotyping procedures failures.

^aThere was no homozygous (A/A) variant of XRCC1 Arg280His among all samples tested.

with those specified for Caucasians in the dbSNP (build 129, April 2008, Table 1) thus attesting to the appropriate data quality.

As seen from Table 4, an association between *ATM* G5557A and PTC, regardless of radiation exposure, was found. The presence of the A allele significantly decreased PTC risk compared with wild-type G allele in the multiplicative model of inheritance (OR=0.69, 95% CI 0.45–0.86, $P=0.03$), which is useful for risk comparison between the groups based on the analysis of allelic frequencies in them.

Main effect on PTC risk appeared also significant for the *XRCC1* gene Arg399Gln polymorphism. The presence of the minor 399Gln allele decreased PTC risk compared with the Arg/Arg genotype (OR=0.66, 95% CI 0.57–0.88, $P=0.02$ and OR=0.70, 95% CI 0.59–0.93, $P=0.03$, in the co-dominant and dominant models respectively).

Analysis of combined *ATM* G5557A and *XRCC1* Arg399Gln genotypes demonstrated that increasing number of minor alleles (i.e. *ATM* 5557A and *XRCC1* 399Gln) significantly decreased PTC risk in corresponding individuals in comparison with those who do not carry minor alleles (Fig. 1).

No other SNP in any gene showed a significant main effect on PTC.

For *ATM* IVS22-77 T>C and *TP53* Arg72Pro, evidence for interaction between radiation exposure and PTC was found (P for interaction 0.04 and 0.01 respectively). As shown in Table 5, the analyses performed in IR-exposed and non-irradiated patients compared respectively, with irradiated and non-exposed controls revealed a significantly increased risk of sporadic PTC for the *ATM* IVS22-77 homozygous CC genotype carriers compared with the TC+TT genotypes (the recessive model of inheritance, OR=1.84, 95% CI 1.10–3.24, $P=0.03$), whereas in the irradiated group an insignificant inverse effect of these genotypes was observed (OR=0.59, 95% CI 0.28–1.27, $P=0.17$). For *TP53* codon 72 polymorphism, in all but the recessive models the increased risk of IR-induced PTC as compared with IR-exposed controls was observed. The highest risk of radiogenic PTC was in the co-dominant model (OR=2.33, 95% CI 1.15–7.21, $P=0.03$). A significant risk was also found in the multiplicative model of inheritance (OR=1.70, 95% CI 1.17–2.46, $P=0.006$). In addition, comparison between IR-exposed and non-exposed controls did not reveal statistically significant difference in adjusted distributions of these polymorphisms. In healthy subjects, the strongest association for the *ATM* IVS22-77 T>C was in the recessive model (OR=1.38, 95% CI 0.84–2.26,

$P=0.21$) and in the multiplicative model for *TP53* Arg72Pro (OR=0.70, 95% CI 0.52–1.19, $P=0.11$) further emphasizing possible role of these SNPs in PTC of different etiology.

Considering multiple pathways for repairing diverse DNA damages induced by endogenous and exogenous carcinogens, genetic variants in different repair pathways may probably have a joint effect on cancer risk. In attempt to search for the stronger associations between PTC and studied SNPs, we performed the analyses of genotype combinations for the *ATM* and *TP53* polymorphisms as these genes are functionally related and three out of four SNPs included in our study showed effects on PTC. Among the possible *ATM/TP53* combinations (rs1801516/rs664677/rs609429/rs1042522) tested, two demonstrated significant differences in the subsets of both groups of PTCs (Fig. 2). Particularly, the combined *ATM/TP53* GG/TC/CG/GC genotype was strongly associated with the IR-induced PTC (OR=2.10, 95% CI 1.17–3.78, $P=0.015$). Another *ATM/TP53* combination, GG/CC/GG/GG, demonstrated a significantly increased risk for sporadic PTC (OR=3.32, 95% CI 1.57–6.99, $P=0.002$).

Discussion

Our study addressed possible associations between SNPs in the genes involved in DNA damage response and the risk of PTC of different etiology. The results demonstrated that the presence of the variant 5557A allele in exon 39 of *ATM* and *XRCC1* 399Gln allele, particularly in the heterozygous state, significantly associated with the decreased risk of PTC. The *ATM* IVS22-77 CC genotype in the non-exposed group and the *TP53* 72Pro allele in the radiation-related one associated with the increased risk of PTC. Moreover, two particular *ATM/TP53* combined genotypes were found with higher frequencies in the IR-induced or sporadic PTC when compared with the controls. Altogether, these data indicate that SNPs in the studied genes may likely modify PTC risk.

A significant association between the *ATM* G5557A and bilateral breast cancer in Caucasian patients has been shown before (Heikkinen *et al.* 2005). Also, this SNP has been reported as a possible modulator of clinical radiosensitivity in cancer. The *ATM* 5557A allele was associated with severe adverse effects of radiation therapy in prostate (Hall *et al.* 1998) and breast cancer patients (Angele *et al.* 2003). Later, an enhanced radiosensitivity of human fibroblasts in the presence of the *ATM* 5557A allele was demonstrated in an experimental work (Alsbeih *et al.* 2007). In contrast

Table 4 OR (95% CI) for papillary thyroid carcinoma (PTC) by gene polymorphism according to different models of inheritance (adjusted for age, gender and radiation exposure). *P* <0.05 in bold

| SNP | Genotype | OR (95% CI) | <i>P</i> |
|------------------------------|--------------------------------|-------------------|-------------|
| ATM G5557A | GG | 1.00 ^a | |
| | GA | 0.75 (0.49–1.15) | 0.31 |
| | AA | 0.61 (0.21–1.77) | 0.45 |
| | Risk per A allele ^b | 0.69 (0.45–0.86) | 0.03 |
| | GA+AA versus GG ^c | 0.73 (0.48–1.10) | 0.13 |
| | AA versus GA+GG ^d | 0.65 (0.23–1.87) | 0.41 |
| ATM IVS22–77 T>C | TT | 1.00 | |
| | TC | 1.03 (0.70–1.50) | 0.74 |
| | CC | 1.19 (0.70–2.04) | 0.47 |
| | Risk per C allele | 1.08 (0.83–1.40) | 0.57 |
| | TC+CC versus TT | 1.06 (0.74–1.53) | 0.75 |
| | CC versus TC+TT | 1.17 (0.72–1.90) | 0.52 |
| ATM IVS48+238 C>G | CC | 1.00 | |
| | CG | 1.10 (0.75–1.62) | 0.55 |
| | GG | 1.14 (0.69–1.89) | 0.84 |
| | Risk per G allele | 1.07 (0.84–1.37) | 0.57 |
| | CG+GG versus CC | 1.11 (0.77–1.60) | 0.57 |
| | GG versus CG+CC | 1.08 (0.69–1.69) | 0.74 |
| XRCC1 Arg280His ^e | GG | 1.00 | |
| | GA | 1.12 (0.62–2.01) | 0.71 |
| | Risk per A allele | 1.15 (0.70–1.87) | 0.61 |
| XRCC1 Arg399Gln | GG | 1.00 | |
| | GA | 0.66 (0.57–0.88) | 0.02 |
| | AA | 0.88 (0.50–1.57) | 0.56 |
| | Risk per A allele | 0.90 (0.69–1.17) | 0.41 |
| | GA+AA versus GG | 0.70 (0.59–0.93) | 0.03 |
| | AA versus GA+GG | 0.98 (0.57–1.69) | 0.94 |
| TP53 Arg72Pro | GG | 1.00 | |
| | GC | 1.02 (0.70–1.47) | 0.89 |
| | CC | 1.16 (0.63–2.14) | 0.38 |
| | Risk per C allele | 1.05 (0.81–1.38) | 0.70 |
| | GC+CC versus GG | 1.04 (0.74–1.48) | 0.82 |
| | CC versus GC+GG | 1.15 (0.64–2.08) | 0.64 |
| XRCC3 Thr241Met | CC | 1.00 | |
| | CT | 0.99 (0.69–1.44) | 0.99 |
| | TT | 0.96 (0.54–1.70) | 0.92 |
| | Risk per T allele | 0.99 (0.76–1.28) | 0.92 |
| | CT+TT versus CC | 0.99 (0.70–1.41) | 0.97 |
| | TT versus CT+CC | 0.96 (0.56–1.64) | 0.88 |
| MTF1 T2193A | TT | 1.00 | |
| | TA | 1.07 (0.73–1.56) | 0.61 |
| | AA | 0.83 (0.49–1.41) | 0.46 |
| | Risk per A allele | 0.94 (0.73–1.21) | 0.63 |
| | TA+AA versus TT | 1.00 (0.70–1.44) | 0.99 |
| | AA versus TA+TT | 0.80 (0.49–1.29) | 0.35 |
| MTF1 G20433A | GG | 1.00 | |
| | GA | 1.14 (0.79–1.63) | 0.43 |
| | AA | 0.76 (0.40–1.43) | 0.21 |
| | Risk per A allele | 0.97 (0.74–1.25) | 0.80 |
| | GA+AA versus GG | 1.05 (0.76–1.49) | 0.76 |
| | AA versus GA+GG | 0.71 (0.39–1.32) | 0.27 |

^aCodominant model of inheritance (wild-type homozygous genotype serves as the reference).

^bMultiplicative model of inheritance (uses allele frequencies).

^cDominant inheritance model (combined heterozygous and homozygous for the minor allele versus wild-type homozygous).

^dRecessive inheritance model (minor allele homozygous versus combined heterozygous and homozygous for the wild-type allele).

^eThe dominant and recessive models are not shown for *XRCC1* Arg280His because of the absence of homozygous (A/A) genotype among 848 samples tested.




Figure 1 Effect of increasing number of minor alleles (MA) for *ATM* G5557A and *XRCC1* Arg399Gln (minor alleles, *ATM* 5557A, and *XRCC1* 399Gln) on PTC risk. The combined genotype with 0 MA was used as a reference. *P* values for genotypes with different MA number: $P_{1MA} < 0.0001$; $P_{2MA} < 0.01$; $P_{3-4MA} < 0.05$. Carriers of three and four minor alleles were combined because of the exceedingly low number of 4 MA carriers in both PTC and control groups.

to these reports, Edvardsen *et al.* (2007) revealed an increasing rate of side effects of radiotherapy with decreasing frequency of this variant allele. Our data are rather in agreement with the latter report and favor the protective role of the *ATM* 5557A allele in PTC development.

The intronic *ATM* polymorphisms IVS22–77 T>C and IVS48 + 238 C>G in the homozygous state have

been associated with increased breast cancer risk and in the heterozygous state with clinical radioprotection (Angele *et al.* 2003). These findings were confirmed in the *in vitro* experiments using lymphoblastoid cell lines established from corresponding patients. Our investigation demonstrated the association between the IVS22–77 CC genotype and increased risk of sporadic PTC in adult patients. By contrast, in the IR-induced PTC group, there was an inverse non-insignificant correlation for this genotype. At the same time, in the IR-induced PTCs, the number of patients heterozygous for IVS22–77 was somewhat, but insignificantly, higher as compared with sporadic PTCs (Table 3). The results for the IVS48 + 238 C>G tended to parallel those for the IVS22–77 T>C remaining below the threshold of significance. At present, the mechanistic and functional basis for the intronic *ATM* SNPs implications in cancer revealed in the previous studies and in ours as well is not fully understood. In a broader sense, however, they may be indicative of a role for the *ATM* gene (or its product) in the development of PTC.

As reviewed by Hu *et al.* (2005), the results of the *XRCC1* gene Arg399Glu investigations vary in different cancers for populations with different ethnicities. In relation to cancer and radiation, the 399Gln allele in combination with 280His was associated with breast cancer risk, and in pair with 194Trp with clinical radiosensitivity in Caucasian women with breast cancer. Also, the 399Gln allele was found to decrease

Table 5 OR (95% CI) for papillary thyroid carcinoma (PTC) of different etiology by *ATM* and *TP53* polymorphisms (adjusted for gender and age). *P* < 0.05 in bold

| SNP | Genotype | IR-induced PTC versus IR-exposed controls | | Sporadic PTC versus non-exposed controls | |
|-------------------------|--------------------------------|---|--------------|--|-------------|
| | | OR (95% CI) | <i>P</i> | OR (95% CI) | <i>P</i> |
| <i>ATM</i> IVS22–77 T>C | TT | 1.00 ^a | | 1.00 | |
| | TC | 1.38 (0.80–2.39) | 0.19 | 0.82 (0.51–1.32) | 0.50 |
| | CC | 0.73 (0.31–1.70) | 0.44 | 1.63 (0.87–3.08) | 0.09 |
| | Risk per C allele ^b | 0.97 (0.66–1.41) | 0.86 | 1.18 (0.86–1.62) | 0.32 |
| | TC+CC versus TT ^c | 1.23 (0.72–2.10) | 0.44 | 0.97 (0.62–1.52) | 0.88 |
| | CC versus TC+TT ^d | 0.59 (0.28–1.27) | 0.17 | 1.84 (1.10–3.24) | 0.03 |
| <i>TP53</i> Arg72Pro | GG | 1.00 | | 1.00 | |
| | GC | 1.68 (1.11–2.75) | 0.03 | 0.84 (0.53–1.33) | 0.52 |
| | CC | 2.33 (1.15–7.21) | 0.03 | 0.84 (0.39–1.79) | 0.73 |
| | Risk per C allele | 1.70 (1.17–2.46) | 0.006 | 0.89 (0.64–1.23) | 0.47 |
| | GC+CC versus GG | 1.80 (1.06–2.36) | 0.01 | 0.84 (0.54–1.29) | 0.43 |
| | CC versus GC+GG | 2.06 (0.79–5.41) | 0.14 | 0.90 (0.44–1.88) | 0.79 |

^aCodominant model of inheritance (wild-type homozygous genotype serves as the reference).

^bMultiplicative model of inheritance (uses allele frequencies).

^cDominant inheritance model (combined heterozygous and homozygous for the minor allele versus wild-type homozygous).

^dRecessive inheritance model (minor allele homozygous versus combined heterozygous and homozygous for the wild-type allele).




Figure 2 The combined *ATM/TP53* genotypes and risk of PTC of different etiology. The combined genotypes were analyzed separately in the IR-exposed and sporadic PTCs versus corresponding control. Six combinations of 3 *ATM* and 1 *TP53* SNPs (rs1801516/rs664677/rs609429/rs1042522) whose frequencies were higher than 5% at least in two of four subgroups are shown. In the numerical codes for any SNP, 0 – the genotype with no MA (i.e. homozygous wild-type); 1 – 1 MA presents (heterozygous genotype); 2 – 2 MA present (homozygous variant genotype); first three numbers correspond to 3 *ATM* SNPs and the last one to *TP53* polymorphism. In the figure, the *GG/TT/CC/GG* genotype is represented by the '0000' numerical code as it does not contain minor alleles; the *GG/TC/CG/GG* corresponds to 0110, *GG/TC/CG/GC* to 0111; *GG/CC/GG/GC* to 0221; *GA/TC/CG/GG* to 1110, and *GG/CC/GG/GG* to 0220. All combinations with frequencies <5% in three or more subgroups are pooled and indicated as 'other'.

the risk of bladder cancer and squamous cell carcinoma of the head and neck.

Interestingly, not only variant but also wild-type allele (i.e. *XRCC1* 399Arg) demonstrated possible role in cancer. High-dose radiation to the chest was more strongly associated with breast cancer among white American women with *XRCC1* Arg399Arg genotype (Duell et al. 2001). Looking for potential biological explanations for these findings, the authors found a higher prevalence of *TP53* deletions in the Arg399Arg cases exposed to occupational radiation compared with exposed patients with the Gln399Gln genotypes or unexposed cases of either genotype. Figueiredo et al. (2004) observed an increased risk of disease among wild-type homozygous (Arg/Arg) and heterozygous Canadian Caucasian women with a family history of breast cancer compared with the individuals without such.

The described above data may be explained, at least in part, by the results of functional study of this polymorphism in which an equal ability for both alleles to suffice single strand break repair by *XRCC1* has been

found (Taylor et al. 2002). The results of our study, taken together with those reported previously, suggest that *XRCC1* polymorphism in particular the Arg399Gln genotype may influence PTC risk, perhaps by modifying the effects of environmental exposure and/or through interaction with other genetic factors.

The *TP53* Arg72Pro polymorphism affects the biological activity of p53. The Arg72 form is more efficient at inducing apoptosis while the Pro72 appears to induce a higher level of G1 arrest (Pim & Banks 2004). Based on these findings, a number of studies have attempted to assess a correlation between *TP53* codon 72 polymorphism and risk of certain types of cancer, however, with inconsistent results, as reviewed by Pietsch et al. (2006). This inconsistency may possibly be explained in part by the coexistence of the codon 72 polymorphism and gain of function mutations in *TP53* in some tumors (Pietsch et al. 2006, Soussi & Wiman 2007).

Several groups have investigated the *TP53* Arg72-Pro polymorphism in PTC. Boltze et al. (2002) found a small number of heterozygotes and no Pro/Pro genotype in differentiated thyroid carcinomas from Germany. By contrast, in ethnically heterogeneous Brazilian population, the Pro/Pro genotype was associated with the higher risk of differentiated thyroid cancer (Granja et al. 2004). The study of codon 72 polymorphism in thyroid tumors from Russian and Ukrainian patients demonstrated a significantly lower frequency of wild-type homozygotes (i.e. Arg/Arg) among adults with IR-induced PTC when compared with sporadic PTC cases and general population (Rogounovitch et al. 2006). Data obtained in the present work, using an independent set of samples, confirm these findings suggesting the modifying role (or as of a marker) of the *TP53* Arg72Pro polymorphism in PTC developed after exposure to IR which is further supported by the absence of significant difference in genotype distributions among our two control groups.

As shown in a genetic study, frequencies of the C allele (encoding 72Pro) do not generally differ in populations of Belarus and Russia (Khrunin et al. 2005). However, East Slavs do not form a single genetic cluster on multidimensional analysis. The 72Pro allele frequency in Belarus is about 0.3; in the two different subpopulations from the Central and Northern regions of the European part of Russia it is 0.24 and 0.32 respectively. The study of healthy population from Poland (bordering with Belarus, linguistically and culturally similar), reported the frequency of 0.28 for the 72Pro allele (Siddique et al. 2005). The 72Pro frequency reported by

Rogounovitch *et al.* (2006) in Russian healthy controls is also 0.28. Thus, the effect of population admixtures in the controls in our investigation could not be completely ruled out. Yet on the other hand, the ratio of Belarusian and Russian subjects in the IR-exposed PTCs and controls was similar (2.24 and 2.30 respectively) suggestive of an unbiased estimate and being an argument in support of *TP53* Arg72Pro polymorphism association with radiation-related PTC.

While many studies established the effect of individual SNPs on cancer, the role of SNP combinations has been less addressed. Several *ATM* and *TP53* haplotypes were associated with clinical radiosensitivity in breast cancer (Angele *et al.* 2003) and brain tumor risk (Malmer *et al.* 2007). Recently, the interactions of SNPs located on different chromosomes were investigated in various malignancies (Yen *et al.* 2008, Yoon *et al.* 2008). One experimental study, in which *ATM* Asp1853Asn, *TP53* Arg72Pro, *XRCC1* Arg399Gln, and *XRCC3* Thr241Met were genotyped, demonstrated that the increasing number of risk alleles enhanced radiosensitivity of human fibroblast cell lines and, potentially, susceptibility to radiation-induced cancers (Alsbeih *et al.* 2007). So far no studies have investigated the joint effect of gene polymorphisms on thyroid cancer. Our observations demonstrated that frequencies of particular combined *ATM/TP53* genotypes were higher in patients with radiogenic or sporadic PTC compared with corresponding control populations.

To some extent these results support the idea that genetic factors may possibly modify predisposition to thyroid cancer. A recent study by Detours *et al.* (2007) reported difference in the expression levels of some genes between Chernobyl PTCs from Ukraine and French sporadic PTCs. Although the mentioned work and the present one are different in molecular approaches, the results of both are suggestive of a possible genetic 'susceptibility signature' that may contribute to the individual predisposition to IR and other carcinogens' effects. These findings are in favor of a 'susceptibility model' that may partly explain why only a minority of the large population exposed to the IR after the Chernobyl disaster developed thyroid cancer (Yamashita & Saenko 2006, Detours *et al.* 2007, 2008).

It is necessary to note that even though nine SNPs were analyzed in our study, no correction for multiple comparisons was applied because of study design and techniques employed. The associations were tested in a one-at-a-time fashion in a limited sample size in the difficult to access groups. The need for correction in such circumstances is still debated (Rothman &

Greenland 1998). Furthermore, since data obtained in this work may be referred to as an initial screening result, non-adjusted presentation enables their inclusion in future meta-analysis. Effects of candidate SNPs that we report need validation in other studies.

In conclusion, the results presented here show that SNPs in *ATM* exon 39 and *XRCC1* exon 10 may be the markers of a decreased PTC risk in adults, whereas the *ATM* IVS22-77 and *TP53* codon 72 SNPs genes may associate with the risk of PTC development in non-irradiated and irradiated individuals. To the best of our knowledge, presented here is the first study of this kind reporting the results of genotyping of candidate DNA damage response genes in irradiated and non-irradiated PTC patients and in corresponding healthy populations. Our data support the paradigm of genetic modifiers of radiation-associated carcinogenesis and perhaps may contribute to genetic determination of PTC-prone subjects. We believe such identification will allow future personalized cancer risk prediction which is of a significant importance in view of the growing thyroid cancer incidence and also because of the relevance to occupational and radiation emergency medicine issues.

Declaration of interest

The authors declare no potential conflict of interest.

Funding

This work was supported in part by Grant-in-Aid for Scientific Research 19256003, 19510058, and 19790651 from Japan Society for the Promotion of Science.

References

- Alsbeih G, El-Sebaie M, Al-Harbi N, Al-Buhairi M, Al-Hadyan K & Al-Rajhi N 2007 Radiosensitivity of human fibroblasts is associated with amino acid substitution variants in susceptible genes and correlates with the number of risk alleles. *International Journal of Radiation Oncology, Biology, Physics* **68** 229–235.
- Angele S, Romestaing P, Moullan N, Vuillaume M, Chapot B, Friesen M, Jongmans W, Cox DG, Pisani P, Gerard JP *et al.* 2003 *ATM* haplotypes and cellular response to DNA damage: association with breast cancer risk and clinical radiosensitivity. *Cancer Research* **63** 8717–8725.
- Bennett B, Repacholi M & Carr Z 2006 *Health Effects of the Chernobyl Accident and Special Health Care Programmes*. Geneva: WHO Press.
- Boltze C, Roessner A, Landt O, Szibor R, Peters B & Schneider-Stock R 2002 Homozygous proline at codon 72

- of p53 as a potential risk factor favoring the development of undifferentiated thyroid carcinoma. *International Journal of Oncology* **21** 1151–1154.
- Bouville A, Likhtarev IA, Kovgan L, Minenko VF, Shinkarev SM & Drozdovitch V 2007 Radiation dosimetry for highly contaminated Belarusian, Russian and Ukrainian populations, and for less contaminated populations in Europe. *Health Physics* **93** 487–501.
- Cardis E, Kesminiene A, Ivanov V, Malakhova I, Shibata Y, Khrouch V, Drozdovitch V, Maceika E, Zvonova I, Vlassov O et al. 2005 Risk of thyroid cancer after exposure to ¹³¹I in childhood. *Journal of the National Cancer Institute* **97** 724–732.
- Davies L & Welch HG 2006 Increasing incidence of thyroid cancer in the United States, 1973–2002. *Journal of the American Medical Association* **295** 2164–2167.
- Davis S, Stepanenko V, Rivkind N, Kopecky KJ, Voilleque P, Shakhtarin V, Parshkov E, Kulikov S, Lushnikov E, Abrosimov A et al. 2004 Risk of thyroid cancer in the Bryansk oblast of the Russian Federation after the Chernobyl Power Station accident. *Radiation Research* **162** 241–248.
- Detours V, Delys L, Libert F, Weiss Solis D, Bogdanova T, Dumont JE, Franc B, Thomas G & Maenhaut C 2007 Genome-wide gene expression profiling suggests distinct radiation susceptibility in sporadic and post-Chernobyl papillary thyroid cancers. *British Journal of Cancer* **97** 818–825.
- Detours V, Verstehey S, Dumont JE & Maenhaut C 2008 Gene expression profiles of post-Chernobyl thyroid cancers. *Current Opinion in Endocrinology, Diabetes, and Obesity* **15** 440–445.
- Duell EJ, Millikan RC, Pittman GS, Winkel S, Lunn RM, Tse CK, Eaton A, Mohrenweiser HW, Newman B & Bell DA 2001 Polymorphisms in the DNA repair gene XRCC1 and breast cancer. *Cancer Epidemiology, Biomarkers & Prevention* **10** 217–222.
- Edvardsen H, Tefre T, Jansen L, Vu P, Haffty BG, Fossa SD, Kristensen VN & Borresen-Dale AL 2007 Linkage disequilibrium pattern of the ATM gene in breast cancer patients and controls; association of SNPs and haplotypes to radio-sensitivity and post-lumpectomy local recurrence. *Radiation Oncology* **2** 25.
- Figueiredo JC, Knight JA, Briollais L, Andrusis IL & Ozcelik H 2004 Polymorphisms XRCC1-R399Q and XRCC3-T241M and the risk of breast cancer at the Ontario site of the Breast Cancer Family Registry. *Cancer Epidemiology, Biomarkers & Prevention* **13** 583–591.
- Granja F, Morari J, Morari EC, Correa LA, Assumpcao LV & Ward LS 2004 Proline homozygosity in codon 72 of p53 is a factor of susceptibility for thyroid cancer. *Cancer Letters* **210** 151–157.
- Hall EJ, Schiff PB, Hanks GE, Brenner DJ, Russo J, Chen J, Sawant SG & Pandita TK 1998 A preliminary report: frequency of A-T heterozygotes among prostate cancer patients with severe late responses to radiation therapy. *Cancer Journal from Scientific American* **4** 385–389.
- Han S, Zhang HT, Wang Z, Xie Y, Tang R, Mao Y & Li Y 2006 DNA repair gene XRCC3 polymorphisms and cancer risk: a meta-analysis of 48 case-control studies. *European Journal of Human Genetics* **14** 1136–1144.
- Heikkinen K, Rapakko K, Karppinen SM, Erkko H, Nieminen P & Winqvist R 2005 Association of common ATM polymorphism with bilateral breast cancer. *International Journal of Cancer* **116** 69–72.
- Hillebrandt S, Streffer C, Demidchik EP, Biko J & Reiners C 1997 Polymorphisms in the p53 gene in thyroid tumours and blood samples of children from areas in Belarus. *Mutation Research* **381** 201–207.
- Hoeijmakers JH 2001 Genome maintenance mechanisms for preventing cancer. *Nature* **411** 366–374.
- Hosking L, Lumsden S, Lewis K, Yeo A, McCarthy L, Bansal A, Riley J, Purvis I & Xu CF 2004 Detection of genotyping errors by Hardy-Weinberg equilibrium testing. *European Journal of Human Genetics* **12** 395–399.
- Hu JJ, Smith TR, Miller MS, Lohman K & Case LD 2002 Genetic regulation of ionizing radiation sensitivity and breast cancer risk. *Environmental and Molecular Mutagenesis* **39** 208–215.
- Hu Z, Ma H, Chen F, Wei Q & Shen H 2005 XRCC1 polymorphisms and cancer risk: a meta-analysis of 38 case-control studies. *Cancer Epidemiology, Biomarkers & Prevention* **14** 1810–1818.
- Jacob P, Bogdanova TI, Buglova E, Chepurnyi M, Demidchik Y, Gavrillin Y, Kenigsberg J, Meckbach R, Schotola C, Shinkarev S et al. 2006 Thyroid cancer risk in areas of Ukraine and Belarus affected by the Chernobyl accident. *Radiation Research* **165** 1–8.
- Khanna KK & Jackson SP 2001 DNA double-strand breaks: signaling, repair and the cancer connection. *Nature Genetics* **27** 247–254.
- Khrunin AV, Tarskaia LA, Spitsyn VA, Lylova OI, Bebyakova NA, Mikulich AI & Limborska SA 2005 p53 polymorphisms in Russia and Belarus: correlation of the 2-1-1 haplotype frequency with longitude. *Molecular Genetics and Genomics* **272** 666–672.
- Likhtarev I, Bouville A, Kovgan L, Luchyanov N, Voilleque P & Chepurnyi M 2006 Questionnaire- and measurement-based individual thyroid doses in Ukraine resulting from the Chernobyl nuclear reactor accident. *Radiation Research* **166** 271–286.
- Malmer BS, Feychting M, Lonn S, Lindstrom S, Gronberg H, Ahlbom A, Schwartzbaum J, Auvinen A, Collatz-Christensen H, Johansen C et al. 2007 Genetic variation in p53 and ATM haplotypes and risk of glioma and meningioma. *Journal of Neuro-oncology* **82** 229–237.
- Nikiforov YE, Nikiforova MN, Gnepp DR & Fagin JA 1996 Prevalence of mutations of ras and p53 in benign and malignant thyroid tumors from children exposed to radiation after the Chernobyl nuclear accident. *Oncogene* **13** 687–693.

- Nikiforov YE, Rowland JM, Bove KE, Monforte-Munoz H & Fagin JA 1997 Distinct pattern of ret oncogene rearrangements in morphological variants of radiation-induced and sporadic thyroid papillary carcinomas in children. *Cancer Research* **57** 1690–1694.
- Pietsch EC, Humbey O & Murphy ME 2006 Polymorphisms in the p53 pathway. *Oncogene* **25** 1602–1611.
- Pim D & Banks L 2004 p53 polymorphic variants at codon 72 exert different effects on cell cycle progression. *International Journal of Cancer* **108** 196–199.
- Rabes HM, Demidchik EP, Sidorow JD, Lengfelder E, Beimfohr C, Hoelzel D & Klugbauer S 2000 Pattern of radiation-induced RET and NTRK1 rearrangements in 191 post-Chernobyl papillary thyroid carcinomas: biological, phenotypic, and clinical implications. *Clinical Cancer Research* **6** 1093–1103.
- Rougounovitch TI, Saenko VA, Ashizawa K, Sedliarou IA, Namba H, Abrosimov AY, Lushnikov EF, Roumiantsev PO, Konova MV, Petoukhova NS *et al.* 2006 TP53 codon 72 polymorphism in radiation-associated human papillary thyroid cancer. *Oncology Reports* **15** 949–956.
- Ron E, Lubin JH, Shore RE, Mabuchi K, Modan B, Pottern LM, Schneider AB, Tucker MA & Boice JD Jr 1995 Thyroid cancer after exposure to external radiation: a pooled analysis of seven studies. *Radiation Research* **141** 259–277.
- Rothman KJ & Greenland S 1998 *Modern Epidemiology*. 2nd edn. Philadelphia: Lipincott-Raven.
- Shiloh Y 2003 ATM and related protein kinases: safeguarding genome integrity. *Nature Reviews. Cancer* **3** 155–168.
- Siddique MM, Barlam C, Fiszler-Maliszewska L, Aggarwal A, Tan A, Tan P, Soo KC & Sabapathy K 2005 Evidence for selective expression of the p53 codon 72 polymorphs: implications in cancer development. *Cancer Epidemiology, Biomarkers & Prevention* **14** 2245–2252.
- Soussi T & Wiman KG 2007 Shaping genetic alterations in human cancer: the p53 mutation paradigm. *Cancer Cell* **12** 303–312.
- Stepanenko VF, Voilleque PG, Gavrilin YuI, Khrouch VT, Shinkarev SM, Orlov MYu, Kondrashov AE, Petin DV, Iaskova EK & Tsyb AF 2004 Estimating individual thyroid doses for a case–control study of childhood thyroid cancer in Bryansk oblast, Russia. *Radiation Protection Dosimetry* **108** 143–160.
- Sturgis EM, Zhao C, Zheng R & Wei Q 2005 Radiation response genotype and risk of differentiated thyroid cancer: a case–control analysis. *Laryngoscope* **115** 938–945.
- Tamura Y, Maruyama M, Mishima Y, Fujisawa H, Obata M, Kodama Y, Yoshikai Y, Aoyagi Y, Niwa O, Schaffner W *et al.* 2005 Predisposition to mouse thymic lymphomas in response to ionizing radiation depends on variant alleles encoding metal-responsive transcription factor-1 (Mtf-1). *Oncogene* **24** 399–406.
- Taylor RM, Thistlethwaite A & Caldecott KW 2002 Central role for the XRCC1 BRCT I domain in mammalian DNA single-strand break repair. *Molecular and Cellular Biology* **22** 2556–2563.
- Thorstenson YR, Roxas A, Kroiss R, Jenkins MA, Yu KM, Bachrich T, Muht D, Wayne TL, Chu G, Davis RW *et al.* 2003 Contributions of ATM mutations to familial breast and ovarian cancer. *Cancer Research* **63** 3325–3333.
- Wang J, Chuang K, Ahluwalia M, Patel S, Umblas N, Mirel D, Higuchi R & Germer S 2005 High-throughput SNP genotyping by single-tube PCR with T_m -shift primers. *BioTechniques* **39** 885–893.
- Williams ED 2006 Chernobyl and thyroid cancer. *Journal of Surgical Oncology* **94** 670–677.
- Yamashita S & Saenko V 2006 Mechanisms of disease: molecular genetics of childhood thyroid cancers. *Nature Clinical Practice. Endocrinology & Metabolism* **3** 422–429.
- Yang T, Namba H, Hara T, Takamura N, Nagayama Y, Fukata S, Ishikawa N, Kuma K, Ito K & Yamashita S 1997 p53 induced by ionizing radiation mediates DNA end-jointing activity, but not apoptosis of thyroid cells. *Oncogene* **14** 1511–1519.
- Yen CY, Liu SY, Chen CH, Tseng HF, Chuang LY, Yang CH, Lin YC, Wen CH, Chiang WF, Ho CH *et al.* 2008 Combinational polymorphisms of four DNA repair genes XRCC1, XRCC2, XRCC3, and XRCC4 and their association with oral cancer in Taiwan. *Journal of Oral Pathology & Medicine* **37** 271–277.
- Yoon YJ, Chang HY, Ahn SH, Kim JK, Park YK, Kang DR, Park JY, Myoung SM, Kim do Y, Chon CY *et al.* 2008 MDM2 and p53 polymorphisms are associated with the development of hepatocellular carcinoma in patients with chronic hepatitis B virus infection. *Carcinogenesis* **29** 1192–1196.

The Usual Ultrasonographic Features of Thyroid Cancer Are Less Frequent in Small Tumors That Develop After a Long Latent Period After the Chernobyl Radiation Release Accident

Valentina M. Drozd,¹ Maxim L. Lushchik,¹ Olga N. Polyanskaya,¹ Michail V. Fridman,¹
Yuri E. Demidchik,¹ Andrej P. Lyshchik,² Johannes Biko,³ Christoph Reiners,³
Yoshisada Shibata,⁴ Vladimir A. Saenko,⁵ and Shunichi Yamashita^{4,5}

Background: The Chernobyl accident resulted in an unprecedented number of radiation-induced thyroid cancers in young individuals as detected by national and international screening programs. The vast majority of thyroid malignancies were papillary carcinomas that, despite being similar by histopathology, displayed large variability in clinical course. The correlations between ultrasound (US) and clinicopathological features in young patients with radiation-induced thyroid cancer, however, have not been well studied. Because of the importance of US for deciding which subjects should have fine-needle aspiration biopsy, we assessed the US features of papillary thyroid carcinoma in patients exposed to Chernobyl fallouts.

Design: We performed a retrospective multivariate logistic regression analysis of US features, clinicopathological data, and the latency period between radiation exposure and the diagnosis of cancer in 94 patients who were 10.6–34.3 years old (16.5 ± 6.2 , mean \pm standard deviation) at the time of diagnosis and 0.1–18.0 years old (5.6 ± 4.2) at the time of the Chernobyl accident.

Results: Nodules greater than 10 mm were associated with the higher frequency of irregular margins ($p = 0.001$), longer period of latency ($p = 0.016$), and bilateral lymph node involvement ($p = 0.025$). Irregular tumor margins correlated with the shorter period of latency ($p = 0.009$) and unilateral nodal disease ($p = 0.010$). Hypoechoic nodules were observed more frequently in female patients ($p = 0.012$), in the absence of halo ($p = 0.003$) or calcifications ($p = 0.005$). Hypoechoic nodules also correlated with the shorter latency ($p = 0.015$) and younger age of patients ($p = 0.048$).

Conclusions: Irregular nodule margins, a usual sign of malignancy, are less useful in detecting thyroid cancers in radiation-exposed patients with tumors less than 10 mm. Thyroid cancers that are detected after longer latent periods display less of the US features characteristic of a malignant process, while benign US features are observed more frequently. Therefore, we recommend fine-needle aspiration biopsy to ensure early diagnosis of thyroid cancer for patients with a history of radiation exposure, even if their nodules are less than 10 mm.

Introduction

CHILDHOOD THYROID CANCER is rare, but early diagnosis is important in pediatric practice. Real-time ultrasound (US) examination is the most useful means of screening for thyroid cancer. US is widely available, has a low cost, requires short examination time, is not painful, and does not expose

the patient to radiation (1). When US is combined with US-guided fine-needle aspiration biopsy (FNAB) this can provide early diagnosis of thyroid disease in high-risk individuals exposed to radiation (2–5).

Among the two million children exposed to Chernobyl fallout in Belarus, over 2000 cases of thyroid cancer have been diagnosed in young patients during the 20-year period

¹Department of Thyroid Disease Research, Belarusian Medical Academy for Postgraduate Education, Minsk, Republic of Belarus.

²Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center, Nashville, Tennessee.

³Clinic and Polyclinic of Nuclear Medicine, University of Wuerzburg, Wuerzburg, Germany.

Departments of ⁴Molecular Medicine and ⁵International Health and Radiation Research, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki, Japan.

TABLE 1. ULTRASOUND FEATURES SUGGESTIVE OF A MALIGNANT OR BENIGN THYROID NODULE

| | Malignant (references) | Benign (references) |
|--------------------------|---|--|
| <i>Number of nodules</i> | | |
| Solitary | Frates <i>et al.</i> (8) | |
| Multiple | Frates <i>et al.</i> (8) | |
| <i>Structure</i> | | |
| Homogeneous | | |
| Heterogeneous | Hegedus (1), Drozd <i>et al.</i> (3) | |
| <i>Echogenicity</i> | | |
| Hypoechoic | Hegedus (1), Drozd <i>et al.</i> (3), Cooper <i>et al.</i> (9), Shimura <i>et al.</i> (10), Tae <i>et al.</i> (11) | Wienke <i>et al.</i> (12) |
| Isoechoic | Drozd <i>et al.</i> (3) | Hegedus (1) |
| Mixed | | |
| <i>Margins</i> | | |
| Regular | Drozd <i>et al.</i> (3) | Hegedus (1), Shimura <i>et al.</i> (10), Tae <i>et al.</i> (11) |
| Irregular | Hegedus (1), Drozd <i>et al.</i> (3), Cooper <i>et al.</i> (9), Shimura <i>et al.</i> (10), Tae <i>et al.</i> (11), Papini <i>et al.</i> (13), Koike <i>et al.</i> (14), Rago <i>et al.</i> (15), Chan <i>et al.</i> (16) | |
| <i>Halo</i> | | |
| Absent | Hegedus (1), Peccin <i>et al.</i> (17) | |
| Present | Drozd <i>et al.</i> (3) | Hegedus (1) |
| <i>Calcifications</i> | | |
| Absent | | Tae <i>et al.</i> (11) |
| Present | Hegedus (1), Cooper <i>et al.</i> (9) | Hegedus (1), Shimura <i>et al.</i> (10), Koike <i>et al.</i> (14) |
| <i>Lymphadenopathy</i> | | |
| Absent | | Hegedus (1) |
| Present | Hegedus (1), Drozd <i>et al.</i> (3), Pellegriti <i>et al.</i> (18) | |
| Hypoechoic | | |
| Isoechoic | Drozd <i>et al.</i> (3) | |

(1986–2006) after the accident (6). The mortality rate has been extremely low. This is probably due, at least in part, to US screening. Had the disaster occurred a decade before US was available, the mortality would probably have been substantially higher (7).

In adults the US features associated with malignancy are hypoechoogenicity, mostly nonhomogenous structure, calcifications, and irregular (microlobulated and blurred) margins (1,3,8–18) as summarized in Table 1. In the pediatric population the potential usefulness of these features is less well studied. Most of the data relating to the US features of thyroid malignancy are based on studies of sporadic cases. Less is known about the US features of radiation-associated thyroid cancer, particularly as it relates to the interval between radiation exposure and US.

In this study we retrospectively analyzed young patients with thyroid cancer who had been exposed to the accidental release of radiation from the Chernobyl reactor in 1986. We found that the US features of malignant nodules in this group were more characteristic of what has been described for thyroid cancer with increasing tumor size and less characteristic of thyroid cancer with increasing duration between radiation exposure and surgical diagnosis.

Materials and Methods

Patients

The study group included 94 patients (47 males and 47 females, sex ratio 1:1) from Belarus with papillary thyroid carcinoma for whom pathological findings, US imaging, and

demographic data were available at the database of Belarusian Medical Academy for Postgraduate Education. Seven additional cases in the database distributed evenly over the period of data acquisition (1990–2003) were not included because of excessively missing clinicopathological information.

The patients' age in the study ranged 0.1–18.0 years with the mean \pm standard deviation (SD) of 5.6 ± 4.2 at the time of the Chernobyl accident, and 10.6–34.3 years with the mean \pm SD of 16.5 ± 6.2 years at diagnosis. The latent period, defined as the interval between radiation exposure (April 26, 1986) and the first thyroid surgery, ranged from 45.6 to 207.6 months with the mean \pm SD of 128.4 ± 49.2 months.

US-guided FNAB followed by cytological analysis was routinely done for all thyroid nodules measuring 5 mm or greater on US. In all cases in this study the preoperative cytological diagnosis was papillary thyroid carcinoma or suspected thyroid carcinoma. All the patients were surgically treated at the Thyroid Cancer Center, Minsk. The cytological diagnosis of papillary thyroid carcinoma was confirmed by pathological examination after surgery.

Clinicopathological data were retrieved from medical records. The study protocol was approved by the Belarusian Medical Academy for Postgraduate Education and Nagasaki University Graduate School of Biomedical Sciences institutional review boards.

US examination

Thyroid nodules were initially detected during annual screenings in most radiocontaminated regions of Belarus.

TABLE 2. CLINICAL, ULTRASONOGRAPHIC, AND PATHOLOGICAL PARAMETERS BY NODULE SIZE

| Parameters | Group | | p-Value ^a |
|---|---------------------------|---------------------------|----------------------|
| | Nodule size ≤10 mm n = 25 | Nodule size >10 mm n = 69 | |
| <i>Clinical characteristics</i> | | | |
| Sex | | | |
| Male | 11 (44.0) ^b | 36 (52.2) | 0.64 |
| Female | 14 (56.0) | 33 (47.8) | |
| Age at the time of accident (years) | 5.0 ± 3.9 ^c | 5.7 ± 4.9 | 0.88 |
| Age at the time of diagnosis (years) | 14.8 ± 5.9 | 17.11 ± 6.6 | 0.14 |
| Latent period (months) | 117.9 ± 46.6 | 133.6 ± 48.7 | 0.089 |
| <i>Ultrasonographic characteristics</i> | | | |
| Nodule volume (mL) | 0.2 ± 0.1 | 1.6 ± 2.5 | <0.001 |
| Number of nodules | | | |
| Solitary | 21 (84.0) | 59 (85.5) | 1.00 |
| Multiple | 4 (16.0) | 10 (14.5) | |
| Structure | | | |
| Homogeneous | 14 (58.3) | 26 (37.7) | 0.097 |
| Heterogeneous | 10 (41.7) | 43 (62.3) | |
| Unknown | 1 | 0 | |
| Echogenicity | | | |
| Hypoechoic | 18 (75.0) | 36 (52.9) | 0.18 |
| Isoechoic | 4 (16.7) | 18 (26.5) | |
| Mixed | 2 (8.3) | 14 (20.6) | |
| Unknown | 1 | 1 | |
| Margins | | | |
| Regular | 10 (43.5) | 10 (14.7) | 0.009 |
| Irregular | 13 (56.5) | 58 (85.3) | |
| Unknown | 2 | 1 | |
| Halo | | | |
| Absent | 22 (88.0) | 56 (85.5) | 1.00 |
| Present | 3 (12.0) | 10 (14.5) | |
| Unknown | 0 | 3 | |
| Calcifications | | | |
| Absent | 25 (100.0) | 59 (85.5) | 0.058 |
| Present | 0 (0.0) | 10 (14.9) | |
| Lymphadenopathy | | | |
| Absent | 15 (60.0) | 39 (56.5) | 0.82 |
| Present | 10 (40.0) | 30 (43.5) | |
| Hypoechoic | 9 (36.0) | 21 (30.4) | |
| Isoechoic | 1 (4.0) | 8 (11.6) | |
| Unknown | 0 | 1 | |
| <i>Pathological characteristics</i> | | | |
| Tumor size (mm) ^d | 7.8 ± 2.6 | 13.2 ± 5.3 | <0.001 |
| T stage | | | |
| T1 | 24 (96.0) | 63 (91.3) | 0.64 |
| T2 | 0 (0.0) | 4 (5.8) | |
| T3 | 1 (4.0) | 2 (2.9) | |
| N stage | | | |
| N0 | 13 (52.0) | 20 (29.4) | 0.003 |
| N1a | 10 (40.0) | 18 (26.5) | |
| N1b | 2 (8.0) | 30 (44.1) | |
| Unknown | 0 | 1 | |
| Histopathological variant | | | |
| Solid | 3 (12.0) | 6 (8.9) | 0.99 |
| Papillary and follicular | 22 (88.0) | 61 (91.1) | |
| Unknown | 0 | 2 | |
| Invasiveness | | | |
| Intrathyroidal | 21 (84.0) | 60 (86.9) | 0.74 |
| Extrathyroidal | 4 (16.0) | 9 (13.1) | |

^aBased on Wilcoxon rank-sum or Kruskal-Wallis test for continuous variables and Fisher's exact test for categorical variables; patients without available information were excluded.

^bNumber of patients with percentage within parentheses.

^cMean ± standard deviation.

^dLargest measurement.

TABLE 3. PARAMETERS ASSOCIATED WITH THE FREQUENCY OF LARGE NODULE SIZE (>10 MM), LOGISTIC REGRESSION ANALYSIS

| Parameters | Comparison | Odds ratio | 95% CI | p-Value ^a |
|-------------------|-----------------------|------------|-----------|----------------------|
| Number of nodules | Singular vs. multiple | 0.24 | 0.05–1.12 | 0.071 |
| Latent period | By 1-month increment | 1.02 | 1.00–1.03 | 0.016 |
| Margins | Irregular vs. regular | 12.75 | 2.99–69.3 | 0.001 |
| N stage | N1a vs. N0 | 0.51 | 0.13–1.92 | 0.331 |
| | N1b vs. N0 | 7.08 | 1.48–48.7 | 0.025 |

^aBased on likelihood ratio test. CI, confidence interval.

Individuals with US thyroid abnormalities underwent health examination, including thyroid ultrasonography, at the hospital. US examination was done with a Toshiba SSA 240A (7.5 MHz sector probe) ultrasonic real-time scanner, with patient in the supine position with the neck slightly hyperextended.

For the purpose of this study, archived US images were reanalyzed by experienced radiologists (V.M.D., M.L.L., and A.P.L.) using identical criteria. Sonograms were interpreted in terms of nodule size (largest dimension), number of nodules (solitary and multiple), echogenicity (hypoechoic, isoechoic, or mixed), echostructure (homogeneous or heterogeneous), margins (regular or irregular), presence or absence of a halo (rim), calcifications, and regional lymphadenopathy. A consensus estimate was reached for every parameter in all cases under study.

Thyroid and nodule volumes (mL) were estimated, using the height (*H*), width (*W*), and length (*L*) of the thyroid lobes or of a nodule measured in the transverse and longitudinal scans (cm) by the formula for ellipsoid volume: $V = H \times W \times L \times \pi/6$ (19).

Histopathology

All cases were pathologically diagnosed as papillary carcinomas. Solid variant of papillary thyroid carcinoma (less mature histotype) was observed in nine (9.6%) cases. Typical papillary morphology and follicular variant of papillary thyroid carcinoma (more differentiated histotypes) were documented in 85 (90.4%) cases. Tumor size varied from 5 to 30 mm with the mean \pm SD of 11.7 ± 3.9 mm. The presence or absence of capsular and extrathyroidal invasion was recorded. Tumor staging was according to the International Union Against Cancer (UICC) TNM Classification of Malignant Tumors (20).

Statistical analysis

All patients were divided into subgroups according to tumor size on ultrasonography (≤ 10 mm and > 10 mm), type of margins (regular or irregular), and echogenicity (hypoechoic, isoechoic, or mixed). Subgroups thus defined were compared with respect to clinical, ultrasonographic, or pathological characteristics by Fisher exact test for categorical

data, and by Wilcoxon rank-sum test or Kruskal-Wallis test for quantitative measurements. Association of these characteristics with tumor size, type of margins, or echogenicity was evaluated by logistic regression analysis, where the most appropriate model was selected by Akaike information criterion starting from the full model. The FREQ, NPAR1WAY, and LOGISTIC procedures in the SAS system (21) were used for the calculations. A *p*-value less than 0.05 was regarded as indicating statistical significance.

Results

Clinical, ultrasonic, and pathological characteristics with respect to nodule size by US

The tumor size on ultrasonography was ≤ 10 mm in 25 patients and > 10 mm in 69 patients (Table 2). No significant difference was observed between the two groups in age at the time of accident, age at diagnosis, or sex ratio. Although not significant, the latent period tended to be longer in the patients with larger nodules.

Irregular margins were detected more frequently in the patients with larger nodules ($p = 0.009$). There was a very strong trend toward the frequency of calcifications being higher in the patients with larger nodules ($p = 0.058$). No significant differences were observed in other US characteristics between the two subgroups. By definition, tumors smaller than 1 cm are classified as pT1. However, 91% of the larger tumors in this series also were classified as pT1 (with diameter smaller than 2 cm). There were no pT4 tumors; the low prevalence of advanced stage tumors is most probably due to extensive screenings. Lymph node involvement (on pathology) significantly correlated with the larger ultrasonic nodule size ($p = 0.003$). No significant difference was observed between the two subgroups regarding the frequency of histopathological variants or tumor invasiveness.

Three factors—latent period, margins, and N stage—were found by logistic regression analysis to be associated with the frequency of larger tumor size (Table 3). The frequency of larger tumors was significantly increased with the latency ($p = 0.016$), was significantly higher among the nodules with irregular margins ($p = 0.001$), and was significantly higher in the patients with bilateral lymph node involvement ($p = 0.025$).

Clinical, ultrasonic, and pathological characteristics with respect to nodule margins

Tumor margins were regular or irregular in 20 and 72 patients, respectively (Table 4). The age at diagnosis was significantly lower ($p = 0.006$) and the latency period was significantly shorter ($p = 0.021$) in the patients with irregular margins of cancer nodules.

A heterogeneous US structure was observed more frequently in nodules with irregular margins ($p = 0.022$). US signs of lymph node involvement as well as N1a tumor stage were significantly more frequent in the patients with nodules with irregular margins ($p = 0.022$ and $p = 0.025$, respectively). The frequency of isoechoic lymph nodes, not considered a suspicious sign, also tended to correlate with the irregular margins of thyroid nodules, although the significance was marginal ($p = 0.064$). An isoechoic pattern and the presence of

TABLE 4. CLINICAL, ULTRASONOGRAPHIC, AND PATHOLOGICAL PARAMETERS BY TUMOR MARGIN TYPE

| Parameters | Group | | p-Value ^a |
|---|------------------------|------------------------|----------------------|
| | Regular margins n=20 | Irregular margins n=72 | |
| <i>Clinical characteristics</i> | | | |
| Sex | | | |
| Male | 11 (55.0) ^b | 36 (50.0) | |
| Female | 9 (45.0) | 36 (50.0) | 0.80 |
| Age at the time of accident (years) | 7.2 ± 6.1 ^c | 5.1 ± 4.2 | 0.27 |
| Age at the time of diagnosis (years) | 20.2 ± 7.5 | 15.3 ± 5.9 | 0.006 |
| Latent period (months) | 155.9 ± 34.3 | 122.7 ± 49.9 | 0.021 |
| <i>Ultrasonographic characteristics</i> | | | |
| Nodule size (mm) ^d | 12.6 ± 5.3 | 15.4 ± 6.7 | 0.052 |
| Nodule volume (mL) | 0.7 ± 0.8 | 1.4 ± 2.5 | 0.019 |
| Number of nodules | | | |
| Solitary | 17 (85.0) | 60 (83.3) | |
| Multiple | 3 (15.0) | 12 (16.7) | 1.00 |
| Structure | | | |
| Homogeneous | 13 (65.0) | 25 (34.7) | |
| Heterogeneous | 7 (35.0) | 47 (65.3) | 0.022 |
| Echogenicity | | | |
| Hypoechoic | 9 (45.0) | 44 (61.1) | |
| Isoechoic | 9 (45.0) | 13 (18.1) | |
| Mixed | 2 (10.0) | 15 (20.8) | 0.051 |
| Halo | | | |
| Absent | 14 (70.0) | 65 (90.3) | |
| Present | 6 (30.0) | 7 (9.7) | 0.029 |
| Calcifications | | | |
| Absent | 18 (90.0) | 64 (88.9) | |
| Present | 2 (10.0) | 8 (11.1) | 1.00 |
| Lymphadenopathy | | | |
| Absent | 16 (80.0) | 37 (51.4) | |
| Present | 4 (20.0) | 35 (48.6) | 0.022 |
| Echogenicity | | | |
| Hypoechoic | 4 (20.0) | 26 (36.1) | |
| Isoechoic | 0 (0.0) | 8 (11.1) | 0.064 |
| Unknown | 0 | 1 | |
| <i>Pathological characteristics</i> | | | |
| Tumor size (mm) ^d | 10.3 ± 3.5 | 12.4 ± 5.6 | 0.15 |
| T stage | | | |
| T1 | 20 (100.0) | 64 (89.0) | |
| T2 | 0 (0.0) | 4 (5.5) | |
| T3 | 0 (0.0) | 4 (5.5) | 0.50 |
| N stage | | | |
| N0 | 12 (60.0) | 20 (28.2) | |
| N1a | 2 (10.0) | 24 (33.8) | |
| N1b | 6 (30.0) | 27 (38.0) | 0.025 |
| Unknown | 0 | 1 | |
| Histopathological variant | | | |
| Solid | 1 (5.0) | 9 (12.5) | |
| Papillary and follicular | 19 (95.0) | 63 (87.5) | 0.70 |
| Invasiveness | | | |
| Intrathyroidal | 18 (90.0) | 60 (83.3) | |
| Extrathyroidal | 2 (10.0) | 12 (16.7) | 0.72 |

^aBased on Wilcoxon rank-sum or Kruskal-Wallis test for continuous variables and Fisher's exact test for categorical variables; patients without available information were excluded.

^bNumber of patients with percentage within parentheses.

^cMean ± standard deviation.

^dLargest measurement.

halo were more frequent in the nodules with regular margins ($p = 0.051$ and $p = 0.029$, respectively).

Three factors, that is, latent period, nodule size, and N stage, were found by logistic regression analysis to be associated with the frequency of irregular margins (Table 5). Irregular

margins were decreased in frequency with longer latency ($p = 0.009$), were significantly frequent in the larger nodules (>10 mm) as compared to smaller ones (≤ 10 mm; $p = 0.042$), and were observed significantly more frequently in the patients with unilateral lymph node involvement ($p = 0.010$).

TABLE 5. FACTORS ASSOCIATED WITH THE FREQUENCY OF IRREGULAR TYPE OF MARGINS, LOGISTIC REGRESSION ANALYSIS

| Factor | Comparison | Odds ratio | 95% CI | p-Value ^a |
|---------------------------|-------------------------------|------------|-----------|----------------------|
| Age at accident | By 1-year increment | 0.95 | 0.84–1.08 | 0.434 |
| Latent period | By 1-month increment | 0.98 | 0.96–0.99 | 0.009 |
| Nodule size on ultrasound | Larger than 10 mm vs. smaller | 1.17 | 1.02–1.37 | 0.042 |
| N stage | N1a vs. N0 | 10.88 | 2.09–89.9 | 0.010 |
| | N1b vs. N0 | | | 0.072 |

^aBased on the likelihood ratio test.
CI, confidence interval.

Clinical, ultrasonic, and pathological characteristics with respect to nodule echogenicity

Tumors were hypoechoic in 54 patients, isoechoic in 22, and mixed in 17 (Table 6). The age at diagnosis was significantly higher ($p = 0.048$) and latent period was significantly longer ($p = 0.015$) in the patients with isoechoic tumors (usually no sign of malignancy) than in those with hypoechoic or mixed-type tumors. The frequency of isoechoic tumors was significantly higher in male patients ($p = 0.003$).

Heterogeneous US structure was significantly more frequent in mixed-type tumors ($p = 0.021$). Irregular margins were more frequent in hypoechoic or mixed-type tumors ($p = 0.051$). The presence of halo was significantly more frequent in isoechoic tumors ($p < 0.001$), while calcifications occurred more frequently in the mixed-type tumors ($p < 0.001$). Tumor size on pathology was significantly larger in the mixed-type tumors ($p = 0.023$).

Three factors—sex, halo, and calcification—were found by logistic regression analysis to be associated with the frequency of hypoechoic tumors (Table 7). It was significantly higher in female patients ($p = 0.012$), in the nodules without halo ($p = 0.003$) or calcifications ($p = 0.005$).

Discussion

Easy detection of thyroid nodules on US, besides of obvious benefits, brings about a variety of associated problems. These are the differential diagnosis, follow-up of a great number of small nodules occurring in 3–50% of population, and establishment of evidence-based criteria for FNAB in different age groups (1,2,7,22). Regarding the latter, the Consensus of Society of Radiologists in US and the American Thyroid Association recommend FNAB for the nodules measuring 1 cm or greater because of "uncertainty as to whether diagnosis of smaller cancers improves life expectancy as well as a concern that assessment of small nodules would lead to an excessive number of biopsies" (9,22). Nodules not exceeding 1 cm in diameter are recommended for follow-up or evaluation in case of suspicious US findings, history of head and neck irradiation, or documented thyroid cancer in consanguine relative(s). Recent guidelines of the American Association of Clinical Endocrinologists and Associazione Medici Endocrinologi also recommend evaluation of thyroid nodules smaller than 1 cm in radiation-exposed patients (23). With regard to the young individuals who were internally irradiated by iodine isotopes, substantiated recommendations for management of small thyroid nodules need further study, in our opinion.

In the present work we specifically focused on a group of young patients with radiation-induced papillary thyroid carcinoma. Nodules in all cases had been initially detected due to the implementation of post-Chernobyl US screening programs in Belarus; all nodules exceeding 5 mm were subjected to FNAB and cytological examination. Unlike in other studies, here we not addressed the differential cancer diagnosis but attempted to determine clinicopathological and ultrasonic criteria with regard to tumor size, ultrasonographic margin type, and nodule echogenicity, that is, with the most important US parameters taken into account for clinical decision making.

The results of analyses demonstrated an association of tumor size (under vs. over 10 mm) with tumor margin type and bilateral lymph node involvement. With increasing nodule size, irregular tumor margins (a usual suspicious sign) and bilateral lymph node involvement (an indicator of tumor aggressiveness) became more frequent. Similar to our findings, Pellegriti *et al.* reported a progressive accumulation of signs of tumor aggressiveness (multifocality, bilaterality, extrathyroidal invasion, and nodal disease) with increasing nodule size at presentation (18). Also, in patients with cancers larger than 1 cm, a close relationship between tumor size and aggressiveness has been demonstrated (24–26). Apparently, smaller tumors display more benign characteristics on US, but these should not be misleading as nodule size *per se* is not a predictor of malignancy (8,13,27) and may not necessarily correlate with an indolent clinical course or lower risk of recurrence (10,18).

Type of margins in our series correlated not only with tumor size, but also with other US features, in particular lymph node involvement and duration of latency. The frequency of irregular margins (a usual suspicious sign) was increasing in parallel with those of heterogeneous structure, hypoechoic or mixed echogenicity (all worrisome), and lymphadenopathy (tumor aggressiveness). Patients with irregular tumor margins were younger at diagnosis and had shorter latent period and more aggressive tumors. Of note, from the molecular studies of Chernobyl thyroid cancers, it is known that mentioned characteristics correlate with the type of genetic abnormality underlying malignancy. RET/PTC3 rearrangements were particularly prevalent in the aggressive tumors developing after the shorter latency; these tumors frequently had solid morphological structure (28,29). In our series, solid morphology was more frequently observed in the tumors with irregular margins (as well as in the hypoechoic ones), but significance of difference was not reached perhaps because of the small number of such cases. It would

TABLE 6. CLINICAL, ULTRASONOGRAPHIC, AND PATHOLOGICAL PARAMETERS BY TUMOR ECHOGENICITY

| Parameters | Group | | | p-Value ^a |
|---|------------------------|------------------|--------------|----------------------|
| | Hypoechoic n = 54 | Isoechoic n = 22 | Mixed n = 17 | |
| <i>Clinical characteristics</i> | | | | |
| Sex | | | | |
| Male | 22 (40.7) ^b | 18 (81.8) | 7 (41.2) | 0.003 |
| Female | 32 (59.3) | 4 (18.2) | 10 (58.8) | |
| Age at the time of accident (years) | 5.2 ± 4.1 ^c | 6.7 ± 6.2 | 5.1 ± 4.2 | 0.87 |
| Age at the time of diagnosis (years) | 14.9 ± 5.8 | 19.4 ± 8.1 | 16.8 ± 5.5 | 0.048 |
| Latent period (months) | 117.7 ± 47 | 151.9 ± 45 | 139.9 ± 47 | 0.015 |
| <i>Ultrasonographic characteristics</i> | | | | |
| Nodule size (mm) ^d | 14.7 ± 7.1 | 14.7 ± 4.8 | 16.6 ± 6.7 | 0.13 |
| Nodule volume, mL | 1.2 ± 1.90 | 1.0 ± 0.87 | 1.9 ± 4.1 | 0.32 |
| Number of nodules | | | | |
| Solitary | 48 (88.9) | 17 (77.3) | 13 (76.5) | 0.38 |
| Multiple | 6 (11.1) | 5 (22.7) | 4 (23.5) | |
| Structure | | | | |
| Homogeneous | 25 (47.2) | 11 (50.0) | 2 (11.8) | 0.021 |
| Heterogeneous | 28 (52.8) | 11 (50.0) | 15 (88.2) | |
| Unknown | 1 | 0 | 0 | |
| Margins | | | | |
| Regular | 9 (17.0) | 9 (40.9) | 2 (11.8) | 0.051 |
| Irregular | 44 (83.0) | 13 (59.1) | 15 (88.2) | |
| Unknown | 1 | 0 | 0 | |
| Halo | | | | |
| Absent | 53 (98.2) | 12 (54.5) | 15 (88.2) | <0.001 |
| Present | 1 (1.8) | 10 (45.5) | 2 (11.8) | |
| Calcifications | | | | |
| Absent | 53 (98.2) | 18 (81.8) | 12 (70.6) | <0.001 |
| Present | 1 (1.8) | 4 (18.2) | 5 (29.4) | |
| Lymphadenopathy | | | | |
| Absent | 27 (50.0) | 15 (68.2) | 11 (64.7) | 0.29 |
| Present | 27 (50.0) | 7 (31.8) | 6 (35.3) | |
| Hypoechoic | 20 (37.0) | 6 (27.3) | 1 (4.5) | |
| Isoechoic | 6 (11.1) | 1 (4.5) | 1 (5.9) | |
| Unknown | 1 | 0 | 4 | |
| <i>Pathological characteristics</i> | | | | |
| Tumor size (mm) ^d | 11.0 ± 5.6 | 11.7 ± 4.1 | 14.4 ± 5.4 | 0.023 |
| T stage | | | | |
| T1 | 50 (92.6) | 22 (100.0) | 13 (76.5) | 0.063 |
| T2 | 3 (5.6) | 0 (0.0) | 1 (5.9) | |
| T3 | 1 (1.8) | 0 (0.0) | 3 (17.6) | |
| N stage | | | | |
| N0 | 23 (43.4) | 6 (27.3) | 4 (23.5) | 0.14 |
| N1a | 17 (32.1) | 5 (22.7) | 4 (23.5) | |
| N1b | 13 (24.5) | 11 (50.0) | 9 (53.0) | |
| Unknown | 1 | 0 | 0 | |
| Histopathological variant | | | | |
| Solid | 7 (13.0) | 1 (4.5) | 2 (11.8) | 0.86 |
| Papillary and follicular | 47 (87.0) | 21 (95.5) | 15 (88.2) | |
| Invasiveness | | | | |
| Intrathyroidal | 46 (85.2) | 19 (86.4) | 14 (82.4) | 0.92 |
| Extrathyroidal | 8 (14.8) | 3 (13.6) | 2 (17.6) | |
| Unknown | 0 | 0 | 1 | |

^aBased on Wilcoxon rank-sum or Kruskal-Wallis test for continuous variables and Fisher's exact test for categorical variables; patients without available information were excluded.

^bNumber of patients with percentage within parentheses.

^cMean ± standard deviation.

^dLargest measurement.