

gests that the core and envelope proteins have only a weak effect on IFN resistance.

In clinical practice, HBV shows high resistance against IFN therapy. This is also the case in the cell culture system, as we showed in this study and has been reported in previous studies [20,28]. The mechanism by which hepatitis viruses resist IFN needs to be clarified in order to develop new and effective therapies for eradication of these viruses.

Acknowledgments

The authors thank Yoshie Yoshida, Kazuyo Hattori, and Rie Akiyama for their excellent technical help.

This study was supported in part by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Labor and Health and Welfare.

References

- [1] Maddrey WC. Hepatitis B: an important public health issue. *J Med Virol* 2000;61:362–366.
- [2] Global surveillance and control of hepatitis C. Report of a WHO Consultation organized in collaboration with the Viral Hepatitis Prevention Board, Antwerp, Belgium. *J Viral Hepat* 1999;6:35–47.
- [3] Beasley RP. Hepatitis B virus. The major etiology of hepatocellular carcinoma. *Cancer* 1988;61:1942–1956.
- [4] Samuel CE. Antiviral actions of interferons. *Clin Microbiol Rev* 2001;14:778–809.
- [5] Sagnelli E, Coppola N, Scolastico C, Filippini P, Santantonio T, Stroffolini T, et al. Virologic and clinical expressions of reciprocal inhibitory effect of hepatitis B, C, and delta viruses in patients with chronic hepatitis. *Hepatology* 2000;32:1106–1110.
- [6] Chiaramonte M, Stroffolini T, Vian A, Stazi MA, Floreani A, Lorenzoni U, et al. Rate of incidence of hepatocellular carcinoma in patients with compensated viral cirrhosis. *Cancer* 1999;85:2132–2137.
- [7] Cacciola I, Pollicino T, Squadrito G, Cerenzia G, Orlando ME, Raimondo G. Occult hepatitis B virus infection in patients with chronic hepatitis C liver disease. *N Engl J Med* 1999;341:22–26.
- [8] Raimondo G, Brunetto MR, Pontisso P, Smedile A, Maina AM, Saitta C, et al. Longitudinal evaluation reveals a complex spectrum of virological profiles in hepatitis B virus/hepatitis C virus-coinfected patients. *Hepatology* 2006;43:100–107.
- [9] Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Gonçales Jr FL, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–982.
- [10] Liu CJ, Lai MY, Chao YC, Liao LY, Yang SS, Hsiao TJ, et al. Interferon alpha-2b with and without ribavirin in the treatment of hepatitis B e antigen-positive chronic hepatitis B: a randomized study. *Hepatology* 2006;43:742–749.
- [11] Gotoh B, Komatsu T, Takeuchi K, Yokoo J. Paramyxovirus strategies for evading the interferon response. *Rev Med Virol* 2002;12:337–357.
- [12] Evans JD, Seeger C. Cardif: a protein central to innate immunity is inactivated by the HCV NS3 serine protease. *Hepatology* 2006;43:615–617.
- [13] Blindenbacher A, Duong FH, Hunziker L, Stutvoet ST, Wang X, Terracciano L, et al. Expression of hepatitis C virus proteins inhibits interferon alpha signaling in the liver of transgenic mice. *Gastroenterology* 2003;124:1465–1475.
- [14] Bode JG, Ludwig S, Ehrhardt C, Albrecht U, Erhardt A, Schaper F, et al. IFN-alpha antagonistic activity of HCV core protein involves induction of suppressor of cytokine signaling-3. *FASEB J* 2003;17:488–490.
- [15] Taylor DR, Shi ST, Romano PR, Barber GN, Lai MM. Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein. *Science* 1999;285:107–110.
- [16] Frese M, Pietschmann T, Moradpour D, Haller O, Bartenschlager R. Interferon-alpha inhibits hepatitis C virus subgenomic RNA replication by an MxA-independent pathway. *J Gen Virol* 2001;82:723–733.
- [17] Christen V, Duong F, Bernsmeier C, Sun D, Nassal M, Heim MH. Inhibition of alpha interferon signaling by hepatitis B virus. *J Virol* 2007;81:159–165.
- [18] Su AI, Pezacki JP, Wodicka L, Brideau AD, Supekova L, Thimme R, et al. Genomic analysis of the host response to hepatitis C virus infection. *Proc Natl Acad Sci USA* 2002;99:15669–15674.
- [19] Wieland S, Thimme R, Purcell RH, Chisari FV. Genomic analysis of the host response to hepatitis B virus infection. *Proc Natl Acad Sci USA* 2004;101:6669–6674.
- [20] Bellecave P, Gouttenoire J, Gajer M, Brass V, Koutsoudakis G, Blum HE, et al. Hepatitis B and C virus coinfection: a novel model system reveals the absence of direct viral interference. *Hepatology* 2009;50:46–55.
- [21] Tsuge M, Hiraga N, Takaishi H, Noguchi C, Oga H, Imamura M, et al. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology* 2005;42:1046–1054.
- [22] Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 2005;11:791–796.
- [23] Hiraga N, Imamura M, Tsuge M, Noguchi C, Takahashi S, Iwao E, et al. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis C virus and its susceptibility to interferon. *FEBS Lett* 2007;581:1983–1987.
- [24] Tateno C, Yoshizane Y, Saito N, Kataoka M, Utoh R, Yamasaki C, et al. Near completely humanized liver in mice shows human-type metabolic responses to drugs. *Am J Pathol* 2004;165:901–912.
- [25] Noguchi C, Ishino H, Tsuge M, Fujimoto Y, Imamura M, Takahashi S, et al. G to A hypermutation of hepatitis B virus. *Hepatology* 2005;41:626–633.
- [26] Rodríguez-Iñigo E, Bartolomé J, Ortiz-Movilla N, Platero C, López-Alcorocho JM, Pardo M, et al. Hepatitis C virus (HCV) and hepatitis B virus (HBV) can coinfect the same hepatocyte in the liver of patients with chronic HCV and occult HBV infection. *J Virol* 2005;79:15578–15581.
- [27] Blight KJ, Kolykhalov AA, Rice CM. Efficient initiation of HCV RNA replication in cell culture. *Science* 2000;290:1972–1974.
- [28] Hayashi Y, Koike K. Interferon inhibits hepatitis B virus replication in a stable expression system of transfected viral DNA. *J Virol* 1989;63:2936–2940.
- [29] Miyamoto M, Kato T, Date T, Mizokami M, Wakita T. Comparison between subgenomic replicons of hepatitis C virus genotypes 2a (JFH-1) and 1b (Con1 NK5.1). *Intervirology* 2006;49:37–43.



Adoptive immunotherapy with liver allograft–derived lymphocytes induces anti-HCV activity after liver transplantation in humans and humanized mice

Masahiro Ohira,^{1,2} Kohei Ishiyama,^{1,2} Yuka Tanaka,^{1,2} Marlen Doskali,^{1,2} Yuka Igarashi,^{1,2} Hirotaka Tashiro,^{1,2} Nobuhiko Hiraga,^{2,3} Michio Imamura,^{2,3} Naoya Sakamoto,⁴ Toshimasa Asahara,^{1,2} Kazuaki Chayama,^{2,3} and Hideki Ohdan^{1,2}

¹Department of Surgery, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, ²Liver Research Project Center, and ³Department of Medicine and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, Minami-ku, Hiroshima, Japan. ⁴Department of Gastroenterology and Hepatology, Tokyo Medical and Dental University, Bunkyo-ku, Tokyo, Japan.

After liver transplantation in HCV-infected patients, the virus load inevitably exceeds pre-transplantation levels. This phenomenon reflects suppression of the host-effector immune responses that control HCV replication by the immunosuppressive drugs used to prevent rejection of the transplanted liver. Here, we describe an adoptive immunotherapy approach, using lymphocytes extracted from liver allograft perfusate (termed herein liver allograft–derived lymphocytes), which includes an abundance of NK/NKT cells that mounted an anti-HCV response in HCV-infected liver transplantation recipients, despite the immunosuppressive environment. This therapy involved intravenously injecting patients 3 days after liver transplantation with liver allograft–derived lymphocytes treated with IL-2 and the CD3-specific mAb OKT3. During the first month after liver transplantation, the HCV RNA titers in the sera of recipients who received immunotherapy were markedly lower than those in the sera of recipients who did not receive immunotherapy. We further explored these observations in human hepatocyte–chimeric mice, in which mouse hepatocytes were replaced by human hepatocytes. These mice unfailingly developed HCV infections after inoculation with HCV-infected human serum. However, injection of human liver–derived lymphocytes treated with IL-2/OKT3 completely prevented HCV infection. Furthermore, an *in vitro* study using genomic HCV replicon–containing hepatic cells revealed that IFN- γ -secreting cells played a pivotal role in such anti-HCV responses. Thus, our study presents what we believe to be a novel paradigm for the inhibition of HCV replication in HCV-infected liver transplantation recipients.

Introduction

Liver failure and hepatocellular carcinoma (HCC) due to chronic hepatitis C infection are the most common indications for liver transplantation (LT), and the incidences of both have been projected to increase further in the future. Recurrent HCV infection of the allograft is universal, occurs immediately after LT, and is associated with accelerated progression to cirrhosis, graft loss, and death (1, 2). This reflects the suppression of those host-effector immune responses that usually control HCV replication, suggesting that the immunosuppressive environment may play a major role in the rapid progression of recurrent HCV infection after LT (3, 4). Further, the immunosuppressive condition described above is considered to increase the incidence of cancer recurrence after LT in HCC patients. We recently proposed the novel strategy of adjuvant immunotherapy for preventing the recurrence of HCC after LT; this immunotherapy involves intravenously injecting LT recipients with activated liver allograft–derived NK cells (5, 6). Since the immunosuppressive regimen currently used after LT reduces the adaptive immune components but effectively maintains the innate components of cellular immunity (7–9), the augmenta-

tion of the NK cell response, which is thought to play a pivotal role in innate immunity, may be a promising immunotherapeutic approach (6). We confirmed that the IL-2/anti-CD3 mAb–treated (IL-2/OKT3–treated) liver allograft–derived NK cells expressed a significantly high level of the tumor necrosis factor–related apoptosis-inducing ligand (TRAIL), which is a critical molecule for tumor cell killing. Further, these cells showed high cytotoxicity against HCC cells, with no such effect on normal cells (5). After obtaining approval from the ethical committee of our institute, we successfully administered adoptive immunotherapy with IL-2/OKT3–treated liver lymphocytes to liver cirrhosis patients with HCC in a phase I trial. Although the long-term benefits of this approach with regard to the control of HCC recurrence after LT remain to be elucidated, this trial provided a unique opportunity to study whether the adoptive administration of IL-2/OKT3–treated liver lymphocytes could also mount an anti-HCV response in HCV-infected LT recipients.

Previous studies have highlighted the important roles of innate lymphocytes in developing immunity against hepatotropic viruses, including HCV (10, 11). In this regard, it is known that patients with chronic HCV infection show diminished NK and NKT cell responses (12–14). In the case of an LT, it has recently been reported that the host CD56⁺ innate lymphocyte population,

Conflict of interest: The authors have declared that no conflict of interest exists.

Citation for this article: *J. Clin. Invest.* doi:10.1172/JCI38374.




Figure 1

Schematic outline of adoptive immunotherapy with lymphocytes extracted from liver allograft perfusate. The therapy involved giving an intravenous injection of IL-2/OKT3–treated liver lymphocytes to LT recipients. The lymphocytes were extracted from the donor liver graft perfusate. After 3 days of culture with IL-2 (100 JRU/ml), the activated liver NK cell–enriched lymphocytes were administered to the LT recipients through venous circulation. OKT3 (1 µg/ml) was added to the culture medium 1 day before this administration in order to prevent GVHD.

consisting of NK and NKT cells, is appreciably associated with the severity of HCV recurrence after LT (15). These insights into the immunopathogenesis of HCV recurrence indicate that the innate immune components mentioned above are potential targets for therapeutic manipulation. In this study, we have demonstrated for the first time to our knowledge that adoptive immunotherapy with IL-2/OKT3–treated liver lymphocytes, including abundant NK and NKT cells, shows anti-HCV activity after LT, even in an immunosuppressive environment.

Results

Adoptive transfer of IL-2/OKT3–treated liver lymphocytes. The human liver contains a significant number of resident lymphocytes. These cells include abundant CD56⁺ NK and NKT cells, many of which differ phenotypically and functionally from the circulating cells (14, 16). In our previous study, we performed ex vivo perfusion of the liver through the portal vein, which was necessary in order to flush blood from the liver graft before implantation. Liver-resident lymphocytes were then extracted from the perfusates (number of lymphocytes extracted from normal liver perfusates, 0.5 ± 0.1 cells per gram of liver weight; $n = 14$) (5). Proportions of CD56⁺CD3⁻NK cells and CD56⁺CD3⁺ NKT cells among the lymphocytes extracted from the liver perfusates (NK cells, $46.4\% \pm 4.2\%$; NKT cells, $17.2\% \pm 2.3\%$; $n = 14$) were significantly ($P < 0.05$) higher than those among the lymphocytes derived from the peripheral blood of the same donors (NK cells, $21.9\% \pm 3.7\%$; NKT cells, $3.8\% \pm 0.9\%$; $n = 14$). Extensive preclinical studies have shown that liver allograft–derived resident NK cells mediate remarkably higher cytotoxic activity against HCC cells than do peripheral blood NK cells (5). On this basis, we undertook a clinical trial of adjuvant immunotherapy with IL-2/OKT3–treated liver lymphocytes for preventing the recurrence of HCC after LT in 14 recipients with HCC (Figure 1 and Tables 1 and 2). The therapy involved administering a single intravenous injection of IL-2/OKT3–treated liver lymphocytes to recipients 3 days after LT ($2\text{--}5 \times 10^8$ cells injected per subject). In order to prevent graft-versus-host disease (GVHD),

i.e., to inactivate CD3⁺ alloreactive T cells, we added an anti-CD3 mAb, OKT3, to the culture medium a day before the inoculation. During the follow-up period (mean, 23.4 months; range, 10.7–32.9 months), neither any remarkable adverse effects nor rejection episodes occurred. All 14 subjects who received the immunotherapy were alive without recurrence of HCC after LT (including 5 patients with HCC exceeding the Milan criteria; ref. 17). At our institute, the survival rate and recurrence rate of historical control patients with HCC exceeding the Milan criteria were 78% (30 of 37) and 10.8% (4 of 37), respectively. The lymphocytes in the peripheral blood of LT recipients who received immunotherapy in the early postoperative period showed significantly enhanced cytotoxicity against an HCC cell line (HepG2) as compared with those in the peripheral blood of LT recipients who did not receive the therapy in the same period (Figure 2A). Although the gross proportions of NK/NKT cells in the peripheral blood of patients treated with immunotherapy did not differ from those in the peripheral blood of untreated patients, the proportions of TRAIL⁺ NK cells significantly increased after immunotherapy in the peripheral blood of the former patients. This increase in the TRAIL⁺ NK cells in the peripheral blood lymphocytes was not observed in untreated patients (Figure 2B). Furthermore, there was a significant correlation between the frequency of TRAIL⁺ NK cells in the peripheral blood lymphocytes and the NK cytolytic activity of the peripheral blood lymphocytes at 7 days after LT (Spearman rank-order correlation coefficient = 0.54, $P = 0.01$; Figure 2C), indicating the anti-HCC effect of adoptively injected TRAIL⁺ NK cells. It would be pertinent to conduct additional clinical trials of this immunotherapy for preventing HCC recurrence after LT.

Anti-HCV activity after adoptive immunotherapy. Of the 14 LT recipients who received the immunotherapy, 7 had chronic HCV infection. During the period of this trial, 5 other HCV-infected LT recipients who did not agree to receive immunotherapy served as controls; the background of the controls, including HCV genotype, age, and immunosuppressive therapy, was similar to that of the immunotherapy recipients (Table 3). It has been reported

Table 1
Recipient and tumor characteristics

Patient no.	Age (yr)	Sex	MELD	Hepatitis virus infection	A	HLA B	C	Milan criteria	AFP (ng/ml)	PIVKA-II (AU/ml)	Tumor no.	Maximum tumor size (mm)	Path. vascular invasion	Path. stage	Postop. months	Outcome
1	67	M	19	B	24,-	13,40	03,-	OUT	-	2,584	5	35	-	III	32.9	Alive
2	53	M	16	B	2603,3303	4002,4403	0304,1403	IN	25.3	43	4	11	-	II	31.0	Alive
3	54	M	7	B	0206,3101	3501,5101	0303,1402	OUT	5.7	213	11	26	-	III	29.4	Alive
4	64	F	16	C	2601,2603	3501,4801	0303,-	IN	5.9	142	-	-	-	-	28.5	Alive
5	59	F	14	B	0206,2601	4002,5502	0102,0304	OUT	<5	65	1	13	b1	II	27.8	Alive
6	47	F	8	C	2402,2601	3501,5201	0303,1202	IN	18	46	3	12	-	II	26.2	Alive
7	57	M	29	B	2402,3101	5101,5201	1202,1402	IN	40.3	514	1	25	-	II	25.4	Alive
8	65	F	18	C	1101,2402	5401,5901	0102,-	IN	-	-	3	6	-	II	24.4	Alive
9	60	F	8	-	1101,3001	1302,4006	0602,0801	OUT	32.8	3,026	2	40	w1	IVA	22.7	Alive
10	56	M	8	C	2402,3303	5201,5801	0302,1202	OUT	-	304	11	22	-	III	19.1	Alive
11	56	M	9	C	0207,-	4601,-	0102,-	IN	47	20	3	25	-	III	17.5	Alive
12	58	M	22	C	1101,3101	1501,3501	0102,0415	IN	-	62	1	17	-	I	16.5	Alive
13	59	M	6	C	1101,2402	1507,1601	0303,0401	IN	202.9	19	3	16	-	II	15.8	Alive
14	51	M	16	B	1101,2601	4002,5401	0102,0304	IN	-	29	-	-	-	-	10.7	Alive

The Milan criteria specifies that liver cancer patients with a single tumor of 5 or fewer centimeters in diameter or 3 or fewer tumors, each no more than 3 cm in diameter, and with no macrovascular invasion, can expect an excellent outcome after LT, with only a 10% risk of cancer recurrence (31). AFP, alpha fetoprotein; F, female; M, male; MELD, model for end-stage liver disease; PIVKA-II, protein induced by vitamin K absence; Path., pathological; Postop., postoperative.

that HCV RNA concentrations sharply decrease a day after LT and increase rapidly thereafter (3). In some of the patients, who did not receive the immunotherapy, HCV RNA titers remained lower than that of the pretransplant titer 1 week after LT, suggesting the individual variation of increasing tempo. However, in almost all patients, HCV RNA titers exceeded the pretransplantation levels by 2 weeks after LT. Notably, HCV infection disappeared in 2 LT recipients after the immunotherapy, but this was not observed in the case of any HCV-infected LT recipients who did not receive the therapy. In one of these patients (who had the lowest HCV RNA levels before LT), HCV RNA has not been detected to date (20 months after LT), even with a qualitative assay. In the other patient, HCV RNA became detectable at 2 months after LT. On the other hand, the 2 patients with the highest HCV viral loads did not respond at all to the immunotherapy. Thus, the effects of immunotherapy were dependent on the HCV virus load before LT, probably because of the proportion of effectors and targets. All patients with HCV viremia are currently being treated with pegylated IFN- α 2b and ribavirin. Nevertheless, during the first month after LT, the HCV RNA titers in the sera of LT recipients who received the immunotherapy were statistically lower than those in the sera of LT recipients who did not receive the therapy ($P < 0.05$) (Figure 3). Among the LT recipients who received the immunotherapy, at 2 weeks after LT, HCV RNA remained undetectable in 4 patients (responders), whereas it was detectable in the other 3 patients (nonresponders). The serum ALT levels did not differ between the responders and nonresponders (Supplemental Figure 1; supplemental material available online with this article; doi:10.1172/JCI38374DS1), suggesting that the immunotherapy did not inhibit HCV RNA by injuring HCV-infected hepatocytes.

In vitro evidence to prove the anti-HCV activity of IL-2/OKT3-treated liver lymphocytes by using HCV replicon-containing hepatic cells. The liver allograft-derived lymphocytes were cultured in complete medium with and without IL-2 for 3 days. This was followed by adding OKT3 to the culture medium 1 day before coculturing the lymphocytes with HCV replicon-containing hepatic cells in a transwell system, at an indicated time. While the freshly isolated liver allograft-derived lymphocytes inhibited HCV replication in the HCV replicon-containing hepatic cells to some extent, the cultivation of these lymphocytes with IL-2/OKT3 markedly promoted anti-HCV activity. Absence of exposure to either IL-2 or OKT3 resulted in reduced anti-HCV activity of the lymphocytes (OKT3 had a more profound influence than IL-2) (Figure 4A). When the lymphocytes were treated with IL-2 alone, the CD56⁺ fraction, including NK and NKT cells, that had been isolated by magnetic cell sorting inhibited HCV replication more strongly than the CD56⁻ fraction; further, the CD3⁺CD56⁺ NK cell and CD3⁺CD56⁺ NKT cell subfractions showed equivalent anti-HCV activity (Figure 4, B and C). On the other hand, when the lymphocytes were treated with both IL-2 and OKT3, the CD56⁺ and CD56⁻ fractions showed similar levels of anti-HCV activity (Figure 4B). After the treatment with IL-2 and OKT3, IFN- γ was the predominant cytokine in the culture supernatant of the lymphocytes (Figure 5A), and intracellular IFN- γ expression was induced in the CD3⁺CD56⁺ NK, CD3⁺CD56⁺ NKT, and CD3⁺CD56⁻ T cells (Figure 5B). There was no difference between the proportions of TRAIL⁺ and TRAIL⁻CD3⁺CD56⁺ NK cells producing IFN- γ (Supplemental Figure 2). Adding mAb against IFN- γ to the coculture of lymphocytes with HCV replicon cells markedly weakened the anti-HCV effects. The incomplete restoration of the anti-HCV effect by anti-IFN- γ treat-



Table 2
Donor and graft characteristics

Donor no.	Donor age (yr)	Donor sex	HLA			Relationship	Graft	Graft weight (g)	No. of cells administered ($\times 10^6$)
			A	B	C				
1	41	M	24,-	07,40	03,07	Offspring	Right	608	172
2	24	M	2402,2603	4002,2603	0304,5201	Offspring	Right	658	38
3	51	F	0201,2402	0702,3901	0702,-	Spouse	Right	670	129
4	34	M	2601,2603	4001,4801	0303,0401	Offspring	Left	414	143
5	31	M	0206,2402	4002,5401	0102,0304	Offspring	Posterior	702	135
6	53	F	2402,-	5201,5401	0102,1202	Sibling	Right	538	411
7	24	M	2601,3101	4006,5201	0801,1202	Offspring	Right	642	350
8	34	M	1101,-	4001,5401	0102,1502	Offspring	Right	846	229
9	37	M	0201,1101	1501,4006	0702,0801	Offspring	Left	402	811
10	28	M	1101,3303	5502,5801	0102,0302	Offspring	Right	686	517
11	28	M	0207,2402	4601,5201	0102,5201	Offspring	Right	558	414
12	27	M	0201,1101	1501,3501	0303,0415	Offspring	Right	628	509
13	54	F	1101,2402	1501,1507	0303,0401	Sibling	Right	650	460
14	21	F	2601,2603	1501,5401	0102,0303	Offspring	Right	436	382

ment suggests the possibility that other inflammatory cytokines may also be responsible for the anti-HCV effect, although we have not defined them at present (Figure 5C). Thus, the vigorous anti-HCV activity of IL-2/OKT3-treated liver lymphocytes was dependent, at least in part, on their IFN- γ -secreting activity.

IFN- γ -secreting activity in LT recipients after adoptive immunotherapy. At 14 days after LT, the number of IFN- γ -secreting cells in the peripheral blood of LT recipients who received adoptive immunotherapy was significantly higher than that in the peripheral blood of LT recipients who did not receive immunotherapy during the trial period (Figure 6). This result was consistent with the results of the in vitro studies showing the crucial role of IFN- γ produced in IL-2/OKT3-treated liver lymphocytes.

In vivo evidence to prove the anti-HCV activity of adoptive immunotherapy by using HCV-infected human hepatocyte-chimeric mice. HCV-infected mice have previously been developed by inoculating HCV-infected human serum into chimeric urokinase-type plasminogen activator-SCID (uPA-SCID) mice with engrafted human hepatocytes (18). This HCV-infected mouse model has been reported to be useful for evaluating anti-HCV drugs such as IFN- α and anti-NS3 protease (19). We also generated a human hepatocyte-chimeric mouse model, in which mouse hepatocytes were almost completely replaced by human hepatocytes (20). These mice consistently developed long-term HCV infections, showing high viral titers after inoculation with HCV genotype 1b-infected human serum (50 μ l/mouse) (Supplemental Figure 3). Intraperitoneal injection of IL-2/OKT3-treated liver lymphocytes (20×10^6 cells/mouse), at 2 weeks after inoculation with the infected serum, consistently prevented the development of HCV infection in

the human hepatocyte-chimeric mice (Figure 7A). Such anti-HCV effects were countered by anti-IFN- γ neutralizing antibodies in some chimeric mice, suggesting the potential role played by IFN- γ in the anti-HCV effects of the immunotherapy. The administration of recombinant human IFN- γ markedly and consistently prevented the development of HCV infection in the human hepatocyte-chimeric mice. Once the HCV RNA became undetectable in the sera of chimeric mice receiving either IL-2/OKT3-treated liver lymphocytes or recombinant IFN- γ , it could not be detected again. The constant levels of human serum albumin in the chimeric mice indicated that neither the immunotherapy nor recombinant IFN- γ administration had significant adverse effects on human hepatocytes in those mice (Figure 7B). Once HCV infection had developed in the human hepatocyte-chimeric mice, who showed high titers of HCV RNA in their sera (over 10^3 copies/ml) 4 weeks after the inoculation of HCV-infected serum, the preventive effects of the adoptive immunotherapy or recombinant IFN- γ on HCV infection were no longer observed (Figure 7C).

Table 3
Characteristics of HCV-infected LT recipients that received and did not receive immunotherapy

No.	Age	Sex	HCV genotype	MELD	Pre-HCV RNA (KIU/ml)	Postoperative months	Immunosuppressant
With immunotherapy							
4	64	F	1b	16	210	29	Basiliximab+FK506+MMF
6	47	F	1b	8	5,000	26	Basiliximab+CsA+MMF
8	65	F	1b	18	2,400	24	Basiliximab+CsA+MMF
10	56	M	1b	8	970	19	Basiliximab+FK506+MMF
11	56	M	1b	9	1,700	17	Basiliximab+FK506+MMF
12	58	M	1b	22	19	17	Basiliximab+FK506+MMF
13	59	M	1b	6	2,200	16	Basiliximab+FK506+MMF
Without immunotherapy							
A	51	M	1b	27	420	42	Basiliximab+FK506+MMF
B	44	M	1b	10	1,600	32	Basiliximab+FK506+MMF
C	54	M	1b	8	180	22	Basiliximab+CsA+MMF
D	56	M	2a	10	470	20	Basiliximab+FK506+MMF
E	57	M	1b	12	3,200	6	Basiliximab+FK506+MMF




Figure 2

Adoptive immunotherapy with IL-2/OKT3-treated liver lymphocytes promoted the cytotoxic activity and TRAIL expression of NK cells in LT recipients. (A) The NK cytotoxic activities of the indicated effectors against their target cells were analyzed by the ⁵¹Cr-release assay. The dot plot represents the NK cytotoxic activities of freshly isolated peripheral blood lymphocytes obtained from recipients who received immunotherapy (+) (*n* = 7) and did not receive immunotherapy (-) (*n* = 5) against HepG2 target cells (effector/target [E/T] ratio, 40:1) 3 and 7 days after LT. NK cytotoxic activities are represented as a proportion (percentage) of the preoperative cytotoxicity in each patient. Horizontal lines indicate the mean. Statistical analyses were performed using the 2-tailed, paired Student's *t* test. **P* < 0.05 for day 7 versus day 3. (B) The frequency of TRAIL⁺ NK cells increased remarkably in the peripheral blood of LT recipients who received the immunotherapy. Horizontal lines indicate the mean. Statistical analyses were performed using the Mann-Whitney *U* test. #*P* = 0.013 for immunotherapy group versus untreated group in postoperative day 7. (C) Correlation between TRAIL⁺ NK cell ratio and NK cytolytic activity after LT (Spearman rank-order correlation coefficient = 0.54, *P* = 0.01). Statistical analyses were performed using the Spearman rank-order correlation coefficient. The diagonal line indicates a linear regression line. Each dot indicates the cytotoxicity and TRAIL⁺ NK cell percentage of each patient. C1, control 1; POD, postoperative day; Pt., patient.

Discussion

The consequences of recurrent hepatitis C on the survival of graft and LT recipients can only be avoided by the development of safe and effective antiviral strategies that can not only prevent initial graft infection but also eradicate established hepatitis C recurrence (3, 4). With regard to initial graft infection, the circulating virions infect the liver graft immediately after LT. HCV RNA concentrations usually increase a few days after LT, reflecting active HCV replication in the liver graft. In general, in such an early phase of a viral infection, the first line of host defense may be effective in removing the virus; however, recent reports have indicated that HCV effectively escapes the innate immune system comprising NK and NKT cells, resulting in persistent infection (21, 22). It has been reported that cross-linking of CD81 on NK cells by the major envelope protein of HCV, HCV-E2, blocks NK cell activation, IFN- γ production, cytotoxic granule release, and proliferation (21). Engagement of CD81 on NK cells blocks tyrosine phosphorylation through a mechanism that is distinct from the negative signaling pathways associated with NK cell inhibitory receptors for major histocompatibility complex class I molecules (22). These

facts prove that HCV-E2-mediated inhibition of NK cells is an efficient HCV evasion strategy, which involves targeting the early antiviral activities of NK cells and allowing the virus to establish itself as a chronic infection.

We have explored whether CD81 cross-linking-induced inhibitory effects occur even in IL-2-stimulated NK cells. CD81 cross-linking by a mAb specific for CD81 inhibited antitumor cytotoxicity and anti-HCV activity mediated by resting NK cells, but this manipulation did not alter both these activities of IL-2-stimulated NK cells (Supplemental Figure 4). This indicated that exposure to IL-2 before CD81 cross-linking abrogates subsequent inhibitory signals in the NK cells. This would be one mechanism whereby the adoptive immunotherapy with IL-2/OKT3-treated liver lymphocytes inhibited HCV replication at the early phase of infection after LT.

Although the role of NK cells in controlling HCV infection and replication has not been completely elucidated, a recent report has indicated that NK cells do not exert a direct cytolytic effect on the HCV replicon-containing hepatic cells but release IFN- γ , suppressing HCV RNA expression (11). The role of IFN- γ in the expression of NK cell-mediated anti-HCV activity has been proved by the observa-




Figure 3

Serial measurement of the HCV RNA titers of LT recipients after LT. The HCV RNA titers in the sera of LT recipients who received immunotherapy were markedly lower than those in the sera of LT recipients who did not receive the therapy during the first month after LT. Each line with a different symbol represents serial HCV RNA titers from an LT recipient who received (+) (A; $n = 7$) and 1 who did not receive (-) (B; $n = 5$) the immunotherapy after LT. KIU, kilo international unit; pre, pre LT; W, week.

tion that NK cell-conditioned media have an enhanced expression of signal transducer and activator of transcription 1, a nuclear factor that is essential in IFN- γ -mediated antiviral pathways. It has also been reported that hepatocytes cultured in NK cell-conditioned media express higher levels of IFN- α/β , IFN regulatory factor 3, and IFN regulatory factor 7, confirming that NK cells play a key role in suppressing HCV infection and replication in human hepatocytes in an IFN-dependent manner (23). Similar to recent reports, in the present study, we demonstrated that the NK cells among the IL-2/OKT3-treated liver lymphocytes released soluble factors, predominantly IFN- γ , thus suppressing HCV replication (Figures 5-7).

In addition to NK cells, NKT cells are thought to be involved in eliciting innate responses against infection; however, the role

of NKT cells in controlling HCV infection/replication remains unclear. One report has indicated that the number of NKT cells in patients with chronic HCV infection does not differ from that in healthy donors; however, activated NKT cells in HCV-infected patients produce higher levels of IL-13 — but comparable levels of IFN- γ — than those in healthy subjects, showing that NKT cells are biased toward T-helper 2-type responses in chronic HCV infection (24). Another recent report has shown that the sustained response of patients with chronic hepatitis C to treatment with IFN- α and ribavirin is closely associated with increased dynamism of NK and NKT cells in the liver, implicating an NKT cell-mediated mechanism in anti-HCV activity (25). Here, we have described that NKT as well as NK cells in the IL-2/




Figure 4

The cultivation of liver lymphocytes with IL-2/OKT3 markedly promoted anti-HCV activity. (A) Activation by IL-2 and OKT3 significantly promoted the anti-HCV effect of the liver allograft-derived lymphocytes that were cultured in complete medium with and without IL-2 (100 JRU/ml) for 3 days. OKT3 (1 $\mu\text{g/ml}$) was then added 1 day before coculturing with HCV replicon cells, at the indicated time. The bar graphs indicate the luciferase activities of the cells in each group. Data are presented as mean \pm SEM ($n = 5$). Statistical analyses were performed using the Mann-Whitney U test with Bonferroni correction after the Kruskal-Wallis H test. $\#P < 0.01$ for OKT3 and IL-2/OKT3 treatment versus no treatment. (B) CD56 $^+$ fraction, including NK and NKT cells, strongly inhibited HCV replication. The culture conditions are described in A. By magnetic cell sorting, CD56 $^+$ and CD56 $^-$ fractions were isolated from the activated lymphocytes and analyzed for anti-HCV activity. The bar graphs indicate the luciferase activities of the cells in each group (IL-2-treated group, white bars; IL-2 plus OKT3-treated group, black bars). Whole, whole lymphocytes. Data are presented as mean \pm SEM ($n = 5$). Statistical analyses were performed using the Mann-Whitney U test. $*P < 0.05$ for CD56 $^+$ fraction versus CD56 $^-$ fraction. (C) Anti-HCV effect of NK cells was almost identical to that of NKT cells after IL-2 activation. The liver allograft-derived lymphocytes were cultured in complete medium with IL-2 (100 JRU/ml) for 3 days. By magnetic sorting, CD3-CD56 $^+$ (NK) and CD3 $^+$ CD56 $^+$ (NKT) fractions were isolated from the activated lymphocytes and analyzed for anti-HCV activity. Data are presented as mean \pm SEM ($n = 6$).




Figure 5

Anti-HCV activity of IL-2/OKT3-treated liver lymphocytes was dependent on their IFN- γ secretion ability. (A) IFN- γ was the major cytokine released from the cultured cells. The bar graphs indicate the concentrations of various cytokines (IFN- γ , TNF- α , IL-2, IL-4, IL-5, and IL-10) detected in the coculture supernatant by CBA. Data are presented as mean \pm SEM ($n = 3$). (B) The effects of IL-2 and OKT3 (100 JRU/ml and 1 μ g/ml, respectively) on IFN- γ production by stimulated CD3-CD56⁺ NK, CD3⁺CD56⁺ NKT, and CD3⁺CD56⁻ T cells were evaluated by a combination of cell surface and cytoplasmic mAb staining and subsequent flow cytometric analysis. Histograms represent the log fluorescence intensities obtained upon staining for IFN- γ after gating of each fraction. Dotted lines represent negative control staining with isotype-matched mAbs. Horizontal lines indicate the gated portion of lymphocytes. GMean, geometric mean fluorescent intensity. (C) Blocking of IFN- γ with mAb (100 μ g/ml) elucidated the marked role played by IFN- γ in producing the anti-HCV effect. The bar graphs indicate the luciferase activities of the cells in each group. Data are presented as mean \pm SEM of a representative triplicate sample.

OKT3-treated liver lymphocytes could play a vital role in controlling HCV replication in hepatic cells via an IFN- γ -associated mechanism (Figures 5 and 6).

Therefore, in the early phase of HCV reinfection after LT, the effects of IFN- γ secretion from adoptively injected liver lymphocytes may include inhibition of HCV virion production, which is probably caused by suppression of viral RNA and protein synthesis without immune lysis of intact hepatic cells. This IFN- γ secretion from both CD3⁺CD56⁺ NKT cells and CD3⁺ T cells was markedly upregulated after treatment with OKT3, which was originally used to prevent GVHD (Figure 5B). This is possibly because of the potent mitogenic activity of OKT3 that induces the activation of CD3⁺CD56⁺ NKT cells and CD3⁺ T cells. However, the administration of OKT3-coated cells in vivo results in the opsonization and subsequent trapping and/or lympholysis of cells by the reticuloendothelial system (26–28). Thus, GVHD is prevented in LT recipients treated with adoptive immunotherapy.

Our finding that the IL-2/OKT3-treated liver lymphocytes controlled HCV replication via an IFN- γ -associated mechanism can lead to the clinical application of recombinant IFN- γ for anti-HCV treatment. However, a clinically applicable dose of recombinant IFN- γ could not induce significant inhibitory effects on HCV

viremia in the previous study (29). Based on the accumulation of adoptively injected IL-2/OKT3-treated liver lymphocytes in the liver of human hepatocyte-chimeric mice (data not shown), the immunotherapy with the liver lymphocytes would provide sufficient IFN- γ to the HCV-infected site.

It has been recently reported that HCV-specific CD8⁺ T cells exert strong antiviral effects by both cytopathic and IFN- γ -mediated noncytopathic effector functions (30). However, in patients with chronic HCV infection, dysfunction and functional restoration of HCV-specific CD8⁺ T cell responses have been reported (31). Since HCV-specific CD8⁺ T cell defects may be important in persistent HCV infections, correcting these defects is considered to our knowledge to be a novel approach to treat HCV infection. Further studies are required to investigate whether activation of NK or NKT cells functionally restores HCV-specific CD8⁺ T cells.

In conclusion, adoptive immunotherapy using IL-2/OKT3-treated liver lymphocytes containing abundant NK and NKT cells could mount remarkable anti-HCV responses in HCV-infected LT recipients, although its effects were incomplete or transient. Treatment-related improvements, such as defining the best schedule and frequency of cell inoculation and developing more potent effectors, could improve clinical benefits.




Figure 6

Adoptive immunotherapy with IL-2/OKT3-treated liver lymphocytes induced the production of IFN-γ in the LT recipients. At 14 days after LT, the number of IFN-γ-secreting cells in the peripheral blood of LT recipients treated with the adoptive immunotherapy (+) with IL-2/OKT3-treated liver lymphocytes, including NK and NKT cells, was significantly higher than that in the peripheral blood of untreated LT recipients (-). Histograms represent the proportion (percentage) of IFN-γ-positive cells among the mononuclear cells obtained from the peripheral blood of the immunotherapy (n = 4) and control group (n = 4) LT recipients. Dotted lines represent negative control staining with isotype-matched mAbs. Horizontal lines indicate the gated portion of lymphocytes. Data are presented as mean ± SEM. Histogram profiles shown are representative of 4 independent experiments. Statistical analyses were performed using the Mann-Whitney U test. *P < 0.05 for immunotherapy group versus control group.

Methods

Subjects. All the human liver samples were collected at Hiroshima University Hospital. Tissue specimens were collected after approval from the Institutional Review Board of Hiroshima University and after written informed consent was obtained from the patients. The use of immunotherapy with IL-2/OKT3-treated liver lymphocytes was approved by the Clinical Institutional Ethical Review Board of Hiroshima University. Written informed consent was

obtained from all of the patients. This approach was successfully used in 14 cirrhotic patients with HCC undergoing clinical LT (Tables 1 and 2). Of these 14 patients, 7 had chronic HCV infection. Five other LT recipients with chronic HCV infection did not agree to receive this immunotherapy during the trial period. HCV RNA was qualitatively detected in the sera of these patients by a standardized qualitative RT-PCR assay (Amplicor HCV monitor, version 2.0; Roche Diagnostics) every week during the first month after LT.




Figure 7

Adoptive immunotherapy with IL-2/OKT3-treated liver lymphocytes prevented HCV infection in human hepatocyte-chimeric mice. (A) Human hepatocyte-chimeric mice were intravenously injected with human serum samples positive for HCV genotype 1b. Two weeks after injecting the infected serum, the mice were intraperitoneally inoculated with IL-2/OKT3-treated liver lymphocytes (20 × 10⁶ cells/mouse; n = 6) for adoptive immunotherapy. When indicated, anti-human IFN-γ mAb was injected intraperitoneally 1 day before the immunotherapy (n = 4). Intraperitoneal injection of recombinant human IFN-γ (rhIFN-γ) was commenced at 2 weeks after injecting the infected serum (n = 5). The untreated mice served as controls (n = 6). The dot plots represent serum HCV RNA titers in each chimeric mouse 4 weeks after the injecting the infected serum. Statistical analyses were performed using the Mann-Whitney U test. *P < 0.01 for immunotherapy group versus control group. (B) The lines represent serial changes in human serum albumin levels in the sera of the mice indicated above. Data are presented as mean ± SEM. (C) IL-2/OKT3-treated liver lymphocytes (20 × 10⁶ cells/mouse) were intraperitoneally inoculated 4 weeks after the injection with the infected serum (n = 5) for adoptive immunotherapy. Intraperitoneal injection of recombinant human IFN-γ was commenced 4 weeks after the injecting the infected serum (n = 9). The untreated mice served as controls (n = 9). The dot plots represent serum HCV RNA titers in each chimeric mouse 6 weeks after injection with the infected serum.



Isolation of lymphocytes from liver allograft perfusate. Donor hepatectomy and the transplantation procedure were performed as described previously (32). After hepatectomy, ex vivo perfusion of the liver allograft was performed through the portal vein. Liver allograft-derived lymphocytes were isolated by gradient centrifugation with Ficoll-Paque (GE Healthcare Bio-Sciences AB).

Adoptive transfer of IL-2/OKT3-treated liver lymphocytes. Liver lymphocytes were cultured with human recombinant IL-2 (100 Japanese reference units/ml [JRU/ml]; Takeda) in complete medium at 37°C in a 5% CO₂ incubator for 3 days. One day before the infusion, 1 µg/ml of OKT3 (Janssen-Kyowa) was added in order to opsonize the CD3⁺ fraction. On the day of infusion, the cells were washed twice with 0.9% sodium chloride and resuspended with 5% human serum albumin in 0.9% sodium chloride for injection (Figure 1). The viability of the cells was assessed by the dye-exclusion test, and the cells were checked twice for possible contamination by bacteria, fungi, and endotoxins.

Cytotoxicity assay. A ⁵¹Cr-release assay was done as previously described (5), using HepG2 tumor cells (Japanese Cancer Research Resources Bank) as targets. Briefly, ⁵¹Cr-labeled target tumor cells were added for 4 hours at 37°C to effector cells in round-bottomed 96-well microtiter plates (BD Biosciences – Discovery Labware). The percentage of specific ⁵¹Cr release was calculated as follows: % cytotoxicity = [(cpm of experimental release – cpm of spontaneous release)/(cpm of maximum release – cpm of spontaneous release)] × 100. All the assays were performed in triplicate.

Flow cytometry. Flow cytometric analyses were performed using a FACSCalibur dual-laser cytometer (BD Biosciences). The following mAbs were used for the surface staining of the lymphocytes: FITC-conjugated anti-CD3 mAb (clone HIT3a; BD Biosciences – Pharmingen); PE-conjugated anti-CD56 mAb (clone B159; BD Biosciences – Pharmingen); and biotinylated anti-TRAIL (biotin-conjugated anti-TRAIL) mAb (clone RIK-2; eBioscience). The biotinylated mAb was visualized using APC-streptavidin (BD Biosciences – Pharmingen). Dead cells identified by light scatter and propidium iodide staining were excluded from the analysis. IFN-γ production in the lymphocytes was measured by a combination of cell surface and cytoplasmic mAb staining and subsequent flow cytometric analysis, as described previously (33).

Isolation of CD56⁺ and CD56⁻ fractions and that of NK and NKT cells. Liver allograft-derived lymphocytes were separated into a CD56⁺ fraction – including NK and NKT cells – and a CD56⁻ fraction by using auto MACS (Miltenyi Biotec) with anti-human CD56 microbeads (Miltenyi Biotec) according to the manufacturer's instructions. The NK and NKT cells were also isolated by magnetic cell sorting, using the human NK cell isolation kit or human CD3⁺CD56⁺ NKT cell isolation kit (Miltenyi Biotec). The purity of the isolated fractions was assessed by flow cytometric analysis, and only the fractions with purities greater than 90% were used for functional studies.

Coculture with HCV replicon-containing hepatic cells. An HCV subgenomic replicon plasmid, pRep-Feo, was derived from pRep-Neo (originally, pHC-Vibneo-delS; ref. 34). The pRep-Feo carries a fusion gene comprising firefly luciferase (*Fluc*) and neomycin phosphotransferase, as described elsewhere (35, 36). After culture in the presence of G418 (Invitrogen), pRep-Feo cell lines stably expressing the replicons were established. For coculture experiments, transwell tissue culture plates (pore size, 1 µm; Costar) were used. HCV replicon-containing hepatic cells (10⁵ cells) were incubated in the lower compartment with different numbers of lymphocytes in the upper compartment. The hepatic cells in the lower compartments were collected 48 hours after coculture for the luciferase assay. Luciferase activities were

measured with a luminometer (Lumat LB9501; Promega), using the Bright-Glo Luciferase Assay System (Promega).

Cytometric bead array. Cytokine (IFN-γ, TNF-α, IL-2, IL-4, IL-5, IL-10) levels in the coculture assay supernatants were measured with the FACSCalibur dual-laser cytometer (BD Biosciences), using a BD Human Th1/Th2 Cytokine Cytometric Bead Array (CBA) Kit according to the manufacturer's instructions.

Generation of human hepatocyte-chimeric mice. Generation of the uPA^{+/+}SCID^{+/+} mice and transplantation of human hepatocytes were performed as described recently by our group (20, 37). Mouse serum concentrations of human serum albumin correlated with the repopulation index (20), and these were measured as described previously (37).

In vivo studies using human hepatocyte-chimeric mice. Human hepatocyte-chimeric mice were intravenously injected with 50 µl of the human serum samples positive for HCV genotype 1b. The serum HCV RNA titer in human hepatocyte-chimeric mice was detected by nested PCR, as previously described (38, 39). All animal protocols described in this study were performed in accordance with the guidelines and with approval of the Ethics Review Committee of Animal Experimentation of the Graduate School of Biomedical Sciences, Hiroshima University. Either 2 or 4 weeks after injecting the infected serum, the mice were intraperitoneally inoculated with IL-2/OKT3-treated liver lymphocytes (20 × 10⁶ cells/mouse) for adoptive immunotherapy. When indicated, anti-human IFN-γ mAb (R&D Systems) (1.5 mg/mouse) was injected intraperitoneally 1 day before the immunotherapy. In a separate experiment, intraperitoneal injection of recombinant human IFN-γ (Imunomax-γ; Shionogi & Co. Ltd.) was commenced at either 2 or 4 weeks after injecting the infected serum. IFN-γ was administered as follows: 1 × 10⁵ IU on the first day and thereafter 2 × 10⁴ IU/day for 13 days.

Statistics. Data are presented as mean ± SEM. The statistical differences of the results were analyzed by 2-tailed, paired Student's *t* test, Mann-Whitney *U* test, and Mann-Whitney *U* test with Bonferroni correction after the Kruskal-Wallis *H* test, using the Stat View program. *P* values of 0.05 or less were considered statistically significant.

Acknowledgments

This work was supported in part by a Grant-in-Aid for Exploratory Research (19659323) from the Japan Society for the Promotion of Science. We thank Kentaro Ide, Toshimitsu Irei, Hiroyuki Tahara, Masataka Banshodani, Nabin Basnet, Hirofumi Tazawa, and Keiko Kajitani for their advice and encouragement and Yuko Ishida for expert technical assistance.

Received for publication December 17, 2008, and accepted in revised form July 29, 2009.

Address correspondence to: Hideki Ohdan, Department of Surgery, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. Phone: 81-82-257-5220; Fax: 81-82-257-5224; E-mail: hohdan@hiroshima-u.ac.jp. Or to: Kazuaki Chayama, Department of Medicine and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. Phone: 81-82-257-5190; Fax: 81-81-257-5194; E-mail: chayama@hiroshima-u.ac.jp.

1. Petrovic, L.M. 2006. Early recurrence of hepatitis C virus infection after liver transplantation. *Liver Transpl.* 12:S32–S37.
2. Brown, R.S. 2005. Hepatitis C and liver transplantation. *Nature.* 436:973–978.

3. Garcia-Retortillo, M., et al. 2002. Hepatitis C virus kinetics during and immediately after liver transplantation. *Hepatology.* 35:680–687.
4. Berenguer, M. 2002. Natural history of recurrent hepatitis C. *Liver Transpl.* 8:S14–S18.

5. Ishiyama, K., et al. 2006. Difference in cytotoxicity against hepatocellular carcinoma between liver and periphery natural killer cells in humans. *Hepatology.* 43:362–372.
6. Ohira, M., et al. 2006. Adoptive transfer of TRAIL-



- expressing natural killer cells prevents recurrence of hepatocellular carcinoma after partial hepatectomy. *Transplantation*. **82**:1712–1719.
7. Miller, J.S., et al. 2005. Successful adoptive transfer and in vivo expansion of human haploidentical NK cells in patients with cancer. *Blood*. **105**:3051–3057.
 8. Hirata, M., et al. 1998. Increase in natural killer cell activity following living-related liver transplantation. *Transpl. Int.* **11**(Suppl. 1):S185–S188.
 9. Harada, N., et al. 2004. IL-12 gene therapy is an effective therapeutic strategy for hepatocellular carcinoma in immunosuppressed mice. *J. Immunol.* **173**:6635–6644.
 10. Golden-Mason, L., and Rosen, H.R. 2006. Natural killer cells: primary target for hepatitis C virus immune evasion strategies? *Liver Transpl.* **12**:363–372.
 11. Li, Y., et al. 2004. Natural killer cells inhibit hepatitis C virus expression. *J. Leukoc. Biol.* **76**:1171–1179.
 12. Deignan, T., et al. 2002. Decrease in hepatic CD56(+) T cells and V alpha 24(+) natural killer T cells in chronic hepatitis C viral infection. *J. Hepatol.* **37**:101–108.
 13. Durante-Mangoni, E., et al. 2004. Hepatic CD1d expression in hepatitis C virus infection and recognition by resident proinflammatory CD1d-reactive T cells. *J. Immunol.* **173**:2159–2166.
 14. Kawarabayashi, N., et al. 2000. Decrease of CD56(+) T cells and natural killer cells in cirrhotic livers with hepatitis C may be involved in their susceptibility to hepatocellular carcinoma. *Hepatology*. **32**:962–969.
 15. Rosen, H.R., et al. 2008. Pretransplantation CD56(+) innate lymphocyte populations associated with severity of hepatitis C virus recurrence. *Liver Transpl.* **14**:31–40.
 16. Doherty, D.G., et al. 1999. The human liver contains multiple populations of NK cells, T cells, and CD3+CD56+ natural T cells with distinct cytotoxic activities and Th1, Th2, and Th0 cytokine secretion patterns. *J. Immunol.* **163**:2314–2321.
 17. Mazzaferro, V., et al. 1996. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N. Engl. J. Med.* **334**:693–699.
 18. Mercer, D.F., et al. 2001. Hepatitis C virus replication in mice with chimeric human livers. *Nat. Med.* **7**:927–933.
 19. Kneteman, N.M., et al. 2006. Anti-HCV therapies in chimeric scid-Alb/uPA mice parallel outcomes in human clinical application. *Hepatology*. **43**:1346–1353.
 20. Tateno, C., et al. 2004. Near completely humanized liver in mice shows human-type metabolic responses to drugs. *Am. J. Pathol.* **165**:901–912.
 21. Tseng, C.T., and Klimpel, G.R. 2002. Binding of the hepatitis C virus envelope protein E2 to CD81 inhibits natural killer cell functions. *J. Exp. Med.* **195**:43–49.
 22. Crotta, S., et al. 2002. Inhibition of natural killer cells through engagement of CD81 by the major hepatitis C virus envelope protein. *J. Exp. Med.* **195**:35–41.
 23. Wang, S.H., et al. 2008. Natural killer cells suppress full cycle HCV infection of human hepatocytes. *J. Viral Hepat.* **15**:855–864.
 24. Kanto, T., and Hayashi, N. 2007. Innate immunity in hepatitis C virus infection: Interplay among dendritic cells, natural killer cells and natural killer T cells. *Hepato. Res.* **37**(Suppl. 3):S319–S326.
 25. Yamagiwa, S., et al. 2008. Sustained response to interferon-alpha plus ribavirin therapy for chronic hepatitis C is closely associated with increased dynamism of intrahepatic natural killer and natural killer T cells. *Hepato. Res.* **38**:664–672.
 26. Van Wauwe, J.P., De Mey, J.R., and Goossens, J.G. 1980. OKT3: a monoclonal anti-human T lymphocyte antibody with potent mitogenic properties. *J. Immunol.* **124**:2708–2713.
 27. Chang, T.W., Kung, P.C., Gingras, S.P., and Goldstein, G. 1981. Does OKT3 monoclonal antibody react with an antigen-recognition structure on human T cells? *Proc. Natl. Acad. Sci. U. S. A.* **78**:1805–1808.
 28. Chatenoud, L., et al. 1990. In vivo cell activation following OKT3 administration. Systemic cytokine release and modulation by corticosteroids. *Transplantation*. **49**:697–702.
 29. Soza, A., et al. 2005. Pilot study of interferon gamma for chronic hepatitis C. *J. Hepatol.* **43**:67–71.
 30. Jo, J., et al. 2009. Analysis of CD8+ T-cell-mediated inhibition of hepatitis C virus replication using a novel immunological model. *Gastroenterology*. **136**:1391–1401.
 31. Penna, A., et al. 2007. Dysfunction and functional restoration of HCV-specific CD8 responses in chronic hepatitis C virus infection. *Hepatology*. **45**:588–601.
 32. Ohdan, H., et al. 2003. Intraoperative near-infrared spectroscopy for evaluating hepatic venous outflow in living-donor right lobe liver. *Transplantation*. **76**:791–797.
 33. Tanaka, Y., Ohdan, H., Onoe, T., and Asahara, T. 2004. Multiparameter flow cytometric approach for simultaneous evaluation of proliferation and cytokine-secreting activity in T cells responding to allo-stimulation. *Immunol. Invest.* **33**:309–324.
 34. Guo, J.T., Bichko, V.V., and Seeger, C. 2001. Effect of alpha interferon on the hepatitis C virus replicon. *J. Virol.* **75**:8516–8523.
 35. Tanabe, Y., et al. 2004. Synergistic inhibition of intracellular hepatitis C virus replication by combination of ribavirin and interferon-alpha. *J. Infect. Dis.* **189**:1129–1139.
 36. Yokota, T., et al. 2003. Inhibition of intracellular hepatitis C virus replication by synthetic and vector-derived small interfering RNAs. *EMBO Rep.* **4**:602–608.
 37. Tsuge, M., et al. 2005. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology*. **42**:1046–1054.
 38. Hiraga, N., et al. 2007. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis C virus and its susceptibility to interferon. *FEBS Lett.* **581**:1983–1987.
 39. Kimura, T., et al. 2008. Establishment of an infectious genotype 1b hepatitis C virus clone in human hepatocyte chimeric mice. *J. Gen. Virol.* **89**:2108–2113.

Original Article

Effects of structural variations of *APOBEC3A* and *APOBEC3B* genes in chronic hepatitis B virus infection

Hiromi Abe,^{1,3} Hidenori Ochi,^{1,3} Toshiro Maekawa,^{1,3} Tsuyoshi Hatakeyama,² Masataka Tsuge,^{1,3,4} Shosuke Kitamura,² Takashi Kimura,² Daiki Miki,² Fukiko Mitsui,² Nobuhiko Hiraga,^{1,3} Michio Imamura,^{1,3} Yoshifumi Fujimoto,^{1,3} Shoichi Takahashi,^{1,3} Yusuke Nakamura,⁵ Hiromitsu Kumada⁶ and Kazuaki Chayama^{1,3}

¹Laboratory for Digestive Diseases, Center for Genomic Medicine, RIKEN, ²Department of Medicine and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences, ³Liver Research Project Center, ⁴Natural Science Center for Basic Research and Development, Hiroshima University, Hiroshima, ⁵Center for Genomic Medicine, RIKEN, Kanagawa, and ⁶Department of Hepatology, Toranomon Hospital, Tokyo, Japan

Aim: Human APOBEC3 deaminases induce G to A hypermutation in nascent DNA strand of hepatitis B virus (HBV) genomes and seem to operate as part of the innate antiviral immune system. We analyzed the importance of APOBEC3A (A3A) and APOBEC3B (A3B) proteins, which are potent inhibitors of adeno-associated-virus and long terminal repeat (LTR)-retrotransposons, in chronic HBV infection.

Methods: We focused on the common deletion polymorphism that spans from the 3' part of A3A gene to the 3' portion of A3B gene. An association study was carried out in 724 HBV carriers and 469 healthy control subjects. We also analyzed hypermutated genomes detected in deletion and insertion (non-deletion) homozygous patients to determine the effect of APOBEC3 gene deletion. Further, we performed functional analysis of A3A gene by transient transfection experiments.

Results: The association study showed no significant association between deletion polymorphism and chronic HBV

carrier state. Context analysis also showed a negligible effect for the deletion. Rather, mild liver fibrosis was associated with APOBEC gene deletion homozygosity, suggesting that A3B deletion is not responsible for chronic HBV infection. Functional analysis of A3A showed that overexpression of A3A induced hypermutation in HBV genome, although the levels of hypermutants were less than those introduced by A3G. However, overexpression of A3A did not decrease replicative intermediates of HBV.

Conclusion: These results suggest that A3A and A3B play little role in HBV elimination through anti-viral defense mechanisms. The significance of hypermutation induced by A3A should be investigated further.

Key words: APOBEC3A, APOBEC3B, APOBEC3G, deaminase, hypermutation, structural variation

INTRODUCTION

APOBEC3 CYTIDINE DEAMINASE family consists of at least seven tandem arrayed genes *APOBEC3A* (A3A), A3B, A3C, A3DE, A3F, A3G, and A3H on

chromosome 22.^{1,2} The anti-viral effect of A3G was initially identified in 2002 when it was found to inhibit the replication of human immunodeficiency virus (HIV).³ Similarly, A3F, A3B and A3DE have been reported to inhibit HIV replication.^{4–8}

APOBEC3 proteins also act on many other viruses such as simian immunodeficiency virus,⁹ adeno-associated virus¹⁰ and retrotransposons.^{11–13} With regard to hepatitis B virus (HBV), A3G was also reported to inhibit HBV replication and induction of hypermutation, although the significance of the latter on viral inactivation is still controversial.^{14–23} Among the APOBEC3 family members, A3B, A3C, A3G and A3F have been

Correspondence: Professor Kazuaki Chayama, Department of Medical and Molecular Science, Division of Frontier Medical Science, Programs for Biomedical Research, Graduate school of Biomedical Science, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan. Email: chayama@hiroshima-u.ac.jp
Received 15 April 2009; revision 17 May 2009; accepted 18 May 2009.

extensively analyzed in these reports for induction of hypermutation and inhibition of replication of HBV. In contrast, the function of A3A on HBV has not been evaluated despite its potent inhibitory effects on adeno-associated virus and retrotransposons.^{9–13} Recently, Henry *et al.*²⁴ reported that, among the APOBEC3 family, A3A is the most efficient editor in induction of hypermutation in the HBV genome. This finding is not consistent with the previous reports. However, the relationship between genomic DNA editing by A3A and its effect on HBV replication have not been elucidated. This background prompted us to examine the effects of A3A on HBV replication and induction of hypermutation.

A recent study²⁵ identified a common deletion polymorphism of APOBEC gene spanning from the 3' end of A3A gene to the 3' portion of A3B gene (the segment extending from exon 5 of A3A to exon 8 of A3B was removed by the deletion, positions 37, 683, 131–37, 712, 716 on chromosome 22). The deletion results in complete elimination of the A3B coding region and the resultant fusion gene has a protein sequence identical to A3A, but has 3' untranslated region of A3B. This polymorphism might modulate the expression levels of A3A peptide because the transcription levels and stability of this fusion mRNA could be altered by replacement of the 3' untranslated region sequences. Analyzing the association between this deletion polymorphism and chronic HBV infection should clarify the effect of A3B on the establishment of chronic HBV carrier state.

The aims of the present study were to determine the association between APOBEC3 gene deletion polymorphism and chronic HBV infection and the effect of A3A, which might be up- or down-regulated by the deletion polymorphism, on HBV replication and induction of hypermutation, by *in-vitro* overexpression experiments.

PATIENTS AND METHODS

Study subjects

BLOOD SAMPLES WERE obtained from 724 patients with chronic HBV infection at the hospitals of the Hiroshima Liver Study Group (<http://home.hiroshima-u.ac.jp/naika1/hepatology/english/study.html>) and Toranomon hospital. We also collected 469 control samples from healthy individuals who agreed to join the BioBank Japan Project at the Institute of Medical Science, the University of Tokyo. The study protocols were approved by the ethics committees of the University of Tokyo and the Center for Genomic Medicine, Riken. All participants were ethnically Japanese and pro-

vided written informed consent. Histological activity and fibrosis was assessed in liver biopsy specimens by the Metavir score.²⁶

HBV markers

We measured DNA polymerase by the method of Robinson *et al.*²⁷ The quantity of HBV DNA was assessed by the following tests. Quantiplex HBV DNA probe assay (Chiron Corporation, Emeryville, CA), PCR (Amplicor Cobas TaqMan HBV Auto; Roche Molecular Diagnostic, Basel), transcription mediated amplification (TMA) assay (Fujirevio Diagnostic, Tokyo). The level of HBV in serum was assessed as high or low according to the following criteria (< 200 or ≥ 200 for DNA polymerase, < 200 or ≥ 200 for probe assay, < 6.0 or ≥ 6.0 for PCR assay, < 6.0 or ≥ 6.0 for TMA assay).

HBV-e antigen (HBeAg) and HBV-e antibody (HBeAb) were measured by commercially available chemiluminescent enzyme immunoassay kit (Abbott Laboratories, Chicago, IL). The cut off levels were 1.0 (cut off index) for HBeAg and 70% for HBeAb.

Genotyping

First, we genotyped genomic samples of 94 individuals by the PCR assay using the Deletion and Insertion specific primer sets reported by Kidd *et al.*²⁵ Since we observed some non-specific amplification, which was confirmed by sequencing analysis, we used the invader probes,²⁸ which specifically recognize A3A and A3B. These probes were designed and synthesized by Third Wave Technologies (Madison, WI). Deletion and two-insertion (non-deletion) PCR assays were performed separately as described previously,²⁵ then pooled (Deletion : Insertion1 : Insertion2 = 3:1:1), and subjected to Invader assay.

Cell culture and transfection

Human liver cancer cell line, HepG2, was purchased from RIKEN Cell Bank (Tsukuba). The cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum at 37°C under 5% CO₂. Cells were seeded to semi-confluence in six-well tissue culture plates. Transient transfection experiments were performed using TransIT-LT1 (Mirus, Madison, WI) according to the instructions provided by the supplier.

Plasmid construction

The expression vector for hemagglutinin (HA)-tagged human A3G was kindly provided by Dr. Takaori (Kyoto University).²⁹ We constructed A3A cDNA expression

plasmid by cloning DNA fragment, which was amplified by PCR from cDNA obtained from lymphocytes of a deletion homozygous patient, into pcDNA3.1/nV5-DEST (Invitrogen, Carlsbad, CA). Construction of the wild-type HBV 1.4 genome length plasmid, pTRE-HB-wt was described previously (Tsuge *et al.*;³⁰ GenBank accession no. AB206816).

Analysis of core-associated HBV DNA

The cells were harvested 4 days after transfection and lysed with 250 μ l lysis buffer [10 mM Tris/HCl, pH 7.4, 140 mM NaCl and 0.5% (v/v) NP-40]. The lysate was then centrifuged for 2 min at 15 000 \times g. The core particles were immunoprecipitated from the supernatant by mouse anti-core monoclonal antibody (anti-HBc determinant α , Institute of Immunology, Tokyo). Genomic DNA was separated from the core particles by SDS/proteinase K digestion followed by phenol extraction and ethanol precipitation. Quantitative analysis was performed using the above HBV DNA by RT-PCR using the RT-PCR system (Applied Biosystems, Foster City, CA). The primers and the probe used were described previously.³¹ The real-time PCR was performed in a 25- μ l reaction volume containing 2 \times TaqMan Gene Expression Master Mix, 0.9 μ M of each primer, 0.25 μ M probe and 1 μ l DNA solution. The thermal profile was 50°C for 2 min, 95°C 10 min, followed by 40 cycles of amplification (denaturation at 95°C for 15 sec, annealing at 55°C for 30 sec and extension at 62°C for 90 sec).

Analysis of hypermutated HBV genomes by 3D-RT-PCR

Hypermutated genomes were detected and quantified by modified 3DRT-PCR using the primers, probe and reagents described previously.³¹ The thermal profile was 50°C for 2 min, 95°C for 10 min followed by initial denaturation at 85°C for 20 min and 45 cycles of amplification (denaturation at 85°C for 15 sec, annealing at 50°C for 30 sec and extension at 62°C for 90 sec).

Detection of A3A-A3B fusion mRNA by RT-PCR

We extracted total RNA from lymphocytes of each allele patients using RNeasy Mini Kit (Qiagen, Hilden) and reverse-transcribed using ReverTra Ace (TOYOBO, Osaka) with random primer in accordance with the instructions supplied by the manufacturer. We then amplified cDNAs by 35 cycles of PCR using primers specific for exon 1 of A3A and 3'-untranslated region of A3B in a 25 μ l reaction volume containing 1 \times KOD-Plus buffer [0.3 μ M each primers, 0.2 mM MgSO₄, 1 μ l DNA

solution and 1 unit of KOD-Plus (TOYOBO Co.)]. The thermal profile was initial denaturation at 98°C for 2 min, followed by 35 cycles of amplification (denaturation at 98°C for 15 sec, annealing at 58°C for 15 sec and extension at 68°C for 60 sec). Nucleotide sequences of the amplified fusion cDNA sequences were confirmed by direct sequencing.

Western blot analysis

Cell lysates prepared as described above were separated by sodium dodecyl sulfate polyacrylamide electrophoresis on a 12% poly acrylamide gel and transferred to polyvinylidene fluoride (Pall Corporation, Pensacola, FL). The membranes were incubated with anti-V5 (Invitrogen), anti-hemagglutinin fusion epitope monoclonal anti-body (Roche) or with anti- β -actin monoclonal antibody (Sigma-Aldrich, St Louis, MO) followed by incubation with horseradish peroxidase-conjugated sheep anti-mouse antibody (GE Healthcare UK, Buckinghamshire). We detected signals using the ECL system (GE Healthcare).

Nucleotide sequencing analysis of hypermutated HBV genomes by 3D-PCR, cloning and nucleotide sequencing

We analyzed hypermutated HBV DNA genomes obtained from serum samples of each genotype patient by 3D PCR (denaturation at 85°C) and cloning and sequencing. The amplified DNA fragments were cloned into pGEM T Easy vector (Promega Corporation, Madison, WI) by TA cloning. Nucleotide sequences were determined using BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems). The nucleotide sequences were compared with those obtained by direct sequencing of amplified PCR products by normal PCR protocol.

Statistical analysis

The allele frequencies was calculated and fit to Hardy-Weinberg equilibrium was tested by the chi-square test between cases and controls using Excel software (Microsoft, Redmond, WA).³² We also compared differences in allele frequency and genotype distribution of the deletion between cases and controls with χ^2 -test. Continuous data were compared by analysis of variance (ANOVA). Differences in categorical data were analyzed by the χ^2 -test. Differences in core-associated HBV and hypermutated HBV genomes per 1 \times 10⁴ copies of HBV genomes, were analyzed by Student's *t*-test.

Table 1 Characteristics of subjects

	Patients	Control	P-value
Number of patients	724	469	–
Sex			NS
Male	499	373	
Female	224	95	
Age (years)	53.1 (20.6–86.4)	55 (18–93)	NS
ALT	66 (5–3634)	–	–
Fibrosis stage		–	–
F0	13		
F1	80		
F2	149		
F3	114		
F4	46		
Activity		–	–
A0	1		
A1	50		
A2	125		
A3	47		
Platelet ($\times 10^4/\text{mm}^3$)	16.5 (2.2–29.8)	–	–
HBV DNA		–	–
High	137		
Middle	108		
Low	156		
HBeAg/HBeAb		–	–
+/-	207		
-/+	184		
Hepatocellular carcinoma	65	–	–

Data are number of patients or median (range) values. Differences in age between case and control were compared by Mann–Whitney *U*-test. The sex ratio was analyzed by the χ^2 -test. ALT, alanine aminotransferase; HBVeAb, hepatitis B virus e antibody; HBVeAg, hepatitis B virus e antigen; NS, not significant.

RESULTS

Association between chronic HBV carriers, clinical parameters and the APOBEC3 gene deletion

TABLE 1 SUMMARIZES the clinicopathological features of the patients and control subjects. If A3B contributes to the prevention of chronic HBV infection, there should be an association between chronic HBV

carrier state and APOBEC gene deletion polymorphism. However, we did not find any association between the two (Table 2). Furthermore, all clinical parameters, with the exception of the extent of liver fibrosis associated with chronic HBV, did not associate with the polymorphism (Tables 3,4). Advanced histopathological stages were associated with insertion homozygosity. These findings also suggest that A3B does not play any important role in anti-viral immunity in the development of chronic HBV infection.

Table 2 Case-control analysis of APOBEC3B deletion

	Frequency (%)		P-value	Additive mode	
	Ins	Del		OR	95% CI
HBV (<i>n</i> = 724)	0.709	0.291	0.599	0.964	0.624–1.489
Control (<i>n</i> = 469)	0.719	0.281			

P-values were calculated from case-control analysis by χ^2 -test. OR, odds ratio; CI, confidence interval; Del, deletion homozygote; Ins, insertion homozygote.

Table 3 Correlation between deletion and clinical parameters

	Genotype			P-value
	I/I	I/D	D/D	
Genotype frequency	0.50	0.42	0.08	NS
Age (years)	54.0 ± 12.8	52.0 ± 12.6	50.4 ± 13.3	NS
ALT	169.0 ± 320.6	149.5 ± 322.9	196.8 ± 309.3	NS
Platelets (×10 ⁴ /mm ³)	16.8 ± 5.2	16.6 ± 6.1	17.0 ± 5.8	NS

Data are number of patients or mean ± SD. Age, ALT and platelet count were compared by ANOVA. ALT, alanine aminotransferase; D/D, deletion homozygote; H, heterozygote; I/I, insertion homozygote; NS, not significant.

Context analysis of hypermutated genomes obtained from deletion homozygous and insertion homozygous patients

The amount of hypermutated genomes was not analyzed in this study because it is known to fluctuate during the clinical course.³³ Instead, we searched for the target context of G to A mutation in hypermutated HBV genomes using serum obtained from patients with deletion homozygotes and with insertion homozygotes. As shown Figure 1, multiple G to A hypermutations were observed in deletion homozygote and insertion homozygote patients. The results of context analysis showed no significant difference between the contexts

obtained from deletion homozygotes and those from non-deletion homozygotes (Fig. 2). In fact, the preferred contexts were similar in all three deletion homozygous patients and one insertion homozygous patient (DD1-3 and II1 in Fig. 2) but slightly different from those of the remaining two (II2 and II3). These results suggest that the effect of deletion is not strong in these preferred context patterns.

Detection of A3A-A3B fusion mRNA

We then analyzed whether the resultant A3A and A3B fusion was actually transcribed. We designed primers specific for exon 1 of A3A and the 3'-untranslated region

Table 4 Association of clinical parameters and APOBEC gene polymorphism (categorical data)

	Genotype frequency			P value	Additive mode	I/I vs I/D, D/D	D/D vs I/I, I/D
	I/I	I/D	D/D				
Sex (Male/Female)					0.76	0.85	0.30
Male (n = 328)	154 (0.47)	143 (0.44)	31 (0.09)	OR	0.75	1.03	0.72
Female (n = 166)	78 (0.47)	74 (0.45)	14 (0.08)	95% CI	0.40–1.41	0.75–1.41	0.40–1.33
Fibrosis stage (F0-F1/F2-F4)					0.0054	0.0019	0.48
F0-F1 (n = 62)	22 (0.35)	34 (0.55)	6 (0.10)	OR	0.51	0.47	0.74
F2-F4 (n = 187)	95 (0.51)	77 (0.41)	15 (0.08)	95% CI	0.21–1.24	0.30–0.76	0.31–1.73
Activity (A0-A1/A2-A3)					0.31	0.46	0.30
A0-A1 (n = 51)	22 (0.43)	23 (0.45)	6 (0.12)	OR	0.56	0.80	0.60
A2-A3 (n = 168)	81 (0.48)	75 (0.45)	12 (0.07)	95% CI	0.20–1.56	0.45–1.44	0.22–1.60
HBV DNA (High/Low)					0.12	0.12	0.47
High (n = 194)	82 (0.42)	94 (0.48)	18 (0.09)	OR	0.66	0.73	0.77
Low (n = 206)	103 (0.50)	88 (0.43)	15 (0.07)	95% CI	0.32–1.40	0.49–1.09	0.38–1.57
HBeAg/HBeAb ((+/-)/(-/+))					0.52	0.34	0.84
+/- (n = 207)	89 (0.43)	99 (0.48)	19 (0.09)	OR	0.96	0.82	1.07
-/+ (n = 184)	88 (0.48)	78 (0.42)	18 (0.10)	95% CI	0.47–1.95	0.55–1.23	0.54–2.11
HCC					0.85	0.89	0.64
(-) (n = 648)	323 (0.50)	266 (0.41)	59 (0.09)	OR	0.73	1.04	0.69
(+) (n = 65)	34 (0.52)	31 (0.47)	0 (0.00)	95% CI	0.25–2.13	0.62–1.73	0.24–1.98

ALT, alanine aminotransferase; CI, confidence interval; D/D deletion homozygote; H, heterozygote; HBeAg, hepatitis B virus e antibody; HBeAg, hepatitis B virus e antigen; HCC, hepatocellular carcinoma; I/I, insertion homozygote; OR, odds ratio.


```

direct_D/D3 2998 CACTGGCCAGAGGC AAATCAGGTAGGAGCGGGAGCATTGGGCCAGGGTCA CCCCACCA 3057
clone1_D/D3 .....A...A..AA.....AA..AA..A.AAA..A.....AAA...AAAT.....
clone2_D/D3 .....A...A..AA.....A.....A.AAA..A.....AA...AAA.....
clone3_D/D3 .....A...A..AA.....AA..AA..A.AAA..A.....AAA...AAAT.....

direct_D/D3 3058 CACGGAGGTCTTTGGGGTGGAGCCCTCAGGCTCAGGGCATATTGACAACAGTGCCAGTA 3117
clone1_D/D3 ...AA..A.....AAAA..AA..A.....AA.....AAA.....A.....A..A...A..
clone2_D/D3 ...AA.....AAAA..AA.....AA.....AAA.....A.....A..A...A..
clone3_D/D3 ...AA..A.....AAAA..AA..A.....AA.....AAA.....A.....A..A...A..

direct_D/D3 3118 GCACCTCCTCCTGCCTCCACCAATCGGCAGTCAGGAAGACAGCCTACTCCATCTCTCCA 3177
clone1_D/D3 A.....A.....AA.....AA.....
clone2_D/D3 A.....AA.....AA..A.....AA.....
clone3_D/D3 A.....AA.....

direct_D/D3 3278 CCTCTAAGAGACAGTCATCCTCAGGCCATGCAATGGAA 3215
clone1_D/D3 .....A..A..T..C.....A.....AA..
clone2_D/D3 .....A..A.....AA.....A..G..AA..
clone3_D/D3 .....A..A..T..C.....A.....AA..

direct_I/I3 2998 CACTGGCCAGAGGC AAATCAGGTAGGAGTGGGAGCATTGGGCCAGGGTCA CCCCACCA 3057
clone1_I/I3 .....A...A..AA.....AA..AA..A.AAA..A.....AAA...AAA.....
clone2_I/I3 ..T..AA...A..AA.....AA..AA..A.AAA..A.....AAA...AAA.....
clone3_I/I3 .....AA...A..AA.....AA..AA..A.AAA..A.....AAA...AAA.....

direct_I/I3 3058 CACGGCGGTCTTTGGGGTGGAGCCCTCAGGCTCAGGGCATATTGACAACAGTGCCAGCA 3117
clone1_I/I3 .....AA..A.....AAAA..A.....AA.....AAA.....A.....A..A...A..
clone2_I/I3 .....AA..AA.....A.....A.....AAA.....A.....A..A...A..
clone3_I/I3 .....A.....AAAA..A.....AA.....AAA.....A.....A..A...A..

direct_I/I3 3118 GCACCTCCTCCTGCCTCCACCAATCGGCAGTCAGGAAGACAGCCTACTCCATCTCTCCA 3177
clone1_I/I3 A.....A.....AA.....AA.....
clone2_I/I3 A.....TA.....AA..A.....A.....
clone3_I/I3 A.....AA.....AA.....

direct_I/I3 3278 CCTCTAAGAGACAGTCATCCTCAGGCCATGCAATGGAA 3215
clone1_I/I3 .....A..A..A.....A.....A..A..AA..
clone2_I/I3 .....A..A.....A.....A..A..AA..
clone3_I/I3 .....A..A..A.....A.....A..AA..

```

Figure 1 Nucleotide sequences of hypermutated genomes detected from deletion homozygous and insertion homozygous patients. Nucleotide sequences of 3D-PCR amplified hepatitis B virus (HBV) DNA clones are compared with those obtained by usual PCR and direct sequencing. Upper panel, nucleotide sequences obtained from a deletion homozygous patient. Lower panel, nucleotide sequences obtained from a homozygous patient. Nucleotide numbers are those from GenBank accession no. AB206816.

of A3B, and performed RT-PCR using cDNAs obtained from patients of each genotype. We obtained amplified DNA fragments of expected size only from deletion homozygotes and heterozygotes (Fig. 3). These results confirmed the transcription of the fusion mRNA with the coding region of A3A and the 3' untranslated region of A3B.

Inhibition of HBV replication and induction of hypermutation by A3A

We then analyzed the antiviral effect and induction of hypermutation by A3A. Although the expression of both A3A and A3G was confirmed by western Blot analysis (Fig. 4A), transient expression of A3A did not reduce the amount of the core-associated HBV DNA in HepG2 cells (Fig. 4B). However, A3A transfection increased the hypermutated genomes of HBV in a dose-dependent manner albeit the level of induction was much lower than that observed when transfected with A3G. These results suggest that A3A has negligible anti-viral effect although it induces hypermutation of HBV genomes.

DISCUSSION

THE MAIN FINDINGS of the present study were: (i) no association between APOBEC3 deletion and chronic HBV infection (Table 2). (ii) Mild liver fibrosis and low alanine amino transferase (ALT) levels were associated with APOBEC gene deletion homozygous genotype. (iii) The absence of A3B is not responsible for chronic HBV carrier status, although A3B is known as a potent inhibitor of adeno-associated virus and retrotransposons.¹² This suggests different antiviral activities for APOBEC proteins against viruses and that A3B plays little role in inhibition of HBV. (iv) The preferred context analysis showed no differences between insertion homozygotes and deletion homozygotes. Only one of the six patients examined showed different context pattern (Fig. 2). These results suggest that A3B protein has only small effect on the formation of hypermutated genomes in the serum of chronic carriers. The protein has been reported to induce hypermutation on the negative and positive strands of HBV.¹⁸ However, our results showed that the effect of A3B is almost negligible in




Figure 2 Context analysis of hypermutated genomes from deletion homozygous and insertion homozygous patients. Context of G to A hypermutation in hypermutated genome clones obtained from three deletion homozygous patients (D/D1, D/D2 and D/D3) and three insertion homozygous patients (I/I1, I/I2 and I/I3) were analyzed. Numbers after each patient represent the number of clones analyzed in each patient. (a) Two letters up-stream, (b) one letter upstream and (c) one letter downstream of mutated G residue were analyzed.

chronic HBV carriers compared to that of A3G. It is assumed that the other APOBEC3 family proteins mainly induce hypermutation of HBV genomes in HBV carriers to compensate for the function of deleted A3B. It is also assumed that the expression pattern of the remaining six APOBEC3 proteins is different from patient to patient.

As discussed above, our results suggest that A3B protein has almost no effect on prevention of chronic HBV infection and induction of hypermutation. It is thus assumed that A3B is not part of the innate anti-viral immune system against HBV. This is consistent with the finding that deletion is commonly seen in normal populations²⁵ irrespective of HBV carrier rates. Other association studies are required to clarify the role of A3B

protein on other pathogens. The functional relevance of other APOBEC3 proteins on HBV infection as anti-viral immunity should be clarified further.

We also found that A3A protein induced hypermutation on the negative strand of HBV. However, the level of induction of hypermutation was much less than that of A3G (Fig. 4). Recent reports showed quite different effects for A3A on induction of hypermutation on HBV genomes. Henry *et al.*²⁴ reported that A3A is the most efficient editor of seven APOBEC3 proteins. In contrast, Zang *et al.*²³ did not detect induction of hypermutation on HBV. Although these different results might come from different cell lines and conditions used in each experiment, our results clearly showed that A3A induced hypermutation on the negative strand of HBV genome.




Figure 3 Detection of A3A-A3B fusion mRNA by PCR. A3A-A3B fusion mRNA was amplified by PCR using primers specific to A3A and A3B (see Materials and Methods) and detected by agarose gel electrophoresis. Lanes 1–3 are those from deletion homozygous patients, lanes 4–6 are from heterozygous patients and 7–9 are from insertion homozygous patients. M, molecular weight size marker (1 kb DNA Ladder; New England BioLabs, Ipswich, MA), N, negative control.






Figure 4 Analysis of inhibition of HBV replication and induction of hypermutation by A3A. HepG2 cells were transiently transfected with expression plasmid of A3A together with 1.4 genome length hepatitis B virus (HBV) expression vector. The indicated amounts of HBV and A3A (1:0, 1:2, 1:6) or A3G (1:6) expression plasmids were transfected into HepG2 cells. All experiments were performed more than twice with similar results. (a) APOBEC3 gene expression levels were detected by western blot analysis. (b) The amounts of core associated replicative intermediates of HBV were measured by RT-PCR. (c) Quantitative measurement of hypermutated genomes by 3D real-time PCR. Data in (B) and (C) are mean \pm SD. * $P < 0.05$; ** $P < 0.01$

The deletion of APOBEC gene spans from 3' end of the A3A gene to the 3' portion of the A3B gene. This deletion results in the formation of a fusion gene that has A3A amino acid sequence and A3B 3' untranslated region. The expression level of this protein may be different from the undeleted A3A due to different stability of RNA or different transcription levels. The results of context analysis also indicated that the effect of this alteration of expression levels of A3A is almost negligible compared with A3G. This is consistent with our results that showed lower levels of induction of hypermutation on HBV genome by A3A compared with A3G.

Our results indicated that the effect of both A3A and A3B is not significant in the development and progression of chronic hepatitis B. The two proteins also have only little influence on the hypermutation state of HBV in chronic HBV carriers. The results also showed higher induction levels of hypermutation by A3G than by A3A.

Sufficient evidence indicates that A3G has anti-viral effects on HBV,^{14–17} suggesting that some but not all APOBEC3 proteins operate as part of the anti-viral immune system against HBV infection. Further study is needed to clarify the functional role of each APOBEC3 protein for innate anti-viral immunity in chronic HBV infection.

ACKNOWLEDGMENT

THE AUTHORS THANK Rie Akiyama, Yoshie Yoshida, Tomomi Murai, Kazuyo Hattori, Mariko Shiota, and Takako Yokogi for the excellent technical assistance, and Yuko Nagai, Junko Sakamiya and Aya Furukawa for the secretarial assistance. We are also grateful to members of Hiroshima Liver Study Group and The Rotary Club of Osaka-Midosuji Direct 2660 Rotary International in Japan for supporting our study. Part of this work was carried out at the Analysis Center of Life Science, Hiroshima University. This work was supported in part by Grants-in-Aid for scientific research and development from the Ministry of Health, Labor and Welfare, Government of Japan.

REFERENCES

- Jarmuz A, Chester A, Bayliss J *et al.* An anthropoid-specific locus of orphan C to U RNA editing enzymes on chromosome 22. *Genomics* 2002; 79: 285–96.
- Wedekind JE, Dance GS, Sowden MP, Smith HC. Messenger RNA editing in mammals: new members of the APOBEC family seeking roles in the family business. *Trends Genet* 2003; 19: 207–16.
- Sheehy AM, Gaddis NC, Choi JD, Malim MH. Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* 2002; 418: 646–50.
- Bishop KN, Holmes RK, Sheehy AM, Davidson NO, Cho SJ, Malim MH. Cytidine deamination of retroviral DNA by diverse APOBEC proteins. *Curr Biol* 2004; 14: 1392–96.
- Liddament MT, Brown WL, Schumacher AJ, Harris RS. APOBEC3F properties and hypermutation preferences indicate activity against HIV-1 in vivo. *Curr Biol* 2004; 14: 1385–91.
- Wiegand HL, Doehle BP, Bogerd HP, Cullen BR. A second human antiretroviral factor, APOBEC3F, is suppressed by the HIV-1 and HIV-2 Vif proteins. *EMBO J* 2004; 23: 2451–8.
- Zheng YH, Irwin D, Kurosui T, Tokunaga K, Sata T, Peterlin BM. Human APOBEC3F is another host factor that blocks human immunodeficiency virus type 1 replication. *J Virol* 2004; 78: 6073–6.
- Dang Y, Wang X, Esselman WJ, Zheng YH. Identification of APOBEC3DE as another antiretroviral factor from the human APOBEC family. *J Virol* 2006; 80: 10522–33.
- Cullen BR. Role and mechanism of action of the APOBEC3 family of antiretroviral resistance factors. *J. Virol* 2006; 80: 1067–76.
- Chen H, Lilley CE, Yu Q *et al.* APOBEC3A is a potent inhibitor of adeno-associated virus and retrotransposons. *Curr Biol* 2006; 16: 480–5.
- Bogerd HP, Wiegand HL, Doehle BP, Lueders KK, Cullen BR. APOBEC3A and APOBEC3B are potent inhibitors of LTR-retrotransposon function in human cells. *Nucleic Acids Res* 2006; 34: 89–95.
- Bogerd HP, Wiegand HL, Hulme AE *et al.* Cellular inhibitors of long interspersed element 1 and Alu retrotransposition. *Proc Natl Acad Sci USA* 2006; 103: 8780–5.
- Chen H, Lilley CE, Yu Q *et al.* APOBEC3A is a potent inhibitor of adeno-associated virus and retrotransposons. *Curr Biol* 2006; 16: 480–5.
- Turelli P, Mangeat B, Jost S, Vianin S, Trono D. Inhibition of hepatitis B virus replication by APOBEC3G. *Science* 2004; 303: 1829.
- Rösler C, Köck J, Malim MH, Blum HE, von Weizsäcker F. Comment on “inhibition of hepatitis B virus replication by APOBEC3G”. *Science* 2004; 305: 1403a.
- Turelli P, Jost S, Mangeat B, Trono D. Response to comment on “inhibition of hepatitis B virus replication by APOBEC3G”. *Science* 2004; 305: 1403b.
- Noguchi C, Ishino H, Tsuge M *et al.* G to A hypermutation of hepatitis B virus. *Hepatology* 2005; 41: 626–33.
- Suspène R, Guétard D, Henry M, Sommer P, Wain-Hobson S, Vartanian JP. Extensive editing of both hepatitis B virus DNA strands by APOBEC3 cytidine deaminases in vitro and in vivo. *Proc Natl Acad Sci* 2005; 102: 8321–6.
- Bonvin M, Achermann F, Greeve I *et al.* Interferon-inducible expression of APOBEC3 editing enzymes in human hepatocytes and inhibition of hepatitis B virus replication. *Hepatology* 2006; 43: 1364–74.
- Xu R, Zhang X, Zhang W, Fang Y, Zheng S, Yu XF. Association of human APOBEC3 cytidine deaminases with the generation of hepatitis virus B x antigen mutants and hepatocellular carcinoma. *Hepatology* 2007; 46: 1810–20.
- Jost S, Turelli P, Mangeat B, Protzer U, Trono D. Induction of antiviral cytidine deaminases does not explain the inhibition of hepatitis B virus replication by interferons. *J Virol* 2007; 81: 10588–96.
- Bonvin M, Greeve J. Effects of point mutations in the cytidine deaminase domains of APOBEC3B on replication and hypermutation of hepatitis B virus in vitro. *J Gen Virol* 2007; 88: 3270–4.
- Zhang W, Zhang X, Tian C *et al.* Cytidine deaminase APOBEC3B interacts with heterogeneous nuclear ribonucleoprotein K and suppresses hepatitis B virus expression. *Cell Microbiol* 2008; 10: 112–21.