

- 2 The Liver Cancer Study Group of Japan. Primary liver cancer in Japan: clinicopathological features and results of surgical treatment. *Ann Surg* 1990; 211: 277–87.
- 3 Wheeler PG, Melia W, Dubbins P *et al.* Non-operative arterial embolization in primary liver tumours. *Br Med J* 1979; 2: 242–4.
- 4 Chuang VP, Wallace S. Hepatic arterial embolization in the treatment of hepatic neoplasms. *Radiology* 1981; 140: 51–8.
- 5 Okamura J, Horikawa S, Fujiyama T *et al.* An appraisal of transcatheter arterial infusion of chemotherapeutic agent for hepatic malignancies. *World J Surg* 1982; 6: 352–7.
- 6 Yamada R, Sato M, Kawabata M, Nakatsuka H, Nakamura K, Takashima S. Hepatic artery embolization in 120 patients with unresectable hepatoma. *Radiology* 1983; 148: 397–401.
- 7 Lin DY, Liaw YF, Lee TY, Lai CM. Hepatic arterial embolization in patients with unresectable hepatocellular carcinoma: a randomized controlled trial. *Gastroenterology* 1988; 94: 453–6.
- 8 Ikeda K, Kumada H, Saitoh S, Arase Y, Chayama K. Effect of repeated transcatheter arterial embolization on the survival time in patients with hepatocellular carcinoma. *Cancer* 1991; 68: 2150–4.
- 9 Llovet JM, Real MI, Montaña X *et al.* Barcelona Liver Cancer Group. Arterial embolisation or chemoembolisation versus symptomatic treatment in patients with unresectable hepatocellular carcinoma: a randomised controlled trial. *Lancet* 2002; 359: 1734–9.
- 10 Lo CM, Ngan H, Tso WK *et al.* Randomized controlled trial of transarterial lipiodol chemoembolization for unresectable hepatocellular carcinoma. *Hepatology* 2002; 35: 1164–71.
- 11 Homma H, Mezawa S, Doi T *et al.* A comparative randomized trial of intermittent intrahepatic arterial carboplatin-versus doxorubicin-lipiodol emulsion in advanced hepatocellular carcinoma (stage IV). *Hepatogastroenterology* 2004; 51: 1135–9.
- 12 Shibata J, Fujiyama S, Sato T, Kishimoto S, Fukushima S, Nakano M. Hepatic arterial injection chemotherapy with cisplatin suspended in an oily lymphographic agent for hepatocellular carcinoma. *Cancer* 1989; 64: 1586–94.
- 13 Sasaki Y, Imaoka S, Kasugai H *et al.* A new approach to chemoembolization therapy for hepatoma using ethiodized oil, cisplatin, and gelatin sponge. *Cancer* 1987; 60: 1194–203.
- 14 Imaoka S, Sasaki Y, Shibata T *et al.* A pre-operative chemoembolization therapy using lipiodol, cisplatin and gelatin sponge for hepatocellular carcinoma. *Cancer Chemother Pharmacol* 1989; 23: S126–8.
- 15 Yamamoto K, Shimizu T, Narabayashi I. Intraarterial infusion chemotherapy with lipiodol-CDDP suspension for hepatocellular carcinoma. *Cardiovasc Intervent Radiol* 2000; 23: 26–39.
- 16 Maeda S, Shibata J, Fujiyama S *et al.* Long-term follow-up of hepatic arterial chemoembolization with cisplatin suspended in iodized oil for hepatocellular carcinoma. *Hepatogastroenterology* 2003; 50: 809–13.
- 17 Ikeda M, Maeda S, Shibata J *et al.* Transcatheter arterial chemotherapy with and without embolization in patients with hepatocellular carcinoma. *Oncology* 2004; 66: 24–31.
- 18 Ando E, Tanaka M, Yamashita F *et al.* Hepatic arterial infusion chemotherapy for advanced hepatocellular carcinoma with portal vein tumor thrombosis: analysis of 48 cases. *Cancer* 2002; 95: 588–95.
- 19 Kaneko S, Urabe T, Kobayashi K. Combination chemotherapy for advanced hepatocellular carcinoma complicated by major portal vein thrombosis. *Oncology* 2002; 62: 69–73.
- 20 Miller AB, Hoogstraten B, Staquet M, Winkler A. Reporting results of cancer treatment. *Cancer* 1981; 47: 207–14.
- 21 Hatanaka K, Kudo M, Fukunaga T *et al.* Clinical characteristics of NonBNonC- HCC: comparison with HBV and HCV related HCC. *Intervirolgy* 2007; 50: 24–31.
- 22 Lokich J, Anderson N. Carboplatin versus cisplatin in solid tumors: an analysis of the literature. *Ann Oncol* 1998; 9: 13–21. Review.
- 23 Go RS, Adjei AA. Review of the comparative pharmacology and clinical activity of cisplatin and carboplatin. *J Clin Oncol* 1999; 17: 409–22. Review.
- 24 Llovet JM, Bruix J. Novel advancements in the management of hepatocellular carcinoma in 2008. *J Hepatol* 2008; 48 (Suppl 1): S20–37. Review.

High Serum Des-gamma-carboxy Prothrombin Level Predicts Poor Prognosis After Radiofrequency Ablation of Hepatocellular Carcinoma

Masahiro Kobayashi, MD, Kenji Ikeda, MD, Yusuke Kawamura, MD, Hiromi Yatsuji, MD, Tetsuya Hosaka, MD, Hitomi Sezaki, MD, Norio Akuta, MD, Fumitaka Suzuki, MD, Yoshiyuki Suzuki, MD, Satoshi Saitoh, MD, Yasuji Arase, MD, and Hiromitsu Kumada, MD

BACKGROUND: Currently, surgical resection is considered the first-line treatment for early stage hepatocellular carcinoma (HCC). Radiofrequency ablation (RFA) has been an alternative choice for unresectable HCC. However, RFA is expected to have similar therapeutic efficacy for early stage HCC with fewer invasions. **METHODS:** The authors retrospectively analyzed 199 patients who underwent surgery and 209 patients who underwent RFA for HCC with a maximum diameter of ≤ 3 cm and tumors numbering ≤ 3 . All patients were complicated with Child-Pugh A cirrhosis. **RESULTS:** The 3- and 5-year survival rates of the resection (90.3%, 79.0%, respectively) and RFA groups were similar (87.4%, 74.8). The 1- and 3-year tumor recurrence-free survival rates of the resection group (83.1%, 51.0%, respectively) were higher than in the RFA group (82.7%, 41.8%; $P = .011$). Multivariate analysis identified prothrombin time $\geq 80\%$ (hazard ratio [HR], 2.72; 95% confidence interval [CI], 1.56-4.74; $P < .001$) as an independent prognostic factor for survival in the resection group. Des-gamma-carboxy prothrombin (DCP) < 100 arbitrary units (AU)/L (HR, 5.49; CI, 2.23-13.5; $P < .001$) and platelet count $\geq 1.0 \times 10^5$ (HR, 2.70; CI, 1.24-5.88; $P = .012$) were significant markers in the RFA group. Among patients with DCP ≥ 100 AU/L, treatment procedure (HR, 1.26; CI, 1.04-1.53; $P = .020$) was a significant prognostic factor for survival. **CONCLUSIONS:** High DCP levels reflect the biologic aggressiveness and progression of HCC tumors. In the aforementioned cases, we recommend surgical resection rather than RFA for such patients. *Cancer* 2009;115:571-80. © 2008 American Cancer Society.

KEY WORDS: hepatocellular carcinoma, DCP, radiofrequency, prognostic factor.

Hepatocellular carcinoma (HCC) is the sixth most common malignancy and third most deadly carcinoma in the world.¹ In Japan, HCC is ranked third among males and fifth among females as the leading causes of cancer death.² Most patients with HCC are infected with either hepatitis B virus (HBV) or hepatitis C virus (HCV), and have complications stemming from underlying chronic liver disease. The importance of liver condition in the treatment of HCC should be clearly discerned.³

Corresponding author: Masahiro Kobayashi, MD, Department of Gastroenterology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-8470, Japan; Fax: (011) 81 (44) 860-1623; mshkobayashi@toranomon.gr.jp

Department of Hepatology, Toranomon Hospital, Tokyo, Japan

Received: February 12, 2008; **Revised:** August 6, 2008; **Accepted:** August 25, 2008

Published online: December 31, 2008, © 2008 American Cancer Society

DOI: 10.1002/cncr.24031, www.interscience.wiley.com

A maximum tumor diameter of ≤ 3 cm and tumors numbering ≤ 3 are good candidates for liver transplantation in patients with Child-Pugh class B and C.^{4,5} However, patients with Child-Pugh class A conditions should be treated curatively.^{5,6} Hepatectomy is currently recommended for patients with single asymptomatic HCC and extremely well-preserved liver function, who have neither clinically substantial portal hypertension nor abnormal bilirubin levels.⁶ However, resection is suitable for only 20% to 35% of patients with HCC because of poor hepatic reserve.^{7,8} Radiofrequency ablation (RFA) was introduced as a minimally invasive therapy for such cirrhotic patients.⁹⁻¹³ RFA was the initial choice for unresectable HCC; however, 2 recent randomized controlled trials concluded that there were no substantial statistical survival differences between resection and RFA.^{9,10} Although the results of these studies have not yet reached a worldwide consensus, some authors recommend RFA as a first-line therapy for such early stage HCC.¹¹⁻¹³

Tumor staging and the decision between possible treatment options are conducted predominantly based on tumor size, number, vascular invasion, and extrahepatic metastasis evaluated by imaging analysis such as ultrasonography or dynamic computed tomography (CT). However, the malignant nature of the tumor as well as other characteristics are not generally considered.^{14,15} Alpha-fetoprotein (AFP) and des-gamma-carboxy prothrombin (DCP) are HCC-specific tumor markers. High levels of serum tumor markers often indicate HCC development in the liver. On the basis of histopathological analysis, serum AFP and DCP levels are also correlated with tumor differentiation, microscopic portal invasion, or intrahepatic metastasis.^{16,17}

This present study is an attempt to evoke discussion on treatment strategies for small HCC measuring ≤ 3 cm by comparing the long-term outcome of patients treated with either hepatectomy or RFA as the first-line treatment for HCC. AFP and DCP were also accounted as indicators in the decision-making and treatment procedure.

MATERIALS AND METHODS

Patients

A total of 1057 patients were admitted to the Department of Hepatology, Toranomon Hospital between 1995 and 2006 for the treatment of initially developed HCC. The major background liver disease was HCV (767 patients,

72.6%), followed by HBV (196 patients, 18.5%), HCV + HBV (8 patients, 0.8%), alcoholic liver diseases (habitual drinking of ethanol at >80 g/day, 48 patients, 4.5%), primary biliary cirrhosis (4 patients, 0.4%), autoimmune hepatitis (2 patients, 0.2%), and cryptogenic liver disease (42 patients, 4.0%). Treatment of HCC included surgical resection in 281 patients, local ablation therapy in 398 patients (RFA, 267 patients; microwave coagulation, 47 patients; ethanol injection, 84 patients), and transarterial chemoembolization in 378 patients. Among these patients, we included patients with Child-Pugh A cirrhosis and HCC measuring ≤ 3 cm in diameter and numbering ≤ 3 tumors who were treated radically by either surgical resection or RFA. Table 1 summarizes the profile of the 199 patients who received resection and 209 patients who received RFA. HBV-related liver diseases were more common among patients who underwent resection, who were younger (62 vs 67 years; $P < .001$) than patients with RFA. The maximum tumor diameter was larger in the resection group than in the RFA group (20 vs 18 mm; $P < .001$). With regard to laboratory tests, serum albumin level, platelet count, and prothrombin time (%) were higher among patients in the resection group, whereas serum aspartate aminotransferase (AST) levels were higher among patients in the RFA group. None of the patients in either group had tumor invasion of the major portal branch or extrahepatic metastasis. Our institution does not require informed consent for retrospective analysis.

Diagnosis of HCC

Diagnosis of HCC was predominantly based on image analysis. If a hepatic nodular lesion was found on screening ultrasonography, the patient underwent dynamic CT and/or dynamic magnetic resonance imaging (MRI). Furthermore, when a liver nodule showed hyperattenuation in the arterial phase of dynamic study and washout in portal or delayed phase, or showed typical hypervascular staining on digital subtraction angiography, the nodule was diagnosed as HCC. According to the American Association for the Study of Liver Disease guidelines, we obtained at least 2 dynamic imaging images before treatment.⁵ When the nodule did not appear in the abovementioned typical imaging features, a fine needle aspiration biopsy was carried out followed by histological examination and diagnosis.

Table 1. Clinical Background of 199 Patients Who Underwent Hepatic Resection and 209 Patients Who Underwent Radiofrequency Ablation of Liver Tumor

Factors	Resection Group (n = 199)	RFA Group (n = 209)	P
Age, y*	62 (29-80)	67 (38-87)	<.001
Sex, men:women	146:53	137:72	NS
HBV:HCV:HBV + HCV:others	60:121:3:15	22:176:1:10	<.001
Habitual alcohol intake, yes:no	29:170	9:200	
Diameter of HCC, mm*	20 (9-30)	18 (8-30)	<.001
No. of HCC, 1:2:3	168:22:9	169:29:11	
Tumor vascularity, present:absent	185:14	156:53	
Albumin, g/dL*	3.7 (2.8-4.7)	3.6 (2.6-4.4)	<.001
Bilirubin, mg/dL*	1.0 (0.3-2.4)	1.0 (0.2-2.4)	NS
AST, IU/L*	43 (13-386)	55 (17-208)	<.001
Platelets, $\times 10^4/\text{mm}^3$ *	13.1 (4.0-27.2)	10.5 (2.7-25.3)	<.001
Prothrombin time, %*	92 (62-115)	88 (57-125)	.006
AFP, ng/mL*	22 (1-7960)	18 (2-1490)	NS
DCP, AU/L*	20 (<10-1650)	17 (<10-1370)	NS

RFA indicates radiofrequency ablation; NS, not significant; HBV, hepatitis B virus; HCV, hepatitis C virus; HCC, hepatocellular carcinoma; AST, aspartate aminotransferase; IU, international units; AFP, alpha-fetoprotein (AFP); DCP, des-gamma-carboxy prothrombin; AU, arbitrary units.

*Data are expressed as median (range).

Method of Treatment

Physicians and surgeons usually discuss together the preferred choice of therapy in individual patients. Hepatic resection was performed under intraoperative ultrasonographic monitoring and guidance. For small and superficial HCCs, arterial and portal vein clamping at hepatic hilum was not usually performed to maintain liver perfusion.

RFA was performed using 3 different devices: the radiofrequency interstitial tumor ablation system (RITA, RITA Medical Systems Inc., Mountain View, Calif), the cool-tip system (Tyco Healthcare Group LP, Burlington, Vt), and the radiofrequency tumor coagulation system (RTC system, Boston-Scientific Japan Co., Tokyo, Japan). In the first 2 systems, treatment procedures were performed according to the protocol advised by the manufacturer. However, treatment using the RTC system was performed by adopting the "stepwise hook extension technique."¹⁸ The needle was inserted into the tumor percutaneously under ultrasonographic guidance. Because the HCC nodule could not be observed by ultrasonography in 6 cases, the needle was inserted under CT assistance. In the case of RFA, dynamic CT was performed 1 to 3 days after therapy, and the ablated area was evaluated. The goal of treatment was to obtain a necrotic area larger than the

original tumor size, with a surrounding treatment margin of ≥ 5 mm in all directions. When this was not achieved or a residual tumor was found, additional ablation was considered. Of 206 total patients, total ablation session was required once in 149 (71.3%) patients, twice in 47 (22.5%) patients, and 3 times or more in 13 patients (6.2%).

Measurement of Serum AFP and DCP

Serum AFP level was measured by chemiluminescent enzyme immunoassay (CLEIA) using a commercial assay kit (Lumipulse Prestoll AFP, Fujirebio Inc., Tokyo, Japan). DCP level was measured by CLEIA (Lumipulse PIVKA II Eisai, Eisai, Tokyo, Japan).

Follow-up Protocol

Physicians examined the patients every 4 weeks after treatment, and liver function tests and tumor markers were also measured once every month. After completion of HCC eradication, recurrence was surveyed with contrast enhancement 3-phase CT every 3 months. Local tumor progression was defined as tumor recurrence adjacent to the resected or ablated area.

Statistical Analysis

Differences in background features and laboratory data between resection and RFA groups were analyzed by the chi-square test and Mann-Whitney *U* test. Survival and recurrence-free survival were analyzed using the Kaplan-Meier technique, and differences in curves were tested using the log-rank test. Independent risk factors associated with survival and recurrence-free survival were studied using stepwise Cox regression analysis.¹⁹ Potential risk factors for survival and recurrence-free survival included the following 14 variables: age, sex, etiology of background liver disease, amount of alcohol intake, serum albumin, bilirubin, AST, platelet count, prothrombin time, AFP, DCP, diameter of the HCC, tumor multiplicity, and tumor vascularity evaluated by dynamic CT or dynamic MRI. A probability of less than .05 was considered significant. Data analysis was performed using SPSS statistical software version 10 (SPSS Inc., Chicago, Ill).

RESULTS

Survival and Recurrence-free Survival Rates

During the median follow-up of 3.3 years (range, 0.1-12.2 years), 112 (56.3%) of 199 patients in the resection group and 120 (57.4%) of 209 patients in the RFA group developed HCC recurrence. HCC recurrences mainly occurred in other sites in the liver. However, in the RFA group, local tumor progression, defined as HCC recurrence adjacent to the treated site, was seen in 18 (8.6%) of 209 patients, but noted in only 1 patient of the resected group. The cumulative local tumor progression rate in the RFA group was 2.7%, 11.3%, and 12.5% at 1, 3, and 5 years, respectively. The tumors were treated with surgical resection in 2 patients, additional tumor ablation in 8 patients, and transcatheter chemoembolization in the remaining 8 patients.

Exactly 64 patients of the resection group and 31 patients of the RFA group died during the follow-up. The cause of death among patients in the resection group was tumor progression in 51, hepatic failure in 10, gastrointestinal bleeding in 1, and other causes in 2. Uniformly, the cause of death in the RFA group was tumor progression in 16, hepatic failure in 13, gastrointestinal bleeding in 1, and other causes in 1. The cumulative survival rates for the resection group at 1, 3, 5, and 7 years were 96.9%,




FIGURE 1. Cumulative recurrence free survival rates of the patients who underwent surgical resection (solid line) and radiofrequency ablation (RFA) (dotted line) are shown. The recurrence-free survival rate of the resection group was higher than of the RFA group ($P = .011$).

90.3%, 79.0%, and 61.5%, whereas those for the RFA group were 99.0%, 87.4%, 74.8%, and 65.4%, respectively. The overall survival rates were not significantly different between the 2 groups.

The tumor recurrence-free survival rates for the resection group at 1, 3, 5, and 7 years were 83.1%, 51.0%, 36.8%, and 23.3%, and for the RFA group were 82.7%, 41.8%, 17.0%, and 5.8%, respectively (Fig. 1). The recurrence-free survival rate was higher in the resection group than in the RFA group ($P = .011$).

Factors Associated With Survival in Patients in the Resection Group

Among 199 patients treated with surgical resection, factors associated with survival were evaluated by both univariate and multivariate analysis (Table 2). Single tumor, serum albumin >3.8 g/dL, and prothrombin time $>80\%$ were significant by Kaplan-Meier analysis, whereas factors such as age, sex, etiology of background liver disease, amount of alcohol intake, bilirubin, AST, platelet count, AFP, DCP, diameter of the HCC, and tumor vascularity were not significantly related to patients' survival. In a multivariate analysis using the Cox proportional hazard model, prothrombin time $>80\%$ (hazard ratio [HR], 2.72; 95% confidence interval [CI], 1.56-4.74; $P < .001$) was the independent prognostic factor for survival.

Table 2. Results of Univariate and Multivariate Analyses of Factors Associated With Survival of Patients Treated by Surgical Resection

Variable	No.	% 5-Year Survival Rate	Hazard Ratio (95% CI)	P
Univariate analysis				
No. of tumors				.012
Single	167	82.4	—	
Multiple	32	59.8	—	
Albumin, g/dL				.020
≥ 3.8	134	91.1	—	
< 3.8	64	73.1	—	
Prothrombin time (%)				.003
≥ 80	168	83.0	—	
< 80	29	60.1	—	
Multivariate analysis				
Prothrombin time, %, ≥ 80 / < 80			2.72 (1.56-4.74)	$< .001$

CI indicates confidence interval.

Table 3. Results of Univariate and Multivariate Analyses of Factors Associated With Survival in Patients Treated With Radiofrequency Ablation

Variable	No.	% 5-Year Survival Rate	Hazard Ratio (95% CI)	P
Univariate analysis				
Platelets, $\times 10^4/\text{mm}^3$.006
≥ 10	111	84.1	—	
< 10	98	65.5	—	
DCP, AU/mL				$< .001$
< 100	187	77.3	—	
≥ 100	13	33.6	—	
Multivariate analysis				
DCP, AU/L, < 100 / ≥ 100			5.49 (2.23-13.5)	$< .001$
Platelets, $\times 10^4/\text{mm}^3$, ≥ 10 / < 10			2.70 (1.24-5.88)	.012

CI indicates confidence interval; DCP, des-gamma-carboxy prothrombin; AU, arbitrary units.

Factors Associated With Survival in Patients in the RFA Group

We also evaluated the factors associated with the survival of 209 patients treated with RFA (Table 3). Platelet count $\geq 1.0 \times 10^5$ and DCP < 100 arbitrary units (AU)/L were significant in univariate analysis, whereas 12 other variables were not associated with survival. Multivariate analysis identified DCP < 100 AU/L (HR, 5.49; 95% CI, 2.23-13.5; $P < .001$) and platelet count $\geq 1.0 \times 10^5$ (HR, 2.70; 95% CI, 1.24-5.88; $P = .012$) as significant and independent determinants of survival.

Factors Associated With Recurrence-free Survival in Patients in the Resection Group

Next, we evaluated the factors associated with recurrence-free survival in patients treated with surgical resection (Table 4). Presence of a single tumor, serum albumin ≥ 3.8 g/dL, platelet count $\geq 1.0 \times 10^5$, and prothrombin time $\geq 80\%$ were significant in the univariate analysis, whereas 10 other variables were not significant factors for recurrence-free survival. In multivariate analysis, single tumor (HR, 2.39; 95% CI, 1.51-3.80; $P < .001$), serum albumin ≥ 3.8 g/dL (HR, 1.54; 95% CI, 1.02-2.32;

Table 4. Results of Univariate and Multivariate Analyses of Factors Associated With Recurrence-free Survival Among Patients Treated With Resection

Variable	No.	% 3-Year Survival Rate	Hazard Ratio (95% CI)	P
Univariate analysis				
No. of tumors				<.001
Single	166	55.4	—	
Multiple	32	24.5	—	
Albumin, g/dL				.009
≥ 3.8	63	71.7	—	
< 3.8	134	42.1	—	
Platelets, $\times 10^4/\text{mm}^3$.025
≥ 10	137	60.3	—	
< 10	60	33.0	—	
Prothrombin time, %				.009
≥ 80	167	55.7	—	
< 80	29	29.0	—	
Multivariate analysis				
No. of tumors, single/multiple			2.39 (1.51-3.80)	<.001
Albumin, g/dL, $\geq 3.8/< 3.8$			1.54 (1.02-2.32)	.040
Platelets, $\times 10^4/\text{mm}^3$, $\geq 10/< 10$			1.47 (1.03-2.12)	.036

CI indicates confidence interval.

$P = .040$), and platelet count $\geq 1.0 \times 10^5$ (HR, 1.47; 95% CI, 1.03-2.12; $P = .036$) were independent prognostic factors for recurrence-free survival.

Factors Associated With Recurrence-free Survival in Patients in the RFA Group

Factors associated with recurrence-free survival were evaluated in patients treated by RFA. The 3-year recurrence-free survival rate was 44.7% in 185 patients with DCP < 100 AU/L, whereas it was 0.0% in 13 patients with DCP ≥ 100 AU/L. Univariate and multivariate analysis identified only DCP < 100 AU/L (HR, 6.82; 95% CI, 3.49-13.3; $P < .001$) as a significant determinant of recurrence-free survival.

Survival and Recurrence-free Survival in Patients With DCP > 100 AU/L

Figure 2 shows the cumulative survival rate, and Figure 3 shows the recurrence-free survival rate based on DCP levels. The survival rate and recurrence-free survival rate were associated with DCP in the RFA group, but they were not associated with DCP in the resection group. AFP > 400 AU/L was associated with neither survival rate nor recur-

rence-free survival rate in either the resection or the RFA group. Therefore, 27 selected patients from the resection group and 13 from the RFA group whose DCP was ≥ 100 AU/L were examined to determine whether the overall survival rate was different between the resection and RFA groups. The backgrounds of the 2 groups based on treatment procedure are shown in Table 5. Treatment procedure (resection), age < 65 years and serum albumin > 3.8 g/dL were significant in the univariate analysis. Multivariate analysis revealed that treatment procedure (HR, 1.26; 95% CI, 1.04-1.53; $P = .020$) was a significant and independent determinant in the overall survival rate (Table 6).

DISCUSSION

Patients with HCC usually have a history of chronic liver disease, especially cirrhosis. Unfortunately, even when curative therapy is performed, tumor recurrence is frequent. For this reason, less invasive treatment procedures are needed to preserve liver function.

The Barcelona Clinic Liver Cancer (BCLC) guideline for the treatment of HCC recommends resection for patients with a single HCC and Child-Pugh A who have no other complications.⁵ The suggested option for RFA includes patients with multiple tumors and associated




FIGURE 2. Cumulative overall survival rate of the patients who underwent surgical resection and radiofrequency ablation based on des-gamma-carboxy prothrombin (DCP) level is shown. Solid line indicates DCP level ≥ 100 AU/L; dotted line, DCP level < 100 AU/L. (A) Cumulative survival rate of patients who underwent resection based on DCP level is shown. (B) Cumulative survival rate of patients who underwent radiofrequency ablation (RFA) based on DCP level is shown. Prognosis of patients who underwent RFA varied according to DCP level, whereas prognosis of patients who underwent resection was independent of DCP level.




FIGURE 3. Cumulative recurrence-free survival (RFS) rate of the patients who underwent surgical resection and radiofrequency ablation based on des-gamma-carboxy prothrombin (DCP) level is shown. Solid line indicates DCP level ≥ 100 AU/L; dotted line, DCP level < 100 AU/L. (A) Cumulative RFS rate of patients who underwent resection based on DCP level is shown. (B) Cumulative RFS rate of patients who underwent RFA based on DCP level is shown. Prognosis of patients who underwent RFA varied according to DCP level.

disease. However, in clinical practice, RFA is widely applied as a curative treatment for variable stages of HCC. Advances in imaging diagnosis have allowed identification of small HCC measuring < 2 cm during the course of chronic liver disease.³ The use of RFA seems to be an excellent option for the aforementioned tumors.

In the present study, we focused on the malignant potential of HCC and examined 2 representative tumor

markers of HCC. AFP has been used as a tumor marker for HCC worldwide, and is considered by some as a predictor of survival or recurrence after RFA.¹² DCP is also useful as a prognostic factor in patients with HCC.²⁰⁻²²

In the present study, serum albumin levels and prothrombin time, which reflect liver function, were significantly associated with survival in patients who undergo resection similar to tumor multiplicity. Likewise, serum

Table 5. Clinical Background of 27 Patients Who Underwent Resection and 13 Patients Who Underwent Radiofrequency Ablation

Factors	Resection Group, n = 27	RFA Group, n = 13	P
Age, y*	60 (35-73)	67 (50-78)	.006
Sex (men:women)	23:4	10:3	NS
HBV:HCV:others	13:12:2	1:9:3	.015
Habitual alcohol intake, yes: no	3:24	2:11	NS
Diameter of HCC, mm*	22 (14-30)	22 (10-30)	NS
No. of HCC, single:multiple	26:1	9:4	.031
Tumor vascularity, present:absent	27:0	12:1	NS
Albumin, g/dL*	3.8 (3.3-4.3)	3.4 (2.6-4.1)	.003
Bilirubin, mg/dL*	0.9 (0.4-1.9)	0.9 (0.4-1.8)	NS
AST, IU/L*	38 (16-240)	49 (17-145)	NS
Platelets, $\times 10^4/\text{mm}^3$ *	15.1 (6.0-24.5)	10.5 (4.5-24.6)	.025
Prothrombin time, %*	94 (79-112)	86 (73-110)	NS

*Data are expressed as median (range).

Table 6. Results of Univariate and Multivariate Analyses of Factors Associated With Survival Among Patients With Serum Des-gamma-carboxy Prothrombin Level ≥ 100

Variable	No.	% 5-Year Survival Rate	Hazard Ratio (95% CI)	P
Univariate analysis				
Age, y				.026
<65	25	87.7	—	
≥ 65	15	45.5	—	
Albumin, g/dL				.024
≥ 3.8	12	100	—	
<3.8	28	58.4	—	
Treatment procedure				.012
Resection	27	83.4	—	
RFA	13	33.6	—	
Multivariate analysis				
Treatment procedure				.020
Resection or RFA			1.26 (1.04-1.53)	

CI indicates confidence interval; RFA, radiofrequency ablation.

albumin level, platelet count, prothrombin time, and presence of multiple tumors were associated with recurrence-free survival. Alternatively, in RFA patients, in addition to platelet count, which indicates severity of portal hypertension, DCP levels were significant predictors of survival. Likewise, DCP levels were also significant predictors in recurrence-free survival. It is noteworthy that both survival and disease-free survival rates of patients who undergo RFA, but not resection, are correlated with DCP levels by multivariate analysis.

It is difficult to explain why DCP influenced survival and disease-free survival in the RFA group but not the resection group. We speculate that a high level of serum tumor marker reflects a high tumor malignant potential. Therefore, for a biologically aggressive tumor like HCC, resection is recommended over RFA because the radical nature of surgical resection may be superior to RFA.

According to previous reports, DCP is related to histological features of HCC.^{21,22} Shirabe et al²¹ examined

218 HCC patients who underwent surgical resection for HCC and concluded that serum DCP level is a predictor of microvascular invasion. They identified microvascular invasion in 44% of their patients with DCP ≥ 100 AU/L, but in only 16% of patients with DCP < 100 AU/L. Shimada et al²² examined explanted liver transplants and reported that serum DCP level is associated with vascular invasion and HCC recurrence. When we evaluated the relationship between clinicopathological features and serum DCP level in the resection group, microvascular invasion was found in 11 (44.0%) of 25 patients with DCP ≥ 100 AU/L, but in only 22 (13.6%) of 162 patients with DCP < 100 AU/L, which was similar to the results in Shirabe et al.²¹

Unfortunately, in the RFA group, histological examination was performed only on nodules that showed atypical image findings. Moreover, it is sometimes difficult to judge microscopic vascular invasion in small specimens obtained by needle biopsy. However, we cannot deny the presence of microscopic vascular invasion in patients with high levels of tumor markers who have been treated with RFA. Furthermore, high levels of DCP are not only a marker of malignancy, but also indicate the biologic aggressiveness and progression of the HCC tumor. Hence, HCCs with high levels of DCP have greater chances of hypervascularity and early infiltration than do HCCs with lower levels of DCP.

In general, microscopic vascular invasion or intrahepatic metastasis is a poor prognostic factor for survival and recurrence-free survival even in patients who undergo surgical resection.²³⁻²⁵ Why was survival and recurrence-free survival not different among the resection group in our study? One of the reasons is that we included patients with an HCC of a maximum diameter of ≤ 3 cm. In contrast, most previous studies included HCC as large as 5 cm in diameter.²³⁻²⁵ The biological features of malignancy might be worse in such large tumors. We speculate that in HCC measuring ≤ 3 cm (median 2.0 cm), minimal microscopic vascular invasion or intrahepatic metastasis adjacent to the main tumor can be curatively resected by surgery, whereas these sometimes become incompletely necrotic even when a sufficient surrounding margin is obtained from treatment with RFA.

Analysis of factors associated with survival in patients with DCP of ≥ 100 AU/L showed that the type of treatment procedure (eg, hepatectomy) significantly

influenced outcome. In contrast, no such relationship was found in patients with DCP of < 100 AU/L. These results indicate that DCP is an important factor in selecting treatment procedure for patients with HCC measuring ≤ 3 cm and numbering ≤ 3 tumors.

In conclusion, DCP levels were significant predictors of both survival and recurrence-free survival in the RFA group. Hence, when the level of DCP is high, hepatic resection should be the treatment of choice even if the maximum tumor diameter is ≤ 3 cm and there are ≤ 3 tumors. If the level is low, RFA should be considered, because it is less invasive.

Because the current study was retrospective in nature, it has certain limitations and potential biases. The baseline characteristics of the 2 groups were quite different. Although we enrolled only Child-Pugh A patients, the resection group was younger and had better liver function. However, etiology of the liver disease was also different; the overall survival rates were not significantly based on etiology in either the resection or the RFA group. Therefore, we believe that the etiology of liver disease could be ignored in these patients. Our study did not uncover the reason for the high risk of mortality and tumor recurrence in patients in the RFA group with high levels of DCP. A cohort validation study is needed to confirm our results. In addition, clinicopathological and molecular analyses are also needed to define the biological significance of the biomarker.

Conflict of Interest Disclosures

Supported in part by a research grant from the Japanese Ministry of Health, Labor, and Welfare, and the Okinaka Memorial Foundation of Toranomon Hospital.

References

1. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin.* 2005;55:74-108.
2. Kiyosawa K, Umemura T, Ichijo T, et al. Hepatocellular carcinoma: recent trend in Japan. *Gastroenterology.* 2004; 127:S17-S26.
3. Ikeda K, Saitoh S, Koida I, et al. A multivariate analysis of risk factors for hepatocellular carcinogenesis: a prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology.* 1993;18:47-53.
4. Mazzaferro V, Regalia E, Doci R, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med.* 1996;334:693-699.

5. Bruix J, Sherman M. Practice Guidelines Committee, American Association for the Study of Liver Diseases. Management of hepatocellular carcinoma. *Hepatology*. 2005;42:1208-1236.
6. Llovet JM, Bru C, Bruix J. Prognosis of hepatocellular carcinoma: the BCLC staging classification. *Semin Liver Dis*. 1999;19:329-338.
7. Cha C, DeMatteo RP, Blumgart LH. Surgery and ablative therapy for hepatocellular carcinoma. *J Clin Gastroenterol*. 2002;35(5 suppl 2):S130-S137.
8. Fan ST, Ng IO, Poon RT, Lo CM, Liu CL, Wong J. Hepatectomy for hepatocellular carcinoma: the surgeon's role in long-term survival. *Arch Surg*. 1999;134:1124-1130.
9. Chen MS, Li JQ, Zheng Y, et al. A prospective randomized trial comparing percutaneous local ablative therapy and partial hepatectomy for small hepatocellular carcinoma. *Ann Surg*. 2006;243:321-328.
10. Lu MD, Kuang M, Liang LJ, et al. Surgical resection versus percutaneous thermal ablation for early-stage hepatocellular carcinoma: a randomized clinical trial [in Chinese]. *Zhonghua Yi Xue Za Zhi*. 2006;86:801-805.
11. Lencioni R, Cioni D, Crocetti L, et al. Early-stage hepatocellular carcinoma in patients with cirrhosis: long-term results of percutaneous image-guided radiofrequency ablation. *Radiology*. 2005;234:961-967.
12. Choi D, Lim HK, Rhim H, et al. Percutaneous radiofrequency ablation for early-stage hepatocellular carcinoma as a first-line treatment: long-term results and prognostic factors in a large single-institution series. *Eur Radiol*. 2007;17:684-692.
13. Livraghi T, Meloni F, Di Stasi M, et al. Sustained complete response and complications rates after radiofrequency ablation of very early hepatocellular carcinoma in cirrhosis: Is resection still the treatment of choice? *Hepatology*. 2008;47:82-89.
14. Vauthey JN, Lauwers GY, Esnaola NF, et al. Simplified staging for hepatocellular carcinoma. *J Clin Oncol*. 2002;20:1527-1536.
15. Liver Cancer Study Group of Japan. Classification of Primary Liver Cancer. First English Edition. Tokyo, Japan: Kanehara Press Co.; 1997.
16. Shimada M, Takenaka K, Fujiwara Y, et al. Des-gamma-carboxy prothrombin and alpha-fetoprotein positive status as a new prognostic indicator after hepatic resection for hepatocellular carcinoma. *Cancer*. 1996;78:2094-2100.
17. Imamura H, Matsuyama Y, Miyagawa Y, et al. Prognostic significance of anatomical resection and des-gamma-carboxy prothrombin in patients with hepatocellular carcinoma. *Br J Surg*. 1999;86:1032-1038.
18. Kobayashi M, Ikeda K, Someya T, et al. Stepwise hook extension technique for radiofrequency ablation therapy of hepatocellular carcinoma. *Oncology*. 2002;63:139-144.
19. Cox DR. Regression models and life tables. *J R Stat Soc*. 1972;34:187-220.
20. Nagaoka S, Yatsushashi H, Hamada H, et al. The des-gamma-carboxy prothrombin index is a new prognostic indicator for hepatocellular carcinoma. *Cancer*. 2003;98:2671-2677.
21. Shirabe K, Itoh S, Yoshizumi T, et al. The predictors of microvascular invasion in candidates for liver transplantation with hepatocellular carcinoma-with special reference to the serum levels of des-gamma-carboxy prothrombin. *J Surg Oncol*. 2007;95:235-240.
22. Shimada M, Yonemura Y, Ijichi H, et al. Living donor liver transplantation for hepatocellular carcinoma: a special reference to a preoperative des-gamma-carboxy prothrombin value. *Transplant Proc*. 2005;37:1177-1179.
23. Sasaki A, Iwashita Y, Shibata K, Matsumoto T, Ohta M, Kitano S. Improved long-term survival after liver resection for hepatocellular carcinoma in the modern era: retrospective study from HCV-endemic areas. *World J Surg*. 2006;30:1567-1578.
24. Grazi GL, Cescon M, Ravaioli M, et al. Liver resection for hepatocellular carcinoma in cirrhotics and noncirrhotics. Evaluation of clinicopathologic features and comparison of risk factors for long-term survival and tumour recurrence in a single centre. *Aliment Pharmacol Ther*. 2003;17(suppl 2):119-129.
25. Hanazaki K, Kajikawa S, Koide N, Adachi W, Amano J. Prognostic factors after hepatic resection for hepatocellular carcinoma with hepatitis C viral infection: univariate and multivariate analysis. *Am J Gastroenterol*. 2001;96:1243-1250.

An Open Pilot Study Exploring the Efficacy of Fluvastatin, Pegylated Interferon and Ribavirin in Patients with Hepatitis C Virus Genotype 1b in High Viral Loads

Hitomi Sezaki^a Fumitaka Suzuki^a Norio Akuta^a Hiromi Yatsuji^a
Tetsuya Hosaka^a Masahiro Kobayashi^a Yoshiyuki Suzuki^a Yasuji Arase^a
Kenji Ikeda^a Yuzo Miyakawa^b Hiromitsu Kumada^a

^aDepartment of Hepatology, Toranomon Hospital, and ^bMiyakawa Memorial Research Foundation, Tokyo, Japan

Key Words

Hepatitis, chronic · Virus, hepatitis C · Statin · Interferon, pegylated · Ribavirin

Abstract

Objective: Response to pegylated (PEG) interferon (IFN) and ribavirin is achieved only in 40–50% of patients infected with hepatitis C virus (HCV) of genotype 1 in high viral loads, which needs to be improved. **Methods:** In an open-label pilot study, fluvastatin (HMG-CoA reductase inhibitor), 20 mg daily, was given along with PEG-IFN/ribavirin to 21 patients with chronic hepatitis C. They were followed for HCV RNA in serum. **Results:** During treatment for 48 weeks, HCV RNA was lost from serum in 93% of the patients. In the 15 patients who received 48-week therapy, a sustained virological response (SVR) with loss of HCV RNA 24 weeks after completion was achieved in 10 (67%), including 7 of the 9 (78%) male and 3 of the 6 (50%) female patients. In the remaining 6 patients who received 72-week therapy, SVR was gained in 4 (67%), including 1 of the 2 male and 3 of the 4 female patients aged 56, 58 and 62 years, respectively. **Conclusion:** Fluvastatin could be used safely to increase the response to PEG-IFN and ribavirin, especially in aged women who respond poorly to combined PEG-IFN/ribavirin.

Copyright © 2009 S. Karger AG, Basel

Introduction

Over the world approximately 190 million people are persistently infected with hepatitis C virus (HCV) [1], and about 30% of them develop serious liver disease such as decompensated cirrhosis and hepatocellular carcinoma (HCC) [2]. Interferon (IFN) is the only drug that can resolve HCV infection. However, using the most advanced treatment currently available, with pegylated (PEG)-IFN and ribavirin for 24–48 weeks, a sustained virological response (SVR) defined by the loss of HCV RNA from serum 24 weeks after completion of therapy is achieved in 40–80% of the patients [3–5]. The response rate is influenced by host factors such as sex, age and ethnicity [6–8], as well as virological factors including genotypes and viral loads [9]. It remains unsatisfactorily low in patients infected with HCV genotype 1 at merely 40–50%, and dismal in women aged 50 years or older at merely 22% [10]. Hence, there is an imminent need to improve the efficacy of antiviral treatment for terminating HCV infection in these patients.

Efforts have been devoted toward increasing the efficacy to PEG-IFN/ribavirin therapy. Inhibitors of viral protease, alone or in combination with the IFN-based treatment, have been found effective in preliminary studies [11, 12]. Recently, drugs that can inhibit the key enzyme

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2009 S. Karger AG, Basel
0300-5526/09/0521-0043\$26.00/0

Accessible online at:
www.karger.com/int

Hitomi Sezaki, MD
Department of Hepatology, Toranomon Hospital
Minato-ku
Tokyo 105-8470 (Japan)
Tel. +81 44 877 5111, Fax +81 44 860 1623, E-Mail hitomis@mx1.harmonix.ne.jp

for controlling the synthesis of cholesterol, 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase, have gained attention due to their potential to decrease the replication of HCV in clinical and experimental settings [13, 14].

More than three quarters of Japanese patients are infected with HCV genotype 1b in high viral loads. They are much older than Western patients owing to the widespread HCV infection which struck Japan about 20 years ago [15]. Owing to this, the response to PEG-IFN/ribavirin in Japanese patients with chronic hepatitis C is poorer than that in Western countries. In an open-label pilot study, 21 patients with chronic hepatitis C, who were infected with HCV-1b at high viral loads, received triple treatment with fluvastatin, PEG-IFN and ribavirin. On-treatment viral dynamics and SVR achieved by the triple therapy could offer hope for improving the response in patients infected with HCV-1 in high viral loads.

Patients and Methods

From December 2005 to March 2006, 21 patients with chronic hepatitis C agreed to receive fluvastatin, in addition to standard PEG-IFN/ribavirin treatment, at the Department of Hepatology, Toranomon Hospital, Metropolitan Tokyo. They all were: (1) positive for antibody to HCV (anti-HCV) and HCV RNA of genotype 1b, and not co-infected with HCV of the other genotypes; (2) negative for hepatitis B surface antigen or antibody to human immunodeficiency virus type-1 (HIV-1); (3) confirmed within the past 2 months to have high HCV RNA levels of ≥ 100 KIU/ml, which is the Japanese definition of high viral loads [16, 17]; (4) platelet counts $>80 \times 10^3/\text{mm}^3$ and without cirrhosis diagnosed by ultrasonography; (5) body weight >40 kg and not pregnant or lactating; (6) a total alcohol intake of <500 kg during the past; (7) without HCC, hemochromatosis, Wilson's disease, primary biliary cirrhosis, alcoholic hepatitis or autoimmune hepatitis; (8) without antiviral or immunosuppressive treatment during the previous 3 months, and (9) with the wish to comply to the treatment protocol for 48–72 weeks.

They were followed for liver function and virological markers at least monthly during treatment and until 24 weeks after completion of triple treatment. Informed consent was obtained from each patient, and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution's human research committee.

Markers of HCV Infection

Anti-HCV was determined by third-generation enzyme-linked immunosorbent assay (ELISA) using commercial kits (Ortho HCV Ab ELISA Test 3; Chiron Cooperation, Emeryville, Calif., USA). HCV RNA was determined quantitatively by polymerase chain reaction (PCR; Cobas Amplicor HCV Monitor ver. 2.0, Roche Diagnostics, Tokyo, Japan), with a dynamic range from 5 to 5,000 KIU/ml, in sera diluted 10-fold at baseline, as well as every 2 weeks until 8 weeks after initiation of therapy and monthly thereafter. Sera negative for HCV RNA (<5 KIU/ml) by the

quantitative assay were tested by qualitative PCR (Amplicor, Roche Molecular Systems, Inc., Branchburg, N.J., USA) with the detection limit at 100 copies/ml.

Fluvastatin Added to Combined PEG-IFN and Ribavirin

Patients received subcutaneous PEG-IFN- $\alpha 2b$ (PEG-Intron, Schering-Plough Corp., Kenilworth, N.J., USA) weekly at a median dose of 1.5 (range 1.3–1.7) $\mu\text{g}/\text{kg}$, along with oral ribavirin daily at a median dose of 12.3 (range 10.2–13.7) mg/kg for 48–72 weeks. The dose of ribavirin was adjusted according to body weight: 600 mg for patients weighing ≤ 60 kg; 800 mg for those between >60 and <80 kg, and 1,000 mg for those ≥ 80 kg. In addition, the patients received 20 mg oral fluvastatin daily for 48–72 weeks.

Results

Patients Who Received Triple Treatment with Fluvastatin, PEG-IFN and Ribavirin

The baseline characteristics of the 21 patients infected with HCV-1b at a high viral load and who received triple therapy with fluvastatin, PEG-IFN and ribavirin are listed in table 1. Their median age was 56 years, and 11 (52%) were men. Their hematological and biochemical values including serum lipids were within normal limits, except for elevated levels of AST and ALT. A lack of mutations affecting the response to IFN in the core protein, i.e., mutations at positions 70 and 91 [16], and amino acid substitutions in the IFN-sensitivity determining region [18] were detected in HCV RNA from 29 and 35% of the patients.

Loss of HCV RNA from Serum in Patients with Triple Treatment

Figure 1 illustrates the cumulative loss of HCV RNA from the serum in 21 patients infected with HCV-1b at high loads. HCV RNA was cleared in 52% of the patients during the first 12 weeks. It decreased slowly thereafter, and was lost in 93% of the patients at 48 weeks. Triple treatment was continued in 6 of the 21 patients for an additional 24 weeks. HCV RNA was cleared from the serum in 5 (83%) of them at completion of the 72-week treatment.

Sustained Virological Response in Patients Receiving Triple Treatment

Of the 21 patients given triple treatment, 19 received it for 48 weeks. Skin rashes and general malaise developed in 2 of them at weeks 17 and 24, respectively, and fluvastatin was withdrawn while PEG-IFN and ribavirin were continued. Six patients continued to receive triple treatment for an additional 24 weeks. Therefore, the loss of HCV RNA from the serum 24 weeks after completion of

Table 1. Baseline characteristics of patients (n = 21) with chronic hepatitis C who received triple treatment with fluvastatin, PEG-IFN and ribavirin

	Normal ranges	Patients
Age, years	NA	56 (32–63)
Men	NA	11 (52%)
Albumin, g/dl	3.9–5.2	3.8 (3.4–4.2)
Hemoglobin, g/dl	11.3–17.0	14.4 (12.1–16.4)
Platelets, $\times 10^3/\text{mm}^3$	141–350	190 (119–240)
AST, IU/l	13–33	50 (21–208)
ALT, IU/l	6–42	64 (23–391)
γ -GTP	9–109	36 (11–301)
ICG ₁₅ , %	<10	12 (2–27)
HCV RNA, KIU/ml	NA	2,000 (14– >5,000)
Past treatments	NA	8 (36%)
Ribavirin dose, mg/kg	NA	12.3 (10.2–13.7)
Serum lipids		
Total cholesterol, mg/dl	122–240	163 (124–273)
LDL-C, mg/dl	86–160	99 (59–187)
HDL-C, mg/dl	35–75	46 (28–73)
Mutations in HCV RNA		
Wild-type at aa70	NA	12 (57%)
Wild-type at aa91	NA	10 (48%)
Double wild, aa71/aa90	NA	6 (29%)
Mutations in ISDR	NA	6/17 (35%)

Data are expressed as the number of patients with percentages in parentheses or median values with ranges in parentheses. AST = Aspartate aminotransferase; ALT = alanine aminotransferase; γ -GTP = γ -glutamyltranspeptidase; ICG₁₅ = retention of indocyanine green at 15 min; LDL-C = low-density lipoprotein cholesterol; HDL-C = high-density lipoprotein cholesterol; NA = not applicable.

therapy was assessed in the remaining 15 patients. Figure 2 shows the SVR in 15 patients. SVR was accomplished in 10 (67%) of them, including 7 of 9 (78%) male patients and 3 of 6 (50%) female patients. Of the 5 female patients aged >50 years, 2 (40%) gained SVR.

Of the 6 patients who received triple treatment for 72 weeks, SVR was accomplished in 4 (67%), including 1 of 2 male and 3 of 4 female patients who were aged 56, 58 and 62 years, respectively.

Discussion

Statins comprise a group of drugs capable of inhibiting HMG-CoA reductase and can regulate the synthesis of cholesterol by competing with the authentic substrate [19,




Fig. 1. Kaplan-Meier table illustrating the cumulative loss of HCV RNA from the serum in patients with chronic hepatitis C. The patients received fluvastatin in addition to PEG-IFN/ribavirin and were followed during 48 weeks of treatment.




Fig. 2. Sustained virological response in patients with chronic hepatitis C to the combined treatment with fluvastatin, PEG-IFN and ribavirin for 48 weeks. SVR was compared among the total, male and female patients, as well as the female patients aged more than 50 years.

20]. They are the best selling drugs in the world, and have been prescribed to more than 30 million patients for lowering serum levels of low-density lipoprotein cholesterol (LDL-C) with the main purpose of preventing cardiovascular accidents [21, 22]. The great virtue of statins is their excellent safety record. Recipients are rarely forced to discontinue statins due to mild side effects including myalgia, headache and skin rashes, thereby allowing their long-term use to be extended to decades and even for life. Liver toxicity is found in less than 1.2% of 49,275 patients with a history of cardiovascular disease [23].

In transfection experiments of cultured hepatocytes, statins inhibited the replication of HCV in synergism with IFN- α [14]. Of particular note, the extent of inhibition differs among various statins, and is highest for lipophilic fluvastatin and lowest for hydrophilic pravastatin. In clinical studies, monotherapy with fluvastatin suppressed serum HCV RNA by log 1.75 at a daily therapeutic dose of <80 mg [13], while no such effects were gained with 20 mg atorvastatin daily [24]. Hence, the ability to inhibit HCV replication may differ among diverse statins through mechanisms not directly associated with lowered levels of LDL-C; fluvastatin decreases total cholesterol and LDL-C levels to a lesser extent than the other statins in current use.

In this open-label pilot study, the response to triple treatment with fluvastatin 20 mg daily supplementing the standard PEG-IFN/ribavirin was examined in 21 patients infected with HCV-1b in high loads. The cumulative loss of HCV RNA from the serum was gained in 93% of the patients after they had received triple treatment for 48 weeks. Only 2 of the 21 (10%) patients did not tolerate fluvastatin due to skin rashes and general malaise. The triple treatment was continued for 72 weeks in 6 patients to determine its long-term effects. SVR was achieved in 10 of 15 (67%) patients on the 48-week schedule, including 7 of 9 (78%) male patients and 3 of 6 (50%) female patients. Of the 5 female patients aged >50 years, 2 (40%) gained SVR. Of the 6 patients who received triple treatment for 72 weeks, 4 (67%) achieved SVR, including 1 of 2 male and 3 of 4 female patients who were aged 56, 58 and 62 years, respectively. SVR gained by triple treatment with fluvastatin, PEG-IFN and ribavirin for 48 weeks (67%) is better than that 80 of 160 (50%) and 55 of 118 (47%) patients who received PEG-IFN and ribavirin for 48 weeks [16, 25]; they all were infected with HCV-1b at high loads.

Sex and age are the two principal host factors influencing the response to IFN-based treatment. Although women respond better to combined IFN and ribavirin than

men in large-scale multinational studies in Western countries [8, 26], target patients were rather young with the median age of about 40–41 years. Sex-dependent responsiveness to IFN is reversed with age, however, and the response to PEG-IFN/ribavirin is much poorer in female than male patients aged 50 years or older (22/100 [22%] vs. 50/94 [53%], $p < 0.001$) [10]. In view of the estradiols that can prohibit fibrosis in experimental cirrhosis induced in rats [27], levels of female sex hormones, which are lower in women than men older than 50 years [28], could be responsible for the sex difference in response to IFN of patients with chronic hepatitis C. Supplementation of aged female patients with estradiols may increase the response, but it can accelerate physiological osteoporosis [29, 30]. Since statin therapy does not increase the risk of severe hepatotoxicity in patients with chronic hepatitis C [31], fluvastatin may provide aged women with an advantage in gaining SVR to PEG-IFN/ribavirin. Furthermore, fluvastatin does not augment the side effects of other drugs, such as cyclosporine A and warfarin, because it is metabolized by CYP2C9 in the liver, unlike other statins that are disposed by CYP3A4 as other drugs [32].

Should statins interfere with the replication of HCV, it remains an open question how they do it. At least three mechanisms may be conceived for the decrease in HCV replication in patients receiving statins. HCV circulates in association with LDL-C [33], and on that basis, the LDL-C receptor has been proposed for infection of hepatocytes [34–36]. HCV RNA replicates in association with lipid droplets in hepatocytes [37, 38], and therefore, the synthesis of cholesterol blocked by statins can reduce the basis on which HCV thrives. However, these two scenarios seem to stand at odds with the response to IFN being better in patients with higher LDL-C levels [39, 40]. Statins prohibit the synthesis of mevalonate that is modified into geranylgeranyl (containing 20 carbons) and then farnesyl (15 carbons) [19]. They both prenylate most cellular proteins to make them lipophilic toward associations with membranes expressing their biological activities [41]. As an extension to these, the geranylgeranylation of host proteins, which may be required for the replication of HCV, is implicated in the capacity of statins to downregulate HCV infection [42–44]. For that matter, most pleiotropic effects of statins, including a decreased incidence of dementia and Alzheimer disease [45–47], could be associated with the inhibition of prenylation rather than lowered levels of LDL-C in the serum [48].

This study is not without limitations. The number of patients receiving fluvastatin along with PEG-IFN and ribavirin is small and fails to gain statistical power in any comparison with those given PEG-IFN/ribavirin in previous studies. The ability of other statins in improving the response to PEG-IFN/ribavirin has not been examined to discover the mechanism of how fluvastatin improves the response to antiviral treatments. The promising results obtained in this study hopefully will invite further interest for planning clinical trials with extended numbers of patients and targeting elderly women in par-

ticular. Meanwhile, it may be worth looking back in the database to examine the response of patients receiving PEG-IFN/ribavirin for chronic hepatitis C, simultaneously with statins for controlling LDL-C levels, to see if they fared better than those not taking statins routinely.

Acknowledgment

This study was supported in part by grants from Okinaka Memorial Foundation in Toranomon Hospital of Japan.

References

- Cohen J: The scientific challenge of hepatitis C. *Science* 1999;285:26–30.
- Seeff LB: Natural history of chronic hepatitis C. *Hepatology* 2002;36:S35–S46.
- Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL Jr, Haussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Lin A, Hoffman J, Yu J: Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–982.
- Hadziyannis SJ, Sette H Jr, Morgan TR, Balan V, Diago M, Marcellin P, Ramadori G, Bodenheimer H Jr, Bernstein D, Rizzetto M, Zeuzem S, Pockros PJ, Lin A, Ackrill AM: Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140:346–355.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, Goodman ZD, Koury K, Ling M, Albrecht JK: Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001;358:958–965.
- Kanwal F, Hoang T, Spiegel BM, Eisen S, Dominitz JA, Gifford A, Goetz M, Asch SM: Predictors of treatment in patients with chronic hepatitis C infection – role of patient versus nonpatient factors. *Hepatology* 2007;46:1741–1749.
- Poynard T, Bedossa P, Opolon P: Natural history of liver fibrosis progression in patients with chronic hepatitis C. *Lancet* 1997;349:825–832.
- Conjeevaram HS, Fried MW, Jeffers LJ, Terrault NA, Wiley-Lucas TE, Afdhal N, Brown RS, Belle SH, Hoofnagle JH, Kleiner DE, Howell CD: Peginterferon and ribavirin treatment in African American and Caucasian American patients with hepatitis C genotype 1. *Gastroenterology* 2006;131:470–477.
- Tsubota A, Chayama K, Ikeda K, Yasuji A, Koida I, Saitoh S, Hashimoto M, Iwasaki S, Kobayashi M, Hiromitsu K: Factors predictive of response to interferon-alpha therapy in hepatitis C virus infection. *Hepatology* 1994;19:1088–1094.
- Sezaki H, Suzuki F, Kawamura Y, Yatsuji H, Hosaka T, Akuta N, Kobayashi M, Suzuki Y, Saitoh S, Arase Y, Ikeda K, Miyakawa Y, Kumada H: Poor response to pegylated interferon and ribavirin in aged women infected with hepatitis C virus of genotype 1b in high viral loads. *Dig Dis Sci*, in press.
- Lawitz E, Rodriguez-Torres M, Muir AJ, Kieffer TL, McNair L, Khunvichai A, McHutchison JG: Antiviral effects and safety of telaprevir, peginterferon alfa-2a, and ribavirin for 28 days in hepatitis C patients. *J Hepatol* 2008;49:163–169.
- Forestier N, Reesink HW, Weegink CJ, McNair L, Kieffer TL, Chu HM, Purdy S, Jansen PL, Zeuzem S: Antiviral activity of telaprevir (VX-950) and peginterferon alfa-2a in patients with hepatitis C. *Hepatology* 2007;46:640–648.
- Bader T, Fazili J, Madhoun M, Aston C, Hughes D, Rizvi S, Seres K, Hasan M: Fluvastatin inhibits hepatitis C replication in humans. *Am J Gastroenterol* 2008;103:1383–1389.
- Ikeda M, Abe K, Yamada M, Dansako H, Naka K, Kato N: Different anti-HCV profiles of statins and their potential for combination therapy with interferon. *Hepatology* 2006;44:117–125.
- Yoshizawa H, Tanaka J, Miyakawa Y: National prevention of hepatocellular carcinoma in Japan based on epidemiology of hepatitis C virus infection in the general population. *Intervirol* 2006;49:7–17.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictors of viral kinetics to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b. *J Med Virol* 2007;79:1686–1695.
- Sezaki H, Suzuki F, Kawamura Y, Yatsuji H, Hosaka T, Akuta N, Kobayashi M, Suzuki Y, Arase Y, Ikeda K, Kumada H: Evaluation of long-term biochemical responses to combination therapy of interferon plus ribavirin in those infected with hepatitis C virus genotype 1b and high baseline viral load. *Hepatol Res* 2007;37:787–792.
- Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Ogura Y, Izumi N, Marumo F, Sato C: Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996;334:77–81.
- Goldstein JL, Brown MS: Regulation of the mevalonate pathway. *Nature* 1990;343:425–430.
- Istvan ES, Deisenhofer J: Structural mechanism for statin inhibition of HMG-CoA reductase. *Science* 2001;292:1160–1164.
- Cannon CP, Braunwald E, McCabe CH, Rader DJ, Rouleau JL, Belder R, Joyal SV, Hill KA, Pfeffer MA, Skene AM: Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350:1495–1504.
- Grundy SM, Cleeman JI, Merz CN, Brewer HB Jr, Clark LT, Hunninghake DB, Pasternak RC, Smith SC Jr, Stone NJ: Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III Guidelines. *J Am Coll Cardiol* 2004;44:720–732.
- de Denu S, Spinler SA, Miller K, Peterson AM: Statins and liver toxicity: a meta-analysis. *Pharmacotherapy* 2004;24:584–591.

- 24 O'Leary JG, Chan JL, McMahon CM, Chung RT: Atorvastatin does not exhibit antiviral activity against HCV at conventional doses: a pilot clinical trial. *Hepatology* 2007;45:895-898.
- 25 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Miyakawa Y, Kumada H: Prediction of response to pegylated interferon and ribavirin in hepatitis C by polymorphisms in the viral core protein and very early dynamics of viremia. *Intervirology* 2007;50:361-368.
- 26 Poynard T, Marcellin P, Lee SS, Niederau C, Minuk GS, Ideo G, Bain V, Heathcote J, Zeuzem S, Trepo C, Albrecht J: Randomised trial of interferon alpha2b plus ribavirin for 48 weeks or for 24 weeks versus interferon alpha2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. *Lancet* 1998;352:1426-1432.
- 27 Yasuda M, Shimizu I, Shiba M, Ito S: Suppressive effects of estradiol on dimethylnitrosamine-induced fibrosis of the liver in rats. *Hepatology* 1999;29:719-727.
- 28 Khosla S, Melton LJ 3rd, Atkinson EJ, O'Fallon WM, Klee GG, Riggs BL: Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bioavailable estrogen. *J Clin Endocrinol Metab* 1998;83:2266-2274.
- 29 Need AG, O'Loughlin PD, Morris HA, Horowitz M, Nordin BE: The effects of age and other variables on serum parathyroid hormone in postmenopausal women attending an osteoporosis center. *J Clin Endocrinol Metab* 2004;89:1646-1649.
- 30 Siris ES, Brenneman SK, Barrett-Connor E, Miller PD, Sajjan S, Berger ML, Chen YT: The effect of age and bone mineral density on the absolute, excess, and relative risk of fracture in postmenopausal women aged 50-99: results from the National Osteoporosis Risk Assessment (NORA). *Osteoporosis Int* 2006;17:565-574.
- 31 Khorashadi S, Hasson NK, Cheung RC: Incidence of statin hepatotoxicity in patients with hepatitis C. *Clin Gastroenterol Hepatol* 2006;4:902-907.
- 32 Corsini A, Bellosta S, Baetta R, Fumagalli R, Paoletti R, Bernini F: New insights into the pharmacodynamic and pharmacokinetic properties of statins. *Pharmacol Ther* 1999;84:413-428.
- 33 Andre P, Komurian-Pradel F, Deforges S, Perret M, Berland JL, Sodayer M, Pol S, Brechot C, Paranhos-Baccala G, Lotteau V: Characterization of low- and very-low-density hepatitis C virus RNA-containing particles. *J Virol* 2002;76:6919-6928.
- 34 Molina S, Castet V, Fournier-Wirth C, Pichard-Garcia L, Avner R, Harats D, Roitelman J, Barbaras R, Graber P, Ghersa P, Smolarsky M, Funaro A, Malavasi F, Larrey D, Coste J, Fabre JM, Sa-Cunha A, Maurel P: The low-density lipoprotein receptor plays a role in the infection of primary human hepatocytes by hepatitis C virus. *J Hepatol* 2007;46:411-419.
- 35 Agnello V, Abel G, Elfahal M, Knight GB, Zhang QX: Hepatitis C virus and other flaviviridae viruses enter cells via low density lipoprotein receptor. *Proc Natl Acad Sci USA* 1999;96:12766-12771.
- 36 Monazahian M, Bohme I, Bonk S, Koch A, Scholz C, Grethe S, Thomssen R: Low density lipoprotein receptor as a candidate receptor for hepatitis C virus. *J Med Virol* 1999;57:223-229.
- 37 Aizaki H, Lee KJ, Sung VM, Ishiko H, Lai MM: Characterization of the hepatitis C virus RNA replication complex associated with lipid rafts. *Virology* 2004;324:450-461.
- 38 Miyazaki Y, Atsuzawa K, Usuda N, Watashi K, Hishiki T, Zayas M, Bartenschlager R, Wakita T, Hijikata M, Shimotohno K: The lipid droplet is an important organelle for hepatitis C virus production. *Nat Cell Biol* 2007;9:1089-1097.
- 39 Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H: Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 2007;46:403-410.
- 40 Gopal K, Johnson TC, Gopal S, Walfish A, Bang CT, Suwandhi P, Pena-Sahdala HN, Clain DJ, Bodenheimer HC Jr, Min AD: Correlation between beta-lipoprotein levels and outcome of hepatitis C treatment. *Hepatology* 2006;44:335-340.
- 41 Zhang FL, Casey PJ: Protein prenylation: molecular mechanisms and functional consequences. *Annu Rev Biochem* 1996;65:241-269.
- 42 Kapadia SB, Chisari FV: Hepatitis C virus RNA replication is regulated by host geranylgeranylation and fatty acids. *Proc Natl Acad Sci USA* 2005;102:2561-2566.
- 43 Wang C, Gale M Jr, Keller BC, Huang H, Brown MS, Goldstein JL, Ye J: Identification of FBL2 as a geranylgeranylated cellular protein required for hepatitis C virus RNA replication. *Mol Cell* 2005;18:425-434.
- 44 Ye J, Wang C, Sumpter R Jr, Brown MS, Goldstein JL, Gale M Jr: Disruption of hepatitis C virus RNA replication through inhibition of host protein geranylgeranylation. *Proc Natl Acad Sci USA* 2003;100:15865-15870.
- 45 Wolozin B, Kellman W, Ruosseau P, Celesia GG, Siegel G: Decreased prevalence of Alzheimer disease associated with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Arch Neurol* 2000;57:1439-1443.
- 46 Jick H, Zornberg GL, Jick SS, Seshadri S, Drachman DA: Statins and the risk of dementia. *Lancet* 2000;356:1627-1631.
- 47 Vaughan CJ: Prevention of stroke and dementia with statins: effects beyond lipid lowering. *Am J Cardiol* 2003;91:23B-29B.
- 48 Liao JK, Laufs U: Pleiotropic effects of statins. *Annu Rev Pharmacol Toxicol* 2005;45:89-118.

Poor Response to Pegylated Interferon and Ribavirin in Older Women Infected with Hepatitis C Virus of Genotype 1b in High Viral Loads

Hitomi Sezaki · Fumitaka Suzuki · Yusuke Kawamura · Hiromi Yatsuji · Tetsuya Hosaka · Norio Akuta · Masahiro Kobayashi · Yoshiyuki Suzuki · Satoshi Saitoh · Yasuji Arase · Kenji Ikeda · Yuzo Miyakawa · Hiromitsu Kumada

Received: 30 March 2008 / Accepted: 22 August 2008 / Published online: 28 October 2008
© Springer Science+Business Media, LLC 2008

Abstract *Background* Response to treatment in patients with chronic hepatitis C, with reference to age and gender, has not been examined fully. *Aim* The influence of gender and age on treatment with pegylated interferon (PEG-IFN) and ribavirin was evaluated in a retrospective study. *Methods* PEG-IFN and ribavirin were given for 48 weeks to 179 men and 121 women infected with hepatitis C virus (HCV) of genotype 1b in high viral loads (>100 kIU/ml). *Results* Sustained virological response at 24 weeks after treatment was poorer in women than men who were aged ≥ 50 years (22% vs 53%, $P < 0.001$). Among the patients aged ≥ 50 years who had received $\geq 80\%$ of the doses of PEG-IFN, ribavirin, or both, women responded less often than men (26% vs 64%, $P < 0.001$; 33% vs 61%, $P = 0.022$; and 32% vs 63%, $P = 0.016$; respectively). In multivariate analysis, male gender, retention of indocyanine green, ribavirin dose and compliance with therapy increased sustained virological response. *Conclusions* Response to combined PEG-IFN and ribavirin is poorer in female than male patients with hepatitis C who are aged ≥ 50 years, irrespective of compliance with treatment. Low estrogen levels in older women could be responsible for their impaired response to PEG-IFN and ribavirin.

Keywords Aging · Women · Chronic hepatitis C · Genotypes · Interferon · Ribavirin

Introduction

There are an estimated 170 million people worldwide that are chronically infected with hepatitis C virus (HCV) [1]. HCV can persist in 70–80% of individuals who have been exposed to it, and it can induce chronic liver disease, through cirrhosis to hepatocellular carcinoma (HCC) in approximately 30% of them until 30–40 years after they were infected [2–4]. A number of viral and host factors influence the velocity of fibrosis progression in chronic hepatitis C. Thus, stage and grade of hepatitis are more severe in patients who are infected with HCV genotype 1 in high viral loads [5–7]. Male gender, age and intake of alcohol accelerate fibrosis, as well [8–10].

Interferon (IFN) combined with ribavirin has been the most effective and favored treatment of chronic hepatitis C to date. The combined treatment with the standard IFN can terminate HCV-1 infection with high viral loads in approximately 20% [11], and that with pegylated IFN (PEG-IFN) in $>40\%$ [12]. Owing to hemolytic side effects, however, women are less tolerant to ribavirin [13]. Although the response to combined treatment has been shown to be better in women than in men in previous studies, there remains a possibility that it could be influenced by age. Hence, there is a need for the comparison of the response between men and women in different age groups.

Virological response to PEG-IFN and ribavirin at the end of a 48-week treatment (ETR), as well as sustained virological response (SVR) 24 weeks after the completion of therapy, was compared between 179 men and 121

H. Sezaki (✉) · F. Suzuki · Y. Kawamura · H. Yatsuji · T. Hosaka · N. Akuta · M. Kobayashi · Y. Suzuki · S. Saitoh · Y. Arase · K. Ikeda · H. Kumada
Department of Hepatology, Toranomon Hospital, Minato-ku, Tokyo 105-8470, Japan
e-mail: hitomis@mx1.harmonix.ne.jp

Y. Miyakawa
Miyakawa Memorial Research Foundation, Tokyo, Japan

women who were infected with HCV-1b in high viral loads. In patients aged ≥ 50 years, both ETR and SVR were poorer in women than in men, irrespective of the total dose of IFN, ribavirin or both.

Methods

Study Population

From December 2001 to February 2006, 490 consecutive patients with chronic hepatitis C received combination therapy with PEG-IFN and ribavirin at the Department of Hepatology in the Toranomon Hospital in Metropolitan Tokyo. The following inclusion criteria were met by 300 (61%) patients: they were (1) positive test results for antibodies to HCV (anti-HCV) and for HCV RNA genotype 1b by qualitative methods, and not co-infected with HCV of other genotypes; (2) negative test results for hepatitis B surface antigen or antibodies to human immunodeficiency virus type-1 (HIV-1); (3) confirmed findings of high HCV RNA levels ≥ 100 kIU/ml, which is the Japanese definition of high viral loads [14, 15], within the past 2 months; (4) no cirrhosis diagnosed by laparoscopy and ultrasonography, and with platelet counts $>80 \times 10^3/\text{mm}^3$; (5) body weight ≥ 40 kg and not pregnant or lactating; (6) total alcohol intake <500 kg in the past; (7) no HCC, hemochromatosis, Wilson's disease, primary biliary cirrhosis, alcoholic hepatitis or autoimmune hepatitis; (8) no treatment with antivirals or immunosuppressants during the previous 3 months; and (9) with the wish to comply with the treatment protocol for 48 weeks. None of them received growth factors before or during the study period.

The 300 patients, comprising 179 men and 121 women, received PEG-IFN and ribavirin for 48 weeks and were followed for at least 24 weeks after completion of this combination therapy. Informed consent was obtained from each patient, and the study protocol conformed to the ethics guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution's human research committee.

Serum Markers of HCV Infection

Anti-HCV was determined by third-generation enzyme-linked immunosorbent assay (ELISA) by commercial kits (Ortho HCV Ab ELISA Test 3; Chiron Cooperation, Emeryville, CA, USA). HCV RNA was determined quantitatively by polymerase chain reaction (PCR) (Cobas Amplicor HCV Monitor ver. 2.0, Roche Diagnostics, Tokyo, Japan) in serum diluted tenfold at the baseline, as well as at least monthly during and after treatment; it has a dynamic range between 5 kIU/ml and 5,000 kIU/ml. Sera

negative for HCV RNA (<5 kIU/ml) by quantitative assay were tested by qualitative PCR (Amplicor, Roche Molecular Systems, Inc., Branchburg, NJ, USA) with a detection limit at 100 copies/ml.

Combined PEG-IFN and Ribavirin Therapy

Patients underwent subcutaneous administration of PEG-IFN- $\alpha 2b$ (PEG-Intron, Schering-Plough Corp, Kenilworth, NJ, USA), weekly, at a median dose of 1.4 $\mu\text{g}/\text{kg}$ (range 0.8–1.9 $\mu\text{g}/\text{kg}$), together with ribavirin orally, at a median daily dose of 11 mg/kg (range 3.7–14.2 mg/kg) for 48 weeks. The dose of ribavirin was adjusted by body weight: 600 mg for patients weighing ≤ 60 kg; 800 mg for those between >60 kg and <80 kg; and 1,000 mg for those ≥ 80 kg. It was tapered in the 99 (33%) patients in whom hemoglobin levels decreased below 10 g/dl during the combination therapy.

Statistical Analysis

Variables were compared between groups by the chi-square test, Fisher's exact probability test and the Mann-Whitney U test. Differences in the loss of HCV RNA from the serum between groups was evaluated with the Kaplan-Meier life table with use of the log rank test. The influence of various factors on the response to PEG-IFN/ribavirin was evaluated by logistic regression in univariate and multivariate analyses. Analysis of all data was performed with the computer program SPSS software (SPSS Inc., Chicago, IL, USA), and a *P* value less than 0.05 was considered significant.

Results

Baseline Characteristics of Male and Female Patients Infected with HCV-1b in High Loads


PEG-IFN and ribavirin were given for 48 weeks to 179 men and 121 women who had been infected with HCV-1b in high viral loads (>100 kIU/ml). Table 1 compares baseline characteristics between them. Women were older, had lower hemoglobin values and platelet counts, and lower levels of albumin, gamma-glutamyl-transpeptidase (γ -GTP) and ferritin, than men. The stage of fibrosis was lower in women than in men, although their alanine aminotransferase (ALT) levels were comparable. Three months before the start of combination therapy, IFN had been given to 75 (42%) of the 179 male patients, comparably frequently to 40 of the 121 (33%) female patients. Age distribution for men and women is shown in Fig. 1. The proportion of patients ≥ 60 years was higher in women than in men (39% vs 19%, *P* < 0.001).

Table 1 Baseline characteristics of 300 patients with chronic hepatitis with high-titers of HCV-1b RNA who had received PEG-IFN and ribavirin for 48 weeks and were followed for 48 weeks or longer

Characteristic ^a	Men (n = 179)	Women (n = 121)	Differences P
Age (years)	50 (19–66)	57 (30–69)	<0.001
Previous IFN treatment	75 (42%)	40 (33%)	0.146
Hemoglobin (g/dl)	15.2 (11.5–17.8)	13.5 (11.2–15.1)	<0.001
Platelets ($\times 10^3/\text{mm}^3$)	176 (88–366)	165 (91–331)	0.025
Albumin (g/dl)	3.9 (3.2–4.6)	3.8 (3.0–4.6)	0.004
ALT (IU/l)	77 (23–504)	68 (19–391)	0.078
γ -GTP (IU/l)	78 (14–409)	37 (11–171)	0.011
LDL (mg/dl)	98 (50–176)	99 (57–168)	0.920
Ferritin (mg/l)	186 (<10–1,327)	95 (<10–4 42)	<0.001
ICG ₁₅ (%)	14 (4–41)	13 (2–31)	0.969
Stage (F0-1/F2-3)	80/66 (50 unknown)	42/55 (57 unknown)	0.050

ALT alanine aminotransferase, γ -GTP gamma-glutamyl transpeptidase, LDL low density lipoprotein, ICG₁₅, retention of indocyanine green at 15 min
^a The means (ranges) are given

Fig. 1 Distribution of ages in the male and female patients with chronic hepatitis C who were infected with HCV-1b in high loads



Virological Response During the 48-Week Treatment with PEG-IFN and Ribavirin

On-treatment response to the combined treatment is compared between men and women in Fig. 2. Through 48 weeks on treatment, women gained a virological response less frequently than did men. ETR was achieved by 58% of women as against 83% of men ($P < 0.001$). Restricted to the patients who gained ETR, women lost HCV RNA from serum later than men did [median (range) 24.1 (2.0–36.4) vs 13.0 (2.0–48.0) weeks, $P < 0.001$]. Figure 3 depicts the on-treatment virological response in patients <50 years and those ≥ 50 years separately. The virological response was no different between men and women <50 years. However, it was poorer in women than




Fig. 2 On-treatment virological responses to PEG-IFN and ribavirin in male and female patients infected with HCV-1b in high viral loads