

Distribution of Hepatitis B Virus Genotypes among Patients with Chronic Infection in Japan Shifting toward an Increase of Genotype A[∇]

Kentaro Matsuura,^{1,2} Yasuhito Tanaka,^{1*} Shuhei Hige,³ Gotaro Yamada,⁴ Yoshikazu Murawaki,⁵ Masafumi Komatsu,⁶ Tomoyuki Kuramitsu,⁷ Sumio Kawata,⁸ Eiji Tanaka,⁹ Namiki Izumi,¹⁰ Chiaki Okuse,¹¹ Shinichi Kakumu,¹² Takeshi Okanoue,¹³ Keisuke Hino,¹⁴ Yoichi Hiasa,¹⁵ Michio Sata,¹⁶ Tatsuji Maeshiro,¹⁷ Fuminaka Sugauchi,² Shunsuke Nojiri,² Takashi Joh,² Yuzo Miyakawa,¹⁸ and Masashi Mizokami^{1,19}

Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan¹; Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan²; Department of Internal Medicine, Hokkaido University Graduate School of Medicine, Sapporo, Japan³; Department of Basic Laboratory Sciences, Kawasaki Medical School, Kawasaki Hospital, Okayama, Japan⁴; Division of Medicine and Clinical Science, Faculty of Medicine, Tottori University, Tottori, Japan⁵; Department of Gastroenterology, Akita City Hospital, Akita, Japan⁶; Kuramitsu Clinic, Akita, Japan⁷; Department of Gastroenterology, Yamagata University School of Medicine, Yamagata, Japan⁸; Department of Medicine, Shinshu University School of Medicine, Matsumoto, Japan⁹; Division of Gastroenterology and Hepatology, Musashino Red Cross Hospital, Tokyo, Japan¹⁰; Department of Internal Medicine, Division of Gastroenterology and Hepatology, St. Marianna University School of Medicine, Kawasaki, Japan¹¹; Department of Gastroenterology, Aichi Medical University School of Medicine, Aichi, Japan¹²; Molecular Gastroenterology and Hepatology, Kyoto Prefectural University of Medicine, Kyoto, Japan¹³; Department of Gastroenterology and Hepatology, Yamaguchi University Graduate School of Medicine, Yamaguchi, Japan¹⁴; Department of Gastroenterology and Metabolism, Ehime University Graduate School of Medicine, Ehime, Japan¹⁵; Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Fukuoka, Japan¹⁶; First Department of Internal Medicine, University Hospital, Faculty of Medicine, University of the Ryukyus, Okinawa, Japan¹⁷; Miyakawa Memorial Research Foundation, Tokyo, Japan¹⁸; and Research Center for Hepatitis and Immunology, Kohnodai Hospital International Medical Center of Japan, Ichikawa, Japan¹⁹

Received 29 October 2008/Returned for modification 17 December 2008/Accepted 2 March 2009

Acute hepatitis B virus (HBV) infection has been increasing through promiscuous sexual contacts, and HBV genotype A (HBV/A) is frequent in patients with acute hepatitis B (AHB) in Japan. To compare the geographic distribution of HBV genotypes in patients with chronic hepatitis B (CHB) in Japan between 2005 and 2006 and between 2000 and 2001, with special attention to changes in the proportion of HBV/A, a cohort study was performed to survey changes in genotypes of CHB patients at 16 hospitals throughout Japan. Furthermore, we investigated the clinical characteristics of each genotype and examined the genomic characteristics of HBV/A isolates by molecular evolutionary analyses. Of the 1,271 patients, 3.5%, 14.1%, and 82.3% were infected with HBV/A, -B, and -C, respectively. In comparison with our previous survey during 2000 and 2001, HBV/A was twice as frequent (3.5% versus 1.7%; $P = 0.02$). The mean age was lower in the patients with HBV/A than in those with HBV/B or -C. Based on phylogenetic analyses of 11 full-length genomes and 29 pre-S2/S region sequences from patients, HBV/A isolates were imported from Europe and the United States, as well as the Philippines and India. They clustered with HBV/A from AHB patients and have spread throughout Japan. HBV/A has been increasing in CHB patients in Japan as a consequence of AHB spreading in the younger generation through promiscuous sexual contacts, aided by a tendency of HBV/A to induce chronic hepatitis. The spread of HBV/A infection in Japan should be prevented by universal vaccination programs.

Hepatitis B virus (HBV), a member of the *Hepadnaviridae*, is a circular, partially double-stranded DNA virus and is one of the major causes of chronic liver diseases, including chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC).

The HBV genome is composed of approximately 3,200 nucleotides. HBV is classified into eight genotypes, designated A to H, based on an intergroup divergence of 8% or more in the complete nucleotide sequence (3, 23, 26, 37). They have dis-

tinct geographical distributions and are associated with differences in clinical and virological characteristics, such as severity of liver disease and response to antiviral therapies (7, 8, 12, 13, 22, 28). Furthermore, subgenotypes have been reported for HBV/A, -B, and -C and named A1 to -3 (17, 38), B1 to -6 (31, 32, 40), and C1 to -6 (20, 31, 45). Equally, other genotypes are classified into subgenotypes. There have been increasing lines of evidence to indicate influences of HBV subgenotypes on the outcome of liver disease and the response to antiviral therapies (1, 39, 44).

In 2001, we reported the geographic distribution of HBV genotypes in Japan (27). Of the 720 Japanese patients with chronic HBV infection (CHB), 12 (1.7%) harbored HBV/A, 88 (12.2%) HBV/B, 610 (84.7%) HBV/C, 3 (0.4%) HBV/D, and 7 (1.0%) mixed genotypes. HBV/C was detected in over 94%

* Corresponding author. Mailing address: Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan. Phone: 81-52-853-8292. Fax: 81-52-842-0021. E-mail: ytanaka@med.nagoya-cu.ac.jp.

[∇] Published ahead of print on 18 March 2009.

of patients on the Japanese mainland, while HBV/B was found in 64% of those in Okinawa, the southernmost islands, and 44% of those in the Tohoku area in the northern part of the mainland.

Recently, acute HBV infection (AHB) has been increasing in Japan, predominantly through promiscuous sexual contacts. In addition, it was reported that HBV/A was more frequent in patients with acute hepatitis than in those with chronic hepatitis (29, 41, 49). Recent studies suggest that the chances for progression to chronic disease may differ among patients acutely infected with HBV of distinct genotypes (21, 25); patients infected with HBV/A run an increased risk of becoming HBV carriers. Hence, it is of utmost concern whether chronic HBV/A infection is increasing in Japan.

In the present study, we compared the geographic distribution of HBV genotypes in Japan during 2005 and 2006 with 2000 and 2001, with special attention to changes in the proportion of HBV/A. Furthermore, we investigated the clinical characteristics of each genotype and examined the genomic characteristics of HBV/A isolates by molecular evolutionary analyses.

MATERIALS AND METHODS

Patients. From September 2005 to October 2006, sera were collected from 1,370 consecutive patients with CHB at 16 representative hospitals that were liver centers in their respective regions throughout Japan for the purpose of investigating the geographic distribution of HBV genotypes in Japan. All of the patients were diagnosed after they had been followed for at least 12 months. Patients diagnosed with AHB were excluded from the study; they had a sudden onset of clinical symptoms of hepatitis, along with high-titer antibody to HBV core antigen of the immunoglobulin M class in serum. Their sera were tested for alanine aminotransferase (ALT), alkaline phosphatase (ALP), γ -glutamyl transpeptidase (γ -GTP), and hepatitis B e antigen (HBeAg), as well as antibody to HBeAg (anti-HBe) (Dinabot, Tokyo, Japan). Four clinical diagnoses were established for them. The inactive carrier state was defined by the presence of HBV surface antigen (HBsAg) with normal ALT levels over 1 year (examined at least four times at 3-month intervals) and without evidence of portal hypertension. Chronic hepatitis was defined by elevated ALT levels (>1.5 times the upper limit of normal [35 IU/liter]) persisting over 6 months (with at least three bimonthly tests). Cirrhosis was diagnosed principally by ultrasonography (coarse liver architecture, nodular liver surface, blunt liver edges, and hypersplenism), platelet counts of $<100,000/\text{cm}^3$, or a combination thereof. Histological confirmation by fine-needle biopsy of the liver was performed as required. HCC was diagnosed by ultrasonography, computerized tomography, magnetic resonance imaging, angiography, tumor biopsy, or a combination thereof.

The study protocol conformed to the 1975 declaration of Helsinki and was approved by the ethics committees of the respective institutions. Every patient or his/her next of kin gave informed consent to the purpose of the study.

Genotypes and subgenotypes of HBV. The six HBV genotypes (A to F) were determined serologically by enzyme immunoassay (EIA) using commercial kits (HBV Genotype EIA; Institutes of Immunology Co., Ltd., Tokyo, Japan). The method depends on the combination of epitopes on pre-S2 region products detected by monoclonal antibodies that were specific for each of them (46, 47). Subgenotypes of HBV/A, designated A1 and A2, were determined by direct sequencing of the pre-S2/S gene, followed by a phylogenetic analysis.

Quantification of HBV DNA and sequencing. HBV DNA levels in sera were quantitated with a commercial kit (Amplicor HBV Monitor; Roche Diagnostics, Basel, Switzerland) with a detection range from 2.6 to 7.6 log copies/ml. Nucleic acids were extracted from 100 μl of serum using the Qiaamp DNA Blood Minikit (Qiagen GmbH, Hilden, Germany). Eleven complete HBV/A genomes and 29 pre-S2/S region sequences were amplified by PCR with appropriate primer sets, as described previously (40). The amplified HBV DNA fragments were directly sequenced using the ABI Prism Big Dye kit version 3.0 (Applied Biosystems, Foster City, CA) in an ABI 3100 automated DNA sequencer (Applied Biosystems). All sequences were analyzed in both forward and reverse directions. Complete and partial HBV genome sequences were aligned using GENETYX version 11.0 (Software Development Co., Ltd., Tokyo, Japan).

TABLE 1. Characteristics of 1,271 CHB patients

Parameter	Value
Characteristic	
Male gender [no. (%)]	766 (60.3)
Age (yr; mean \pm SD)	51.4 \pm 14.0
Diagnosis	
Inactive carrier state [no. (%)]	206 (16.2)
Chronic hepatitis [no. (%)]	786 (61.8)
Cirrhosis [no. (%)]	175 (13.8)
HCC [no. (%)]	104 (8.2)
Antiviral treatment [no. (%)]	577 (45.4)
Blood tests	
Platelets ($10^4/\text{mm}^3$)	21.4 \pm 30.2
ALT (IU/liter)	59.8 \pm 103.0
ALP (IU/liter)	270.4 \pm 136.0
γ -GTP (IU/liter)	47.4 \pm 66.1
HBV markers	
HBeAg [no. (%)]	399 (31.4)
HBV DNA (median [range] [log copies/ml])	4.2 (<2.6 to >7.6)

Molecular evolutionary analysis of HBV. Reference sequences were retrieved from the DDBJ/EMBL/GenBank databases with their accession numbers for identification. To investigate the relationship between HBV isolates from patients with chronic and acute hepatitis B in Japan, HBV/A isolates (AH1 to -10) were randomly retrieved from them and sequenced in our previous study (29). Nucleotide sequences of HBV DNA were aligned by the program CLUSTAL X, and genetic distance was estimated by the six-parameter method (10) in the Hepatitis Virus Database (36). Based on these values, phylogenetic trees were constructed by the neighbor-joining method (30) with the midpoint rooting option. To confirm the reliability of the phylogenetic trees, bootstrap resampling tests were performed 1,000 times.

Statistical analysis. Categorical variables were compared between groups by the χ^2 test or Fisher's exact test and noncategorical variables by the Mann-Whitney U test. A *P* value of less than 0.05 was considered significant.

Nucleotide sequence accession numbers. The DDBJ/EMBL/GenBank accession numbers of the complete genome sequences of HBV isolates JPN_CH1 to -11 are AB453979 to AB453989.

RESULTS

Distribution of HBV genotypes among patients with CHB.

Of the 1,370 serum samples, the genotype could not be determined for 99 (7.2%) by EIA due to low HBsAg levels, leaving 1,271 for analysis in this study (Table 1). Of these, 206 (16.2%) were inactive carriers, 786 (61.8%) had chronic hepatitis, 175 (13.8%) cirrhosis, and 104 (8.2%) HCC. They had a mean age of 51.4 \pm 14.0 years and included 766 (60.3%) men. They had a median HBV DNA level of 4.2 log copies/ml, and 399 (31.4%) of them were positive for HBeAg. Antiviral treatment had been given to 577 (45.4%) of them with interferon, lamivudine, adefovir pivoxil, or entecavir.

The genotypes were HBV/A in 44 (3.5%), HBV/B in 179 (14.1%), HBV/C in 1,046 (82.2%), and HBV/D in 2 (0.2%) (Table 2). In comparison with our previous report on the distribution of genotypes in Japan in 2001 (27), HBV/A was more frequent in this study (3.5% versus 1.7%; *P* = 0.02). Of the 16 hospitals in this study, 10 overlapped with those in our previous report from 2001. In these 10 hospitals, HBV/A was more frequent in the present than in the previous survey (3.6% versus 1.7%; *P* = 0.04).

The distribution of HBV genotypes in Japan differed by

TABLE 2. Distribution of HBV Genotypes

Genotype	No. (%)	
	2005–2006 (n = 1,271)	2000–2001 ^a (n = 720)
A	44 (3.5 ^b)	12 (1.7)
B	179 (14.1)	88 (12.2)
C	1,046 (82.3)	610 (84.7)
D	2 (0.2)	3 (0.4)
Mixed	0 (0.0)	7 (1.0)

^a From Orito et al. (27).^b $P = 0.02$.

geographic location (Fig. 1). HBV/C was the most prevalent in the majority of areas. In the Tohoku area, the northern part of the Japanese mainland (Honshu), HBV/B was more prevalent than in the other areas of the Japanese mainland. In Okinawa, the southernmost islands of Japan, HBV/B was predominant. Of note, HBV/A was more frequent in the Kanto area (9.5%), the metropolitan area, and Okinawa (9.1%) than in the other areas.

Clinical differences among HBV/A, -B, and -C. Clinical backgrounds were compared among the patients infected with HBV/A, -B, and -C (Table 3). HBeAg was significantly less prevalent in the patients infected with HBV/B than in those infected with HBV/A or -C ($P < 0.01$ for each). When the positivity of HBeAg was stratified by age, HBeAg was markedly less common in patients infected with HBV/B than in those infected with HBV/A or -C who were older than 40 years of age (7/157 [4.5%] versus 4/19 [21.1%] [$P < 0.05$] or 215/755 [28.5%] [$P < 0.01$]) (Fig. 2). There were no significant differences in HBV DNA levels among patients infected with the three genotypes. As antiviral treatments might have influenced the severity of liver disease, clinical states were compared among patients infected with HBV/A, -B, and -C who did and

did not receive it; antiviral treatments did not affect the above-mentioned trends represented in Table 3 in age, diagnosis, and HBeAg, as well as ALT and HBV DNA levels (data not shown).

Additionally, we compared the distributions of age and liver diseases in patients infected with HBV/A, -B, and -C. In patients infected with HBV/C, the prevalence of cirrhosis and HCC increased in those older than 50 years of age compared to younger patients (Fig. 3), whereas in the patients infected with HBV/B, cirrhosis and HCC were rare in elderly patients. The proportion of patients younger than 40 years of age was higher in those infected with HBV/A than in those infected with HBV/B or -C (25/44 [56.8%] versus 22/179 [12.3%] or 288/1,046 [27.5%]; $P < 0.01$ for each), while cirrhosis and HCC were also found in those older than 50 years of age infected with HBV/A.

Coinfection with human immunodeficiency virus type 1 (HIV-1) was found in 6 of the 44 (13.6%) patients infected with HBV/A compared to only 3 of the 1,046 (0.3%) patients infected with HBV/C ($P < 0.0001$); it occurred in none of the 179 patients infected with HBV/B.

Phylogenetic analyses. Among the 44 HBV/A isolates, the complete genome was sequenced successfully in 11 (JPN_CH1 to -11). Seven of them were classified as HBV/A2 and four as HBV/A1. A phylogenetic tree was constructed based on the complete genome sequences of these 11 isolates, along with those from two patients with AHB and those from 40 HBV/A isolates retrieved from the database (Fig. 4). Of the seven HBV/A2 isolates, the four from patients with CHB in this study formed a cluster with the Japanese isolates retrieved from the database and two from patients with AHB. Of the other three isolates, JPN_CH5 clustered with French and U.S. isolates, JPN_CH6 with German isolates, and JPN_CH7 with




FIG. 1. Geographic distribution of HBV genotypes in patients with chronic HBV infection in Japan during 2005 and 2006.

TABLE 3. Clinical characteristics of individuals chronically infected with HBV of different genotypes

Parameter	Value for genotype:		
	A (n = 44)	B (n = 179)	C (n = 1,046)
Male gender [no. (%)]	32 (72.7)	112 (62.6)	621 (59.4)
Age (yr [mean \pm SD])	41.3 \pm 14.9 ^a	55.8 \pm 13.7 ^b	48.8 \pm 13.3
Diagnosis			
Inactive carrier state [no. (%)]	13 (29.5) ^c	63 (35.2) ^b	129 (12.3)
Chronic hepatitis [no. (%)]	26 (59)	103 (57.5)	656 (62.7)
Cirrhosis [no. (%)]	3 (6.8)	10 (5.6) ^b	162 (15.5)
HCC [no. (%)]	2 (4.5)	3 (1.7) ^b	99 (9.5)
Anti viral treatment [no. (%)]	13 (29.5) ^d	48 (26.8) ^b	516 (49.3)
Blood tests			
Platelet (10 ⁴ /mm ³)	23.3 \pm 21.9	25.9 \pm 35.9 ^e	20.6 \pm 29.5
ALT (IU/liter)	56.2 \pm 83.8	42.2 \pm 104.2 ^e	63.0 \pm 103.3
ALP (U/liter)	247.1 \pm 123.0	255.5 \pm 97.9	273.9 \pm 141.9
γ -GTP (U/liter)	39.6 \pm 34.6	49.3 \pm 63.4	47.5 \pm 67.6
HBV markers			
HBeAg [positive rate(%)]	15 (34.0) ^f	17 (9.5) ^b	367 (35.1)
HBV DNA (median [range]) (log copies/ml)	4.2 (<2.6-7.6)	4.1 (<2.6-7.6)	4.2 (<2.6-7.6)

^a $P < 0.01$, A versus B or C.

^b $P < 0.01$, B versus C.

^c $P < 0.01$, A versus C.

^d $P < 0.05$, A versus C.

^e $P < 0.05$, B versus C.

^f $P < 0.01$, A versus B.

Spanish and Italian isolates. All four HBV/A1 isolates in this study formed a cluster with Philippine and Indian isolates.

In addition, the pre-S2/S region sequences of a total of 29 isolates were determined, including the 11 isolates whose complete genomes were sequenced. Of these, 21 (72%) were classified as HBV/A2 and the remaining 8 as HBV/A1. A phylogenetic tree was constructed based on the pre-S2/S region sequences from the 29 isolates, along with those from 10 patients with AHB infected with HBV/A and 47 HBV/A isolates retrieved from the database (Fig. 5). The 21 HBV/A2 isolates in the present study formed a cluster with Japanese, American, and European isolates retrieved from the database and those from patients with acute hepatitis. In addition, some of them were highly homologous with each other. Likewise, HBV/A1 isolates from eight patients with chronic hepatitis in this study

were highly homologous with those from two patients with acute hepatitis and isolates from the Philippines and India. Based on the phylogenetic analyses, HBV/A isolates were imported from Europe and the United States, as well as the Philippines and India, and had infiltrated throughout Japan.

DISCUSSION

Perinatal transmission from carrier mothers to their babies has been the principal route for establishing persistent HBV infection in Asian countries (19). In Japan, passive and active immunoprophylaxis with HBV immune globulin and vaccine has been mandated for babies born to HBeAg-positive carrier mothers since 1986; this was extended to HBeAg-negative carrier mothers in 1995. As a result, HBsAg has become rare in Japanese born after 1986; it was detected in only 0.2% of first-time blood donors younger than 19 years of age in 2000 (24). However, AHB has been increasing in Japan, predominantly through promiscuous sexual contacts.

In Japan, HBV/A is detected rarely among patients with CHB but is frequent in those with acute hepatitis (14, 25, 29, 41, 43). Yotsuyanagi et al. reported the distribution of genotypes in 145 Japanese patients with AHB and found HBV/A in 27 (19%), HBV/B in 8 (5%), and HBV/C in 109 (75%) (49). HBV/A is more frequent in metropolitan areas than other areas. The majority of patients with HBV/A infection in metropolitan areas have had extramarital sexual contacts with multiple irregular partners, through which they could have contracted infection. In support of this view, among men who have sex with men (MSM) who are coinfecting with HBV and HIV-1 in Tokyo, most were infected with HBV/A (15, 35).

In Japan, AHB in adulthood becomes chronic in only ~1%




FIG. 2. Prevalence of HBeAg among patients infected with HBV of different genotypes stratified by the age.




FIG. 3. Distribution of HCC, cirrhosis, chronic hepatitis, and inactive carrier state among the 1,271 patients infected with HBV of different genotypes stratified by the age.

of cases. This is much less than the progression to chronic disease (close to 10%) in Europe and the United States, where HBV/A prevails (34). Recent studies have suggested that the chances for persistence may differ among patients acutely infected with HBV of distinct genotypes (21, 25). In particular, acute infection with HBV/A may bring about an increased risk of progression to chronic disease. Therefore, an increase of acute infection with HBV/A would result in a surge of HBV/A among patients with CHB in Japan. In actuality, in comparison with our previous results during 2000 and 2001 (27), HBV/A was twice as frequent in this study (3.5% versus 1.7%; $P = 0.02$). HBV/A has been increasing in patients with CHB in the Kanto area, where HBV/A in patients with acute hepatitis is more frequent than in the other areas. In the islands of Okinawa, also, HBV/A was found to be prevalent in this study. Of the four patients infected with HBV/A there, two were coinfecting with HIV-1. They were both MSM, and they were sus-




FIG. 4. Phylogenetic tree constructed based on the complete genome sequences of HBV/A isolates. Those from 11 patients with chronic infection in this study are shown in boldface italic (JPN_CH1 to -11), along with two isolates (JPN_AH1 and -2) from patients with acute hepatitis in Japan reported in our previous study (17). Representative isolates were retrieved from the DDBJ/EMBL/GenBank databases, including 21 HBV/Ae, 10 HBV/Aa, and 2 HBV/Ac isolates, along with 7 HBV isolates representative of the other seven genotypes. Isolates from the databases are identified by accession numbers, followed by the country of origin. The bar at the bottom spans 0.01 nucleotide substitutions per site.

pected to have been infected with HIV through sexual contacts on the Japanese mainland. It has been reported that HIV infection increases the probability that AHBs will become chronic (2, 11, 33, 48). Because they share routes of transmission and the risk for HIV-1 and HBV infections, approximately




FIG. 5. Phylogenetic tree constructed based on pre-S2/S region sequences of HBV/A isolates. Those from 29 patients with chronic infection in this study are shown in boldface italic (*JPN_CH1 to -29*), along with 10 isolates (*JPN_AH1 to -10*) from patients with acute hepatitis in Japan reported in our previous study (17). Representative isolates were retrieved from the DDBJ/EMBL/GenBank databases, including 28 HBV/Ae, 10 HBV/Aa, and 2 HBV/Ac isolates and 7 HBV isolates representative of the other seven genotypes. Isolates from the databases are identified by accession numbers, followed by the country of origin. The bar at the bottom spans 0.01 nucleotide substitutions per site.

90% of patients with AIDS have markers of past or ongoing HBV infection (18). Thus, HBV carriers are more frequent in the HIV-1-positive than in the HIV-1-negative population (4, 9). Among patients with HIV infection in Japan, 6.3% are HBsAg positive, in particular, 8.3% of HIV-infected MSM (16). In this study, coinfection with HIV was found in 6 of the 44 (13.6%) patients infected with HBV/A. All of them were men. Their median age was 27.7 ± 4.1 years, and five patients were positive for HBeAg. Thus, there is a possibility that HIV-1 and HBV/A coinfections are increasing among young people in Japan, and the high rate of HBeAg positivity may be influenced by immune suppression due to HIV infection.

In the phylogenetic analysis, the HBV/A2 isolates recovered in this study were homologous to those from Europe and the United States, and some of them clustered with the Japanese isolates. On the other hand, there were HBV/A1 isolates that formed a cluster with those from the Philippines and India. Furthermore, some isolates from patients with acute hepatitis who were infected with HBV/A in Japan were highly homologous to HBV/A isolates from patients with chronic hepatitis. This invites speculation that some HBV/A isolates were introduced into Japan from foreign countries, while others have already settled down there and spread from patients with chronic infection to their contacts. HBV/A would have been infiltrating throughout Japan by these two different routes.

Clinical differences among patients infected with HBV/A, -B, and -C were observed. The mean age was lower in the patients infected with HBV/A than in those infected with HBV/B or -C. As mentioned above, AHB patients infected with HBV/A have been increasing in the younger generation in Japan, and around 10% of them would have progressed to chronic infection. This is one of the reasons why the patients infected with HBV/A are younger than those infected with HBV/B or -C. Most patients infected with HBV/B were negative for HBeAg, while a high proportion of the patients infected with HBV/A and -C had it. In particular, this difference was remarkable in the patients who were older than 40 years of age. Thus, the seroconversion rate for the loss of HBeAg among younger people may be higher in infection with HBV/B than in that with HBV/A or -C. Inactive carriers were commoner in HBV/A than in HBV/C infection, as well.

These lines of evidence indicate that the activity of hepatitis is lower in HBV/B than HBV/C infection, and patients with HBV/B seroconvert from HBeAg to anti-HBe at young ages. In addition, cirrhosis and HCC were less frequent in the patients infected with HBV/B than in those infected with HBV/C. Therefore, the prognosis would be better in the patients infected with HBV/B than in those infected with HBV/C. These results are in accord with previous reports (5, 13, 28, 42). There have been few reports on the clinical features of patients with chronic hepatitis infected with HBV/A in Japan. Chu et al. have reported the distribution of HBV genotypes with reference to clinical characteristics in the United States (6). They have shown that HBV/A and HBV/C infections are accompanied by a higher frequency of HBeAg than HBV/B infection, while HBV/B is associated with a lower rate of hepatic decompensation than HBV/A and -C. In our study, inactive carriers were commoner, while cirrhosis and HCC were found less often in HBV/A than in HBV/C infection. HBeAg was more prevalent in the patients infected with HBV/A than in those

infected with HBV/B who were older than 40 years of age. Therefore, it can be said that the prognosis is better for patients infected with HBV/A than for those infected with HBV/C; it may be poorer than for those infected with HBV/B.

In conclusion, HBV/A has been increasing among CHB patients in Japan. On the basis of phylogenetic analyses, some HBV/A isolates appear to have been imported from foreign countries. They clustered with HBV/A from AHB patients and have infiltrated throughout Japan. It is very likely that acute and chronic infections with HBV/A have been increasing in Japan. Obviously, immunoprophylaxis of perinatal HBV infection, implemented since 1986 on a national basis, has been insufficient to prevent horizontal HBV/A infection diffusing among high-risk groups by transmission routes shared by HIV infection. The foreseeable spread of HBV/A infection in Japan should be prevented by universal vaccination programs extended to high-risk groups or the general population.

ACKNOWLEDGMENTS

The study was supported in part by a grant-in-aid from the Ministry of Health, Labor and Welfare of Japan and a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology.

We thank T. Kimura and K. Sato, Institutes of Immunology Co., Ltd. (Tokyo, Japan), for determining HBV genotypes in this study and Takashi Saito, Yamagata University Hospital; Akihiro Matsumoto, Shinshu University Hospital; Yasuhiro Asahina, Musashino Red Cross Hospital; Yoshito Ito, University Hospital, Kyoto Prefectural University of Medicine; Keiko Hoshino, Tottori University Hospital; Morikazu Onji, Ehime University Hospital; Tatsuya Ide, Kurume University Hospital; and Hiroshi Sakugawa, Hospital, University of the Ryukyus, for their help throughout this work.

Kentaro Matsuura wrote the study protocol and the first draft of the manuscript and performed the experiments and statistical analysis. Yasuhito Tanaka contributed to the experimental work and the final version of the manuscript. Shuhei Hige, Gotaro Yamada, Yoshikazu Murawaki, Masafumi Komatsu, Tomoyuki Kuramitsu, Sumio Kawata, Eiji Tanaka, Namiki Izumi, Chiaki Okuse, Shinichi Kakumu, Takeshi Okanoue, Keisuke Hino, Yoichi Hiasa, Michio Sata, and Tatsuji Maeshiro contributed to the collection of the samples and clinical data from patients and to the final version of the manuscript. Fuminaka Sugauchi, Shunsuke Nojiri, Takashi Joh, and Yuzo Miyakawa contributed to the final version of the manuscript. Masashi Mizokami had the original idea and did the planning of the study and contributed to the final version of the manuscript. All of the authors have seen and approved the final draft of the manuscript.

REFERENCES

1. Akuta, N., F. Suzuki, M. Kobayashi, A. Tsubota, Y. Suzuki, T. Hosaka, T. Someya, S. Saitoh, Y. Arase, K. Ikeda, and H. Kumada. 2003. The influence of hepatitis B virus genotype on the development of lamivudine resistance during long-term treatment. *J. Hepatol.* **38**:315-321.
2. Alter, M. J. 2006. Epidemiology of viral hepatitis and HIV co-infection. *J. Hepatol.* **44**:S6-S9.
3. Arauz-Ruiz, P., H. Norder, B. H. Robertson, and L. O. Magnius. 2002. Genotype H: a new Amerindian genotype of hepatitis B virus revealed in Central America. *J. Gen. Virol.* **83**:2059-2073.
4. Bodsworth, N. J., D. A. Cooper, and B. Donovan. 1991. The influence of human immunodeficiency virus type 1 infection on the development of the hepatitis B virus carrier state. *J. Infect. Dis.* **163**:1138-1140.
5. Chu, C. J., M. Hussain, and A. S. Lok. 2002. Hepatitis B virus genotype B is associated with earlier HBeAg seroconversion compared with hepatitis B virus genotype C. *Gastroenterology* **122**:1756-1762.
6. Chu, C. J., E. B. Keeffe, S. H. Han, R. P. Perrillo, A. D. Min, C. Soldevilla-Pico, W. Carey, R. S. Brown, Jr., V. A. Luketic, N. Terrault, and A. S. Lok. 2003. Hepatitis B virus genotypes in the United States: results of a nationwide study. *Gastroenterology* **125**:444-451.
7. Chu, C. J., and A. S. Lok. 2002. Clinical significance of hepatitis B virus genotypes. *Hepatology* **35**:1274-1276.
8. Ding, X., M. Mizokami, G. Yao, B. Xu, E. Orito, R. Ueda, and M. Nakanishi. 2001. Hepatitis B virus genotype distribution among chronic hepatitis B virus carriers in Shanghai, China. *Intervirology* **44**:43-47.

9. Fujii, T., H. Taguchi, H. Katano, S. Mori, T. Nakamura, N. Nojiri, K. Nakajima, K. Tadokoro, T. Juji, and A. Iwamoto. 1999. Seroprevalence of human herpesvirus 8 in human immunodeficiency virus 1-positive and human immunodeficiency virus 1-negative populations in Japan. *J. Med. Virol.* 57:159-162.
10. Gojobori, T., K. Ishii, and M. Nei. 1982. Estimation of average number of nucleotide substitutions when the rate of substitution varies with nucleotide. *J. Mol. Evol.* 18:414-423.
11. Hadler, S. C., F. N. Judson, P. M. O'Malley, N. L. Altman, K. Penley, S. Buchbinder, C. A. Schable, P. J. Coleman, D. N. Ostrow, and D. P. Francis. 1991. Outcome of hepatitis B virus infection in homosexual men and its relation to prior human immunodeficiency virus infection. *J. Infect. Dis.* 163:454-459.
12. Kao, J. H. 2002. Clinical relevance of hepatitis B viral genotypes: a case of *deja vu*? *J. Gastroenterol. Hepatol.* 17:113-115.
13. Kao, J. H., P. J. Chen, M. Y. Lai, and D. S. Chen. 2000. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology* 118:554-559.
14. Kobayashi, M., Y. Arase, K. Ikeda, A. Tsubota, Y. Suzuki, S. Saitoh, F. Suzuki, N. Akuta, T. Someya, M. Matsuda, J. Sato, K. Takagi, Y. Miyakawa, and H. Kumada. 2002. Viral genotypes and response to interferon in patients with acute prolonged hepatitis B virus infection of adulthood in Japan. *J. Med. Virol.* 68:522-528.
15. Koibuchi, T., A. Hitani, T. Nakamura, N. Nojiri, K. Nakajima, T. Jyuji, and A. Iwamoto. 2001. Predominance of genotype A HBV in an HBV-HIV-1 dually positive population compared with an HIV-1-negative counterpart in Japan. *J. Med. Virol.* 64:435-440.
16. Koike, K., Y. Kikuchi, M. Kato, J. Takamatsu, Y. Shintani, T. Tsutsumi, H. Fujie, H. Miyoshi, K. Moriya, and H. Yotsuyanagi. 2008. Prevalence of hepatitis B virus infection in Japanese patients with HIV. *Hepatol. Res.* 38:310-314.
17. Kurbanov, F., Y. Tanaka, K. Fujiwara, F. Sugauchi, D. Mbanya, L. Zekeng, N. Ndembu, C. Ngansop, L. Kaptue, T. Miura, E. Ido, M. Hayami, H. Ichimura, and M. Mizokami. 2005. A new subtype (subgenotype) Ac (A3) of hepatitis B virus and recombination between genotypes A and E in Cameroon. *J. Gen. Virol.* 86:2047-2056.
18. Lebovics, E., B. M. Dworkin, S. K. Heier, and W. S. Rosenthal. 1988. The hepatobiliary manifestations of human immunodeficiency virus infection. *Am. J. Gastroenterol.* 83:1-7.
19. Lok, A. S. 1992. Natural history and control of perinatally acquired hepatitis B virus infection. *Dig. Dis.* 10:46-52.
20. Lusida, M. I., V. E. Nugrahaputra, Soetjipto, R. Handajani, M. Nagano-Fujii, M. Sasayama, T. Utsumi, and H. Hotta. 2008. Novel subgenotypes of hepatitis B virus genotypes C and D in Papua, Indonesia. *J. Clin. Microbiol.* 46:2160-2166.
21. Mayerat, C., A. Mantegani, and P. C. Frei. 1999. Does hepatitis B virus (HBV) genotype influence the clinical outcome of HBV infection? *J. Viral Hepat.* 6:299-304.
22. Miyakawa, Y., and M. Mizokami. 2003. Classifying hepatitis B virus genotypes. *Intervirology* 46:329-338.
23. Norder, H., B. Hammas, S. Lofdahl, A. M. Courouce, and L. O. Magnus. 1992. Comparison of the amino acid sequences of nine different serotypes of hepatitis B surface antigen and genomic classification of the corresponding hepatitis B virus strains. *J. Gen. Virol.* 73:1201-1208.
24. Noto, H., T. Terao, S. Ryou, Y. Hirose, T. Yoshida, H. Ookubo, H. Mito, and H. Yoshizawa. 2003. Combined passive and active immunoprophylaxis for preventing perinatal transmission of the hepatitis B virus carrier state in Shizuoka, Japan during 1980-1994. *J. Gastroenterol. Hepatol.* 18:943-949.
25. Ogawa, M., K. Hasegawa, T. Naritomi, N. Torii, and N. Hayashi. 2002. Clinical features and viral sequences of various genotypes of hepatitis B virus compared among patients with acute hepatitis B. *Hepatol. Res.* 23:167-177.
26. Okamoto, H., F. Tsuda, H. Sakugawa, R. I. Sastrosoewignjo, M. Imai, Y. Miyakawa, and M. Mayumi. 1988. Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J. Gen. Virol.* 69:2575-2583.
27. Orito, E., T. Ichida, H. Sakugawa, M. Sata, N. Horiike, K. Hino, K. Okita, T. Okanoue, S. Iino, E. Tanaka, K. Suzuki, H. Watanabe, S. Hige, and M. Mizokami. 2001. Geographic distribution of hepatitis B virus (HBV) genotype in patients with chronic HBV infection in Japan. *Hepatology* 34:590-594.
28. Orito, E., M. Mizokami, H. Sakugawa, K. Michitaka, K. Ishikawa, T. Ichida, T. Okanoue, H. Yotsuyanagi, S. Iino, et al. 2001. A case-control study for clinical and molecular biological differences between hepatitis B viruses of genotypes B and C. *Hepatology* 33:218-223.
29. Ozasa, A., Y. Tanaka, E. Orito, M. Sugiyama, J. H. Kang, S. Hige, T. Kuramitsu, K. Suzuki, E. Tanaka, S. Okada, H. Tokita, Y. Asahina, K. Inoue, S. Kakumu, T. Okanoue, Y. Murawaki, K. Hino, M. Onji, H. Yatsuhashi, H. Sakugawa, Y. Miyakawa, R. Ueda, and M. Mizokami. 2006. Influence of genotypes and precore mutations on fulminant or chronic outcome of acute hepatitis B virus infection. *Hepatology* 44:326-334.
30. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406-425.
31. Sakamoto, T., Y. Tanaka, E. Orito, J. Co, J. Clavio, F. Sugauchi, K. Ito, A. Ozasa, A. Quino, R. Ueda, J. Sollano, and M. Mizokami. 2006. Novel subtypes (subgenotypes) of hepatitis B virus genotypes B and C among chronic liver disease patients in the Philippines. *J. Gen. Virol.* 87:1873-1882.
32. Sakamoto, T., Y. Tanaka, J. Simonetti, C. Osioy, M. L. Borresen, A. Koch, F. Kurbanov, M. Sugiyama, G. Y. Minuk, B. J. McMahon, T. Joh, and M. Mizokami. 2007. Classification of hepatitis B virus genotype B into 2 major types based on characterization of a novel subgenotype in Arctic indigenous populations. *J. Infect. Dis.* 196:1487-1492.
33. Salmon-Ceron, D., C. Lewden, P. Morlat, S. Bevilacqua, E. Jougla, F. Bonnet, L. Heripret, D. Costagliola, T. May, and G. Chene. 2005. Liver disease as a major cause of death among HIV infected patients: role of hepatitis C and B viruses and alcohol. *J. Hepatol.* 42:799-805.
34. Sherlock, S. D. J. 1997. Virus hepatitis, p. 265-392. *In* S. D. J. Sherlock (ed.), *Diseases of the liver and biliary system*, 10th ed. Blackwell Scientific Publications, London, United Kingdom.
35. Shibayama, T., G. Masuda, A. Ajisawa, K. Hiruma, F. Tsuda, T. Nishizawa, M. Takahashi, and H. Okamoto. 2005. Characterization of seven genotypes (A to E, G and H) of hepatitis B virus recovered from Japanese patients infected with human immunodeficiency virus type 1. *J. Med. Virol.* 76:24-32.
36. Shin, I. T., Y. Tanaka, Y. Tateno, and M. Mizokami. 2008. Development and public release of a comprehensive hepatitis virus database. *Hepatol. Res.* 38:234-243.
37. Stuyver, L., S. De Gendt, C. Van Geyt, F. Zoulim, M. Fried, R. F. Schinazi, and R. Rossau. 2000. A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J. Gen. Virol.* 81:67-74.
38. Sugauchi, F., H. Kumada, S. A. Acharya, S. M. Shrestha, M. T. Gamutan, M. Khan, R. G. Gish, Y. Tanaka, T. Kato, E. Orito, R. Ueda, Y. Miyakawa, and M. Mizokami. 2004. Epidemiological and sequence differences between two subtypes (Ae and Aa) of hepatitis B virus genotype A. *J. Gen. Virol.* 85:811-820.
39. Sugauchi, F., E. Orito, T. Ichida, H. Kato, H. Sakugawa, S. Kakumu, T. Ishida, A. Chutaputti, C. L. Lai, R. G. Gish, R. Ueda, Y. Miyakawa, and M. Mizokami. 2003. Epidemiologic and virologic characteristics of hepatitis B virus genotype B having the recombination with genotype C. *Gastroenterology* 124:925-932.
40. Sugauchi, F., E. Orito, T. Ichida, H. Kato, H. Sakugawa, S. Kakumu, T. Ishida, A. Chutaputti, C. L. Lai, R. Ueda, Y. Miyakawa, and M. Mizokami. 2002. Hepatitis B virus of genotype B with or without recombination with genotype C over the precore region plus the core gene. *J. Virol.* 76:5985-5992.
41. Sugauchi, F., E. Orito, T. Ohno, Y. Tanaka, A. Ozasa, J. H. Kang, J. Toyoda, T. Kuramitsu, K. Suzuki, E. Tanaka, Y. Akahane, T. Ichida, N. Izumi, K. Inoue, H. Hoshino, S. Iino, H. Yotsuyanagi, S. Kakumu, E. Tomita, T. Okanoue, S. Nishiguchi, Y. Murawaki, K. Hino, M. Onji, H. Yatsuhashi, M. Sata, Y. Miyakawa, R. Ueda, and M. Mizokami. 2006. Spatial and chronological differences in hepatitis B virus genotypes from patients with acute hepatitis B in Japan. *Hepatol. Res.* 36:107-114.
42. Sumi, H., O. Yokosuka, N. Seki, M. Arai, F. Imazeki, T. Kurihara, T. Kanda, K. Fukui, M. Kato, and H. Saisho. 2003. Influence of hepatitis B virus genotypes on the progression of chronic type B liver disease. *Hepatology* 37:19-26.
43. Suzuki, Y., M. Kobayashi, K. Ikeda, F. Suzuki, Y. Arase, N. Akuta, T. Hosaka, S. Saitoh, T. Someya, M. Matsuda, J. Sato, S. Watabiki, Y. Miyakawa, and H. Kumada. 2005. Persistence of acute infection with hepatitis B virus genotype A and treatment in Japan. *J. Med. Virol.* 76:33-39.
44. Tanaka, Y., I. Hasegawa, T. Kato, E. Orito, N. Hirashima, S. K. Acharya, R. G. Gish, A. Kramvis, M. C. Kew, N. Yoshihara, S. M. Shrestha, M. Khan, Y. Miyakawa, and M. Mizokami. 2004. A case-control study for differences among hepatitis B virus infections of genotypes A (subtypes Aa and Ae) and D. *Hepatology* 40:747-755.
45. Tanaka, Y., E. Orito, M. F. Yuen, M. Mukaide, F. Sugauchi, K. Ito, A. Ozasa, T. Sakamoto, F. Kurbanov, C. L. Lai, and M. Mizokami. 2005. Two subtypes (subgenotypes) of hepatitis B virus genotype C: a novel subtyping assay based on restriction fragment length polymorphism. *Hepatol. Res.* 33:216-224.
46. Usuda, S., H. Okamoto, H. Iwanari, K. Baba, F. Tsuda, Y. Miyakawa, and M. Mayumi. 1999. Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J. Virol. Methods* 80:97-112.
47. Usuda, S., H. Okamoto, T. Tanaka, K. Kidd-Ljunggren, P. V. Holland, Y. Miyakawa, and M. Mayumi. 2000. Differentiation of hepatitis B virus genotypes D and E by ELISA using monoclonal antibodies to epitopes on the preS2-region product. *J. Virol. Methods* 87:81-89.
48. Weinbaum, C. M., K. M. Sabin, and S. S. Santibanez. 2005. Hepatitis B, hepatitis C, and HIV in correctional populations: a review of epidemiology and prevention. *AIDS* 19(Suppl. 3):S41-S46.
49. Yotsuyanagi, H., C. Okuse, K. Yasuda, E. Orito, S. Nishiguchi, J. Toyoda, E. Tomita, K. Hino, K. Okita, S. Murashima, M. Sata, H. Hoshino, Y. Miyakawa, and S. Iino. 2005. Distinct geographic distributions of hepatitis B virus genotypes in patients with acute infection in Japan. *J. Med. Virol.* 77:39-46.

A Genetic Variant of Hepatitis B Virus Divergent from Known Human and Ape Genotypes Isolated from a Japanese Patient and Provisionally Assigned to New Genotype J^{V†}

Kanako Tatematsu,¹ Yasuhito Tanaka,^{1*} Fuat Kurbanov,¹ Fuminaka Sugauchi,²
Shuhei Mano,³ Tatsuji Maeshiro,⁴ Tomokuni Nakayoshi,⁵ Moriaki Wakuta,⁶
Yuzo Miyakawa,⁷ and Masashi Mizokami^{1,8}

Department of Clinical Molecular Informative Medicine¹ and Department of Gastroenterology and Metabolism,² Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan; Nagoya City University Graduate School of Natural Sciences, Nagoya, Japan³; Control and Prevention of Infectious Diseases, Department of Medicine and Therapeutics, Faculty of Medicine, University of the Ryukyus, Okinawa, Japan⁴; Heart Life Hospital, Okinawa, Japan⁵; Wakusan Clinic, Okinawa, Japan⁶; Miyakawa Memorial Research Foundation, Tokyo, Japan⁷; and Research Center for Hepatitis and Immunology, International Medical Center of Japan Kohnodai Hospital, Chiba, Japan⁸

Received 5 March 2009/Accepted 24 July 2009

Hepatitis B virus (HBV) of a novel genotype (J) was recovered from an 88-year-old Japanese patient with hepatocellular carcinoma who had a history of residing in Borneo during the World War II. It was divergent from eight human (A to H) and four ape (chimpanzee, gorilla, gibbon, and orangutan) HBV genotypes, as well as from a recently proposed ninth human genotype I, by 9.9 to 16.5% of the entire genomic sequence and did not have evidence of recombination with any of the nine human genotypes and four nonhuman genotypes. Based on a comparison of the entire nucleotide sequence against 1,440 HBV isolates reported, HBV/J was nearest to the gibbon and orangutan genotypes (mean divergences of 10.9 and 10.7%, respectively). Based on a comparison of four open reading frames, HBV/J was closer to gibbon/orangutan genotypes than to human genotypes in the P and large S genes and closest to Australian aboriginal strains (HBV/C4) and orangutan-derived strains in the S gene, whereas it was closer to human than ape genotypes in the C gene. HBV/J shared a deletion of 33 nucleotides at the start of preS1 region with C4 and gibbon genotypes, had an S-gene sequence similar to that of C4, and expressed the *ayw* subtype. Efficient infection, replication, and antigen expression by HBV/J were experimentally established in two chimeric mice with the liver repopulated for human hepatocytes. The HBV DNA sequence recovered from infected mice was identical to that in the inoculum. Since HBV/J is positioned phylogenetically in between human and ape genotypes, it may help to trace the origin of HBV and merits further epidemiological surveys.

Worldwide, an estimated 400 million people are infected with hepatitis B virus (HBV) persistently, of whom three quarters live in the Southeast and Far East Asia, and one million die of decompensated cirrhosis and/or hepatocellular carcinoma (HCC) annually (8, 15). HBV is the smallest animal DNA virus and has a genome made of approximately 3,200 nucleotides (nt) that contains four open reading frames for P, C, S, and X genes; they code for DNA polymerase/reverse-transcriptase, core protein, surface protein, and X protein, respectively (49). The S gene is divided into preS1 and preS2 regions and the small S gene, and the C gene splits into PreC and C.

Eight genotypes of HBV have been recognized by a sequence divergence of >8% in the entire genome and named by capital alphabet letters (A to H) in the order of discovery (3, 26, 29, 42). HBV genotypes are further classified into subgenotypes, such as B1/Bj and B2-5/Ba (44), as well as C1/Cs, C2/Ce,

and C3-5 (36). A systematic nomenclature is proposed for designating HBV subgenotypes using Arabic numbers, such as A1, A2, and A3 (25). HBV genotypes have distinct geographical distribution (16, 23). Genotype A is prevalent in Africa, Europe and India, genotypes B and C are common in Asia, and genotype E is common in sub-Saharan Africa. Genotypes F and H are restricted to Central and South American continents, whereas genotype D is distributed all over the world. HBV genotypes have clinical application, and they influence severity and progression of liver disease and the response to antiviral therapies. Previous reports indicate that HCC is more frequent in the patients infected with genotype C than B (7, 47), and interferon is more effective in those infected with genotype B than C in Asia and more effective in those infected with genotype A than D in Europe (18, 34, 51).

Recently, a ninth genotype (I) was tentatively proposed for HBV strains detected in Laos (31). These strains are phylogenetically similar to aberrant Vietnamese strains that display complex recombination over the genome (10). In the present study, an HBV isolate was recovered from a Japanese patient with HCC, who was involved in military actions in Borneo during the World War II. The isolated strain was compared against eight human (A to H) and four ape (chimpanzee, gorilla, gibbon, and orangutan) genotypes and was provisionally designated genotype J. The new genotype was assigned based on a sequence diver-

* Corresponding author. Mailing address: Department of Clinical Molecular Informative Medicine, Nagoya, City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan. Phone: (81) 52-853-8292. Fax: (81) 52-842-0021. E-mail: ytanaka@med.nagoya-cu.ac.jp.

† Supplemental material for this article may be found at <http://jvi.asm.org/>.

^V Published ahead of print on 29 July 2009.

TABLE 1. Nucleotide divergence in the full-genome sequence estimated from pairwise comparison between the Ryukyu 34 strain of a provisional genotype J and 1,440 HBV strains from the database entered by September 2008

Genotype	No. of strains	Divergence (%)		
		Range	Mean	SD
A	202	12.1–15.9	13.0	0.4
B	309	11.1–13.6	11.9	0.5
C	396	11.2–13.1	11.9	0.5
D	264	12.6–15.0	13.4	0.2
E	90	12.3–13.4	12.7	0.3
F	56	15.2–16.5	15.6	0.2
G	23	12.8–14.6	13.7	0.3
H	21	15.4–16.3	15.7	0.3
I	16	11.4–12.0	11.7	0.2
Chimpanzee	14	11.6–12.7	12.1	0.3
Gorilla	1	12.2		
Gibbon	34	9.9–11.7	10.9	0.5
Orangutan	12	10.4–11.2	10.7	0.4
Woolly monkey	2	27.2–27.4	27.3	0.1

gence of 10.7 to 15.7% from other genotypes, a unique phylogenetic position between human and ape genotypes, and the absence of strong evidence of recombination.

MATERIALS AND METHODS

Patient. A Japanese man, 88 years old, developed HCC in 2006. He had a history of residing in Borneo during the World War II. No HBV infections were recorded in his family members. In October 1996, he was diagnosed with chronic hepatitis B. Hepatitis B surface antigen (HBsAg) was detected in serum, and the aspartate aminotransaminase and alanine aminotransferase levels were elevated to 83 and 73 U/liter, respectively (normal levels, <30 U/liter for both). Thereafter, the transaminase levels were normalized, and he had been monitored as an asymptomatic HBV carrier. In August 2000, the level of a tumor marker (des-γ-carboxy prothrombin) was elevated to 52 mAU/ml (normal, <40 mAU/ml), while another tumor marker (alpha-fetoprotein) remained within normal range (<10 ng/ml) as alanine aminotransferases. In October 2006, a tumor (4.3 by 4.1 cm) was detected in the liver by ultrasonography, and he received treatment with transarterial embolization. Des-γ-carboxy prothrombin was elevated to 419 mAU/ml, while the aminotransferase levels remained within normal limits. Hepatitis B e antigen (HBeAg) was negative, and the corresponding antibody (anti-HBe) was detected in his serum. The subtype of HBsAg in this serum was *ayw*.

HBV DNA was extracted from his serum specimen obtained in 2006, and the full-length genome sequence was determined for phylogenetic and biological analyses. An informed consent had been obtained from the patient, and the study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution's human research committee.

Markers of HBV infection. HBeAg and anti-HBe were determined by enzyme-linked immunosorbent assay (ELISA) with commercial kits (HBeAg EIA; Institute of Immunology, Tokyo, Japan), and subtypes of HBsAg by ELISA with commercial kits (HBsAg Subtype EIA; Institute of Immunology). Hepatitis B core-related antigen (HBcrAg) was determined by chemiluminescence enzyme immunoassay (13). The method allows more sensitive detection of core protein and, as was shown in previous studies, HBcrAg levels reflect HBV DNA loads and well correlate with intrahepatic covalently closed circular DNA (cccDNA) levels. The measurement of serum HBcrAg is a useful noninvasive tool for monitoring intrahepatic HBV viral status (52). HBV DNA was quantified by the S gene-targeted real-time detection PCR with a sensitivity of 100 copies/ml (equivalent to 20 IU/ml) (1). However, due to small volumes of sera available from the challenged mice, HBV DNA was extracted from 10-fold-diluted specimens, resulting in reduced assay sensitivity in the present study (1,000 copies/ml [200 IU/ml]).

Determination of the complete nucleotide sequence of HBV/J isolate. HBV DNA was extracted by using the QIAamp DNA blood kit (Qiagen, GmbH, Hilden, Germany) from 100 μl of serum that had been stored at -80°C. The complete genome sequence of an HBV/J isolate recovered from the patient was determined by the strategy previously reported (43). In brief, two sets of primers were designed to amplify overlapping fragments (A and B) covering the entire

HBV genome (stat not shown). Nested PCR was carried out for 35 cycles (95°C, 30 s; 57°C, 30 s; and 72°C, 2 min) using TaKaRa LA *Taq* polymerase (Takara Biochemicals, Kyoto, Japan). Amplified fragments were inserted into the pGEM-T Easy vector (Promega, Madison, WI), and cloned in DH5a cells (Toyobo, Osaka, Japan). Obtained HBV DNA clones were confirmed to have the sequence identical to the major-clone consensus sequence determined directly on PCR products by Prism BigDye (Applied Biosystems, Foster City, CA) in the ABI 3100 automated sequencer.

Phylogenetic analysis. Full-length sequences of HBV isolates were aligned with use of the CLUSTAL W software program (48) (available at www.ebi.ac.uk), and the alignment was confirmed by visual inspection. Genetic distances were estimated by the six-parameter method, and phylogenetic trees were constructed with the neighbor-joining method (35). To confirm the reliability of phylogenetic trees, bootstrap resampling and reconstruction were carried out 1,000 times using the program




FIG. 1. Phylogenetic tree constructed on the entire genome sequences of 44 HBV isolates representing four ape and eight human genotypes. A woolly monkey HBV isolate serves as an outgroup. The HBV/J isolate (JRB34) is indicated by an arrowhead, and the genetic distance is indicated by a bar below.




FIG. 2. Complete genome scanning carried by PHYLIP, the phylogeny inference package implemented in the Simmonic software, for the JRB34 strain versus 228 selected nonrecombinant HBV genotypes (HBV/Ba and HBV/I not included) reference strains grouped by genotype. Kimura two-parameter distance model (A) and grouping scan (B) were determined with a 300-nt size window sliding by an increment of 15 nucleotides. The x axis indicates the genome position (corresponding to the midpoint of the scanning fragment), and the y axis indicates the mean distances between JRB34 and reference groups (A). Phylogenetic association (y axis) was evaluated throughout entire HBV genome (x axis) with the same window and step size parameters (B). The association value below 0.5 was considered to represent an outgroup. The open reading frame map is shown schematically at the top of the figure.

of the Hepatitis Virus Database (39). All 1,440 complete genomes available in the DDBJ/GenBank served as references for the initial alignment in the present study. Divergence in the nucleotide sequence between a strain of provisional genotype J and previously reported strains was estimated by using MEGALIGN v.6.00 (Lazer-gene package; DNASTAR, Inc., Madison, WI).

Examination of recombination evidence. Evidence of possible recombination was investigated by using the software packages Simmonic 2005 v1.6 and SimPlot v3.5.1, both implementing PHYLIP (Phylogeny Inference Package v3.68; J. Felsenstein, Department of Genome Sciences, University of Washington, Seattle [distributed by the authors]) (19, 40).

Inoculation of chimeric mice with the liver repopulated for human hepatocytes. Severe combined immunodeficiency mice transgenic for the urokinase-type plasminogen activator gene (uPA^{+/+}/SCID^{+/+} mice) with the liver repopulated with human hepatocytes (chimeric mice) were purchased from Phoenix Bio Co., Ltd. (Hiroshima, Japan). Human serum albumin was measured by ELISA with commercial assay kits (Eiken Chemical Co., Ltd., Tokyo, Japan) for estimating the extent of repopulation. The research complied with all relevant federal guidelines and institutional policies.

Immunofluorescence. Freshly prepared liver tissues were snap-frozen in isopentane precooled in liquid nitrogen. Frozen specimens were cut at 5 to 6 μ m by cryostat, mounted on glass slides, air dried, and fixed in 100% acetone at room

temperature for 10 min. Sections were blocked with antibody diluent (Dako, Tokyo, Japan) and stained for hepatitis B core antigen (HBcAg). They were incubated with rabbit anti-HBc (Dako) at room temperature for 1 h, washed in phosphate-buffered saline, and then incubated with goat anti-rabbit immunoglobulin G conjugated with Cy3 (Chemicon International, Inc., Temecula, CA) or goat anti-human albumin antibody labeled with fluorescein isothiocyanate (Bethyl Laboratories, Inc., Montgomery, TX). Sections were washed with phosphate-buffered saline and observed in a fluorescence microscope (Eclipse E800M; Nikon, Tokyo, Japan).

Nucleotide sequence accession numbers. The nucleotide sequence data reported in the present study will appear in the DDBJ/EMBL/GenBank databases under accession no. AB486012.

RESULTS

Composition of the HBV genome of genotype J. HBV DNA was extracted from serum of a patient with HCC. It was named JRB34 ("J" for Japanese; "R" after the southernmost island [Ryukyu] where the patient has spent most of his life now




FIG. 3. Phylogenetic tree constructed on the preS/S gene (A) and C gene (B) sequences of 44 HBV isolates representing four ape and eight human genotypes. A woolly monkey HBV isolate serves as an outgroup. The HBV/J isolate (JRB34) is indicated by an arrowhead, and an HBVC4 isolate from Australian aborigine is indicated by an open triangle. The genetic distance is indicated by a bar below.

exceeding 90 years; and “B” for Borneo where he is suspected to have contracted the HBV infection). The entire nucleotide sequence was determined for the JRB34 isolate of genotype J (HBV/J). It had a genomic length of 3,182 nt, which consisted of envelope gene containing preS1 region (nt 2848 to 3171, coding for 108 amino acids [aa]), preS2 region (nt 3172 to 154 [55 aa]), and the small S gene (nt 155 to 835 [226 aa]), X gene (nt 1374 to 1838 [154 aa]), preC region (nt 1814 to 1897 [27 aa]), C gene (nt 1901 to 2452 [183 aa]), and P gene (nt 2307 to 1623 [832 aa]).

Sequence divergence of the JRB34 strain from other genotypes. The complete genome sequence of the JRB34 strain obtained in the present study was compared against those of 1,440 HBV genomes registered in the Viral Hepatitis Database

(39). Estimated nucleotide sequence divergence of the JRB34 strain from four ape and nine human genotypes is summarized in the Table 1. The mean divergence by genotypes ranged from 10.7 and 10.9% (from orangutan and gibbon, respectively) to 15.6 and 15.7% (from genotypes F and H, respectively). Surprisingly, the minimum divergence of 9.9% was observed in comparison with a nonhuman HBV isolate from *Hilobates agilis* gibbon confiscated in Taiwan in 1993 (AY330917) (41). Since the sequence divergence from any documented genotypes, including recently proposed genotype I, exceeded 8%, the JRB34 strain was tentatively classified into a novel genotype J of HBV.

Phylogenetic analysis of the entire genomic sequence. In the phylogenetic tree constructed on 1,440 complete genome




FIG. 4. Comparison of the amino acid sequence in the preC gene and carboxy-terminal amino acid sequences in the C gene of HBV isolates of various genotypes. The sequence of the HBV/J isolate (JRB34) is indicated at the top. Dots represent amino acids shared by JRB34, and a dash indicates the deletion of an amino acid. The sequence of the arginine-rich domain bearing the binding site with HBV DNA is boxed.

EMBL/DDBJ/GenBank database entries, the HBV/J strain was positioned distinctively from all known human genotypes (data not shown). It was closest to the cluster formed by gibbon- and orangutan-derived strains. However, including recombinant strains in such analyses may significantly affect the overall phylogenetic topology. This possibility was ruled out by reconstruction of the phylogeny using nonrecombinant HBV strains that further confirmed the phylogenetic peculiarity of the studied JRB34 strain (see Fig. S1 in the supplemental material). A total of 44 representative reference strains were further selected for establishing the consistency. Thus, phylogenetic topology indicating genotype-specific clustering is shown in the Fig. 1. Hence, using various sets of references, we confirmed that genotype J undoubtedly differed phylogenetically from all other known genotypes.

Lack of significant evidence of recombination with other human or ape genotypes in genotype J. To investigate possible recombination in the JRB34 genome, a window scanning analysis of aligned HBV genomes was performed by means of Simplot and Simmonics software packages. Both Bootscanning

by SimPlot and GroupScanning by Simmonics showed similar output results. However, the methodological approach is different between these two software packages; GroupScanning provides more robust analysis of the phylogenetic relation between the examined strain and clusters of reference strains, whereas SimPlot does this comparison between the examined strain and parametrically generated consensus of the reference strains. The results obtained by SimPlot therefore can be significantly affected by selected parameters for the generation of consensus. This is especially undesirable when a new genotype strain (for which no references are available among known genotypes) is being analyzed (40). Figure 2 shows genome-wide distance scanning and GroupScanning plots for the JRB34 strain in comparison with a reference set consisting of 228 nonrecombinant HBV isolates retrieved from the public database (the phylogenetic tree is shown in Fig. S1 in the supplemental material). It is evident that the JRB34 strain was divergent from all known genotypes, and the closest genetic neighbors were estimated by distance and phylogenetic association scanning were the gibbon genotype (in preS, S, and P




FIG. 5. Comparison of amino acid sequences of the preS/S gene among HBV isolates of various genotypes. The sequence of the HBV/J isolate (JRB34) is indicated at the top. Dots represent amino acids shared by JRB34, and a dash indicates the deletion of an amino acid. The sequence from positions 101 to 156 forming loops, bearing the common antigenic determinants of HBsAg, is boxed.

genes) and genotype C (in the core gene). However, no significant evidence of recombination between these two ape and human genotypes was revealed by the used methods. Homology scan carried out by SimPlot using the same set of reference sequences gave concordant results.

Phylogenetic analyses of the four open reading frames. Phylogenetic relationship between the JRB34 strain and other genotypes was further analyzed in four open reading frames. In the small S gene, subgenotype C4 recovered from Australian aborigines (43) changed its phylogenetic topology from the branch of human genotypes to a branch intermediate between orangutan and gibbon strains (Fig. 3A). Remarkably, genotype J and C4 strains joined together to create a clade between orangutan and gibbon strains. In contrast, genotype J clustered with human genotypes in the phylogenetic analysis of the C gene and was closely related to genotype C; it took a position outside genotype I strains, however (Fig. 3B). Genotype J was closer to gibbon and orangutan genotypes in the phylogenetic trees constructed on P and large S genes (data not shown), demonstrating its topology similar to that in the analysis of the entire genome (Fig. 1).

Amino acid sequence of the HBV/J isolate. The amino acid sequence of HBV/J was compared against those of other genotypes over three different areas of the genome. The amino

acid sequence in the preC gene and arginine-rich domain in the carboxy-terminal sequence in the C gene were well conserved by genotype J (Fig. 4). In the preS1 region, genotype J had a deletion of 11 aa as gibbon and chimpanzee genotypes (Fig. 5). This deletion was shared by one of the two HBV/C4 isolates from Australian aborigines, as well as all HBV/D isolates. Amino acid sequence in the S gene of genotype J was the same as those of aborigine isolates of subgenotype C4; they would share antigenic epitopes of HBsAg. Amino acids at codons 122 and 160 were arginine (with G as nt 365) and lysine (with G as nt 479), respectively, which was consistent with subtype *ayw* of HBsAg from this patient (27).

Five domains (A to E) of DNA polymerase/reverse transcriptase in the P gene were preserved well in HBV/J, and it did not have mutations in the Tyr-Met-Asp-Asp motif in the domain C that determines the sensitivity to lamivudine (data not shown). HBV/J possessed A1762T/G1764A double mutations in the core promoter and G1896A stop codon mutation in the preC region, which was compatible with an HBeAg-minus phenotype of HBV recovered from the patient positive for anti-HBe.

Infection with HBV/J in chimeric mice with the liver repopulated for human hepatocytes. Two chimeric mice that had been transplanted with human hepatocytes were inoculated with 10⁴ HBV DNA copies of genotype J. In both mice, HBV




FIG. 6. Markers of HBV infection in two chimeric mice inoculated with the HBV/J isolate (JRB34). The levels of HBV DNA are illustrated in panel A, and those of HBsAg and HBcrAg are illustrated in panel B. Values represent the means for two mice.

DNA in a high titer (10^5 copies/ml) appeared in the circulation at week 7, plateaued at high levels (10^6 to 10^8 copies/ml), and stayed detectable until 22 weeks of observation after the inoculation (Fig. 6A). HBsAg and HBcrAg became detectable at week 7 and kept increasing in concentrations until week 15 when they reached a plateau at high levels (Fig. 6B). HBV strains recovered from mice at the last day of follow-up were identical in the complete genome sequence to the JRB34 strain used for inoculation.

The liver from chimeric mice infected with HBV/J was stained for HBcAg by immunofluorescence (Fig. 7A). The staining for HBcAg was confined to areas where mouse liver had been replaced for human hepatocytes, and the same areas were stained for human albumin (Fig. 7B). Colocalization of HBcAg and human hepatocytes was demonstrated by double staining for HBcAg and human albumin (Fig. 7C). Finally, expression and replication of the JRB34 strain were confirmed by successful detection of cccDNA and HBV RNA in the liver tissue from both sacrificed mice (see Fig. S2A and B in the supplemental material).

DISCUSSION

An HBV isolate (JRB34) was recovered from a male, 88-year-old Japanese patient with HCC and sequenced over the entire genome. In the full-genome sequence, the JRB34 strain

had 10.9 to 15.7% divergence from 1,440 HBV strains retrieved from the DDBJ/EMBL/GenBank. The divergence exceeds 8% that has been defined originally for distinguishing between four genotypes (A to D) (29) and later for an additional four genotypes (E to H) (3, 26, 42). Phylogenetically, the sequence of JRB34 was closer to ape than human HBV genotypes. No significant evidence of recombination with eight known human and four ape genotypes was revealed by the GroupScanning analysis (40) and phylogenetic analyses. These lines of evidence have qualified the JRB34 strain to represent a possible new HBV genotype. To further confirm the epidemiological significance of this strain, capable of establishing new infections, two chimeric mice were each inoculated with 10^4 copies of JRB34 HBV DNA. They both were successfully infected with sharp increases in HBV DNA and HBsAg in serum several weeks after the inoculation. Replication in the chimeric mice was also confirmed by detection of cccDNA and HBV RNA in their liver tissues.

Recently, an HBV isolate from Vietnam (VH24 [accession no. AB231908]) was reported as a ninth human genotype (I) (12). However, VH24 differed by only $7.0\% \pm 0.4\%$ from HBV isolates of genotype C and possessed complex recombination with genotypes A and G in three genomic areas. A number of sporadic HBV isolates have been reported to date that contain recombination between human genotypes (4, 24, 40), as well as between human and ape genotypes (21). Only a few recombinant variants, however, became widely spread in human populations, developing their own specific distributions and epidemiologies. This is particularly demonstrated for the B/C recombinant designated as a distinct subgenotype; Ba/B2-5 now accounts for the majority of genotype B strains in mainland Asia (44). Likewise, the C/D recombinant prevails in Tibet and northern China (50). To avoid assigning a new genotype for every newly discovered sporadic recombinant HBV variant, evidence of intergenotypic recombination should be carefully eliminated (14). However, in some cases, designation of a new genotype is proposed by a potential epidemiological significance of a novel genetic variant. Recently, a study carried out in Laos described a number of strains closely related phylogenetically with the Vietnamese genotype I strains, thereby suggesting their epidemiological significance (31). The JRB34 strain documented in the present study was genetically and phylogenetically distinct from any previously published strains, including those of genotype I from Vietnam and Laos. To avoid possible misconceptions in the future, the strain is provisionally designated genotype J.

HBV of distinct genotypes can infect great apes in the wild, including chimpanzee, gorilla, orangutan and gibbons (9, 20, 37, 51). HBV genotypes of chimpanzee and gorilla, as well as those of orangutan and gibbon, cocluster in agreement with their geographical distribution in Africa and Southeast Asia, respectively (41). Genotype J represented by the JRB34 strain clustered with gibbon/orangutan genotypes. In a phylogenetic analysis of the S region/gene sequence, JRB34 belonged to a nonhuman HBV group but was closely related to an HBV isolate of subgenotype C4 (AB048704) recovered from an Australian aborigine; C4 is most divergent from other subgenotypes of genotype C (43). In the phylogenetic analysis of the C gene, however, JRB34 clustered with human genotypes and closely related to genotype C, including C4, and was positioned




FIG. 7. (A and B) Immunofluorescent staining of a frozen liver section of a chimera mouse inoculated with the HBV/J isolate (JRB34). HBeAg is stained in panel A, and human albumin is stained in panel B. (C) Colocalization of HBeAg and human albumin is revealed by double staining. (D to F) HBV-uninfected mouse liver shows that only human albumin is stained.

outside genotype I strains (Fig. 4). Taken together, genotype J is phylogenetically close to gibbon/orangutan genotypes in the entire genome and to genotype C (C4 in particular) in the S and C genes. However, despite observed interchangeable relatedness with gibbon and genotype C/I strains, no strong evidence of recombination was confirmed in the JRB34.

In the sequence of C gene, carboxyl-terminal arginine-rich region, required for binding with HBV DNA, was preserved in JRB34. It had the G1896A stop codon in the precore region that aborts the translation of HBeAg (5, 30) and A1762T/G1764A double mutations in the core promoter that interfere with the transcription of HBeAg by downregulating preC mRNA (28, 45); they are compatible with the HBeAg⁻ anti-HBe⁺ phenotype of the patient from whom JRB34 was isolated. Since the double mutations are detected frequently in HBV DNA sequences from patients with HCC (17, 33), it could be implicated in hepatocarcinogenesis of the patient from whom JRB34 was isolated. It is not certain, however, if precore and core-promoter mutations had existed in HBV transmitted to the patient who is presumed to have been infected 60 years ago. Since amino acid sequences constituting antigenic loops of HBsAg (6) were the same as those of Australian aborigine isolates of C4, they would share antigenic epitopes of HBsAg. The amino acids at codons 122 and 160 were arginine (with G at nt 365) and lysine (with G at nt 479),

respectively (27), in agreement with subtype *ayw* of HBsAg from this patient. Five domains (A to E) of DNA polymerase/reverse transcriptase in the P gene were preserved well in HBV/J, and it did not have mutations in the Tyr-Met-Asp-Asp motif in the domain C that determines the sensitivity to lamivudine (2).

How and when the patient contracted infection with HBV/J is not certain. It is very unlikely, however, that he acquired infection in Japan via perinatal or horizontal transmission. There are no wild primates in Okinawa, where the patient was originally from, and the prevalent human HBV genotypes are limited to B (60%), C (39%), and sporadic cases of A (1%) (32). Furthermore, HBV/J was not found among patient's family members who are currently alive (data not shown). The phylogenetic position within open reading frames of JRB34 in between gibbon/orangutan genotypes and human genotype C gives a clue where and when the patient had contracted HBV infection. He was drafted to Borneo during World War II (1939 to 1945); the island in the Southeast Asia is inhabited by gibbons and orangutans and has a local population mainly infected with genotypes B or C. Zoonotic infection of HBV has been previously reported (11, 46), and HBV of genotype E was recovered from a chimpanzee captured in West Africa where this genotype is common. There is a possibility that JRB34 of

genotype J had been transmitted to the study patient in Borneo during the war (38).

The origin of genotype J in gibbon/orangutan or human inhabitants in Borneo is not certain but very likely. HBV DNA and/or HBsAg was detected in 26% (55/213) and 20% (58/297) of gibbons and orangutans, respectively, captured in Southeast Asia (38). HBV is also endemic in people living there, with a prevalence of HBsAg at 2 to 8%. There would be high chances for cross-species transmission of HBV where it prevails both in human beings and nonhuman primates. Phylogenetic analysis for close relationship between human and nonhuman HBV genotypes has indicated geographical influence rather than association with particular species (41).

It remains to be determined whether genotype J and ape-derived strains originate from species-specific convergent evolution of distant strains or whether they have diverged from a single common ancestor sometime in the past and evolved independently thereafter. The validity of cross-species infection or species-specific evolution for genotype J would be verified by sequence analysis of HBV DNA from gibbons and humans living in Borneo. If they turn out to be the same, cross-species infection will be justified. Should genotype J be restricted to human beings, in converse, species-specific infection will be confirmed.

In conclusion, a novel HBV genotype was identified in the Ryukyu isolate and provisionally named genotype J. Phylogenetic analyses over the full-length sequence and open reading frames indicate a close relationship of genotype J with gibbon/orangutan genotypes and human genotype C. The index patient would have been infected with HBV/J while he resided in Borneo inhabited by gibbons and orangutans. Although only one HBV isolate of genotype J (JRB34) has been identified, this may be only the tip of an iceberg. It would be worthwhile to examine the genotype of HBV infecting people and gibbons, as well as orangutans, living in Borneo and neighboring countries for mapping the epidemiology of genotype J and finding any clinical relevance.

ACKNOWLEDGMENTS

This study was supported in part by a grant-in-aid from the Ministry of Health, Labor and Welfare of Japan and a grant-in-aid from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

REFERENCES

- Abe, A., K. Inoue, T. Tanaka, J. Kato, N. Kajiyama, R. Kawaguchi, S. Tanaka, M. Yoshida, and M. Kohara. 1999. Quantitation of hepatitis B virus genomic DNA by real-time detection PCR. *J. Clin. Microbiol.* 37:2899–2903.
- Allen, M. I., M. Deslauriers, C. W. Andrews, G. A. Tipples, K. A. Walters, D. L. Tyrrell, N. Brown, L. D. Condreay, et al. 1998. Identification and characterization of mutations in hepatitis B virus resistant to lamivudine. *Hepatology* 27:1670–1677.
- Arauz-Ruiz, P., H. Norder, B. H. Robertson, and L. O. Magnius. 2002. Genotype H: a new Amerindian genotype of hepatitis B virus revealed in Central America. *J. Gen. Virol.* 83:2059–2073.
- Bollyky, P. L., and E. C. Holmes. 1999. Reconstructing the complex evolutionary history of hepatitis B virus. *J. Mol. Evol.* 49:130–141.
- Carman, W. F., M. R. Jacyna, S. Hadziyannis, P. Karayiannis, M. J. McGarvey, A. Makris, and H. C. Thomas. 1989. Mutation preventing formation of hepatitis B e antigen in patients with chronic hepatitis B infection. *Lancet* ii:588–591.
- Carman, W. F., A. R. Zanetti, P. Karayiannis, J. Waters, G. Manzillo, E. Tanzi, A. J. Zuckerman, and H. C. Thomas. 1990. Vaccine-induced escape mutant of hepatitis B virus. *Lancet* 336:325–329.
- Fung, S. K., and A. S. Lok. 2004. Hepatitis B virus genotypes: do they play a role in the outcome of HBV infection? *Hepatology* 40:790–792.
- Ganem, D., and A. M. Prince. 2004. Hepatitis B virus infection—natural history and clinical consequences. *N. Engl. J. Med.* 350:1118–1129.
- Grethe, S., J. O. Heckel, W. Rietschel, and F. T. Hufert. 2000. Molecular epidemiology of hepatitis B virus variants in nonhuman primates. *J. Virol.* 74:5377–5381.
- Hannoun, C., H. Norder, and M. Lindh. 2000. An aberrant genotype revealed in recombinant hepatitis B virus strains from Vietnam. *J. Gen. Virol.* 81:2267–2272.
- Hu, X., A. Javadian, P. Gagneux, and B. H. Robertson. 2001. Paired chimpanzee hepatitis B virus (ChHBV) and mtDNA sequences suggest different ChHBV genetic variants are found in geographically distinct chimpanzee subspecies. *Virus Res.* 79:103–108.
- Huy, T. T. T., T. N. Trinh, and K. Abe. 2008. New complex recombinant genotype of hepatitis B virus identified in Vietnam. *J. Virol.* 82:5657–5663.
- Kimura, T., A. Rokuhara, Y. Sakamoto, S. Yagi, E. Tanaka, K. Kiyosawa, and N. Maki. 2002. Sensitive enzyme immunoassay for hepatitis B virus core-related antigens and their correlation to virus load. *J. Clin. Microbiol.* 40:439–445.
- Kurbanov, F., Y. Tanaka, A. Kramvis, P. Simmonds, and M. Mizokami. 2008. When should “I” consider a new hepatitis B virus genotype? *J. Virol.* 82:8241–8242.
- Lee, W. M. 1997. Hepatitis B virus infection. *N. Engl. J. Med.* 337:1733–1745.
- Lindh, M., A. S. Andersson, and A. Gusdal. 1997. Genotypes, nt 1858 variants, and geographic origin of hepatitis B virus: large-scale analysis using a new genotyping method. *J. Infect. Dis.* 175:1285–1293.
- Liu, C. J., B. F. Chen, P. J. Chen, M. Y. Lai, W. L. Huang, J. H. Kao, and D. S. Chen. 2006. Role of hepatitis B viral load and basal core promoter mutation in hepatocellular carcinoma in hepatitis B carriers. *J. Infect. Dis.* 193:1258–1265.
- Liu, C. J., J. H. Kao, and D. S. Chen. 2005. Therapeutic implications of hepatitis B virus genotypes. *Liver Int.* 25:1097–1107.
- Lole, K. S., R. C. Bollinger, R. S. Paranjape, D. Gadkari, S. S. Kulkarni, N. G. Novak, R. Ingersoll, H. W. Sheppard, and S. C. Ray. 1999. Full-length human immunodeficiency virus type 1 genomes from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J. Virol.* 73:152–160.
- MacDonald, D. M., E. C. Holmes, J. C. Lewis, and P. Simmonds. 2000. Detection of hepatitis B virus infection in wild-born chimpanzees (*Pan troglodytes verus*): phylogenetic relationships with human and other primate genotypes. *J. Virol.* 74:4253–4257.
- Magiorkinis, E. N., G. N. Magiorkinis, D. N. Paraskevis, and A. E. Hatzakis. 2005. Re-analysis of a human hepatitis B virus (HBV) isolate from an East African wild born *Pan troglodytes schweinfurthii*: evidence for interspecies recombination between HBV infecting chimpanzee and human. *Gene* 349:165–171.
- Reference deleted.
- Miyakawa, Y., and M. Mizokami. 2003. Classifying hepatitis B virus genotypes. *Intervirology* 46:329–338.
- Morozov, V., M. Pisareva, and M. Groudinin. 2000. Homologous recombination between different genotypes of hepatitis B virus. *Gene* 260:55–65.
- Norder, H., A. M. Courouze, P. Coursaget, J. M. Echevarria, S. D. Lee, I. K. Mushahwar, B. H. Robertson, S. Locarnini, and L. O. Magnius. 2004. Genetic diversity of hepatitis B virus strains derived worldwide: genotypes, subgenotypes, and HBsAg subtypes. *Intervirology* 47:289–309.
- Norder, H., A. M. Courouze, and L. O. Magnius. 1994. Complete genomes, phylogenetic relatedness, and structural proteins of six strains of the hepatitis B virus, four of which represent two new genotypes. *Virology* 198:489–503.
- Okamoto, H., M. Imai, F. Tsuda, T. Tanaka, Y. Miyakawa, and M. Mayumi. 1987. Point mutation in the S gene of hepatitis B virus for a *dy* or *w/r* subtypic change in two blood donors carrying a surface antigen of compound subtype *adyr* or *adwr*. *J. Virol.* 61:3030–3034.
- Okamoto, H., F. Tsuda, Y. Akahane, Y. Sugai, M. Yoshida, K. Moriyama, T. Tanaka, Y. Miyakawa, and M. Mayumi. 1994. Hepatitis B virus with mutations in the core promoter for an e antigen-negative phenotype in carriers with antibody to e antigen. *J. Virol.* 68:8102–8110.
- Okamoto, H., F. Tsuda, H. Sakugawa, R. I. Sastrosoewignjo, M. Imai, Y. Miyakawa, and M. Mayumi. 1988. Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J. Gen. Virol.* 69(Pt. 10):2575–2583.
- Okamoto, H., S. Yotsumoto, Y. Akahane, T. Yamanaka, Y. Miyazaki, Y. Sugai, F. Tsuda, T. Tanaka, Y. Miyakawa, and M. Mayumi. 1990. Hepatitis B viruses with precore region defects prevail in persistently infected hosts along with seroconversion to the antibody against e antigen. *J. Virol.* 64:1298–1303.
- Olinger, C. M., P. Jutavittum, J. M. Hubschen, A. Yousukh, B. Samoury, T. Thammavong, K. Toriyama, and C. P. Muller. 2008. Possible new hepatitis B virus genotype, southeast Asia. *Emerg. Infect. Dis.* 14:1777–1780.
- Orito, E., T. Ichida, H. Sakugawa, M. Sata, N. Horiike, K. Hino, K. Okita, T. Okanoue, S. Iino, E. Tanaka, K. Suzuki, H. Watanabe, S. Hige, and M. Mizokami. 2001. Geographic distribution of hepatitis B virus (HBV) genotype in patients with chronic HBV infection in Japan. *Hepatology* 34:590–594.

33. Orito, E., M. Mizokami, H. Sakugawa, K. Michitaka, K. Ishikawa, T. Ichida, T. Okanoue, H. Yotsuyanagi, and S. Iino. 2001. A case-control study for clinical and molecular biological differences between hepatitis B viruses of genotypes B and C. *Hepatology* 33:218–223.
34. Palumbo, E. 2007. Hepatitis B genotypes and response to antiviral therapy: a review. *Am. J. Ther.* 14:306–309.
35. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406–425.
36. Sakamoto, T., Y. Tanaka, E. Orito, J. Co, J. Clavio, F. Sugauchi, K. Ito, A. Ozasa, A. Quino, R. Ueda, J. Sollano, and M. Mizokami. 2006. Novel subtypes (subgenotypes) of hepatitis B virus genotypes B and C among chronic liver disease patients in the Philippines. *J. Gen. Virol.* 87:1873–1882.
37. Sall, A. A., S. Starkman, J. M. Reynes, S. Lay, T. Nhim, M. Hunt, N. Marx, and P. Simmonds. 2005. Frequent infection of *Hylobates pileatus* (pileated gibbon) with species-associated variants of hepatitis B virus in Cambodia. *J. Gen. Virol.* 86:333–337.
38. Sa-nguanmoo, P., C. Thongmee, P. Ratanakorn, R. Pattanarangsarn, R. Boonyarittichaikij, S. Chodapisitkul, A. Theamboonlers, P. Tangkijvanich, and Y. Poovorawan. 2008. Prevalence, whole genome characterization and phylogenetic analysis of hepatitis B virus in captive orangutan and gibbon. *J. Med. Primatol.* 37:277–289.
39. Shin-I, T., Y. Tanaka, Y. Tateno, and M. Mizokami. 2008. Development and public release of a comprehensive hepatitis virus database. *Hepatology Res.* 38:234–243.
40. Simmonds, P., and S. Midgley. 2005. Recombination in the genesis and evolution of hepatitis B virus genotypes. *J. Virol.* 79:15467–15476.
41. Starkman, S. E., D. M. MacDonald, J. C. Lewis, E. C. Holmes, and P. Simmonds. 2003. Geographic and species association of hepatitis B virus genotypes in non-human primates. *Virology* 314:381–393.
42. Stuyver, L., S. De Gendt, C. Van Geyt, F. Zoulim, M. Fried, R. F. Schinazi, and R. Rossau. 2000. A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J. Gen. Virol.* 81:67–74.
43. Sugauchi, F., M. Mizokami, E. Orito, T. Ohno, H. Kato, S. Suzuki, Y. Kimura, R. Ueda, L. A. Butterworth, and W. G. Cooksley. 2001. A novel variant genotype C of hepatitis B virus identified in isolates from Australian Aborigines: complete genome sequence and phylogenetic relatedness. *J. Gen. Virol.* 82:883–892.
44. Sugauchi, F., E. Orito, T. Ichida, H. Kato, H. Sakugawa, S. Kakumu, T. Ishida, A. Chutaputti, C. L. Lai, R. Ueda, Y. Miyakawa, and M. Mizokami. 2002. Hepatitis B virus of genotype B with or without recombination with genotype C over the precore region plus the core gene. *J. Virol.* 76:5985–5992.
45. Takahashi, K., K. Aoyama, N. Ohno, K. Iwata, Y. Akahane, K. Baba, H. Yoshizawa, and S. Mishiro. 1995. The precore/core promoter mutant (T1762A1764) of hepatitis B virus: clinical significance and an easy method for detection. *J. Gen. Virol.* 76(Pt. 12):3159–3164.
46. Takahashi, K., B. Brotman, S. Usuda, S. Mishiro, and A. M. Prince. 2000. Full-genome sequence analyses of hepatitis B virus (HBV) strains recovered from chimpanzees infected in the wild: implications for an origin of HBV. *Virology* 267:58–64.
47. Tanaka, Y., and M. Mizokami. 2007. Genetic diversity of hepatitis B virus as an important factor associated with differences in clinical outcomes. *J. Infect. Dis.* 195:1–4.
48. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673–4680.
49. Tiollais, P., P. Charnay, and G. N. Vyas. 1981. Biology of hepatitis B virus. *Science* 213:406–411.
50. Wang, Z., Z. Liu, G. Zeng, S. Wen, Y. Qi, S. Ma, N. V. Naoumov, and J. Hou. 2005. A new intertype recombinant between genotypes C and D of hepatitis B virus identified in China. *J. Gen. Virol.* 86:985–990.
51. Wiegand, J., D. Hasenclever, and H. L. Tillmann. 2008. Should treatment of hepatitis B depend on hepatitis B virus genotypes? A hypothesis generated from an explorative analysis of published evidence. *Antivir. Ther.* 13:211–220.
52. Wong, D. K., Y. Tanaka, C. L. Lai, M. Mizokami, J. Fung, and M. F. Yuen. 2007. Hepatitis B virus core-related antigens as markers for monitoring chronic hepatitis B infection. *J. Clin. Microbiol.* 45:3942–3947.

ユニバーサル HB ワクチン接種に関する提言

厚生労働科学研究補助金 肝炎等克服緊急対策研究事業

「肝炎ウイルス感染防御を目指したワクチン接種の基盤構築」班

平成 21 年 3 月

研究代表者

国立感染症研究所 血液・安全性研究部

水 落 利 明

目 次

提言	-----	3
研究組織	-----	5
提言作成の背景	-----	7
世界におけるユニバーサル HB ワクチン接種(HB-UV)の現状	-----	8
国内における HBV 感染者の実態	-----	10
HB-UV の効果	-----	12
HB ワクチンの選択	-----	14
HB ワクチンの追加接種	-----	16
HB ワクチンの副作用	-----	18
HB-UV の対費用効果	-----	20
HB-UV の是非について： 日本肝臓学会評議員を対象に実施されたアンケート調査結果	-----	22
研究班各班員による HB-UV についての意見	-----	23
まとめ／謝辞	-----	29
参考資料（文献）	-----	31