

hepatitis B specimens was divided into two distinct patterns. In one liver specimen with moderate fibrosis and inflammation from patient 2 in Table 2 who was diagnosed with HCC 9 years later, intense pSmad3L immunostaining was present in the nuclei of all hepatocytes throughout the liver lobules; C-terminal phosphorylation of Smad3 was strongly suppressed in hepatocytic nuclei (Fig. 1A and Supplementary Fig. 1). In another specimen with similar fibrotic stage and necroinflammatory activity from patient 44 in Table 2 who had not developed HCC, many hepatocytes retained phosphorylation at Smad3C but showed scarce phosphorylation at Smad3L (Fig. 1B). Among 37 patients with chronic hepatitis B who had strong pSmad3L positivity, 24 patients showed little Smad3C phosphorylation, and only 13 patients

**Table 3. Correlation Between pSmad3L and pSmad3C in Chronic Hepatitis B Specimens**

	pSmad3C Positivity *		Total
	Low (1 and 2)	High (3 and 4)	
pSmad3L positivity *			
Low (1 and 2)	11	22	33
High (3 and 4)	24	13	37
Total	35	35	70

Abbreviations: pSmad3C, C-terminally phosphorylated Smad3; pSmad3L, linker-phosphorylated Smad3.

\*Hepatocytic Smad3 phosphorylation is scored as follows: 0, no phosphorylation; 1, <25% Smad3 phosphorylation; 2, 25% to 50% Smad3 phosphorylation; 3, 50% to 75% Smad3 phosphorylation; 4, >75% Smad3 phosphorylation.

showed abundant Smad3C phosphorylation (64.9% versus 35.1% [ $P = 0.03$ ]) (Table 3). In contrast, 22 patients with little Smad3L phosphorylation (scores 0 to 2) versus only 13 patients with abundant Smad3L phosphorylation (scores 3 to 4) showed strong pSmad3C positivity (62.9% versus 37.1% [ $P = 0.04$ ]). Because the extent of Smad3L phosphorylation increased as fibrotic stage and necroinflammatory activity progressed in chronic hepatitis C, Smad3L showed little phosphorylation in early chronic hepatitis C.<sup>20</sup> In contrast, degree of linker phosphorylation of Smad3 in hepatocytic nuclei remained high (staining scored as 3 or 4) in 21 of 47 patients with chronic hepatitis B (F1 to F2) (Fig. 1C). These results indicate differential mechanisms of HBV- and HCV-





Fig. 1. Two distinct hepatocytic Smad3 signaling pathways in early chronic hepatitis B: pSmad3L- and pSmad3C-dominant types. (A) Smad3 in the nuclei of hepatocytes was phosphorylated sparsely at the C-terminal region ( $\alpha$  pSmad3C column) but intensely at the linker region ( $\alpha$  pSmad3L column). The liver specimen showing moderate fibrosis and inflammation was obtained from patient 2 in Table 2 diagnosed with HCC 9 years later. Bar = 50  $\mu$ m. (B) In patient 44 in Table 2 who had not developed HCC, hepatocytes retained phosphorylation at Smad3C ( $\alpha$  pSmad3C column) but showed little phosphorylation at Smad3L ( $\alpha$  pSmad3L column). The specimen showed degrees of fibrosis and necroinflammatory activity similar to those in (A). Formalin-fixed, paraffin-embedded liver sections were stained with anti-pSmad3L Ab ( $\alpha$  pSmad3L column) and anti-pSmad3C Ab ( $\alpha$  pSmad3C column). The pSmad3C section was paired with an adjacent section stained using anti-pSmad3L Ab. Abs were then bound by goat anti-rabbit IgG conjugated with peroxidase-labeled polymer. Peroxidase activity was detected with 3,3'-diaminobenzidine tetrahydrochloride. All sections were counterstained with hematoxylin (blue). Brown color indicates specific Ab reactivity. Bar = 50  $\mu$ m. (C) Degrees of Smad3 phosphorylation were stable in hepatocytic nuclei in early chronic hepatitis B specimens (F1 to F2), whereas pSmad3L increased and pSmad3C decreased as chronic hepatitis B (F3) progressed through cirrhosis to HCC. Smad3 phosphorylation in hepatocytes did not change between F1 and F2 stages. Phosphorylation of Smad3L and Smad3C in hepatocytes of cirrhotic liver (F4) and HCC was greater and less than that in livers with grade F1 and F2 fibrosis, respectively. Extent of Smad3 phosphorylation: □, 0; □, 1; ■, 2; ■, 3; ■, 4. \* $P < 0.05$ .

associated carcinogenesis, especially in the early stages of chronic hepatitis.

**pSmad3L Prominence in Hepatocytic Nuclei in Proportion to Plasma HBV DNA Levels.** Because HCC risk is related to plasma HBV DNA levels and chronic inflammation,<sup>24</sup> we next investigated the correlation of hepatocytic pSmad3L positivity with plasma HBV DNA levels and necroinflammatory activity in chronic hepatitis B patients (Fig. 2). Positivity of hepatocytic nuclei for pSmad3L in chronic hepatitis B specimens gradually increased in proportion to amounts of HBV DNA.



Sixteen sera samples of 35 patients with abundant Smad3L phosphorylation (scores 3 to 4) but only seven sera samples of 35 patients with little Smad3L phosphorylation (scores 0 to 2) contained more than 5.0 log copies/mL (45.7% versus 20.0% [ $P = 0.02$ ]) (Fig. 2A). However, 23 of 35 chronic hepatitis B patients with abundant Smad3L phosphorylation and 22 of 35 patients with little Smad3L phosphorylation showed a high level of inflammatory activity (A 2 to 3) (65.7% versus 62.9% [ $P = 0.80$ ]) (Fig. 2B). These results indicated that HBV itself could up-regulate the hepatocytic phosphorylation at Smad3L, but inflammation could not strongly affect linker phosphorylation.

To further confirm the direct effects of HBV, besides chronic inflammation, on phosphorylation at Smad3L in early chronic hepatitis B, we examined the degrees of pSmad3L and pSmad3C in a group of patients with little fibrosis (F1), little inflammation (A1), and high plasma HBV DNA. Smad3L was highly phosphorylated in hepatocytic nuclei, whereas the phosphorylation at Smad3C was suppressed (Fig. 2C). Of 20 chronic hepatitis B samples, 12 samples showed abundant Smad3L phosphorylation (scores 3 to 4), but only five samples had abundant Smad3C phosphorylation (60.0% versus 25.0% [ $P = 0.03$ ]) (Table 4).

**HBx Protein Involvement in c-Myc-Mediated Oncogenic Activity via the pSmad3L Pathway in Human Chronic Hepatitis B.** Integrated viral sequences produce HBx protein, which brings about up-regulation of c-Myc oncoprotein.<sup>25</sup> We therefore investigated whether HBx protein affected Smad3L phosphorylation and expression of c-Myc in biopsy specimens from HBV-infected livers by immunostaining sections for

Fig. 2. In proportion to plasma HBV DNA levels, JNK-dependent pSmad3L became prominent in the nuclei of hepatocytes in human early chronic hepatitis B. (A) Positivity for pSmad3L in hepatocytic nuclei in chronic hepatitis B specimens was greater in proportion to plasma HBV DNA levels. Patients with strong pSmad3L positivity in hepatocytic nuclei (staining scored as 3 or 4) had more HBV DNA in plasma than patients with weak pSmad3L positivity (staining scored as 0 to 2). Hepatocytic Smad3 phosphorylation in chronic hepatitis B specimens is scored as follows: 0, no phosphorylation; 1, <25%; 2, 25% to 50%; 3, 50% to 75%; 4, >75%. (B) Degree of Smad3 phosphorylation at the linker region did not strongly correlate with necroinflammatory activity of chronic hepatitis B. Hepatocytic Smad3 phosphorylation at the linker region in livers with necroinflammatory activities of A2 to A3 was essentially similar to phosphorylation in those with activities of A0 to A1. Extent of necroinflammatory activity: □, 0; ◻, 1; ◼, 2; ◼, 3. (C) Smad3 in the nuclei of hepatocytes was phosphorylated intensely at linker region (α pSmad3L column) but sparsely at the C-terminal region (α pSmad3C column). The liver specimen showing minimal fibrosis (F1) and inflammation (A1) was obtained from patient 10 in Table 2 who showed high plasma HBV DNA and was diagnosed with HCC 4 years later.

**Table 4. Clinicopathologic Features, Smad3L/C Phosphorylation, and Plasma HBV DNA Levels in Specimens from Patients with Early Chronic Hepatitis B**

Patient No.	Sex	Age	pSmad3L staining*	pSmad3C staining*	Fibrotic Stage†	Inflammatory Activity†	HBV DNA (log copies/mL)
1	M	46	4	2	1	1	5.6
2	M	45	4	2	1	1	5.4
3	F	38	3	1	1	1	6.2
4	F	49	3	1	1	1	5.2
5	F	52	3	1	1	1	6.1
6	F	40	3	2	1	1	5.8
7	F	28	3	2	1	1	5.6
8	M	38	3	2	1	1	5.5
9	M	44	3	2	1	1	5.3
10	F	40	3	2	1	1	5.4
11	M	55	3	2	1	1	5.1
12	F	43	3	3	1	1	5.8
13	M	34	2	1	1	1	5.2
14	M	28	1	1	1	1	5.2
15	F	35	1	2	1	1	5.1
16	M	30	1	2	1	1	5.3
17	M	45	1	3	1	1	5.2
18	M	38	1	4	1	1	5.6
19	M	54	1	4	1	1	5.2
20	M	48	1	4	1	1	5.4

Abbreviations: F, female; HBV, hepatitis B virus; M, male; pSmad3C, C-terminally phosphorylated Smad3; pSmad3L, linker-phosphorylated Smad3.

\*Hepatocytic Smad3 phosphorylation is scored as follows: 0, no phosphorylation; 1, <25% Smad3 phosphorylation; 2, 25% to 50% Smad3 phosphorylation; 3, 50% to 75% Smad3 phosphorylation; 4, >75% Smad3 phosphorylation.

†Necroinflammatory activity and fibrotic stage are determined histologically according to Desmet's classification.

pSmad3L, paired with sections immunostained for HBx and c-Myc.

In specimens from patient 3 in Table 2 with chronic hepatitis B, pSmad3L, HBx, and c-Myc were distributed in hepatocytes throughout liver lobules (Fig. 3A and Supplementary Fig. 2). Double immunofluorescence studies in chronic hepatitis B specimens confirmed that pSmad3L was colocalized in HBx- and c-Myc-immunoreactive hepatocytes (Fig. 3B). HBx and c-Myc expression increased in hepatocytes of hepatitis B specimens as Smad3 showed more phosphorylation at the linker region (Fig. 3C).

**Increased JNK/pSmad3L/c-Myc Oncogenic Signaling and Impaired pSmad3C/p21<sup>WAF1</sup> Tumor-Suppressive Signaling as Chronic Hepatitis B Progresses From Cirrhosis to HCC.** We further investigated tumor-suppressive and oncogenic Smad3 signaling in biopsy specimens during HBV-related hepatocarcinogenesis by staining sections using anti-pSmad3L Ab and anti-pSmad3C Ab, paired with sections stained for anti-c-Myc Ab and anti-p21<sup>WAF1</sup> Ab. Although pSmad3L accelerates tumor growth by up-regulating c-Myc, pSmad3C participates in tumor suppression by up-regulating p21<sup>WAF1</sup> transcription.<sup>16, 20</sup>

In specimens from a patient with chronic hepatitis B, the distribution of pSmad3L fit well with the pattern shown by c-Myc immunolabeling (Fig. 4A, chronic hep-

atitis panel): both were strong in hepatocytes throughout liver lobules. Linker phosphorylation and c-Myc staining increased further as chronic liver disease progressed through cirrhosis to HCC (Fig. 4A, cirrhosis and HCC panels).

Distribution of pSmad3C resembled the pattern obtained by p21<sup>WAF1</sup> staining in chronic hepatitis B specimens (Fig. 4B, chronic hepatitis panel). As with pSmad3C distribution, hepatocytes showed increased p21<sup>WAF1</sup> staining in nuclei. In contrast to intense staining for pSmad3L and c-Myc, pSmad3C and p21<sup>WAF1</sup> staining decreased in hepatocytic nuclei in cirrhotic liver (Fig. 4B, cirrhosis panel). Nuclear pSmad3C and p21<sup>WAF1</sup> immunostaining showed only a scattered distribution throughout HCC specimens (Fig. 4B, HCC panel). Semiquantitative analyses of positivity for pSmad3L, pSmad3C, and c-Myc in HBV-related chronic liver disease showed increasing pSmad3L/c-Myc and decreasing pSmad3C as chronic hepatitis B progressed from cirrhosis (F4) to HCC (Table 1).

We next quantified the extent of phosphorylation at Smad3L and Smad3C by immunoblotting with domain-specific Abs against phosphorylated Smad3 in tissue samples representing various stages of HBV-related chronic liver disorders. The linker region of Smad3 showed very little phosphorylation in normal liver (Fig. 4C,  $\alpha$  pSmad3L panel). Remarkable up-




Fig. 3. HBx might be involved in c-Myc-mediated oncogenic activity in human chronic hepatitis B via the pSmad3L pathway. (A) Hepatocytes of chronic hepatitis B specimens from patient 3 in Table 2 showed diffuse immunostaining for pSmad3L, HBx, and c-Myc. All sections were counterstained with hematoxylin (blue). Brown color indicates specific Ab reactivity. Bar = 50  $\mu$ m. (B) pSmad3L in hepatocytic nuclei of chronic hepatitis B specimens was colocalized with HBx and c-Myc proteins. Sections of chronic hepatitis B tissues were stained for immunofluorescence to simultaneously detect pSmad3L (red) and HBx or c-Myc (green). Yellow color indicates overlap of proteins. Hepatocytes immunoreactive for pSmad3L showed colocalization of HBx (upper column) and c-Myc (lower column). Bar = 50  $\mu$ m. (C) HBx and c-Myc expression increased in hepatocytes of chronic hepatitis B specimens as Smad3 was increasingly phosphorylated at the linker region. HBx and c-Myc expression was greater in hepatocytes with high phosphorylation at Smad3L (staining scored as 3 or 4) than in hepatocytes with staining scored as 0 to 2. The extent of HBx and c-Myc expression is indicated as that of pSmad3L positivity: □, 0; ◻, 1; ◼, 2; ◼, 3; ◼, 4. \* $P < 0.05$ .

regulation of pSmad3L was seen with progression of hepatic fibrosis and carcinogenesis. In cirrhotic liver and HCC, pSmad3L was far more abundant than in chronic hepatitis. In contrast, pSmad3C gradually decreased as disease stages progressed toward HCC (Fig. 4C,  $\alpha$  pSmad3C panel).

We previously reported that Smad3L served as a substrate for JNK.<sup>12</sup> To address the functional relationship between activated JNK and Smad3L phosphorylation during hepatocarcinogenesis, we presently assayed kinase activity *in vitro*. Although JNK from normal liver showed little ability to phosphorylate Smad3 at the linker region, JNK from livers involved by chronic hepatitis B, cirrhosis, and HCC could directly phosphorylate Smad3L (Fig. 4D). These results suggested that JNK in preneoplastic liver tissues and HCC directly phosphorylated the linker region of Smad3.

Collectively, JNK/pSmad3L/c-Myc oncogenic signaling in hepatocytes came to predominate while the tumor-suppressive pSmad3C/p21<sup>WAF1</sup> pathway became quiescent as chronic hepatitis B progressed to cirrhosis and then HCC.

**Selective Blockade of Linker Phosphorylation Abolishes pSmad3L-Mediated Cell Growth in HBx-Expressing Hepatocytes.** pSmad3L, HBx, and c-Myc were colocalized in preneoplastic lesions including chronic hepatitis and cirrhosis (Fig. 3). These findings suggest that HBx oncoprotein might alter hepatocytic TGF- $\beta$  signaling in chronic hepatitis B. We investigated this hypothesis using HBx-expressing hepatocytes. Selective blockade of linker phosphorylation by a mutant Smad3 lacking the JNK-dependent linker phosphorylation sites abolished pSmad3L-mediated cell growth in HBx-expressing hepatocytes (Supplementary Figs. 3-5). These results suggest that HBx activated the JNK/pSmad3L pathway, further promoting cell proliferation by up-regulating c-Myc transcription (Fig. 5).

**Activation of the pSmad3L/c-Myc Pathway as HBx Transgenic Mouse Livers Progress Through Hyperplasia to HCC.** We further investigated localization of pSmad3L, HBx, and c-Myc during HBx-induced hepatocarcinogenesis in HBx transgenic mouse livers. Beginning at the age of 2 months, HBx transgenic mouse liver showed centrilobular foci of cellular alteration with cytoplasmic vacuolation surrounding the central veins where bromodeoxyuridine was uptaken into the hepatocytes.<sup>6</sup>

In this hyperplastic mouse liver, phosphorylation at Smad3L was observed in hepatocytic nuclei in the centrilobular region, and distribution of pSmad3L was similar to those of HBx and c-Myc (Fig. 6A). pSmad3L, HBx,




Fig. 4. As chronic hepatitis B progressed through cirrhosis to HCC, JNK/pSmad3L/c-Myc oncogenic signaling started to increase, whereas the tumor-suppressive pSmad3C/p21<sup>WAF1</sup> pathway decreased. (A) pSmad3L and c-Myc increased as human chronic hepatitis B progressed through cirrhosis to HCC. (B) pSmad3C and p21<sup>WAF1</sup> decreased as human chronic hepatitis B progressed through cirrhosis to HCC. All sections in A and B were counterstained with hematoxylin (blue). Brown staining indicates specific Ab reactivity. Bar = 50  $\mu$ m. (C) Immunoblotting of pSmad3L and pSmad3C in HBV-related chronic liver diseases. Cell lysates obtained from hepatocellular carcinoma (HCC) and surrounding nonneoplastic liver tissues including chronic hepatitis B (CH) or liver cirrhosis (LC) as well as uninvolved normal liver tissues from a patient with a metastatic liver tumor were subjected to anti-Smad3 immunoprecipitation (IP) and were then immunoblotted with each anti-pSmad3 Ab (upper panels). Relative amounts of endogenous Smad3 were determined via immunoblotting using anti-Smad3 Ab (bottom panel). (D) JNK in human HBV-related chronic liver tissue directly phosphorylated Smad3 at the linker region. Cell lysates obtained from HCC and surrounding nonneoplastic liver tissue including chronic hepatitis (CH) and liver cirrhosis (LC) from HBV-infected patients, as well as uninvolved normal liver tissue from a patient with a liver metastasis, were subjected to anti-phospho-JNK1/2 immunoprecipitation (IP), and were then mixed with bacterially expressed GST-Smad3 and GST-Smad2. After *in vitro* kinase assay, phosphorylation of Smad3L was analyzed via immunoblotting using anti-pSmad3L antibody (upper panel). Total Smad3 and Smad2 were determined via immunoblotting using anti-Smad2/3 Ab (lower panel).

and c-Myc were distributed diffusely in HCC specimens (Fig. 6B, HCC panel). Semiquantitative analyses of positivity for these molecules in HBx transgenic mouse livers also revealed that hepatocytic pSmad3L, HBx, and c-Myc increased as mouse liver progressed through hyperplasia to HCC (Fig. 6C). Double immunofluorescence studies in hyperplastic specimens confirmed that pSmad3L was colocalized in HBx- and c-Myc-immunoreactive hepatocytes (Fig. 6D).

Success in comparative study of HBx, pSmad3L, and c-Myc positivity during human and mouse hepatocarcinogenesis identified pSmad3L as a key regulatory element

that offers a general framework for understanding the origins of HBV-related HCC.

**Chronic Hepatitis B Patients with Hepatocytes Positive for pSmad3L and Negative for pSmad3C Increased Risk of HCC Development.** We finally investigated whether phosphorylation levels of Smad3 could affect the risk of neoplastic evolution in the patients with chronic hepatitis B (Table 2). To compare HCC incidence, patients were classified into those with abundant (scores 3 to 4) and limited (scores 0 to 2) Smad3 phosphorylation in hepatocytic nuclei. HCC developed in six of 28 patients with abundant Smad3L phosphorylation




Fig. 5. Reversibility of Smad3-dependent signaling between tumor suppression and oncogenesis in HBx-expressing hepatocytes. Hepatocytes exhibit TGF- $\beta$ -dependent Smad3 phosphorylation at the C-terminal region, which results in growth inhibition by repression of c-Myc. High expression of HBx protein in hepatocytes tends to shut down pSmad3C-mediated signaling and favor acquisition of constitutively active JNK-mediated pSmad3L signaling, which fosters cell growth by up-regulating c-Myc. Selective blockade of linker phosphorylation by a mutant Smad3 lacking the JNK-dependent linker phosphorylation sites (Smad3E9PSM) restores the TGF- $\beta$ -dependent tumor-suppressive response involving pSmad3C that is shown by parental hepatocytes.

lation, but in only one of 32 patients with limited Smad3L phosphorylation (log-rank = 0.03) (Fig. 7A). In contrast, HCC developed only in the patients with limited Smad3C phosphorylation, and no patients with abundant hepatocytic pSmad3C developed HCC (log-rank = 0.009) (Fig. 7B).

Several studies have analyzed risk factors for HCC occurrence in patients with HBV-related chronic liver disease, including elevated plasma HBV DNA<sup>24</sup> and seropositivity for hepatitis B e antigen.<sup>26</sup> In the univariate analysis, HCC occurrence in high pSmad3L positivity ( $P = 0.01$ ), low pSmad3C positivity ( $P = 0.03$ ), and plasma HBV-DNA levels of more than 5.0 log copies/mL ( $P = 0.05$ ) showed  $P$  values less than 0.10, thus being significantly associated with HCC (Table 5). All variables with statistical significance in the univariate analysis were entered in the multivariate analysis, and high pSmad3L and low pSmad3C positivity were considered significantly predictive of HCC development within 12 years. Hepatocytic positivity for pSmad3L and pSmad3C should allow us to distinguish chronic hepatitis B patients at high and low risk for the development of HCC in near future.

## Discussion

In patients with chronic hepatitis B, persistent HBV infection is clearly the primary inducer of HCC.<sup>1-7</sup> Com-

parative studies that seek to identify conserved oncogenic signaling common to HCC in both humans and experimental animals will help to eventually identify the molecular pathways that drive the development of HCC.<sup>27</sup> Much is known about the morphologic changes of cells and tissues that precede and accompany development of HCC in humans, allowing earlier diagnosis in some instances.<sup>28</sup> A variety of molecular alterations have been detected in fully developed HCC and to a lesser extent in morphologically defined preneoplastic precursor lesions.<sup>29</sup> Our current studies compared pSmad3L- and pSmad3C-mediated signaling in biopsy specimens of chronic hepatitis, cirrhosis, or HCC from 90 patients with chronic HBV infection versus signaling in preneoplastic and neoplastic liver lesions of HBx transgenic mice. Taken together with the results of *in vitro* experiments using HBx-expressing hepatocytes, our findings indicate that the HBx oncoprotein participates directly in hepatocarcinogenesis by shifting hepatocytic Smad3-mediated signaling from tumor suppression to oncogenesis in patients with early chronic hepatitis B (Fig. 7C). According to the two-step model of carcinogenesis (initiation and promotion), tumor formation can be explained by permanent HBx-dependent activation of the JNK/pSmad3L cascade that has a tumor promoter-like action.

HCC is a human neoplasm associated with viral infection.<sup>1,3</sup> At present, hepatitis virus-associated carcinogenesis can be seen as a multifactorial process that includes both direct and indirect mechanisms.<sup>19</sup> A major factor in the process of HCC development is the host immune system.<sup>30</sup> Chronic inflammation, degeneration, and regeneration induced by the host cellular immune response are common to a variety of human liver diseases, and subsequent cellular proliferation might increase the risk of cancer. We previously reported that increased phosphorylation of Smad3L and decreased phosphorylation of Smad3C were associated with an increased risk of HCV-related HCC.<sup>20</sup> Similarly to HCV-related chronic liver disease, strong pSmad3L positivity was observed in the late stages of HBV-related chronic liver disease (F3 to F4) (Fig. 1C). Considering the development of HCC in HCV core gene-transgenic mice,<sup>31</sup> hepatitis viruses themselves together with the host immune response might promote human hepatocarcinogenesis via the JNK/pSmad3L pathway during the late stage of the carcinogenic process in both HBV- and HCV-related chronic liver disease.

However, HBV and HCV have different roles in human hepatocarcinogenesis when early chronic hepatitis (F1 to F2) is considered. The histological severity of HCV-related liver disease correlates closely with the risk




Fig. 6. The pSmad3L/c-Myc pathway was activated as HBx transgenic mouse liver progressed through hyperplasia to HCC. (A) Distribution of pSmad3L, HBx, and c-Myc in hyperplastic specimens from HBx transgenic mouse liver. (B) Distributions of pSmad3L, HBx, and c-Myc in normal liver, hyperplasia, and HCC specimens from HBx transgenic mice. Immunostaining for pSmad3L, HBx, and c-Myc was present in hyperplastic hepatocytes surrounding central veins in HBx transgenic mouse liver [(A) and (B), hyperplasia panels], and was distributed diffusely in HCC specimens [(B), HCC panel]. All sections were counterstained with hematoxylin (blue). Brown color indicates specific Ab reactivity. Bar = 50  $\mu$ m. (C) Hepatocytic pSmad3L, HBx, and c-Myc increased as HBx transgenic mouse liver progressed from hyperplasia to HCC. Staining for pSmad3L, HBx, and c-Myc was detected minimally in normal mouse livers, but was strongly up-regulated in neoplastic livers. In HCC, pSmad3L, HBx, and c-Myc were significantly greater than in livers with hyperplasia. \* $P < 0.05$ . Extent of pSmad3L, HBx, and c-Myc: □, 0; ◻, 1; ◼, 2; ◽, 3; ◼, 4. (D) Hepatocytic pSmad3L in hyperplastic specimens from HBx transgenic mouse liver was colocalized with HBx and c-Myc. Hyperplasia sections of HBx transgenic mouse livers were stained for immunofluorescence to simultaneously detect pSmad3L (red) and HBx or c-Myc (green). Yellow color indicates overlap of proteins. Hepatocytes immunoreactive for pSmad3L showed colocalization of HBx (upper column) and c-Myc (lower column). Bar = 50  $\mu$ m.

of HCC.<sup>32</sup> In contrast, HCC occasionally develops in healthy HBV surface antigen carriers, who are persistently infected with HBV but have normal liver function parameters and no necroinflammation.<sup>33</sup> This indicates that HBV itself has a direct influence on hepatocarcinogenesis in early chronic hepatitis B. Although integration of the viral genome into chromosomal DNA has not been reported in patients with HCV infection, integration of HBV has been detected in almost all cases of chronic hepatitis B,<sup>3</sup> leading to activation of the HBx-mediated oncogenic pathway.<sup>4</sup> It is noteworthy that HCC developed in patient 10 (Table 2), who showed strong pSmad3L positivity of hepatocytic nuclei but had minimal necroinflammatory activity (A1) or fibrosis (F1). In summary, HCV contributes indirectly to the development of HCC through chronic inflammation in early

chronic hepatitis C. In contrast, HBV directly triggers the JNK/pSmad3L oncogenic pathway in early chronic hepatitis B, thus playing a role beyond mere stimulation of the host immune response.

Our findings also open up a new avenue to understanding the development and progression of hepatic fibrogenesis.<sup>34</sup> Whereas HSCs have traditionally been considered as the principal source of liver fibrosis, mature hepatocytes can acquire a mesenchymal phenotype and perform the functions of activated HSC—that is, they can contribute to fibrogenesis.<sup>35,36</sup> In support of this notion, pSmad3L-mediated signaling promotes liver fibrosis by hepatocytes as well as activated HSCs during long-standing carcinogenesis.<sup>13,18,20</sup> In this manner, either HBV- or HCV-related chronic hepatitis progresses through fibrogenesis to HCC.




Fig. 7. Chronic hepatitis B patients with hepatocytes positive for pSmad3L and negative for pSmad3C increased risk of HCC development. (A) HCC occurred subsequently among patients whose hepatocytes in chronic hepatitis B specimens were strongly positive for pSmad3L. Incidence of HCC was significantly higher in patients with abundant Smad3L phosphorylation (scores 3 to 4, solid line) in hepatocytic nuclei versus those with sparse Smad3L phosphorylation (scores 0 to 2, dotted line). (B) HCC did not occur subsequently among patients whose hepatocytes in chronic hepatitis B specimens were strongly positive for pSmad3C. HCC occurred only in patients with sparse Smad3C phosphorylation (scores 0 to 2, solid line) in hepatocytic nuclei, while no patients with abundant Smad3C phosphorylation (scores 3 to 4, dotted line) have developed HCC. Cumulative rates of HCC occurrence from chronic hepatitis B were compared between cases with high and low phosphorylation of Smad3L and Smad3C (Kaplan-Meier analysis and log-rank test). (C) HBx protein shifted hepatic TGF- $\beta$  signaling from the tumor-suppressive pSmad3C pathway to the oncogenic JNK-dependent pSmad3L pathway in early stages of chronic hepatitis B. Normal hepatocytes exhibited TGF- $\beta$ -dependent Smad3 phosphorylation at the C-terminal region, which is related to growth inhibition by up-regulation of p21<sup>WAF1</sup>. HBx protein activates JNK, promoting the oncogenic pSmad3L signaling, which fosters cell growth by up-regulating c-Myc, in a mean time reducing tumor-suppressive pSmad3C-mediated signaling.

The general biomedical approach to HCC is shifting away from population risk assessment and empirical treatment of patients to predictive personalized medicine based on molecular classification and targeted therapy.<sup>29</sup> Better knowledge of the risk factors associated with the occurrence of HCC can improve the effectiveness of surveillance programs. Our approach has identified pSmad3L and pSmad3C as prognostic markers that may prove to be clinically useful. Such predictive markers could allow us to select patients with chronic hepatitis B who have a high or low risk of developing HCC. Although the latter group could be followed up on an annual basis, the patients with a high risk require targeted surveillance measures to allow early diagnosis of HCC.

Phosphorylation of many transcription factors is controlled by the dynamic interplay between kinases and phosphatases. In this regard, we studied the kinetics of both linker and C-terminal phosphorylation of Smad3 in parental and HBx-expressing hepatocytes in response to TGF- $\beta$  (unpublished observation). In parental hepatocytes, the levels of linker and C-terminal phosphorylation peaked at 30 minutes after the start of exposure to TGF- $\beta$  and then gradually declined. However, HBx-expressing hepatocytes showed constitutive phosphorylation at Smad3L during continuous exposure to TGF- $\beta$ . Several lines of evidence have identified small C-terminal domain phosphatase (SCP1-3) and protein phosphatase magnesium 1A (PPM1A) as the linker and C-terminal phosphatases, respectively.<sup>37,38</sup> Accordingly, SCP1-3 and PPM1A may reverse domain-specific phosphorylation in normal hepatocytes. In contrast, HBx-expressing hepatocytes may not show induction or activation of SCP1-3. Alternatively, linker phosphorylation in HBx-expressing hepatocytes might be resistant to SCP1-3.

Many researchers have been seeking key transcription factors regulating tumor-suppressive pathways that are altered in cancer. Our current model of JNK/pSmad3L signaling during HBV-related chronic liver disease suggests that specific inhibitors of the JNK/pSmad3L pathway might inhibit the progression of HCC. With respect to molecular targeting therapy for human HCC, pSmad3L and pSmad3C should be assessed as biomarkers to evaluate the benefit from specific inhibition of the JNK/pSmad3L pathway.

**Acknowledgment:** We thank Dr. Rik Derynck (University of California at San Francisco) and Dr. Seishi Murakami (Kanazawa University) for providing us with complementary DNAs encoding human Smad3 and HBx. We also thank Chiaki Kitano for assistance to construct ecotropic retrovirus and Natsuko Ohira for assistance with immunoblotting.



**Table 5. Variables with Independent Predictive Value for HCC in Univariate and Multivariate Analyses**

Characteristics	n	No. of Patients with HCC (%)	Univariate Analysis		Multivariate Analysis	
			Hazard Ratio (95% CI)	P Value	Hazard Ratio (95% CI)	P Value
pSmad3L positivity*						
Low (1 and 2)	32	1 (3)	1.00		1.00	
High (3 and 4)	28	6 (21)	3.8 (1.4-10.6)	0.01	14.8 (1.8-118.5)	0.01
pSmad3C positivity*						
High (3 and 4)	30	0 (0)	1.00		1.00	
Low (1 and 2)	30	7 (23)	2.8 (0.001-7.0)	0.03	16.4 (1.0-125.0)	0.04
Fibrotic stage†						
Low (F1 and F2)	39	4 (10)	1.00		1.00	
High (F3)	21	3 (14)	1.9 (0.7-5.4)	0.24	3.9 (0.4-38.6)	0.24
Inflammatory activity‡						
Low (A0 and A1)	23	1 (4)	1.00		1.00	
High (A2 and A3)	37	6 (16)	1.8 (0.7-4.8)	0.27	0.2 (0.02-1.1)	0.06
HBV DNA (copies /mL)						
<10 <sup>5</sup>	42	3 (7)	1.00		1.00	
>10 <sup>5</sup>	18	4 (22)	1.9 (1.0-3.5)	0.05	2.5 (0.9-6.9)	0.08
HBeAg						
Negative	42	4 (10)	1.00		1.00	
Positive	18	3 (17)	2.1 (0.5-9.5)	0.32	9.9 (1.1-89.3)	0.03

Abbreviations: CI, confidence interval; HBeAg, hepatitis B e antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; pSmad3C, C-terminally phosphorylated Smad3; pSmad3L, linker-phosphorylated Smad3.

\*Hepatocytic Smad3 phosphorylation in chronic hepatitis B specimens is scored as follows: 0, no phosphorylation; 1, <25% Smad3 phosphorylation; 2, 25% to 50% Smad3 phosphorylation; 3, 50% to 75% Smad3 phosphorylation; 4, >75% Smad3 phosphorylation.

†Necroinflammatory activity and fibrotic stage are determined histologically according to Desmet's classification.

## References

- Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 2003; 362:1907-1917.
- El-Serag HB. Hepatocellular carcinoma: recent trends in the United States. *Gastroenterology* 2004;127(Suppl):27S-34S.
- Brechot C, Pourcel C, Louise A, Rain B, Tiollais P. Presence of integrated hepatitis B virus DNA sequences in cellular DNA of human hepatocellular carcinoma. *Nature* 1980;286:533-535.
- Kim CM, Koike K, Saito I, Miyamura T, Jay G. HBx gene of hepatitis B virus induces liver cancer in transgenic mice. *Nature* 1991;351:317-320.
- Yu DY, Moon HB, Son JK, Jeong S, Yu SL, Yoon H, et al. Incidence of hepatocellular carcinoma in transgenic mice expressing the hepatitis B virus X-protein. *J Hepatol* 1999;31:123-132.
- Koike K, Moriya K, Iino S, Yotsuyanagi H, Endo Y, Miyamura T, et al. High-level expression of hepatitis B virus HBx gene and hepatocarcinogenesis in transgenic mice. *HEPATOLOGY* 1994;19:810-819.
- Benn J, Su F, Doria M, Schneider RJ. Hepatitis B virus HBx protein induces transcription factor AP-1 by activation of extracellular signal-related and c-Jun N-terminal mitogen-activated protein kinases. *J Virol* 1996;70:4978-4985.
- Roberts AB, Sporn MB. The transforming growth factor- $\beta$ s. In: Sporn MB, Roberts AB, eds. *Peptide Growth Factors and Their Receptors*. Berlin: Springer-Verlag, 1990:419-472.
- Heldin CH, Miyazono K, ten Dijke P. TGF- $\beta$  signaling from cell membrane to nucleus through SMAD proteins. *Nature* 1997;390:465-471.
- Massagué J. TGF- $\beta$  signal transduction. *Annu Rev Biochem* 1998;67:753-791.
- Kretzschmar M, Doody J, Timokhina I, Massagué J. A mechanism of repression of TGF- $\beta$ /Smad signaling by oncogenic Ras. *Genes Dev* 1999; 13:804-816.
- Mori S, Matsuzaki K, Yoshida K, Furukawa F, Tahashi Y, Yamagata H, et al. TGF- $\beta$  and HGF transmit the signals through JNK-dependent Smad2/3 phosphorylation at the linker regions. *Oncogene* 2004;23:7416-7429.
- Furukawa F, Matsuzaki K, Mori S, Tahashi Y, Yoshida K, Sugano Y, et al. p38 MAPK mediates fibrogenic signal through Smad3 phosphorylation in rat myofibroblasts. *HEPATOLOGY* 2003;38:879-889.
- Matsuura I, Denissova NG, Wang G, He D, Long J, Liu F. Cyclin-dependent kinases regulate the antiproliferative function of Smads. *Nature* 2004;430:226-231.
- Yamagata H, Matsuzaki K, Mori S, Yoshida K, Tahashi Y, Furukawa F, et al. Acceleration of Smad2 and Smad3 phosphorylation via c-Jun NH(2)-terminal kinase during human colorectal carcinogenesis. *Cancer Res* 2005; 65:157-165.
- Sekimoto G, Matsuzaki K, Yoshida K, Mori S, Murata M, Seki T, et al. Reversible Smad-dependent signaling between tumor suppression and oncogenesis. *Cancer Res* 2007;67:5090-5096.
- Arany PR, Rane SG, Roberts AB. Smad3 deficiency inhibits v-ras-induced transformation by suppression of JNK MAPK signaling and increased farnesyl transferase inhibition. *Oncogene* 2008;27:2507-2512.
- Yoshida K, Matsuzaki K, Mori S, Tahashi Y, Yamagata H, Furukawa F, et al. Transforming growth factor- $\beta$  and platelet-derived growth factor signal via c-Jun N-terminal kinase-dependent Smad2/3 phosphorylation in rat hepatic stellate cells after acute liver injury. *Am J Pathol* 2005;166:1029-1039.
- Block TM, Mehta AS, Fimmel CJ, Jordan R. Molecular viral oncology of hepatocellular carcinoma. *Oncogene* 2003;22:5093-5107.
- Matsuzaki K, Murata M, Yoshida K, Sekimoto G, Uemura Y, Sakaida N, et al. Chronic inflammation associated with hepatitis C viral infection perturbs hepatic TGF- $\beta$  signaling, promoting cirrhosis and hepatocellular carcinoma. *HEPATOLOGY* 2007;46:48-57.
- Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Scheuer PJ. Classification of chronic hepatitis: diagnosis, grading and staging. *HEPATOLOGY* 1994;19:1513-1520.
- Cox DR. Regression models and life-tables. *J R Stat Soc (B)* 1972;34:187-220.
- Pardali K, Moustakas A. Actions of TGF- $\beta$  as tumor suppressor and prometastatic factor in human cancer. *Biochim Biophys Acta* 2007;1775:21-62.

24. Chen CJ, Yang HI, Su J, Jen CL, You SL, Lu SN, et al. REVEAL-HBV study group. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006;295:65-73.
25. Feltel MA. c-Myc overexpression in hepatocarcinogenesis. *Human Pathology* 2004;35:1299-1302.
26. Yang HI, Lu SN, Liaw YF, You SL, Sun CA, Wang LY, et al. Taiwan community-based cancer screening project group. Hepatitis B e antigen and the risk of hepatocellular carcinoma. *N Engl J Med* 2002;347:168-174.
27. Thorgeirsson SS, Lee JS, Grisham JW. Functional genomics of hepatocellular carcinoma. *HEPATOLOGY* 2006;43:145-150.
28. Theise ND, Park YN, Kojiro M. Dysplastic nodules and hepatocarcinogenesis. *Clin Liver Dis* 2002;6:497-512.
29. Thorgeirsson SS, Grisham JW. Molecular pathogenesis of human hepatocellular carcinoma. *Nat Genet* 2002;31:339-346.
30. Chusari FV, Klopchin K, Moriyama T, Pasquinelli C, Dunsford HA, Sell S, et al. Molecular pathogenesis of hepatocellular carcinoma in hepatitis B virus transgenic mice. *Cell* 1989;59:1145-1156.
31. Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, et al. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065-1067.
32. Di Bisceglie AM. Hepatitis C and hepatocellular carcinoma. *HEPATOLOGY* 1997;26 (Suppl):34S-38S.
33. Popper H, Shafritz DA, Hoofnagle JH. Relation of the hepatitis B virus carrier state to hepatocellular carcinoma. *HEPATOLOGY* 1987;7:764-772.
34. Inagaki Y, Okazaki I. Emerging insights into transforming growth factor  $\beta$  Smad signal in hepatic fibrogenesis. *Gut* 2007;56:284-292.
35. Kaumori A, Potter J, Kaumori JY, Wang C, Mezey E, Koteish A. Transforming growth factor- $\beta$ 1 induces an epithelial-to-mesenchymal transition state in mouse hepatocytes in vitro. *J Biol Chem* 2007;282:22089-22101.
36. Weng HL, Ciucian L, Liu Y, Hamzavi J, Godoy P, Gantantz H, et al. Profibrogenic transforming growth factor- $\beta$ /activin receptor-like kinase 5 signaling via connective tissue growth factor expression in hepatocytes. *HEPATOLOGY* 2007;46:1257-1270.
37. Lin X, Duan X, Liang YY, Su Y, Wrighton KH, Long J, et al. PPM1A functions as a Smad phosphatase to terminate TGF $\beta$  signaling. *Cell* 2006;125:915-928.
38. Wrighton KH, Willis D, Long J, Liu F, Lin X, Feng XH. Small C-terminal domain phosphatases dephosphorylate the regulatory linker regions of Smad2 and Smad3 to enhance transforming growth factor- $\beta$  signaling. *J Biol Chem* 2006;281:38365-38375.

# A Single Amino Acid of Toll-like Receptor 4 That Is Pivotal for Its Signal Transduction and Subcellular Localization\*

Received for publication, April 22, 2008, and in revised form, October 29, 2008. Published, JBC Papers in Press, December 8, 2008, DOI 10.1074/jbc.M803086200

Shintaro Yanagimoto<sup>†§¶</sup>, Keita Tatsuno<sup>§</sup>, Shu Okugawa<sup>§</sup>, Takatoshi Kitazawa<sup>§</sup>, Kunihiisa Tsukada<sup>§</sup>, Kazuhiko Koike<sup>§</sup>, Tatsuhiko Kodama<sup>¶</sup>, Satoshi Kimura<sup>||</sup>, Yoshikazu Shibasaki<sup>‡1</sup>, and Yasuo Ota<sup>\*\*1,2</sup>

From the <sup>†</sup>Center for Structuring Life Sciences, Graduate school of Arts and Sciences, University of Tokyo, Meguro-ku, Tokyo 153-8903, the <sup>§</sup>Department of Infectious Diseases, Graduate School of Medicine, University of Tokyo, Bunkyo-ku, Tokyo 113-8655, the <sup>¶</sup>Laboratory for Systems Biology and Medicine, Research Center for Advanced Science and Technology, University of Tokyo, Meguro-ku, Tokyo 153-8904, the <sup>||</sup>Tokyo Teishin Hospital, Fujimi, Chiyoda-ku, Tokyo 102-8798, and the <sup>\*\*</sup>Department of Medicine, Teikyo University School of Medicine, 2-11-1, Kaga, Itabashi-ku, Tokyo 173-8605, Japan

Toll-like receptor 4 (TLR4) is essential for recognizing a Gram-negative bacterial component, lipopolysaccharide (LPS). A single amino acid mutation at position 712 of murine TLR4 leads to hyporesponsiveness to LPS. In this study we determined that an amino acid, a leucine at position 815 of human TLR4, is also pivotal for LPS responsiveness and subcellular distribution. By replacing the leucine with alanine, the mutant TLR4 lost responsiveness to LPS and did not localize on the plasma membrane. In addition, it does not coprecipitate with myeloid differentiation-2, an accessory protein that is necessary for TLR4 to recognize LPS. These results suggest that the leucine at position 815 is required for the normal maturation of TLR4 and for formation of the TLR4-MD-2 complex.

Toll-like receptors (TLRs)<sup>3</sup> play essential roles in both innate and adaptive immunity (1). Thirteen members of the TLR family have been identified in mammals. TLRs have leucine-rich-repeats in their extracellular domains and a Toll/Interleukin-1 receptor (TIR) in their cytoplasmic domains, the latter of which mainly mediates intracellular signaling. Signaling pathways of TLRs, except for TLR3, depend on an adapter protein, MyD88 (myeloid differentiation factor 88), which interacts with the TIR domain of TLRs. This pathway leads to the activation of the transcription fac-

tor NF- $\kappa$ B and production of cytokines such as tumor necrosis factor- $\alpha$  and interleukin-6. Another important signaling pathway mediated by TLR3 and TLR4 that exploits the TIR domain is the MyD88-independent pathway. This pathway involves different adapter proteins, such as the TIR domain-containing adaptor inducing interferon- $\beta$  (TRIF) and TRIF-related adaptor molecule (2–4), and is essential for production of type I interferon through activation of interferon regulatory factor-3.

TLRs recognize as ligands several microbial pathogen-associated molecular patterns. One such pathogen-associated molecular pattern is lipopolysaccharide (LPS), which is recognized by TLR4. LPS triggers severe immunologic reactions by the host in Gram-negative bacterial infections and has drawn attention in many clinical situations. TLR4 is the first mammalian TLR to be discovered in the context of immunology. TLR4 was identified in the search for the genes responsible for LPS hyporesponsiveness (5, 6). The defect was found to stem from a single amino acid mutation, replacement of proline with histidine at position 712, in the cytoplasmic tail of murine TLR4. The study led to the discovery of the importance of TLR4 in innate immunity.

A variety of cells are activated by LPS stimulation through TLR4. TLR4 forms a receptor complex with an accessory protein, myeloid differentiation-2 (MD-2). MD-2 first associates with TLR4 in the endoplasmic reticulum (ER) and *cis*-Golgi, and both proteins move together to the plasma membrane (7, 8). Upon recognition of LPS, the TLR4-MD-2 complex receives LPS on the cell surface and initiates intracellular signaling. The expression of TLR4 in the absence of MD-2 does not confer full responsiveness to LPS stimuli in experimental cell lines (9). An analysis of MD-2 knockout mice revealed that MD-2 is important not only for LPS sensing but also for cellular distribution of TLR4.

In this study we hypothesized that the cytoplasmic tail of TLR4 contains regions that control both localization and signaling. Using truncation and mutation analysis, and paying particular attention to the TIR domain, we identified a single amino acid that is pivotal for both TLR4 signaling and subcellular distribution. The site we found was on the C-terminal portion of the TIR domain for which no specific function has been yet determined.

\* This work was partly supported by the Program of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation, by the Focus 21 project of the New Energy and Industrial Technology Development Organization, and by the Special Coordination Fund for Science and Technology from the Ministry of Education, Culture, Sports, Science and Technology. This study was also partly supported by a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology (to Y. O.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Both authors contributed equally to this work.

<sup>2</sup> To whom correspondence should be addressed. Tel.: 81-3-3964-1211 (ext. 1756); Fax: 81-3-3579-6310; E-mail: yasuo-ota-ty@umin.ac.jp.

<sup>3</sup> The abbreviations used are: TLR, Toll-like receptor; TIR, Toll/Interleukin-1 receptor; TRIF, TIR domain-containing adaptor inducing interferon- $\beta$ ; LPS, lipopolysaccharide; MD-2, myeloid differentiation-2; ER, endoplasmic reticulum; GFP, green fluorescent protein; EGFP, enhanced GFP; RLA, relative luciferase activity; Sulfo-NHS-SS-Biotin, sulfosuccinimidyl-2-(biotinamido)ethyl-1,3-dithiopropionate.

## An Important Amino Acid of TLR4 for Its Function

### EXPERIMENTAL PROCEDURES

**Reagents and Other Materials**—Lipopolysaccharide (LPS) from *Escherichia coli* O55:B5 was purchased from Sigma-Aldrich and applied without repurification. FLAG- and hexa-histidine (His<sub>6</sub>)-tagged human TLR4 expression plasmid (pEFBOS/humanTLR4<sub>flaghis</sub>) and FLAG- and His<sub>6</sub>-tagged human MD-2 expression plasmid (pEFBOS/humanMD-2<sub>flaghis</sub>) were generous gifts from Dr. Kensuke Miyake (Institute of Medical Science, University of Tokyo, Japan). Human CD14 cDNA plasmid (pCMV6-XL5/humanCD14) was purchased from OriGene (Rockville, MD). Fluorescent protein expression vector pEGFP-N3 was purchased from Clontech (Mountain View, CA). Anti-TLR4 monoclonal antibody (clone HTA125) was purchased from Abcam (Cambridge, MA). Anti-FLAG monoclonal antibody (clone M2) was purchased from Sigma-Aldrich. Anti-*A.v.* (GFP) monoclonal and polyclonal antibodies were purchased from Clontech. Control immunoglobulins for immunoprecipitation were purchased from BD Biosciences (San Jose, CA). Horseradish peroxidase-labeled anti-immunoglobulins antibodies were purchased from Dako (Glostrup, Denmark). BlockAce (DS Pharma Biomedical, Osaka, Japan) solution was used as blocking buffer for Western blotting.

**Cell Culture**—Human embryonic kidney (HEK) 293T cells were maintained in Dulbecco's modified Eagle's medium (Sigma-Aldrich) containing 10% heat-inactivated fetal bovine serum supplemented with penicillin-streptomycin solution (Invitrogen). FuGENE 6 transfection reagent (Roche Applied Science) was used for transient cotransfection according to the manufacturer's instructions. Culture dishes or plates were prepared to 70% confluence prior to transfection. Cells were used for experiments 36 h later. The transfection conditions were optimized for microscopic observation of the expressed fluorescent protein and were kept unchanged in other experiments.

**Expression Vector Subcloning and Mutagenesis**—Wild-type TLR4 cDNA was excised from pEFBOS/humanTLR4<sub>flaghis</sub> and subcloned into pEGFP-N3 so that when expressed enhanced green fluorescent protein (EGFP) would be fused at the C terminus of TLR4 (pEGFP-N3/humanTLR4). All mutations were introduced into pEFBOS/humanTLR4<sub>flaghis</sub> and pEGFP-N3/humanTLR4 using the QuikChange site-Directed mutagenesis kit (Stratagene, La Jolla, CA) according to the manufacturer's instructions and were confirmed by sequencing. For the truncation analysis, two identical unique restriction sites were prepared in the TLR4-coding region of pEFBOS/humanTLR4 using a QuikChange kit, and the DNA fragment to be removed, which was a part of the C terminus of TLR4, was excised enzymatically. After agarose gel purification, the linear double-stranded DNA was ligated to re-form a circular plasmid. Restriction sites were designed so as not to cause a frameshift between TLR4 and EGFP.

**Confocal Laser Scanning Microscopy of Cells**—Samples were fixed in 3% paraformaldehyde-phosphate-buffered saline at 37 °C for 10 min. Fluorescence images of fixed samples were recorded using a FluoView FV1000 Confocal Microscope (an inverted confocal laser scanning microscope, Olympus, Tokyo, Japan).

**Immunoprecipitation**—Transfected cells were lysed in lysis buffer (50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 0.1% Triton X-100, 1 mM 1,4-dithiothreitol, and proteinase inhibitor mixture), sonicated, and centrifuged at 4 °C. Antibody was added to the supernatant, and the sample was rotated 1 h at 4 °C followed by the addition of protein G-Sepharose (GE Healthcare Life Sciences, Piscataway, NJ) and an additional 8-h incubation at 4 °C. Bound protein was washed three times in lysis buffer. Proteins were eluted by boiling in SDS sample buffer.


**Biotinylation and Purification of Cell Surface Proteins**—Prior to surface biotinylation, HEK 293T cells plated in a 100-mm dish were transiently transfected as described above. Surface biotinylation and subsequent purification of biotinylated proteins were performed using a Cell Surface Protein Biotinylation and Purification Kit (Pierce) following the manufacturer's instructions. Briefly, membrane-impermeable sulfo-succinimidyl-2-(biotinamido)ethyl-1,3-dithiopropionate (Sulfo-NHS-SS-Biotin) was added to cell monolayers in the culture dishes and covalently bound to amines in proteins exposed on the cell surface. The affinity resin that binds to the biotin end of Sulfo-NHS-SS-Biotin was used to collect the biotinylated proteins. Reduction by 1,4-dithiothreitol causes cleavage of the disulfide bond in Sulfo-NHS-SS-Biotin, and the elute contains the biotinylated cell surface proteins. Each final sample obtained was considered to contain proteins from an equal amount of cells, because all culture plates were treated equally and grown to full confluence. All samples were sonicated and subjected to SDS-PAGE and Western blotting. The membrane to which protein was transferred was blocked in blocking buffer for 1 h. Then the membrane was incubated with a primary antibody, followed by incubation with horseradish peroxidase-labeled anti-immunoglobulins antibody. The protein bands were then visualized by using a chemiluminescence reagent, Immobilon Western Chemiluminescent HRP Substrate (Millipore, Billerica, MA), according to the manufacturer's instructions.

**Cell Stimulation Assays**—HEK293T cells were plated and transiently transfected for assays. Thirty-six hours after the transfection, LPS was added to fresh culture medium in each well of the culture plates at the stated concentration. The duration of LPS stimulation was 7 h.

**Dual Luciferase Reporter Assays for NF- $\kappa$ B Activation**—HEK293T cells were plated in 12-well culture plates ( $4 \times 10^4$  cells/well), and experimental cDNA plasmids were transiently transfected 36 h later using the FuGENE 6 transfection reagent with 0.5  $\mu$ g of NF- $\kappa$ B reporter plasmid expressing firefly luciferase (pNF- $\kappa$ B-Luc, Stratagene) and 0.05  $\mu$ g of constitutively active *Renilla* luciferase reporter plasmid (pRL-TK, Promega, Madison, WI) in addition to 0.5  $\mu$ g each of TLR4-EGFP plasmid and MD-2 plasmid. Stimulation experiments were performed 36 h later. Firefly luciferase and *Renilla* luciferase activities were measured using the Dual-Luciferase Reporter Assay System (Promega) and the Genelight55 luminometer (Microtech, Chiba, Japan). Relative luciferase activity (RLA) was obtained as the ratio of firefly luciferase activity to *Renilla* luciferase activity. Results are expressed as the ratio of RLA with LPS stimulation to RLA without LPS stimulation ( $[\text{RLA LPS+}]/[\text{RLA LPS-}]$ ). This ratio should ideally approach 1 when no response to LPS stimulation is observed.



**FIGURE 1. Alignment of the cytoplasmic domains of EGFP fusion TLR4 truncation mutants used in this study.** TLR4 (766tr) signifies the mutant truncated at position 766. Others are named in the same manner. The amino acids are colored based on their physicochemical properties: pink, basic; blue, acidic; green, polar and neutral; and orange, hydrophobic. The black underline represents the TIR domain. Colored overlines indicate amino acid sequences identical to known sorting signal motifs except for two LLs, which are dileucine motif-like sequences in that they consist of solely two consecutive leucines without preceding aspartate or glutamate. Capital letters on the line signify the single-letter code for amino acids: E, glutamic acid; L, leucine; R, arginine; and Y, tyrosine. X signifies any amino acid, and Ø signifies an amino acid residue with a bulky hydrophobic side chain.



**FIGURE 2. LPS responsiveness measured by NF- $\kappa$ B luciferase assay.** HEK293T cells were transfected with plasmids containing the gene for wild-type TLR4 or a truncated human TLR4-EGFP fusion protein, in addition to a luciferase reporter and human MD-2 plasmid (A) or unmodified plasmids (control) (B). After 36 h, cells were stimulated with LPS (10 ng/ml) for 7 h, and luciferase reporter gene activity was measured. All results were expressed as the ratio of relative luciferase activity with LPS stimulation to that without stimulation. The data were from three independent experiments. Small bars indicate 95% confidence intervals of the mean ( $p$  values for \* are: TLR4 (WT)-EGFP/MD-2 (+),  $p = 0.002$ ; TLR4 (826tr)-EGFP/MD-2 (+),  $p = 0.016$ ).

**Statistical Analyses**—All quantitative experiments were repeated three times, and each experiment was done in triplicate. The ratio of relative luciferase activity of LPS+ to LPS- was calculated as the index of the responsiveness to the stimuli as explained above. When positive response is observed, the ratio should significantly exceed one. The means of the ratio were represented in bar graphs. The 95% confidence interval of the mean of the ratio was calculated and indicated on each bar in the graph, and  $p$  values were calculated using Student's  $t$  distribution compared with the hypothetical mean, one.

## RESULTS

**Truncation Analysis of TLR 4**—To identify amino acid sequences in the cytoplasmic tail of TLR4 that are involved in


both signal transduction and subcellular distribution, first we generated five truncation mutants of TLR4 with a fluorescent protein (EGFP) at the C terminus of TLR4.

Although there are no known definite sorting signal motifs in the cytoplasmic tail of TLR4, some amino acid sequences are similar or identical to known general sorting signal motifs as shown in Fig. 1. YXXØ, a form of tyrosine-based sorting signal, and EXXXLL, a form of dileucine (LL)-based sorting signal, both control protein internalization, lysosomal targeting, and basolateral targeting (10), where "X" represents any amino acid, "Ø" stands for an amino acid residue with a bulky hydrophobic side chain, and other letters are single-letter abbreviations for the amino acids. "Diacidic" signals such as DXE mediate export from the ER (11). RR or RXR is another example of an ER export signal (12). Truncation sites were chosen so that some of these amino acid sequences were deleted in each mutant. Because the TIR domain, which is essential in TLR4 signaling and possibly subcellular localization (13), spans most of the cytoplasmic domain of TLR4, four out of five mutants have involvement in the TIR domain, which we hypothesized could result in impaired signal transduction and a change in subcellular distribution. Part of the cytoplasmic portion of the amino acid sequence of the truncation mutants is shown in Fig. 1. The five truncation mutant proteins lost their C-terminal tails at positions 826, 815, 802, 788, and 766, respectively, and were conjugated with EGFP *in vitro*. Actual truncation and ligation sites of all actual mutants were confirmed to have the designed DNA alignment by sequencing.

We utilized the luciferase reporter assay to assess NF- $\kappa$ B transcription activity as an indicator of TLR4 response to LPS stimuli. MD-2 is reported to be essential for this response (9). However, because it is not known whether MD-2 is necessary for transduction of the truncated TLR4 signal as well, we performed the assays with and without MD-2. The index of cell responsiveness to the stimulation was measured as the ratio between RLA with LPS stimulation and RLA without LPS stimulation. Only cells transfected with TLR4 (826tr)-EGFP in combination with MD-2 retained responsiveness to LPS stimulation. One exception was wild-type TLR4-EGFP (Fig. 2A). HEK293T cells transfected with TLR4 but without MD-2 did not respond to LPS stimuli regardless of the TLR4-EGFP genotype (Fig. 2B).

Next, we compared the localization of wild-type and truncated mutants of TLR4-EGFP in HEK293T by fluorescence microscopy (Fig. 3A). The wild-type TLR4 cotransfected with MD-2 was expressed on the plasma membrane and also in the

## An Important Amino Acid of TLR4 for Its Function



**FIGURE 3. Residues 815–826 of TLR4 contain a region necessary for plasma membrane localization.** *A*, cells were cultured on coverslips in 12-well plates and transfected as in Fig. 2. EGFP-tagged TLR4 was visualized by laser confocal microscopy. Fluorescence from EGFP was observed in green. Each genotype of TLR4-EGFP was cotransfected with a human MD-2 plasmid or empty vector. Bar, 20  $\mu$ m. *B*, TLR4 (826tr)-EGFP with or without coexpression of MD-2 were tagged by biotinylation of the cell surface proteins and affinity-purified. TLR4 was visualized by immunoblotting using an anti-GFP monoclonal antibody. Samples from both combinations of DNAs were prepared from the same number of cells.

perinuclear area. These findings were consistent with observations by others (14, 15). TLR4 is reported to localize in the Golgi apparatus as well as on the plasma membrane. Our observation of TLR4-EGFP accumulation in the perinuclear area does not contradict the report that TLR4 partly localizes in the Golgi apparatus (14).

TLR4-EGFP truncation mutants, 815tr, 802tr, 788tr, and 766tr apparently did not localize at the plasma membrane. No particular fluorescence pattern that might be characteristic of localization to a specific intracellular compartment was observed. Only TLR4 (826tr)-EGFP, which has the shortest truncation, was expressed on the plasma membrane and in the perinuclear area, and the fluorescence pattern was similar to that of wild-type (Fig. 3A). No TLR4 genotypes, including wild-type TLR4-EGFP, clearly localized on the plasma membrane in the absence of MD-2 (Fig. 3A). MD-2 is reported to be necessary for localization of wild-type TLR4 at the plasma membrane (15), which is consistent with our observation. Intracellular distribution of mutant TLR4 varied depending on the genotype, but no particular cellular structure was identified as an alternative target site. Furthermore, we examined the plasma membrane expression of TLR4 (826tr)-EGFP by cell surface protein biotinylation. The expression level of TLR4 (826tr)-EGFP was markedly decreased without coexpression of MD-2 (Fig. 3B), which is compatible with the microscope observation.

Removal of the C-terminal segment of TLR4 at residue 826 does not qualitatively affect LPS responsiveness and subcellular distribution. However, when more residues, up to position 815, were removed, both signal transduction and plasma membrane localization were impaired. These results suggest that residues 815–826 of TLR4 contain at least one segment that is critical for those functions.

**Amino Acid Sequence Replacement Analysis**—To identify critical amino acid sequences in this region, we generated an amino acid replacement mutant of TLR4 instead of truncation mutants. As shown in Fig. 1, although it is not a canonical sequence, leucine-leucine at 815–816 partially fits a known sorting signal motif, a dileucine motif, (D/E)XXXL(L/I) or DXXLL, which plays an important role in internalization of plasma membrane protein or sorting from the *trans*-Golgi network (10). Thus, as has been done in a similar study (16), a mutant was generated in which alanines were substituted for both leucines at positions 815 and 816.

We measured the NF- $\kappa$ B activity of TLR4 (L815A/L816A)-EGFP, the mutant in which both leucines were replaced with alanines, under LPS stimulation (Fig. 4A). This mutant protein did not respond to LPS stimuli. Microscopic observation revealed that TLR4 (L815A/L816A)-EGFP was not expressed on the plasma membrane regardless of whether MD-2 was cotransfected (Fig. 4B). The phenotype of this doubly substituted mutant appeared to be the same as that of the truncation mutants. These results imply that the leucines in positions 815 and 816 play an important role in TLR4 plasma membrane localization.

**Analysis of Single Amino Acid Substitution Mutants**—As previously mentioned, the amino acid sequence leucine-leucine at positions 815 and 816 does not completely match the dileucine motif, *i.e.* it lacks a preceding acidic amino acid. Therefore it




FIGURE 4. Leucines at positions 815–816 of TLR4 are responsible for impairment of LPS responsiveness and plasma membrane expression. A, the LPS stimulation assay was done for TLR4 (L815A/L816A)-EGFP as in Fig. 2. The data were from three independent experiments. Small bars indicate 95% confidence intervals of the mean (*p* value for \* are: TLR4 (WT)-EGFP/MD-2 (+), *p* = 0.002). B, TLR4 (L815A/L816A)-EGFP expression in HEK293T cells was observed by laser confocal microscopy. Bar, 20  $\mu$ m.



FIGURE 5. Alignment of the cytoplasmic domain of EGFP fusion TLR4 amino acid-replacement mutants used in this study. TLR4 (L813A) signifies a mutant with leucine replaced with alanine at position 813. Others are named in the same manner. The amino acids are colored as in Fig. 1. All amino acids are designated using the single-letter code.

was reasonable to explore whether leucines 815 and 816 need to be adjacent to each other. We created five genotypes of single amino acid mutants of TLR4: TLR4 (K813A)-EGFP, TLR4 (L815A)-EGFP, TLR4 (L816A)-EGFP, and TLR4 (D817A)-EGFP. We excluded the amino acid at position 814 from the analysis, because the amino acid in position 814 of wild-type TLR4 is alanine. The amino acid sequence alignment of wild-type TLR4 and the single amino acid replacement mutants is shown in Fig. 5. DNA sequences were confirmed by sequencing.


As was done with truncation mutants, we measured NF- $\kappa$ B activity of wild-type TLR4-EGFP, TLR4 (K813A)-EGFP, TLR4 (L815A)-EGFP, TLR4 (L816A)-EGFP, and TLR4 (D817A)-EGFP in response to LPS stimulation. All mutants except TLR4 (L815A)-EGFP showed responsiveness to LPS stimulation with coexpression of MD-2 (Fig. 6A). Without MD-2, no genotype of TLR4-EGFP responded to LPS stimulation (Fig. 6B). LPS stimulation was performed in an identical manner as with truncation mutants.

We analyzed the subcellular distribution of single amino acid mutants of TLR4-EGFP with and without MD-2 coexpression by fluorescence microscopy. TLR4 (K813A)-EGFP and TLR4 (D817A)-EGFP showed a similar fluorescence pattern to the wild-type, which localized at the plasma membrane when coexpressed with MD-2. No genotypes of TLR4-EGFP localized on the plasma membrane without MD-2 (Fig. 7). The cells transfected with TLR4 (L815A)-EGFP coexpressed with MD-2 did not show plasma membrane fluorescent pattern. Also, TLR4 (L815A)-EGFP showed comparatively weaker fluorescence than other mutants, possibly due to lower expression of the protein. Fluorescence of TLR4 (L816A)-EGFP with MD-2 was ambiguous as for the plasma membrane expression. Some kind of membranous structure was observed in the cytoplasmic area, but the intensity of the plasma membrane green fluorescence was obscure. Together with the results from the LPS stimulation experiment, the leucines at positions 815 and 816 are considered to play important roles in signal transduction and/or subcellular distribution of TLR4.

Because EGFP consists of 239 amino acids, which is about one-third the size of the complete TLR4 protein, the experimental results obtained using TLR4-EGFP could have been influenced by the presence of the EGFP fused at the C terminus of TLR4. To rule out this possibility, we tested the functional integrity of both TLR4 (L815A) and TLR4 (L816A) with and without EGFP at the C terminus. Reporter assays were performed under the same conditions except that the shorter tag, FLAG-His<sub>6</sub>, which has only 21-amino acid tags at the C terminus, was fused to TLR4 in place of EGFP. There was no difference

was obscure. Together with the results from the LPS stimulation experiment, the leucines at positions 815 and 816 are considered to play important roles in signal transduction and/or subcellular distribution of TLR4.

## An Important Amino Acid of TLR4 for Its Function




**FIGURE 6. Leucine at position 815 of TLR4 is pivotal for LPS responsiveness as measured by NF- $\kappa$ B luciferase assay.** A, HEK293T cells were transfected with single amino acid replacement mutants of the human TLR4-EGFP fusion protein plasmid, human MD-2 plasmid, and luciferase reporter and control plasmids. After 36 h, cells were stimulated with LPS (10 ng/ml) for 7 h, and luciferase reporter gene activity was measured. B, instead of MD-2, an empty vector was cotransfected with TLR4-EGFP plasmid and reporter assay vectors. LPS stimulation was done as in A. All results were expressed in the ratio of relative luciferase activity with LPS stimulation to that without the stimulation as in Fig. 2. The data were from three independent experiments. Small bars indicate 95% confidence intervals of the mean ( $p$  values for \* are: TLR4 (WT)-EGFP/MD-2 (+),  $p = 0.002$ ; TLR4 (K813A)-EGFP/MD-2 (+),  $p = 0.000$ ; TLR4 (L816A)-EGFP/MD-2 (+),  $p = 0.018$ ; and TLR4 (D817A)-EGFP/MD-2 (+),  $p = 0.007$ ).

between EGFP-tagged proteins and FLAG-His<sub>6</sub>-tagged proteins in the relative pattern of responsiveness against LPS stimulation (Fig. 8A). Because CD14 is also important for LPS recognition by TLR4, we examined the effect of CD14 coexpression on the phenotypic changes of the mutants (17, 18). Coexpression of CD14 did not change the phenotypes of wild-type TLR4, TLR4 (L815A), and TLR4 (L816A) in terms of LPS responsiveness (data not shown).

Cell surface expressions of the wild-type, L815A mutant, and L816A mutant TLR4-FLAG-His<sub>6</sub> fusion proteins were also examined. Live cells transfected with wild-type TLR4, the L815A mutant or the L816A mutant as well as human MD-2 and CD14 were biotinylated on the cell surface, and the biotinylated proteins were affinity-purified and subjected to Western blotting. Fig. 8B shows the marked difference in cell surface expression of wild-type and mutants L815A and L816A. Note that biotinylated proteins have additional residues on every amine of the extracellular domain, which leads to a band shift during electrophoresis. Although both mutants were detected far less than the wild-type on the cell surface, comparatively more L816A mutant was expressed on the plasma membrane than L815A mutant, and the amount of L815A mutant seemed to be negligible compared with the wild type. These results may clarify the ambiguity of the microscopic observation of TLR4 (L815A) and TLR4 (L816A). Plasma membrane expression of TLR4 was impaired when the leucine at 815 or 816 was replaced to alanine. But the leucine at 815 is more critical, and the mutant L816A may show the weaker phenotypic change.

To further investigate the characteristics of the TLR4 (L815A) mutant, we performed an immunoprecipitation assay of wild-type and mutant TLR4. Cells were transfected with a human MD-2-FLAG-His<sub>6</sub> expression vector and either the wild-type or the mutant (L815A) TLR4-EGFP expression vector. Anti-TLR4 monoclonal antibody (clone HTA125), anti-GFP polyclonal antibody, or anti-FLAG monoclonal antibody



**FIGURE 7. Leucines at the position 815 and 816 of TLR4 are responsible for full plasma membrane expression.** Cells were cultured on coverslips in 12-well plates and transfected as in Fig. 2. EGFP-tagged TLR4 was visualized by laser confocal microscopy. Each genotype of TLR4-EGFP was cotransfected with human MD-2 plasmid or empty vector. Bar, 20  $\mu$ m.

was added to the lysate and precipitated with Protein G-Sepharose beads. Collected proteins were eluted and subjected to Western blotting. The results are shown in Fig. 8C. TLR4 (L815A) was not immunoprecipitated with anti-TLR4 antibody (HTA125). HTA125 antibody was raised against TLR4-expressing cells (9) and recognizes the extracellular portion of TLR4. This result suggests that the amino acid replacement at position 815 may cause a change in the extracellular portion of TLR4 and/or that the replacement may also inhibit cell surface expression of the mutant protein. On the other hand, both wild-type TLR4-EGFP and mutant TLR4-EGFP were immunoprecipitated with anti-GFP polyclonal antibody, which recognized EGFP. However, of the two bands of TLR4, the heavier band seems to be somewhat faint in the mutant, whereas in the wild type the heavier band is at least as dense as the lighter one. TLR4 can be detected as two separate bands in a Western blot (19), especially under transient transfection conditions. The difference in proportion of the heavy and light bands between wild-type and mutant TLR4 may suggest that there is some difference in glycosylation. Furthermore, wild-type TLR4 was coprecipitated with MD-2-FLAG-His<sub>6</sub>, but the mutant TLR4 could not be detected (Fig. 8C, lanes 4 and 8). Because MD-2 is



## An Important Amino Acid of TLR4 for Its Function

associated with TLR4 (9), it is logical to expect that immunoprecipitating MD-2-FLAG-His<sub>6</sub> with anti-FLAG antibody should cause TLR4 to be coprecipitated with it. It is suggested by the result here that the association of the TLR4 mutant with MD-2 is impaired.

### DISCUSSION


In this research, we performed mutagenesis analyses of particular amino acid residues in TLR4 to explore the mechanisms of TLR4 intracellular signal transduction and subcellular distribution. We found the candidate residues by analyzing truncation mutants of TLR4 in the cytoplasmic region, in which both signaling and normal subcellular distribution of TLR4 are disturbed. Because we are focusing on a common mechanism for the impaired signaling and distribution, we finally picked a single amino acid mutant that does not respond to LPS stimuli, as measured with NF- $\kappa$ B reporter luciferase assay, and one that does not localize on the plasma membrane. TLR4 (L815A) is a mutant that meets these conditions, and our results suggest that the leucine at position 815 of TLR4 is required for both signal transduction and plasma membrane localization.

The best known single amino acid mutant of TLR4 is TLR4 (P712H) known as the *Lps<sup>d</sup>* mutation in the C3H/HeJ mouse, which corresponds to position 714 in this study of human TLR4 (5, 6, 20). Mice carrying this mutation opened up the rediscovery of TLR4 as a key player in innate immunity. Because this proline residue at this position is within the TIR domain and is conserved among TLRs or TLR4s of other species, it is assumed that the residue plays an important role in TLR4 function. The association of TLR4 (P712H) with its adapter proteins is reported to be intact, and the explanation for the functional impairment of TLR4 (P712H) is not clear (21–23).

Some single amino acid variants are found in humans, and these are related to the incidence or prognosis of some infections and other diseases. A growing body of data suggests that the ability of certain individuals to respond properly to TLR4 ligands may be impaired by single-nucleotide polymorphisms within TLR4 genes (24). The D299G and T399I alleles of the TLR4 gene have been associated with increased risk of severe infections (25).

By clarifying the subcellular component where the mutant protein is retained, or by clarifying to which compartment the mutant is not delivered, the abnormal intracellular sorting that is caused by the mutation in TLR4 (L815A) could be elucidated more precisely. Usually a sorting signal motif is comprised of several amino acids. In this regard, if the leucine at position 815 is a part of a motif, there should be other amino acids that are also members of the motif. Although replacement of leucine with alanine at position 816 did not cause an apparent signal transduction impediment, plasma membrane expression of TLR4 (L816A) was impaired to a certain extent. Positive

respectively, were prepared from the same number of cells as for the biotinylation experiment. C, immunoprecipitation with antibodies further reveals the characteristics of TLR4 (L815A). Anti-TLR4 monoclonal antibody (HTA125) does not precipitate the mutant TLR4, whereas anti-GFP polyclonal antibody (pAb) precipitates both wild-type and mutant TLR4. Mutant TLR4 was not coprecipitated with MD-2-FLAG-His<sub>6</sub>. Lysates were prepared from cells transiently expressing wild-type or mutant TLR4-EGFP and MD-2-FLAG-His<sub>6</sub>.



**FIGURE 8.** A, TLR4 mutants L815A and L816A with and without EGFP fusion exhibit the same phenotypes in LPS responsiveness and plasma membrane expression. HEK293T cells were transfected with the wild-type, the L815A or L816A mutant TLR4flaghis plasmid plus the human MD-2 plasmid and luciferase reporter, or control plasmids. After 36 h, cells were stimulated with LPS (10 ng/ml) for 7 h, and luciferase reporter gene activity was measured. The data were from three independent experiments. Small bars indicate 95% confidence intervals of the mean (*p* values for \* are: TLR4 (WT) flaghis/MD-2 (+), *p* = 0.046; TLR4 (L816A) flaghis/MD-2 (+), *p* = 0.003). N.S.: not significant. B, wild-type and mutant TLR4s L815A and L816A were tagged by biotinylation of the cell surface proteins and affinity-purified. Human MD-2 and CD14 were coexpressed. TLR4 was visualized by immunoblotting using an anti-FLAG monoclonal antibody (mAb). Faint bands below 100 kDa are considered to be unbiotinylated intracellular TLR4 proteins that were not washed off during the process. Samples from TLR4 (WT), TLR4 (L815A), and TLR4 (L816A),

## An Important Amino Acid of TLR4 for Its Function

response to LPS stimulation by TLR4(L816A) could be attributable to this small amount of expression on the plasma membrane. Mutagenesis analyses of neighboring amino acids of the leucine at 815 were not definitive, but the results could be suggestive that the adjacent leucine at 816 may work together with the leucine at 815. Leucines at position 815 and 816 could be in the same motif, and the leucine at position 816 may be less critical.

Several proteins have been reported to be involved in TLR4 cell surface expression. Heat shock protein gp96 is necessary for TLR4 association with MD-2 in the ER and for subsequent cell surface expression (26). PRAT4A and PRAT4B are associated with TLR4 and regulate TLR4 cell surface expression (27, 28). In embryonic fibroblasts of MD-2 knockout mice, TLR4 localization on the cell surface is severely impaired, and most TLR4 is retained in the ER or Golgi apparatus (15). MD-2 binds to TLR4 at its extracellular domain and is essential for LPS recognition by TLR4 (29). Although proteins such as CD14 and LPS-binding protein are reported to have important roles in LPS recognition by TLR4, in an *in vitro* setting HEK293T cells gain LPS responsiveness by introducing only TLR4 and MD-2 genes when measured by NF- $\kappa$ B reporter assay (9, 30). Without transfection, HEK293 cells do not express TLR4, MD-2, or CD14, which are involved in LPS-induced intracellular signaling (31, 32). In this study, we show that the association of the TLR4 mutant and MD-2 is impaired (Fig. 8C).

Post-translational modification is another important factor for TLR4 function. Asparagine residues in the extracellular portion of TLR4 need to be glycosylated for plasma membrane expression of TLR4 (15, 19, 33). TLR4:MD-2 association is necessary for this glycosylation as well. The difference in the proportion of the heavy band to lighter band between wild-type and L815A mutant TLR4 immunoprecipitated with anti-GFP polyclonal antibody suggests that there may be some difference in glycosylation between wild-type and L815A mutant TLR4 (Fig. 8C). Although leucine at position 815 is located in the cytoplasmic tail of TLR4, we speculated that substitution of leucine at position 815 may cause a conformational change in the extracellular portion of the protein, which may interfere with the association between L815A mutant TLR4 and MD-2, leading to inhibition of glycosylation and cell surface expression of the mutant protein. Further investigation may reveal the mechanism involved in this phenotypic change in TLR4 (L815A), which would lead to better understanding of the mechanism of wild-type TLR4 signaling and trafficking.

**Acknowledgment**—We greatly appreciate the gift of human TLR4 and MD-2 cDNA from Dr. Kensuke Miyake (Institute of Medical Science, University of Tokyo, Japan).

### REFERENCES

- Takeda, K., Kaisho, T., and Akira, S. (2003) *Annu. Rev. Immunol.* **21**, 335–376
- Hoebe, K., Du, X., Georgel, P., Janssen, E., Tabet, K., Kim, S. O., Goode, J., Lin, P., Mann, N., Mudd, S., Crozat, K., Sovath, S., Han, J., and Beutler, B. (2003) *Nature* **424**, 743–748
- Oshiumi, H., Sasai, M., Shida, K., Fujita, T., Matsumoto, M., and Seya, T. (2003) *J. Biol. Chem.* **278**, 49751–49762
- Yamamoto, M., Sato, S., Mori, K., Hoshino, K., Takeuchi, O., Takeda, K., and Akira, S. (2002) *J. Immunol.* **169**, 6668–6672
- Qureshi, S. T., Lariviere, L., Leveque, G., Clermont, S., Moore, K. J., Gros, P., and Malo, D. (1999) *J. Exp. Med.* **189**, 615–625
- Poltorak, A., He, X., Smirnova, I., Liu, M. Y., Van Huffel, C., Du, X., Birdwell, D., Alejos, E., Silva, M., Galanos, C., Freudenberg, M., Ricciardi-Castagnoli, P., Layton, B., and Beutler, B. (1998) *Science* **282**, 2085–2088
- Visintin, A., Mazzoni, A., Spitzer, J. A., and Segal, D. M. (2001) *Proc. Natl. Acad. Sci. U. S. A.* **98**, 12156–12161
- Nishitani, C., Mitsuzawa, H., Hyakushima, N., Sano, H., Matsushima, N., and Kuroki, Y. (2005) *Biochem. Biophys. Res. Commun.* **328**, 586–590
- Shimazu, R., Akashi, S., Ogata, H., Nagai, Y., Fukudome, K., Miyake, K., and Kimoto, M. (1999) *J. Exp. Med.* **189**, 1777–1782
- Bonifacio, J. S., and Traub, L. M. (2003) *Annu. Rev. Biochem.* **72**, 395–447
- Nishimura, N., and Balch, W. E. (1997) *Science* **277**, 556–558
- Nufer, O., and Hauri, H. P. (2003) *Curr. Biol.* **13**, R391–393
- Slack, J. L., Schooley, K., Bonnert, T. P., Mitcham, J. L., Qwarnstrom, E. E., Sims, J. E., and Dower, S. K. (2000) *J. Biol. Chem.* **275**, 4670–4678
- Latz, E., Visintin, A., Lien, E., Fitzgerald, K. A., Monks, B. G., Kurt-Jones, E. A., Golenbock, D. T., and Espevik, T. (2002) *J. Biol. Chem.* **277**, 47834–47843
- Nagai, Y., Akashi, S., Nagafuku, M., Ogata, M., Iwakura, Y., Akira, S., Kitamura, T., Kosugi, A., Kimoto, M., and Miyake, K. (2002) *Nat. Immunol.* **3**, 667–672
- Hein, C., and Andre, B. (1997) *Mol. Microbiol.* **24**, 607–616
- Beutler, B. (2000) *Curr. Opin. Immunol.* **12**, 20–26
- Akashi, S., Ogata, H., Kirikae, F., Kirikae, T., Kawasaki, K., Nishijima, M., Shimazu, R., Nagai, Y., Fukudome, K., Kimoto, M., and Miyake, K. (2000) *Biochem. Biophys. Res. Commun.* **268**, 172–177
- Ohnishi, T., Muroi, M., and Tanamoto, K. (2003) *Clin. Diagn. Lab. Immunol.* **10**, 405–410
- Xu, Y., Tao, X., Shen, B., Horng, T., Medzhitov, R., Manley, J. L., and Tong, L. (2000) *Nature* **408**, 111–115
- Dunne, A., Ejdeback, M., Ludidi, P. L., O'Neill, L. A., and Gay, N. J. (2003) *J. Biol. Chem.* **278**, 41443–41451
- Fitzgerald, K. A., Palsson-McDermott, E. M., Bowie, A. G., Jefferies, C. A., Mansell, A. S., Brady, G., Brint, E., Dunne, A., Gray, P., Harte, M. T., McMurray, D., Smith, D. E., Sims, J. E., Bird, T. A., and O'Neill, L. A. (2001) *Nature* **413**, 78–83
- Horng, T., Barton, G. M., and Medzhitov, R. (2001) *Nat. Immunol.* **2**, 835–841
- Schroder, N. W., and Schumann, R. R. (2005) *Lancet Infect. Dis.* **5**, 156–164
- Agnese, D. M., Calvano, J. E., Hahn, S. J., Coyle, S. M., Corbett, S. A., Calvano, S. E., and Lowry, S. F. (2002) *J. Infect. Dis.* **186**, 1522–1525
- Random, F., and Seed, B. (2001) *Nat. Cell Biol.* **3**, 891–896
- Wakabayashi, Y., Kobayashi, M., Akashi-Takamura, S., Tanimura, N., Konno, K., Takahashi, K., Ishii, T., Mizutani, T., Iba, H., Kouro, T., Takaki, S., Takatsu, K., Oda, Y., Ishihama, Y., Saitoh, S., and Miyake, K. (2006) *J. Immunol.* **177**, 1772–1779
- Konno, K., Wakabayashi, Y., Akashi-Takamura, S., Ishii, T., Kobayashi, M., Takahashi, K., Kusumoto, Y., Saitoh, S., Yoshizawa, Y., and Miyake, K. (2006) *Biochem. Biophys. Res. Commun.* **339**, 1076–1082
- Nishitani, C., Mitsuzawa, H., Sano, H., Shimizu, T., Matsushima, N., and Kuroki, Y. (2006) *J. Biol. Chem.* **281**, 38322–38329
- Akashi, S., Shimazu, R., Ogata, H., Nagai, Y., Takeda, K., Kimoto, M., and Miyake, K. (2000) *J. Immunol.* **164**, 3471–3475
- Espevik, T., Latz, E., Lien, E., Monks, B., and Golenbock, D. T. (2003) *Scand. J. Infect. Dis.* **35**, 660–664
- Muta, T., and Takeshige, K. (2001) *Eur. J. Biochem.* **268**, 4580–4589
- da Silva Correia, J., and Ulevitch, R. J. (2002) *J. Biol. Chem.* **277**, 1845–1854



ELSEVIER

Contents lists available at ScienceDirect

## Atherosclerosis

journal homepage: [www.elsevier.com/locate/atherosclerosis](http://www.elsevier.com/locate/atherosclerosis)

## Association between metabolic syndrome and carotid atherosclerosis in individuals without diabetes based on the oral glucose tolerance test

Nobukazu Ishizaka<sup>a,\*</sup>, Yuko Ishizaka<sup>b</sup>, Minoru Yamakado<sup>b</sup>, Eiichi Toda<sup>b</sup>, Kazuhiko Koike<sup>c</sup>, Ryozi Nagai<sup>a</sup>

<sup>a</sup> Department of Cardiovascular Medicine, University of Tokyo, Graduate School of Medicine, Hongo 7-3-1 Bunkyo-ku, Tokyo 113-8655, Japan

<sup>b</sup> Center for Multiphasic Health Testing and Services, Mitsui Memorial Hospital, Tokyo, Japan

<sup>c</sup> Department of Infectious Diseases, University of Tokyo, Graduate School of Medicine, Tokyo, Japan

## ARTICLE INFO

## Article history:

Received 31 July 2008

Received in revised form 20 October 2008

Accepted 21 October 2008

Available online 30 October 2008

## Keywords:

Metabolic syndrome

Carotid artery

Atherosclerosis

Risk factors

Glucose metabolism

## ABSTRACT

**Introduction:** Whether or not metabolic syndrome is predictive of atherosclerotic disorders may depend on the population studied. We investigated whether metabolic syndrome is associated with carotid atherosclerosis in individuals who were shown not to have diabetes mellitus based on results of the 75-g oral glucose tolerance test (OGTT).

**Methods and results:** Between 1994 and 2003, 3904 individuals underwent general health screening that included the OGTT. Among these 3904 individuals, 3679 had a fasting plasma glucose of <126 mg/dL (subgroup 1), and 3488 had a 2-h post-OGTT glucose value of <200 mg/dL (subgroup 2). In both subgroups, metabolic syndrome was found to be a risk factor for carotid plaque and for carotid intima-media thickening in men, and tended to be a risk factor for carotid plaque in women after adjustment for age. Among 3473 individuals who had both a fasting plasma glucose value of <126 mg/dL and a 2-h post-OGTT glucose of <200 mg/dL, 2440 did not have hypertension, which was defined as systolic and diastolic blood pressure of <140/90 mmHg and absence of use of anti-hypertensive medication. In these non-diabetic non-hypertensive individuals, the association between metabolic syndrome and carotid plaque or carotid intima-media thickening was not statistically significant even with adjustment only for age.

**Conclusions:** In men who did not have impaired fasting glycemia and/or in those without impaired glucose tolerance, metabolic syndrome was a predictor of carotid atherosclerosis after age adjustment, although metabolic syndrome was not found to be a predictor of carotid atherosclerosis when hypertensive individuals were excluded from the study population.

© 2008 Elsevier Ireland Ltd. All rights reserved.

### 1. Introduction

Metabolic syndrome (MetS) is a cluster of metabolic and hemodynamic abnormalities linked with insulin resistance. Since components of MetS also represent risk factors for atherosclerotic disorders, it is natural that individuals with this syndrome have an increased risk for ischemic heart disease [1] and stroke [2,3]. On the other hand, the clinical utility of MetS may depend on whether the risk conveyed by this syndrome is higher than the sum of each component utilized as diagnostic criteria for MetS [4,5].

Carotid artery intima-media thickness has been reported to be a discriminator as a surrogate of cardiovascular mortality in community-dwelling Japanese people [6] and, conversely, aggre-

gation of established major coronary risk factors has been reported to strongly influence the presence of carotid atherogenesis in the general Japanese population [7]. Previously, we reported that the presence of MetS may not increase the risk for carotid atherosclerosis in individuals without hypertension, with hypertension defined as systolic blood pressure (SBP) of  $\geq 140$  mmHg, diastolic blood pressure (DBP) of  $\geq 90$  mmHg, or the use of anti-hypertensive medication [8]. This observation suggested that the properties of MetS that present a risk for atherosclerotic diseases may differ according to the populations selected. Consistent with this idea, it was reported that MetS was not found to be associated with cardiovascular mortality in non-diabetic non-hypertensive Chinese individuals [9], and that MetS did not significantly increase the risk of mortality from cardiovascular disease in non-diabetic Mexican Americans and non-Hispanic whites [10]. In the current study, we investigated whether MetS was associated with carotid atherosclerosis in Japanese individuals who did not have diabetes mellitus based on results of the 75-g oral glucose tolerance test (OGTT).

\* Corresponding author. Tel.: +81 3 3815 5411x37156; fax: +81 3 5842 5586.

E-mail address: [nobuishizka-tky@umin.ac.jp](mailto:nobuishizka-tky@umin.ac.jp) (N. Ishizaka).




Fig. 1. Flow chart showing selection of the four subgroups.

## 2. Methods

### 2.1. Study subjects and selection of subgroups

The study was approved by The Ethical Committee of Mitsui Memorial Hospital and University of Tokyo, Faculty of Medicine. Between September 1994 and December 2003, 8143 subjects underwent general health screening including carotid ultrasonography at the Center for Multiphasic Health Testing and Services, Mitsui Memorial Hospital. Of the 8143 subjects, 95 were treated as having diabetes, and of the remaining 8048 individuals, 3904 underwent an OGTT. Among these 3904 individuals, three subgroups were sequentially selected based on various parameters (Fig. 1). Those with a fasting plasma glucose (FPG) value of  $<126$  mg/dL were designated as subgroup 1, and those with a 2-h post-OGTT plasma glucose (2-h PG) value of  $<200$  mg/dL were designated as subgroup 2. Subgroup 3 was comprised of subjects who met all the following conditions: FPG of  $<126$  mg/dL, 2-h PG of  $<200$  mg/dL, and not having hypertension. Hypertension was defined as SBP  $\geq 140$  mmHg, DBP  $\geq 90$  mmHg, or the use of anti-hypertensive medication. We also selected individuals without impaired glucose tolerance (IGT), i.e., individuals with a 2-h PG value of  $<140$  mg/dL.

At our institute, several types of health screening programs are available, and some general health screening programs include carotid ultrasonography and/or OGTT, while others do not. However, the decision on the type of health screening was made by the individuals and/or their companies and was not decided upon or recommended by any attending physician.

### 2.2. Definition of MetS

MetS was defined as the presence of three or more of the following: (1) fasting glucose  $\geq 110$  mg/dL; (2) SBP/DBP  $\geq 130/85$  mmHg or taking anti-hypertensive medication; (3) triglycerides  $\geq 150$  mg/dL mmol/L; (4) HDL cholesterol  $<40$  mg/dL in men and  $<50$  mg/dL in women; and (5) body mass index  $\geq 25$  kg/m<sup>2</sup> [11].

### 2.3. Carotid ultrasonography

Carotid artery status was studied using high resolution B-mode ultrasonography (Sonolayer SSA270A, Toshiba, Japan) equipped with a 7.5 MHz transducer as described previously [12]. Plaque was defined to be present when there is one or more clearly isolated focal thickening(s) of the intima-media layer with thickness of  $\geq 1.3$  mm at the common or internal carotid artery or the carotid bulb. Carotid wall intima-media thickening was said to be present when intima-media thickness which was measured at the far wall of the distal 10 mm of the common carotid artery was  $\geq 1.0$  mm [12].

### 2.4. Statistical analysis

Logistic regression analysis was used to obtain adjusted odds ratios and their 95% confidence intervals (CIs) to predict the presence of carotid plaque or carotid intima-media thickening. Statistical analyses were carried out by using Dr. SPSS II (SPSS Inc., Chicago, IL). Results are expressed as the mean  $\pm$  standard deviation (SD). A value of  $p < 0.05$  was taken to be statistically significant.