

needle with an internal diameter of 2 mm (Tohoku University style, Kakinuma Factory, Tokyo, Japan), fixed in 10% formalin, and stained with hematoxylin-eosin, Masson's trichrome, silver impregnation, and periodic acid-Schiff after diastase digestion. The size of specimens for examination was more than six portal areas. Histopathological interpretations of specimens were made independently by experienced liver pathologists (YA and HK) who had no clinical information. Baseline liver histology of chronic hepatitis prior to IFN therapy was classified according to the extent of fibrosis, into three stages: mild, periportal expansion; moderate, portoportal septa; and severe, portocentral linkage, or bridging fibrosis [Desmet et al., 1994]. Patients with severe fibrosis were considered as pre-cirrhosis or cirrhosis.

Statistical Analysis

Baseline characteristics and treatment differences of both groups were analyzed using Fisher's exact test or Kruskal Wallis test. HCC appearance rates were analyzed by the log rank test. A Cox proportional hazards model was used to analyze the factors contributing to the rate of development of HCC: factors examined included age, gender, histologic findings, HCV genotype, HCV load, aspartate aminotransferase (AST), ALT, and AFP. A *P*-value of <0.05 were considered statistically significant. The SPSS software package (SPSS 11.0 for windows; SPSS, Inc., Chicago, IL) was used for analyses.

RESULTS

Pretreatment Clinical Characteristics

Table I shows the characteristics of the patients with and without IFN treatment. There were no significant differences between the two groups with regard to sex ratio, age, histopathological stage of the liver, serum HCV-RNA level, and AST, ALT, AFP, and blood cell counts. In the control group, eighty-two patients started IFN during follow-up.

Changes in Serum AST, ALT, and AFP Activity After Treatment

Figure 1 shows the serum AST and ALT levels after initiation of treatment. The serum AST and ALT levels declined to normal levels after initiation of treatment with IFN. Transaminase levels at 6, 12, 18, and 24 months after in the IFN group were lower than that of patients not treated with IFN group with statistical significance. Figure 2 shows change in serum AFP level after initiation of treatment. Serum AFP level decreased after the initiation of IFN therapy compared the untreated group with statistical significance.

Loss of HCV-RNA

Of 120 patients treated with IFN, 18 patients lost serum HCV-RNA during IFN treatment. Of the 96 patients who stopped IFN therapy, 8 patients lost HCV-RNA in the serum 6 month after the termination of IFN treatment.

Cumulative Rates of HCC

HCC was diagnosed in four patients in the IFN group and 38 in patients not treated with IFN. Figure 3 shows the cumulative HCC development rates in both groups. The 5- and 10-year cumulative rates of HCC were 5.9 and 13.7%, and 17.1 and 32.8%, for IFN- and no-IFN groups, respectively. The cumulative rate of development of HCC in the IFN group was significantly lower than that not treated with IFN (*P* = 0.045).

Risk Factors for the Development of HCC

The rate of development of HCC after initiation of treatment, Cox regression analysis was performed using several variables. Univariate analysis showed that the following four factors affected significantly the cumulative development of HCC in all patients: histopathological staging (*P* < 0.0001), serum AFP level at 0.5 year after IFN therapy (*P* = 0.005), sex (*P* = 0.006), and IFN therapy (*P* = 0.045) (Table II). The variables were correlated mutually and multivariate Cox regression

TABLE I. Clinical Profiles Before Treatment

Characteristic	IFN group	Non-IFN group	<i>P</i> -value
N ^a	120	240	
Sex (M/F)	65/55	130/110	1
Age (years) ^b	63(60-75)	63(60-75)	1
Liver fibrosis (mild/moderate/severe) ^a	36/40/44	82/70/88	0.965
Genotype (1/2) ^a	85/31	170/61	0.782
HCV-RNA (KIU/ml) ^b	680(10-5,000)	720(5-5,000)	0.176
AST (IU/L) ^b	71(26-446)	67(20-355)	0.493
ALT (IU/L) ^b	86(38-699)	78(46-374)	0.101
AFP (ng/ml) ^b	10(3-316)	9(2-190)	0.342
Hemoglobin (g/dl) ^b	13.7(10.9-17.0)	14.0(11.8-17.0)	0.167
Platelet ($\times 10^4/\text{mm}^3$) ^b	12.8(5.2-25.6)	13.1(5.6-23.2)	0.275
WBC ($\times 10^3/\text{mm}^3$) ^b	4.0(2.1-11.3)	4.0(2.6-7.9)	0.570

AFP, alpha-fetoprotein; ALT, alanine aminotransferase; AST, aspartate aminotransferase; HCV, hepatitis C virus; WBC, white blood count; Normal reference ranges ≤ 10 ng/ml for AFP, 6-50 IU/L for ALT, 11-38 IU/L for AST.

^aData are number of patients.

^bData are median (range).




Fig. 1. Changes in serum AST (panel A) and ALT (panel B) after initiation of treatment. Data were expressed as mean ± standard deviation. *Year after initiation of treatment, †differences of both group by Kruskal Wallis test.

analysis was carried out with the four statistically significant variables in the model (Table III). The development of HCC occurred significant by when: (1) histological staging was advanced, (2) serum AFP level after the initiation of treatment was >10 ng/ml, and (3) IFN was not given. The relative risk of HCC in patients of the IFN group was 0.3 times of that in patients of no-IFN-group (Fig. 3). The relative risk for the development of HCC in patients with severe fibrosis was 3.9 compared to patients with mild or moderate fibrosis. Figure 4 shows the rate of development of HCC based on the difference of treatment and histological staging. IFN therapy could reduce significantly the development of HCC in severe fibrosis (Figure 4, Panel B).

Safety and Tolerance of IFN

Of the 120 patients included in this study, 9 discontinued IFN therapy because of adverse events: 3 cases of general fatigue, 2 cases of psychiatric disorder, 2 cases of aggravation of diabetes mellitus, 1 patient each with thrombocytopenia, pneumonia, and Parkinson's syndrome. The cumulative dropout rate because of IFN-related side effects is plotted in Figure 5. The onset of IFN-related side effects ranged from 204 to 1,569 days after initiation of IFN therapy. These side effects in nine patients disappeared 1 month after cessation of IFN therapy. None of the other patients developed serious side effects that required discontinuation of IFN.




Fig. 2. Changes in serum AFP after initiation of treatment. *Year after initiation of treatment, †differences between IFN group and no-IFN group by the Kruskal Wallis test.




Fig. 3. Cumulative rate of development of hepatocellular carcinoma (HCC) based on the difference in treatment.

TABLE II. Factors Associated With the Development of HCC After Initiation of Treatment by Univariate Cox Regression Analysis

Factor	Category	Odds ratio	95% confidence interval	P-value
AFP after treatment (ng/ml) ^a	≤10	1	2.34–7.00	<0.0001
	>10	4.05		
Liver fibrosis	Mild or moderate	1	2.86–7.24	<0.0001
	Severe	4.55		
Sex	Male	1	0.34–0.84	0.006
	Female	0.532		
IFN therapy	–	1	0.13–0.98	0.045
	+	0.35		
AFP (ng/ml) ^b	≤10	1	0.45–36.90	0.209
	>10	2.23		
ALT after treatment ^a	≤50	1	0.53–6.29	0.342
	>50	1.82		
ALT (IU/L) ^b	≤100	1	0.73–2.34	0.371
	>100	1.31		
Age (years) ^b	≤65	1	0.23–11.78	0.615
	>65	1.65		
HCV-RNA after treatment ^a	–	1	0.29–14.88	0.474
	+	2.06		
BMI ^b	≤25	1	0.08–7.29	0.800
	>25	0.74		
HCV-RNA (KIU/ml) ^b	>100	1	0.05–4.21	0.490
	≤100	0.57		
HCV genotype ^b	1	1	0.45–2.49	0.907
	2	1.05		

AFP, alpha-fetoprotein; ALT, alanine aminotransferase; BMI, body mass index; HCC, hepatocellular carcinoma; IFN, interferon.

^aSerum level 6 months after initiation of treatment.

^bMeasurement at initiation of treatment.

DISCUSSION

Several findings from the present study have direct implications for long-term IFN treatment in aged patients with chronic hepatitis or cirrhosis. First, the AFP baseline was decreased after initiation of IFN therapy in most patients. Second, the cumulative HCC development rate in patients whose serum level of AFP was within normal limits after initiation of IFN therapy was lower than that of patients with high level of AFP despite of IFN therapy. These suggest that AFP is a suitable indicator in long-term IFN therapy for protecting against HCC. If long-term IFN therapy could maintain normalization of serum AFP level, HCC development could be prevented in HCV patients.

AFP is a glycoprotein produced by the liver or yolk sac in fetal life in vertebrates, and it is not normally present in the serum of adults and is used commonly as a tumor marker for HCC [Otsuru et al., 1988]. Many reports have cited elevated AFP baselines as an independent HCC risk factor together with age, gender, liver histology stage, and ethnicity in patients infected with HCV [Ikeda et al., 1993; Tsukuma et al., 1993]. Elevation of AFP has been observed after a rise in transaminase in acute hepatitis, fulminant hepatitis, and acute exacerbation of chronic hepatitis. This type of AFP elevation is explained as a result of hepatocyte regeneration accompanied by necroinflammatory changes [Hu et al., 2004]. On the other hand, Yoshida et al. [2002] have reported that the HCV-coding core

TABLE III. Factors Associated With the Development of HCC After Initiation of Treatment by Multivariate Cox Regression Analysis

Factor	Category	Odds ratio	95% confidence interval	P-value
Liver fibrosis	Mild or moderate	1	1.69–6.13	<0.0001
	Severe	3.89		
AFP after treatment (ng/ml) ^a	≤10	1	1.45–4.84	0.002
	>10	2.65		
IFN therapy	–	1	0.11–0.86	0.025
	+	0.304		

AFP, alpha-fetoprotein; HCC, hepatocellular carcinoma; IFN, interferon.

^aSerum AFP level 6 months after initiation of treatment.




Fig. 4. Cumulative rate of development of hepatocellular carcinoma (HCC) based on the difference of treatment and histological staging (Panel A); patients with mild or moderate fibrosis. Panel B: patients with severe fibrosis.

protein is related to the cell cycle and cell proliferation at the transcriptional level in hepatocytes. This might mean that the HCV-coding core protein upregulate AFP production in hepatocytes.

AFP has the following functions associated with the hepatocarcinogenesis. First, AFP plays a role in the carrier-transport of various ligands and binds to a large variety of ligands such as fatty acids and estrogens. The action of AFP ligands can lead to cells toward multiplication or differentiation [Parmelee et al., 1978; Jacobson et al., 1990; Deutsch, 1991]. Second, AFP could control cell growth. AFP synergies growth factors such as epidermal growth factor and insulin like growth factor to cause proliferation of granulose cells. AFP has been found to regulate the proliferation of human

mammary tumor cells [Wang and Alpert, 1995]. Wang and Xu [1998] have reported that human AFP can enhance the mouse hepatoma H-22 and human hepatoma SNMC-7721 cells in vitro. Third, AFP has been found to have immunosuppressive activity. AFP also suppresses the natural killer cell activity and induces suppressor T cells. AFP prevents the expression of MHC-II class molecule on macrophages [Lester et al., 1976].

Subsequently, Wang et al. [2001] has reported that antisense phosphorothioate oligodeoxyribonucleotide targeted to AFP genes inhibit the growth of human hepatoma cells and solid hepatoma, which is related to their cell apoptosis induction. As described above, elevated AFP might be associated with the hepatocarcinogenesis. Murashima et al. [2006] have reported that the competing action of IFN against HCV-related protein may cause decrease the production of AFP. Thus, normalization of serum AFP level by prolonged-IFN therapy could also protect against the development of HCC.

IFN therapy could reduce significantly the development of HCC in severe fibrosis. Long-term IFN treatment can be associated with serious side effects and is costly. Accordingly, careful selection of patients for long-term IFN therapy is important. The development rate of HCC is high in aged patients with severe liver fibrosis and elevated AFP. On the other hand, the development of HCC is low in no-aged patients with non-severe fibrosis and low level of AFP. Therefore, long-term IFN treatment for protection against HCC could be




Fig. 5. Cumulative dropout rates due to IFN-related side effect.

recommended for patients with elevated AFP level and/or severe liver fibrosis who can tolerate IFN-related side effects. Considering cost-effectiveness, it seems reasonable to select aged patients with elevated AFP and/or severe fibrosis for long-term IFN therapy for protecting against the development of HCC.

ACKNOWLEDGMENTS

The present work was supported in part by grants-in-aid from Okinaka Memorial Institute for Medical Research and Japanese Ministry of Health, Labour And Welfare.

REFERENCES

- Albadalejo J, Alonso R, Antinozzi R, Bogard N, Bourgalet AM, Colucci G, Fenner T, Petersen H, Sala F, Vinceleto J, Young C. 1998. Multicenter evaluation of the COBAS AMPLICOR HCV assay, an integrated PCR system for rapid detection of hepatitis C virus RNA in the diagnostic laboratory. *J Clin Microbiol* 36:862-865.
- Arase Y, Ikeda K, Murashima N, Chayama K, Tsubota A, Koida I, Suzuki Y, Saitoh S, Kobayashi M, Kumada H. 1997. The long term efficacy of glycyrrhizin in chronic hepatitis C patients. *Cancer* 79:1494-1500.
- Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Sheuer PJ. 1994. Classification of chronic hepatitis: Diagnosis, grading, and staging. *Hepatology* 19:1513-1520.
- Deutsch HF. 1991. Chemistry and biology of alpha-fetoprotein. *Adv Cancer Res* 56:253-312.
- Doglio A, Laffont C, Caroli-Bose FX, Rochet P, Lefebvre J. 1999. Second generation of the automated Cobas Amplicor HCV assay improves sensitivity of hepatitis C virus RNA detection and yields results that are more clinically relevant. *J Clin Microbiol* 37:1567-1569.
- Dusheiko G, Schmilovitz-Weiss H, Brown D, McOmish F, Yap PL, Sherlock S, McIntyre N, Simmonds P. 1994. Hepatitis C virus genotypes; an investigation of type-specific differences in geographic origin and disease. *Hepatology* 19:13-18.
- Fogler WE, Volker K, McCormick KL, Watanabe M, Ortaldo JR, Wiltrout RH. 1994. NK cell infiltration into lung, liver, and subcutaneous B16 melanoma is mediated by VCAM-1/VLA-4 interaction. *J Immunol* 196:4707-4714.
- Gutterman JU. 1994. Cytokine therapeutics: Lessons from interferon alpha. *Proc Natl Acad Sci* 91:1198-1205.
- Hagiwara H, Hayashi N, Mita E, Ueda K, Takehara T, Kasahara A, Fusamoto H, Kamada T. 1992. Detection of hepatitis C virus RNA in serum of patients with chronic hepatitis C treated with interferon-alpha. *Hepatology* 15:37-41.
- Harrington DP, Fleming TR. 1983. A class of rank test procedures for censored survival data. *Biomedica* 62:553-566.
- Hu KQ, Kyulo NL, Lim N, Elhazin B, Hillebrand DJ, Bock T. 2004. Clinical significance of elevated alpha-fetoprotein (AFP) in patients with chronic hepatitis C, but not hepatocellular carcinoma. *Am J Gastroenterol* 99:860-865.
- Ikeda K, Saitoh S, Koida I, Arase Y, Tsubota A, Chayama K, Kumada H, Kawanishi M. 1993. A multivariate analysis of risk factors for hepatocellular carcinogenesis: A prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology* 18:47-53.
- Ikeda K, Arase Y, Saitoh S, Kobayashi M, Suzuki Y, Suzuki F, Tsubota A, Chayama K, Murashima N, Kumada H. 2000. Interferon beta prevents recurrence of hepatocellular carcinoma after complete resection or ablation of primary tumor: A prospective randomized study of hepatitis C virus-related liver cancer. *Hepatology* 32:228-232.
- Imai Y, Kawata S, Tamura S, Yabuuchi I, Noda S, Inada M, Maeda Y, Shirai Y, Fukuzaki T, Kaji I, Ishikawa H, Matsuda Y, Nishikawa M, Seki K, Matsuzawa Y. 1998. Relation of interferon therapy and hepatocellular carcinoma in patients with chronic hepatitis C. Osaka Hepatocellular Carcinoma Prevention Study Group. *Ann Intern Med* 129:94-99.
- Jacobson HI, Bennett JA, Mizejewski GJ. 1990. Inhibition of estrogen-dependent breast cancer growth by a reaction product of alpha-fetoprotein and estradiol. *Cancer Res* 50:415-420.
- Lester EP, Miller JB, Baron JM, Yachnin S. 1976. Inhibition of human lymphocyte transformation by human alpha-fetoprotein (HAFF): Studies on the mode of HAFF action and the role of HAFF polymorphism. *Immunology* 34:189-194.
- Lindner DJ. 2002. Interferons as antiangiogenic agents. *Curr Oncol Rep* 4:510-514.
- Lindsay KL, Trepo C, Heintges T, Shiffman ML, Gordon SC, Hoefs JC, Schiff ER, Goodman ZD, Laughlin M, Yao R, Albrecht JK. Hepatitis Interventional Therapy Group. 2001. A randomized, double-blind trial comparing pegylated interferon alfa-2b to interferon alfa-2b as initial treatment for chronic hepatitis C. *Hepatology* 34:395-403.
- Lu CY, Changelian PS, Unanue ER. 1984. Alpha-fetoprotein inhibits macrophage expression of Ia antigen. *J Immunol* 132:1722-1727.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, Goodman ZD, Koury K, Ling M, Albrecht JK. 2001. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment for chronic hepatitis C: A randomized trial. *Lancet* 358:958-965.
- McHutchison JG, Gordon SC, Schiff ER, Shiffman ML, Lee WM, Rustgi VK, Goodman ZD, Ling MH, Cortes, Albrecht JK. 1998. Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. Hepatitis Interventional Therapy Group. *N Engl J Med* 339:1485-1492.
- Murashima S, Tanaka M, Haramaki M, Yutani S, Nakashima Y, Harada K, Ide T, Kumashiro R, Sata M. 2006. A decrease in AFP level related to administration interferon in patients with chronic hepatitis C and a high level of AFP. *Dig Dis Sci* 51:808-812.
- Murphy D, Detjen KM, Welzel M, Wiedenmann B, Rosewicz S. 2001. Interferon-alpha delays S-phase progression in human hepatocellular carcinoma cells via inhibition of specific cyclin-dependent kinases. *Hepatology* 33:346-356.
- Nishiguchi S, Kuroki T, Nakatani S, Morimoto H, Takeda T, Nakajima S, Shiomi S, Seki S, Kobayashi K, Otani S. 1995. Randomised trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 346:1051-1055.
- Otsuru A, Nagataki S, Koji T, Tamaoki T. 1988. Analysis of alpha-fetoprotein gene expression in hepatocellular carcinoma and liver cirrhosis by situ hybridization. *Cancer* 62:1105-1112.
- Parmelee M, Evenson M, Deutsch H. 1978. The presence of fatty acids in human alpha-fetoprotein. *J Biol Chem* 253:2114-2119.
- Poynard T, Marcellin P, Lee SS, Niederau C, Minuk GS, Ideo G, Bain V, Heathcote J, Zeuzem S, Trepo C, Albrecht J. 1998. Randomised trial of interferon alpha2b plus ribavirin for 48 weeks or for 24 weeks versus interferon alpha2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. International Hepatitis Interventional Therapy Group (IHIT). *Lancet* 52:1426-1432.
- Reichard O, Norkrans G, Fryden A. 1998. Randomised, double-blind, placebo-controlled trial of interferon alpha 2b with and without ribavirin for chronic hepatitis C. The Swedish Study Group. *Lancet* 351:83-87.
- Scarpa S, Giuffrida A, Palumbo C, Vasaturo F, Signorelli P, Fomi G, Modesti M, Ferrantini M, Belardelli F, Musiani P, Modesti A. 1997. Extracellular matrix remodelling in a murine mammary adenocarcinoma transfected with the interferon-alpha 1 gene. *J Pathol* 181:116-123.
- Schalm SW, Hansen BE, Chemello L, Bellobuon A, Brouwer JT, Weiland O, Cavalletto L, Schvarcz R, Ideo G, Alberti A. 1997. Ribavirin enhances the efficacy but not the adverse effects of interferon in chronic hepatitis C. Meta-analysis of individual patient data from European Centers. *J Hepatol* 26:961-966.
- Takano S, Ito Y, Yokosuka O, Ohto M, Uchiyumi K, Hirota K, Omata M. 1994. A multicenter randomized controlled dose study of ursodeoxycholic acid for chronic hepatitis C. *Hepatology* 20:558-564.
- Tsubota A, Kumada H, Arase Y, Chayama K, Saitoh S, Ikeda K, Kobayashi M, Suzuki Y, Murashima N. 1999. Combined ursodeoxycholic acid and glycyrrhizin therapy for chronic hepatitis C virus infection: A randomized controlled trial in 170 patients. *Eur J Gastroenterol Hepatol* 11:1077-1083.
- Tsukuma H, Hiyama T, Tanaka S, Nakao M, Yabuuchi T, Kitamura T, Nakanishi K, Fujimoto I, Inoue A, Yamazaki H, et al. 1993. Risk factors for hepatocellular carcinoma among patients with chronic liver disease. *N Engl J Med* 428:1797-1801.
- Wang W, Alpert E. 1995. Downregulation of phorbol 12-myristate 13-acetate-induced tumor necrosis factor-alpha and interleukin-1beta production and gene expression in human monocytic cells by human alpha-fetoprotein. *Hepatology* 22:921-928.

- Wang XW, Xu B. 1998. Stimulation of tumor-cell growth by alpha-fetoprotein. *Int J Cancer* 75:596-599.
- Wang XW, Yuan JH, Zhang RG, Guo LX, Xie Y, Xie H. 2001. Antihepatoma effect of alpha-fetoprotein antisense phosphorothioate oligodeoxyribonucleotides in vitro and in mice. *World J Gastroenterol* 7:345-351.
- Yoshida H, Shiratori Y, Moriyama M, Arakawa Y, Ide T, Sata M, Inoue O, Yano M, Tanaka M, Fujiyama S, Nishiguchi S, Kuroki T, Iwazaki F, Yokosuka O, Kinoyama S, Yamada G, Omata M. 1999. Interferon therapy reduces the risk of hepatocellular carcinoma: National surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHTT Study Group. Inhibition of hepatocarcinogenesis by interferon therapy. *Ann Intern Med* 131:174-181.
- Yoshida T, Hanada T, Tokuhisa T, Kosai K, Sata M, Kohara M, Yoshimura A. 2002. Activation of STAT3 by the hepatitis C virus core protein leads to cellular transformation. *J Exp Med* 196:641-653.
- Zuzem S, Feinman SV, Rasenack J, Heathcote EJ, Lai MY, Gane E, O'Grady J, Reichen J, Diago M, Lin A, Hoffiman J, Brunda MJ. 2000. Peginterferon alfa-2a in patients with chronic hepatitis C. *N Engl J Med* 343:1666-1672.

© John Wiley & Sons, Ltd.

The Publisher and other involved in compiling the content of this publication make no warranty as to the accuracy or completeness of any information and accept no responsibility or liability for any inaccuracy or errors or omissions. Please check all prescribing information directly with the manufacturer.



Reprinted with permission by Content'Ed Net Inc.
1-14-24-401, Kyomachibori, Nishi-ku, Osaka, JAPAN
Tel: (06)-6445-1301 Fax: (06)-6445-1334 E-mail: info@contentednet.co.jp

Original Article

Evaluation of long-term biochemical responses to combination therapy of interferon plus ribavirin in those infected with hepatitis C virus genotype 1b and high baseline viral load

Hitomi Sezaki, Fumitaka Suzuki, Yusuke Kawamura, Hiromi Yatsuji, Tetsuya Hosaka, Norio Akuta, Masahiro Kobayashi, Yoshiyuki Suzuki, Yasuji Arase, Kenji Ikeda and Hiromitsu Kumada

Department of Hepatology, Toranomon Hospital, Tokyo, Japan

Aim: The aim of this study was to determine the long-term effects in non-responders (NRs) to 48-week interferon (IFN) and ribavirin combination treatment in patients infected with hepatitis C virus (HCV) genotype 1b and high baseline viral loads.

Methods: We measured serum alanine aminotransferase (ALT) and HCV-RNA levels in 52 consecutive patients infected with HCV genotype 1b and high viral loads who received combination therapy for 48 weeks.

Results: Sustained virologic response (SVR) was noted in 30 patients (57.7%). Virologic response (VR), that is serum HCV-RNA negativity by the end of treatment and positivity during follow-up, was noted in nine patients (17.3%). Thirteen (25.0%) patients were NRs. Significantly lower serum albumin ($P = 0.007$) and ribavirin doses according to body weight ($P = 0.021$) and higher gamma glutamyl transpeptidase (GGT,

$P = 0.038$) were noted at baseline in the NR group than in the SVR and VR groups. ALT normalization rates at six months after the completion of treatment were 55.6% (5/9) in VR and 61.5% (8/13) in NRs. Sustained ALT normalization at two years after the completion of treatment was noted in 55.6% (5/9) and 58.3% (7/12), respectively.

Conclusion: Our study indicates a high rate of ALT normalization in patients infected with HCV genotype 1b and high baseline viral loads who received combination therapy and that such a rate could be maintained after the completion of therapy, even in NRs. Our results suggest that combination therapy should be continued in NRs who show ALT normalization in order to prevent potential hepatocarcinogenesis.

Key words: ALT, chronic hepatitis C, combination therapy, non-responder, ribavirin

INTRODUCTION

HEPATITIS C VIRUS (HCV) usually causes chronic infection and is a major cause of chronic hepatitis, cirrhosis and hepatocellular carcinoma (HCC).^{1–3} The aim of any treatment for chronic hepatitis C is to delay the progression of liver fibrosis and inhibit the development of HCC by the eradication of HCV, and normalization of alanine aminotransferase (ALT), even if viral clearance cannot be achieved.⁴

The combination therapy of interferon (IFN) α -2b or pegylated-IFN (PEG IFN) α -2b plus ribavirin is the first-line therapy for patients with chronic hepatitis C. The addition of ribavirin to IFN or PEG IFN is reported to enhance the virological response even in "IFN-resistant" patients.^{5–7} The achieved sustained virological response (SVR) rate by 48-week combination therapy was approximately 50% and significantly higher than in patients who received 24- or 48-week IFN monotherapy.^{8,9} However, some patients, particularly those with genotype 1b and high viral load, fail to respond to treatment and cannot achieve viral clearance even following combination therapy.

Recently, Iino *et al.*¹⁰ reported that sustained ALT normalization (ALT normal at 24 weeks after the end of treatment) following combination therapy was 28.1%,

Correspondence: Dr Hitomi Sezaki, Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-8470, Japan. Email: hitomis@mx1.harmonix.ne.jp
Received 9 November 2006; revision 27 March 2007; accepted 2 April 2007.

which was higher than the 10.5% with IFN monotherapy. Their results suggested that combination therapy might inhibit the progression to HCC, based on the high SVR rate and high rate of sustained ALT normalization.¹⁰

The aims of the present study were to determine the non-virological response (no-response, NR) rate and evaluate ALT normalization during therapy and at long-term follow-up in patients with high viral load and chronic infection with HCV genotype 1b who received combination therapy of IFN α -2b or PEG IFN α -2b plus ribavirin.

METHODS

Patients

FROM DECEMBER 2001 to February 2006, a total of 425 Japanese patients with chronic HCV infection caused by genotype 1b and high baseline viral loads received combination therapy of IFN- α -2b or PEG IFN α -2b plus ribavirin for 48 weeks at Toranomon Hospital, Tokyo, Japan. Among them, 52 patients who could be followed up for more than one year after the completion of 48 week-combination therapy were selected for the present retrospective study. The inclusion criteria were: (i) a positive test for anti-HCV antibody; (ii) HCV genotype 1b (confirmed by a PCR-based method¹¹); (iii) serum HCV-RNA levels more than 100 KIU/mL by quantitative PCR assay (Amplicor GT-HCV Monitor version 2.0; Roche Diagnostic Systems, Pleasanton, CA) within the preceding 12 weeks (defined as "high" viral load); (iv) persistently high serum ALT concentrations (the upper limit of normal for ALT is 50 IU/L) during the preceding 12 weeks; (v) a diagnosis of chronic hepatitis on liver biopsy specimen obtained within the preceding one year of enrollment; (vi) hemoglobin concentration of ≥ 12.0 g/dL; (vii) platelet count of $\geq 100 \times 10^3/\mu\text{l}$; and (viii) signing a consent form of the study protocol that had been approved by Human Ethics Review Committee of Toranomon Hospital. Patients with the following conditions were excluded from the study: (i) other forms of liver disease (e.g. primary biliary cirrhosis, alcoholic liver disease and autoimmune liver disease); (ii) treatment with any other antiviral or immunomodulatory agents administered within the preceding 24 weeks; (iii) patients with hepatitis B surface antigen or hepatitis B core antibody; (iv) coinfection with human immunodeficiency virus; and (v) women who were pregnant or lactating. The selected subjects

included 33 males and 19 females, aged 19–65 years, with a median age of 52 years.

Study protocol

The combination treatment was provided for 48 weeks, with a subsequent follow-up period of more than one year. In 21 (40.3%) patients, IFN α -2b (Schering-Plough, Osaka, Japan) was injected intramuscularly at 6 million units (MU)/day for the initial two weeks, followed by three times per week for 46 weeks. In the remaining 31 (59.6%) patients, 1.5 $\mu\text{g}/\text{kg}/\text{week}$ of PEG IFN α -2b (Schering-Plough, Osaka, Japan) was injected subcutaneously for 48 weeks. All patients received ribavirin (Schering-Plough, Osaka, Japan) at a dose adjusted to body weight (600 mg for individuals with body weight of ≤ 60 kg, 800 mg for weight of 60–80 kg, and 1000 mg for weight ≥ 80 kg).

Biochemical and virological responses to treatment were assessed during the 48-week treatment period and during the subsequent follow-up period. Biochemical response was defined as normalization of serum ALT activity (the upper limit of normal for ALT is 50 IU/L) by the end of treatment. Virological response (VR) was defined as undetectable serum HCV-RNA at the end of the 48-week treatment, as confirmed by a qualitative PCR assay (Amplicor HCV version 2.0; Roche Molecular Systems, Belleville, NJ), but reappearance of HCV-RNA during the one-year follow-up period. A sustained virological response (SVR) was defined as disappearance of serum HCV-RNA after the completion of treatment until the end of the follow-up period. A non-virological response (NR) was defined as persistent presence of HCV-RNA during treatment.

Blood tests

Routine biochemical and hematological tests were performed at least once every month during and after treatment. Serum HCV-RNA levels were measured using a quantitative PCR assay with a lower detection limit of quantification of 0.5 KIU/mL (Amplicor HCV Monitor version 2.0; Roche Diagnostics, Pleasanton, CA). The presence or absence of serum HCV-RNA was assessed using a qualitative PCR assay (Amplicor HCV version 2.0; Roche Diagnostics, Pleasanton, CA) with a lower detection limit of 100 copies/mL.

Liver histopathological examination

Histopathological staging of liver biopsy specimens obtained at baseline and during treatment was performed according to the classification of Desmet *et al.*¹²

Table 1 Comparison of baseline clinical profiles of patients infected by HCV genotype 1b and a high viral load with or without viral disappearance during combination therapy

	Total (n = 52)	SVR + VR (n = 39)	NR (n = 13)	P-value
Age (y)†	19-65 (52)	19-64 (51)	37-65 (53)	0.808
Sex (M/F)	33/19	26/13	7/6	0.501
Body weight (kg)	42.3-120 (62.4)	46-77 (63.1)	42.3-120 (60)	0.759
Ribavirin dose/kg weight (mg/kg)	6.7-14.2 (11.2)	10.1-13.1 (11.4)	6.7-14.2 (10.7)	0.021
Histopathological staging (F1/2/3)	28/18/6	22/14/3	6/4/3	0.322
Histopathological grading (A1/2/3)	28/24/0	22/17/0	6/7/0	0.541
ALT (IU/L)	29-276 (98)	29-276 (101)	50-135 (71)	0.131
GGT (IU/L)	16-240 (62)	16-240 (60)	40-121 (75)	0.038
Hemoglobin (g/dL)	12.0-17.4 (14.4)	12.0-17.4 (14.5)	12.5-16.3 (14.4)	0.674
Platelet count ($\times 10^3/\mu\text{L}$)	101-309 (177)	101-309 (178)	111-237 (173)	0.550
Fe ($\mu\text{g/dL}$)	46-308 (139)	46-308 (143)	70-214 (130)	0.393
Ferritin ($\mu\text{g/L}$)	<10-644 (176)	<10-644 (188)	52-335 (125)	0.290
ICG R15 (%)	7-41 (15)	7-33 (13)	7-41 (17)	0.842
Albumin (g/dL)	3.4-4.5 (3.8)	3.4-4.5 (3.9)	3.4-3.9 (3.7)	0.007
HCV-RNA (KIU/mL)	49-3500 (795)	110-3500 (810)	49-2800 (940)	0.512
Follow-up period (month)	18-43 (38)	18-43 (38)	18-42 (39)	0.617

†Data are ranges (median).

ALT, alanine aminotransferase; GGT, gamma glutamyl transpeptidase; ICG R15, indocyanine green retention rate at 15 min; NR, non-responders; SVR, sustained virological response; VR, virological response.

Statistical analysis

Nonparametric tests, including the χ^2 , Fisher's exact probability and Mann-Whitney *U*-tests, were used to analyze the baseline clinical profile of patients. Univariate and multivariate logistic regression analyses were used to determine the factors that significantly contributed to a VR. We also calculated the odds ratios and 95% confidence intervals (95% CI). A *P*-value < 0.05 by the two-tailed test was considered statistically significant. All analyses were performed using SPSS version 10.1 (SPSS, Chicago, IL).

RESULTS

Baseline characteristics

THE BASELINE CLINICAL profile of the patients at commencement of the combination therapy is shown in Table 1. The follow-up period ranged from 18 to 43 months (median, 38 months). The baseline viral load ranged from 49 to more than 5000 KIU/mL (median, 795 KIU/mL).

Virological response rates

Of 52 patients, 39 (75.0%) achieved viral disappearance during the combination therapy, 30 (57.7%)

patients attained an SVR and nine (17.3%) patients had a reappearance of HCV-RNA during the follow-up period, that is VR. The remaining 13 (25.0%) were NRs (Table 2). We compared the baseline clinical profile of SVR plus VR group and NR group (Table 1). There were no differences between the two groups with respect to age, sex, body weight, histopathological findings, serum ALT, hemoglobin, platelets count and HCV-RNA levels at the commencement of the combination therapy. Serum albumin concentrations (*P* = 0.007) and ribavirin dose according to body weight (*P* = 0.021) were significantly lower and γ -glutamyl transpeptidase (GGT, *P* = 0.038) was higher in the NR group than in the other group. We then analyzed the data to determine those factors that could predict a VR. Univariate analysis identified three parameters that significantly influenced the VR: serum albumin concentrations ≥ 3.9 g/dL (*P* = 0.012), GGT < 50 IU/L (*P* = 0.048) and ribavirin dose according to body weight ≥ 11.2 (*P* = 0.009). Multivariate analysis using variables including sex, age, serum albumin, ALT, GGT and ribavirin dose according to body weight, identified three parameters that independently influenced the virological response: male (*P* = 0.035), ribavirin dose according to body weight ≥ 11.2 (*P* = 0.012) and GGT < 50 IU/L (*P* = 0.010).

Table 2 ALT normalization in patients infected with genotype 1b and a high baseline viral load, who received 48-week combination therapy of IFN plus ribavirin

	ALT normalization after completion of treatment			
	End of treatment	6 months	1 year	2 years
VR (<i>n</i> = 9)	7	5	5	5/9
Follow up period; 17–42 (40) months	(77.8%)	(55.6%)	(55.6%)	(55.6%)
NR (<i>n</i> = 13)	10	8	6	7/12†
Follow up period; 19–42 (36) months	(76.9%)	(61.5%)	(46.2%)	(58.3%)
Total		13/21 (59.1%)		12/21† (57.1%)

†One patient could not be followed up for 2 years after completion of treatment.
ALT, alanine aminotransferase; NR, non-responders; VR, virological response.

ALT normalization rate

ALT normalization rate during the 48-week combination therapy is shown in Table 2. We determined the ALT normalization rate for non-SVR, VR and NR, because all SVR cases exhibited ALT normalization after the completion of treatment. ALT normalization rates in patients who showed VR and NR were 77.8% (7/9) and 76.9% (10/13) at the end of treatment, respectively; and 55.6% (5/9) and 61.5% (8/13) at six months after the completion of treatment, respectively. For the same two groups, the rates of sustained ALT normalization at two-years after the completion of treatment were 55.6% (5/9) and 58.3% (7/12), respectively. The overall ALT normalization rate at two years after the completion of treatment is shown in Table 3. Because of the high rate of sustained ALT normalization among the SVR cases (*n* = 30), the overall ALT normalization rate at two years after completion of treatment for all patients was 82.4% (42/51).

Table 3 Rates of SVR, VR and NR, and overall ALT normalization in patients infected with genotype 1b and a high baseline viral load who received 48-week combination therapy of IFN plus ribavirin

	N	Rate	ALT normalization at 2 years after completion of treatment	
SVR	30	57.7%	30/30	100%
VR	9	17.3%	5/9	55.6%
NR	13	25.0%	7/12†	57.1%
Total	52		42/51†	82.4%


†One patient could not be followed up for 2 years after completion of treatment.

ALT, alanine aminotransferase; NR, non-responders; SVR, sustained virological response; VR, virological response.

Figure 1 shows the clinical course of a patient who showed the persistent presence of HCV-RNA during treatment but achieved sustained ALT normalization after the completion of combination therapy. The HCV-RNA level decreased only 1 log relative to baseline viral load at the end of the combination therapy. However, serum ALT normalized rapidly after the commencement of treatment and was persistently normal over more than three years after the completion of the treatment.

DISCUSSION

THE PRIMARY PURPOSE of treatment of patients with chronic hepatitis C is the eradication of HCV, that is SVR. However, patients who are infected with HCV genotype 1b and have a high baseline viral load are considered refractory to IFN, and the SVR rate in such patients is less than 10%.⁹ However, the use of ribavirin in combination with IFN has markedly improved the SVR rate.^{5–7,9,13} Early disappearance of HCV-RNA in the serum samples is a predictor of SVR to IFN-based therapy, and the most appropriate time-point for determining the outcome of combination treatment is week

**Figure 1** Clinical course of a patient with sustained ALT normalization who showed the persistent presence of HCV-RNA.

12 of such therapy.^{14,15} We reported previously that when the loss of HCV occurs at more than 24 weeks, the negative predictive value for SVR was 100%.¹⁴ Consequently, it is not clear how to treat such patients (i.e. those infected with genotype 1b and had a high baseline viral load who are HCV-positive at 24 weeks after the initiation of combination therapy of IFN plus ribavirin).

In the present study, we investigated cases who were HCV-RNA positive during the combination therapy, that is NRs. Compared with patients who were HCV-RNA negative during the combination therapy, our patients had significantly lower serum albumin and higher GGT (Table 1). In multivariate analysis to determine those factors that could predict a virological response, although serum albumin did not independently influence the VR ($P = 0.062$), we consider this result approximately similar to our previous reports, in which we identified serum albumin as a predictor of non-VR in patients on 48 weeks of the same combination therapy.^{16,17} The results that GGT independently influenced the virological response ($P = 0.010$) confirm the findings of previous studies that GGT levels correlate with sustained virological response.^{18,19} The low level of serum albumin reflects the deterioration of the ability of the liver to synthesize serum proteins due to progression of liver fibrosis. This finding suggests that eradication of HCV-RNA is difficult in patients with advanced liver fibrosis, even when treated with IFN monotherapy or combination therapy.

However, our results showed that even when patients with genotype 1b and high viral load do not achieve HCV-RNA negativity during combination therapy with IFN and ribavirin, they achieved a high rate of ALT normalization (76.9%) at the end of therapy and could maintain a normal ALT level over a long period of time after the completion of combination therapy (58.3% at two years after the completion of such therapy) (Table 2). In IFN monotherapy, the ALT normalization rate reported in genotype 1 patients ranged from 10 to 32%.^{4,20,21} We also reported that among 1654 patients treated with IFN alone, 266 (16.1%) showed normal levels of ALT without loss of HCV-RNA for ≥ 6 months after the completion of IFN monotherapy. Considered together, these results suggest that a higher ALT normalization rate can be achieved by combination therapy.

On the other hand, the SVR rate improved to approximately 57.7% in our 48-week combination therapy, and ALT normalization was noted in approximately 60% of non-SVR cases at six months and 57.1% at two years after completion of combination therapy (Table 2). In other words, the overall ALT normalization rate was

extremely high (82.4%). These results are similar to those of a previous study,¹⁰ which showed a higher rate of sustained ALT normalization (normal ALT levels at 24 weeks after the end of treatment) with combination therapy than with IFN monotherapy.

The natural history of chronic hepatitis C includes cirrhosis and hepatocellular carcinoma. Previous studies reported that the predictive factors of progression to cirrhosis from chronic hepatitis C were male sex, heavy alcohol consumption, elevated serum ALT levels and histology of high grade necroinflammatory activity.²² In this regard, we reported previously that normalization of ALT levels after IFN therapy without loss of serum HCV-RNA was associated with decreased incidence of hepatocarcinogenesis.^{23,24} In the present study, we could not investigate whether normalization of ALT levels after combination therapy was associated with reduced incidence of hepatocarcinogenesis because the median follow-up period was 38 months. However, in view of the previous and present results, we consider that maintenance of ALT normalization over a long time after the completion of the combination therapy seems to suppress progression of liver fibrosis and future development of hepatocellular carcinoma. Accordingly, we recommend that even NRs to combination therapy should continue combination therapy for 48 weeks, especially when they achieve ALT normalization during the therapy, to further maintain ALT normalization after the completion of the combination therapy.

In this study, to compare it with past studies, we set the upper limit of ALT to 50 IU/L as a definition of ALT normalization. To review correct ALT normalization, we should lower the upper limit of ALT and study further large-scale populations in the future. And, because we investigated only cases that completed 48 weeks of IFN plus ribavirin treatment, we consider our treatment results about the SVR and the sustained ALT normalization rate were good in comparison with former results. Unfortunately, however, there were cases that had to stop the treatment because of adverse events. Therefore, we consider that even NRs to combination therapy should continue the combination therapy for 48 weeks if they achieve the ALT normalization during the therapy, and then by further maintaining ALT normalization after the completion of the combination therapy, they may suppress the progression of liver fibrosis and prevent future development of hepatocellular carcinoma.

In conclusion, we have demonstrated that combination therapy for patients infected with HCV genotype 1b and a high baseline viral load achieved a high rate of ALT normalization that could be maintained after the

completion of therapy, even in patients who failed to show HCV-RNA eradication. Thus, our results suggest that NRs with ALT normalization should continue the combination therapy to prevent potential future hepatocarcinogenesis.

REFERENCES

- Niederau C, Lange S, Heintges T *et al.* Prognosis of chronic hepatitis C: results of a large, prospective cohort study. *Hepatology* 1998; 28: 1687–95.
- Ikeda K, Saitoh S, Suzuki Y *et al.* Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: a prospective observation of 2215 patients. *J Hepatol* 1998; 28: 930–8.
- Kenny-Walsh E. Clinical outcomes after hepatitis C infection from contaminated anti-D immune globulin. Irish Hepatology Research Group. *N Engl J Med* 1999; 340: 1228–33.
- Ikeda K, Saitoh S, Arase Y *et al.* Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C: a long-term observation study of 1,643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* 1999; 329: 1124–30.
- Manns MP, McHutchison JG, Gordon SC *et al.* Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001; 358: 958–65.
- McHutchison JG, Gordon SC, Schiff ER *et al.* Interferon alpha-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. *N Engl J Med* 1998; 339: 1485–92.
- Poynard T, Marcellin P, Lee SS *et al.* Randomised trial of interferon alpha 2b plus ribavirin for 48 weeks or for 24 weeks versus interferon alpha 2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. *Lancet* 1998; 352: 1426–32.
- Reichard O, Norkrans G, Frydén A *et al.* Randomised double-blind, placebo-controlled trial of interferon α -2b with and without ribavirin for chronic hepatitis C. *Lancet* 1998; 351: 83–7.
- Lindsay KL, Trepo C, Heintges T *et al.* A randomized, double-blind trial comparing pegylated interferon alfa-2b to interferon alfa-2b as initial treatment for chronic hepatitis C. *Hepatology* 2001; 34: 395–403.
- Iino S, Tomita E, Kumada H *et al.* Impact of daily high-dose IFN α -2b plus ribavirin combination therapy on reduction of ALT levels in patients with chronic hepatitis C with genotype 1 and high HCV-RNA levels. *Hepatol Res* 2005; 31: 88–94.
- Chayama K, Tsubota A, Arase Y *et al.* Genotypic subtyping of hepatitis C virus. *J Gastroenterol Hepatol* 1993; 8: 150–6.
- Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Scheuer PJ. Classification of chronic hepatitis: diagnosis, grading and staging. *Hepatology* 1994; 19: 1513–20.
- Shehab TM, Fontana RJ, Oberhelman K, Marrero JA, Su GL, Lok ASF. Effectiveness of interferon α -2b and ribavirin combination therapy in the treatment of naïve chronic hepatitis C patients in clinical practice. *Clin Gastroenterol Hepatol* 2004; 2: 425–31.
- Tsubota A, Arase Y, Someya T *et al.* Early viral kinetics and treatment outcome in combination of high-dose interferon induction vs. pegylated interferon plus ribavirin for naïve patients infected with hepatitis C virus of genotype 1b and high viral load. *J Med Virol* 2005; 75: 27–34.
- Davis GL, Wong JB, McHutchison JG, Manns MP, Harvey J, Albrecht J. Early virologic response to treatment with peginterferon α -2b plus ribavirin in patients with chronic hepatitis C. *Hepatology* 2003; 38: 645–52.
- Akuta N, Suzuki F, Sezaki H *et al.* Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 2005; 48: 372–80.
- Akuta N, Suzuki F, Tsubota A *et al.* Efficacy of interferon monotherapy to 394 consecutive naïve cases infected with hepatitis C virus genotype 2a in Japan: therapy efficacy as consequence of tripartite interferon of viral, host and interferon treatment-related factors. *J Hepatol* 2002; 37: 831–6.
- Berg T, Sarrazin C, Herrmann E *et al.* Prediction of treatment outcome in patients with chronic hepatitis C: significance of baseline parameters and viral dynamics during therapy. *Hepatology* 2003; 37: 600–9.
- Berg T, Von Wagner M, Nasser S *et al.* Extended treatment duration for hepatitis C virus type 1: Comparing 48 versus 72 weeks of peginterferon-alfa-2a plus ribavirin. *Gastroenterology* 2006; 130: 1086–97.
- Iino S. *Incidence of Progression to Hepatocellular Carcinoma Following Interferon Treatment of Chronic Hepatitis C Using a Survey by Questionnaire*. Annual report from Non-A, Non-B Hepatitis Research Group sponsored ministry of Health and Welfare, Tokyo. 1997; 49–52 (in Japanese).
- Kasahara A, Hayashi N, Mochizuki K *et al.* Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998; 27: 1394–402.
- Freeman AJ, Law MG, Kaldor JM, Dore GJ. Predicting progression to cirrhosis in chronic hepatitis C virus infection. *J Viral Hepat* 2003; 10: 285–93.
- Ikeda K, Arase Y, Saitoh S *et al.* Anticarcinogenic impact of interferon on patients with chronic hepatitis C: a large-scale long-term study in a single center. *Intervirology* 2006; 49: 82–90.
- Akuta N, Suzuki F, Suzuki Y *et al.* Long-term follow-up of interferon monotherapy in 454 consecutive naïve patients infected with hepatitis C virus: multi-course interferon therapy may reduce the risk of hepatocellular carcinoma and increase survival. *Scand J Gastroenterol* 2005; 40: 688–96.

Original Article

Efficacy and anticarcinogenic activity of interferon for hepatitis C virus-related compensated cirrhosis in patients with genotype 1b low viral load or genotype 2

Eiko Hasegawa, Masahiro Kobayashi, Yusuke Kawamura, Hiromi Yatsuji, Hitomi Sezaki, Tetsuya Hosaka, Norio Akuta, Fumitaka Suzuki, Yoshiyuki Suzuki, Yasuji Arase, Kenji Ikeda and Hiromitsu Kumada

Department of Hepatology, Toranomon Hospital, Tokyo, Japan

Background: We assessed the efficacy and anticarcinogenic effects of interferon (IFN) therapy in patients with hepatitis C virus (HCV)-related cirrhosis.

Methods: The study subjects were 123 Japanese patients with HCV-related cirrhosis with genotype 1b low viral load or genotype 2 who received IFN from 1989 to 2005 (18 patients continue to receive IFN therapy). They included 81 men and 42 women aged 29–74 years (median, 56 years).

Results: Univariate analysis identified four parameters that significantly influenced SVR; viral load (low HCV concentration, $P < 0.001$), duration of IFN therapy (≥ 52 weeks, $P = 0.029$), daily dose of IFN (≥ 6 million units, $P = 0.018$), induction therapy (presence, $P = 0.010$) and choline esterase ($> 1.0 \Delta\text{pH}$, $P = 0.037$). Multivariate analysis identified viral load (risk ratio = 6.329, $P < 0.001$) and daily dose of IFN (risk ratio = 2.62, $P = 0.042$) as two independent parameters that

influenced SVR. During the observation period, newly developed hepatocellular carcinoma (HCC) was detected in 22 patients. The rates of development of HCC in patients with SVR were 5.8% at the fifth year and 10.3% at the 10th year, compared with 25.8% at the fifth year and 42.5% at the 10th year in non-SVR patients. Multivariate analysis showed that IFN efficacy (SVR) was the only independent factor of hepatocarcinogenesis (hazard ratio: 0.185, 95% confidence interval: 0.042–0.810, $P = 0.025$).

Conclusion: Among patients with HCV-related cirrhosis, the rate of development of HCC is significantly less in patients with SVR.

Key words: cancer prevention, cirrhosis, hepatitis C virus, hepatocellular carcinoma, interferon

INTRODUCTION

HEPATITIS C VIRUS (HCV)-associated hepatocellular carcinoma (HCC) typically develops through a sequence of events that progresses from chronic inflammation through fibrosis and cirrhosis accompanied by dysplasia and ultimately HCC. In a cohort study of Japanese patients with HCV-related cirrhosis,¹ the cumulative appearance rates of HCC at 5, 10 and 15 years were 32.5, 59.6 and 77.4%, respectively. Furthermore, the life expectancy of patients with HCV-related cirrhosis is

largely influenced by the development of HCC during the clinical course.² Because an effective and curative therapy for HCC remains limited at best, primary prevention of HCC in patients with chronic liver disease is of great importance at present.

Interferon (IFN) is effective in eliminating HCV and reducing the serum level of alanine aminotransferase (ALT) in some patients with chronic hepatitis C.^{3–5} The response to IFN therapy is related to various factors such as HCV subtype, serum concentration of HCV, IFN treatment and liver histology.^{6–8} Kasahara *et al.* reported that the presence of a persistently normal level of serum aminotransferase after IFN therapy was associated with significant reduction in the incidence of HCC in patients with chronic hepatitis C.⁹ Furthermore, reduced incidence of HCC in HCV-associated cirrhosis by IFN has been reported by many investigators;^{10–13} in fact, only a few studies have failed to find its benefit.^{14,15}

Correspondence: Dr Masahiro Kobayashi, Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-8470, Japan. Email: mshkobayashi@toranomon.gr.jp

Received 11 December 2006; revision 28 February 2007; accepted 7 April 2007.

In order to evaluate whether IFN can reduce the rate of hepatocarcinogenesis and to analyze the pretreatment predictive factors associated with response to IFN and carcinogenesis in patients with HCV-related cirrhosis, we retrospectively analyzed 123 patients with HCV-related cirrhosis.

PATIENTS AND METHODS

Study population

A TOTAL OF 634 patients were diagnosed with HCV-related cirrhosis from 1989 to 2005 at the Department of Hepatology at Toranomon Hospital, Tokyo, Japan. Of these, 267 (42.1%) patients were treated with IFN. They included 140 patients with genotype 1b-high HCV concentration, 24 patients with genotype 1b-low HCV concentration, 38 patients with genotype 2a-high HCV concentration, 42 patients with genotype 2a-low HCV concentration, 18 patients with genotype 2b-high HCV concentration, and one patient with genotype 2b-low HCV concentration. A total of 123 patients with HCV-related cirrhosis with genotype 1b low viral load or genotype 2 were enrolled in this analysis, whose initial sera tested negative for hepatitis B surface antigen by radioimmunoassay (Ausria; Dainabot, Tokyo, Japan) and positive for anti-HCV by the second or third-generation enzyme-linked immunosorbent assay (Dainabot). They included 81 men and 42 women aged 29–74 years (median, 56 years). The diagnosis of liver cirrhosis was based on clinical features, laboratory tests, and peritoneoscopy or liver biopsy. In order to investigate hepatocarcinogenesis in HCV-related cirrhosis, patients coinfecting with HBV were excluded. Our institution does not require informed consent for retrospective studies.

Background and laboratory findings

Table 1 shows demographic profiles and results of laboratory tests for the 123 patients at baseline (before treatment with IFN). Quantitative analysis of HCV-RNA was performed using a branched DNA probe assay (bDNA probe assay, version 2.0; Chiron, Dai-ichi Kagaku, Tokyo) and polymerase chain reaction (PCR)-based assay using the protocol provided by the manufacturer (Amplicor HCV Monitor assay version 2.0; Roche Diagnostics, Tokyo, Japan). HCV genotype was classified by PCR, using a mixture of primers for six subtypes known to exist in Japan, as reported previously.¹⁶

Table 1 Demographics and baseline characteristics of 105 patients in the present study

Age (years)	56 (29–74)†
Sex (M/F)	81/42
BMI (kg/m ²)	23.9 (16.9–35.7)†
Albumin (g/dL)	3.7 (1.8–4.7)†
AST (IU/L)	70 (26–338)†
ALT (IU/L)	80 (11–434)†
Cholesterol (mg/dL)	159 (93–272)†
Choline esterase (Δ pH)	0.8 (0.3–1.5)†
AFP (μ g/L)	11 (2–631)†
Ferritin (μ g/L)	178 (<10–2076)†
Hyaluronic acid (μ L)	184 (30–1000)†
FBS (mg/mL)	94 (65–338)†
Platelet ($\times 10^4/\mu$ L)	9.8 (2.5–22.3)†
HCV genotype	
1b	24
2a	80
2b	19
HCV-RNA	
High viral load	56
Low viral load	67

†Data expressed as median (range).

High viral load > 100 KIU/mL or > 1 Meq/mL; low viral load < 100 KIU/mL or < 1 Meq/mL.

AFP, alpha fetoprotein; ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index; FBS, fasting blood sugar; HCV, hepatitis C virus.

IFN treatment and evaluation of response to therapy

Among the 123 patients, 85 (69.1%) received IFN for the first time, while the remaining 38 (30.9%) patients had received IFN prior to this protocol. In the present study, IFN treatment involved the use of natural or recombinant IFN- α ($n = 83$), natural IFN- β ($n = 38$), or both ($n = 2$). The dosage of IFN varied in this study; 22 (17.9%) patients received 3–9 million units (MU) IFN daily for 4–8 weeks; 38 (30.9%) patients received 3–9 MU IFN daily for 2–8 weeks followed by two or three times per week; 57 (46.3%) patients received intermittent IFN two to three times per week; six patients received pegylated IFN. Among the 123 patients, 10 (8.1%) were treated with both IFN and ribavirin. The median dose of IFN was 399 MU (18–14 778 MU) during a median period of 25 weeks (1.9–602 weeks), and the daily dose was < 6 MU ($n = 59$) and ≥ 6 MU ($n = 58$). In this study, the initial daily administration of IFN for two or more weeks was defined as induction therapy.

The response to IFN was evaluated by clearance of HCV-RNA from serum and serum levels of ALT. Sustained virological response (SVR) was defined as persistent disappearance of HCV-RNA after therapy, while biochemical response (BR) was defined as normalization of ALT levels without elimination of HCV-RNA for at least 6 months after therapy. No response (NR) was defined as elevation or a transient decrease in serum ALT levels with persistent HCV-RNA levels in the serum.

Follow up of patients and diagnosis of HCC

Patients were followed up monthly after diagnosis of liver cirrhosis in our outpatient clinic and monitored clinically by hematological, biochemical and virological tests. In addition to admission to receive IFN treatment, biweekly or monthly follow up was performed in almost all patients who received IFN. Imaging studies were conducted every 3 months in the majority of patients using ultrasonography or computed tomography (CT). Angiography was considered only when HCC was suspected on ultrasonography or CT. The diagnosis of HCC was made by characteristic hypervascular stain on hepatic angiography. When the hepatic nodule did not show hypervascular stain, a fine-needle biopsy was carried out to exclude or diagnose HCC.

Statistical analysis

We used univariate and multivariate logistic regression analyses to determine those factors that contributed to SVR. We also calculated the odds ratios and 95% confidence intervals (95% CI). All *P*-values of less than 0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance ($P < 0.05$) or marginal significance ($P < 0.15$) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent factors. Potential predictive factors associated with SVR included the following 20 variables: age, sex, body mass index (BMI), serum albumin, cholinesterase, total cholesterol, platelet count, α -fetoprotein (AFP), indocyanine green retention rate at 15 min (ICG R15), fasting blood glucose, aspartate aminotransferase (AST), ALT, level of viremia, genotype, combination therapy with ribavirin, duration of IFN therapy, total dose of IFN, daily dose of IFN, method of IFN administration and type of IFN. The incidence of hepatocarcinogenesis was calculated by the Kaplan–Meier method; it was based on the duration between the start of IFN therapy and detection of HCC. Differences in slopes of carcinogenesis curves were evaluated by the log–rank test. Independent factors associated with the development of HCC were studied

using stepwise Cox regression analysis. The following 21 variables were analyzed: age, sex, BMI, serum albumin, cholinesterase, total cholesterol, platelet count, AFP, ICG R15, fasting blood glucose, AST, ALT, level of viremia, genotype, combination therapy with ribavirin, duration of IFN therapy, total dose of IFN, daily dose of IFN, method of IFN administration, type of IFN and the effect of IFN (SVR). Statistical analysis was conducted by using SPSS software (version 10; SPSS, Chicago, IL, USA).

RESULTS

Response to IFN

AMONG 123 PATIENTS who received IFN therapy, the response to IFN therapy could be evaluated in 105 who completed the treatment protocol until December 2005, while the remaining 18 patients continue to receive IFN therapy. The dose of IFN was reduced from 6 to 3 MU per day in 20 patients. In 14 of the 20 patients, the IFN dose was reduced according to the study protocol; it was reduced at 1 week after starting IFN in 10 patients and at two weeks in four patients. In the other six patients, the dose was reduced due to thrombocytopenia. Among the 105 patients, 48 (45.7%) showed SVR, 14 (13.3%) showed BR and 43 (41%) were NR.

Efficacy of IFN treatment according to baseline viral load and genotype

For this part of the study, pretreatment viral load was measured in 123 cases and subjects were divided into two groups using a cut-off viral load of 1 Meq/mL or 10^6 copies/mL. Table 2 shows the treatment efficacy estimated by baseline (pretreatment) viral load and genotype among the 105 cases. Of 21 patients with genotype 1b (low HCV concentration), nine (42.9%) showed SVR. Among 31 patients with genotype 2a (high HCV concentration), eight (25.8%) showed SVR and six (19.4%) showed BR. Among 35 patients with genotype 2a (low HCV concentration), 26 (74.3%) showed SVR and two (5.7%) showed BR. Among 17 patients with genotype 2b (high HCV concentration), four (23.5%) showed SVR and two (11.8%) showed BR. The single patient with genotype 2b (low HCV concentration) showed SVR. In summary, among 57 patients with a low viremia level, 36 (63.2%) achieved SVR and six (10.9%) achieved BR, while of 48 patients with a high viremia level, 12 (25.0%) showed SVR and eight (16.7%) patients showed BR.

Table 2 Proportion of patients with SVR among the 105 patients who were treated with IFN for HCV-related cirrhosis

	HCV genotype			Total
	1b	2a	2b	
HCV-RNA high viral load	–	8/31 (25.8)	4/17 (23.5)	12/48 (25.0)
HCV-RNA low viral load	9/21 (42.9)	26/35 (74.3)	1/1 (100.0)	36/57 (63.2)
Total	8/20 (42.9)	34/66 (51.5)	5/18 (27.8)	48/105 (48.0)

Numbers in parentheses are percentages of patients.

High viral load > 100 KIU/mL or > 1 Meq/mL; low viral load < 100 KIU/mL or < 1 Meq/mL.

HCV, hepatitis C virus; IFN, interferon; SVR, sustained virological response.

Side-effects

Almost all patients treated with IFN showed a variable degree of fever, chills, myalgia, headache, and general malaise after the first injection of IFN. Most patients developed a variable degree of leukocytopenia and thrombocytopenia. IFN therapy was discontinued due to anemia in one patient, thrombocytopenia in one patient, skin eruption in one patient, worsening of diabetes mellitus in one patient, retinopathy in one patient, bleeding from the ocular fundus in one patient, and interstitial pneumonia in one patient (total seven patients, 5.7%).

Predictive factors associated with SVR in multivariate analysis

We then analyzed the data for the entire population sample to determine those factors that could predict SVR. Univariate analysis identified four parameters that significantly influenced the SVR. These included viral load (low HCV concentration, $P < 0.001$), daily dose of IFN (≥ 6 MU, $P = 0.018$), induction therapy (present, $P = 0.010$) and choline esterase level (> 1.0 ΔpH, $P = 0.037$). AFP (< 20 μg/L, $P = 0.058$) and duration of IFN therapy (≥ 52 weeks, $P = 0.064$) were marginally associated with SVR (Table 3). Multivariate analysis identified two parameters that independently influenced SVR, including viral load (risk ratio = 6.99, $P < 0.001$) and daily dose of IFN (risk ratio = 2.62, $P = 0.042$) (Table 4).

Crude rates of hepatocarcinogenesis

Four of the 123 patients received IFN therapy after removal of HCC by either surgical resection or locoregional ablation. Therefore, these four patients were excluded from the following analysis. During the observation period (median: 4.6 years, range: 0.3–14.0 years), HCC developed in 22 (18.5%) of the 119 patients. Of these, three patients showed SVR, 16

patients showed NR and the remaining three patients developed HCC while still receiving IFN therapy and their ALT were below the upper limit of normal. One patient continued IFN therapy after the diagnosis of HCC. The crude rates of hepatocarcinogenesis were 16.8% at the fifth year, 29.1% at the 10th year and 34.2% at the 15th year. The rates of hepatocarcinogenesis in patients with SVR were 5.8% at the fifth year, and 10.3% at the 10th year, and in patients with non-SVR were 25.8% at the fifth year, and 42.5% at the 10th year (Fig. 1). Hepatocarcinogenesis was significantly less frequent in patients with SVR than in patients with non-SVR. (log-rank test, $P = 0.007$).

Predictive factors of hepatocarcinogenesis

Univariate analysis identified three factors that correlated significantly with hepatocarcinogenesis (Table 5). They were the response to IFN therapy (SVR, $P = 0.007$), serum albumin level (> 4.0 g/dL, $P = 0.043$) and choline esterase (> 1.0 ΔpH, $P = 0.009$). Age (> 56 years, $P = 0.080$) and daily dose of IFN (> 6 MU, $P = 0.100$) were marginally associated with hepatocarcinogenesis. Multivariate analysis showed that efficacy of IFN therapy independently influenced the development of HCC in the cohort; SVR was associated with a significant decrease in risk of hepatocarcinogenesis (hazard ratio: 0.185, 95%CI: 0.042–0.810), compared with non-SVR (Table 6).

DISCUSSION

PROGNOSIS OF PATIENTS with HCV-related cirrhosis is greatly affected by the development of HCC, especially during the compensation period.² Kasahara *et al.* reported previously that the development of HCC could be suppressed by IFN therapy and elimination of HCV in patients with chronic hepatitis C.⁹ Likewise, hepatocarcinogenesis is significantly inhibited by IFN

Table 3 Results of univariate analysis for SVR to IFN therapy in patients with HCV-related cirrhosis

Factor	Category	SVR (n = 48)	Non-SVR (n = 58)	P-value
Age	> 56 years	29 (60.4%)	32 (55.2%)	NS
Sex	Male	31 (64.6%)	38 (65.5%)	NS
BMI	≥ 25 kg/m ²	14 (29.2%)	17 (29.3%)	NS
Albumin	> 4.0 g/dL	17 (35.4%)	18 (31.0%)	NS
AST	> 76 IU/l	23 (47.9%)	29 (50.0%)	NS
ALT	> 100 IU/l	14 (29.2%)	22 (37.9%)	NS
Cholesterol	> 160 mg/dL	26 (54.2%)	27 (46.6%)	NS
Choline esterase	> 1.0 ΔpH	22 (46.8%)	14 (25.5%)	0.037
AFP	> 20 μg/L	11 (23.9%)	24 (43.6%)	0.058
FBS	> 126 mg/mL	10 (23.3%)	6 (11.1%)	NS
Platelets	> 10 × 10 ⁴ /mL	27 (56.3%)	27 (46.6%)	NS
HCV genotype	1b	9 (18.8%)	12 (20.7%)	NS
HCV-RNA level	High	12 (25.0%)	36 (62.1%)	< 0.001
Total dose of IFN†	≥ 400 MU	21 (45.7%)	26 (44.8%)	NS
Daily dose of IFN†	≥ 6 MU	31 (67.4%)	25 (43.1%)	0.018
Duration of IFN	≥ 52 weeks	4 (8.3%)	13 (22.4%)	0.064
Induction therapy†	Yes	31 (67.4%)	24 (41.4%)	0.010
Type of IFN	Alpha	29 (60.4%)	38 (65.5%)	NS
Combination with ribavirin	Yes	5 (10.4%)	2 (3.4%)	NS

†Patients who were treated with peglated IFN were excluded from analysis.

High viral load ≥ 100 KIU/mL or ≥ 1 Meq/mL.

AFP, alpha fetoprotein; ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index; FBS, fasting blood sugar; HCV, hepatitis C virus; IFN, interferon; NS, not significant; SVR, sustained virological response.

therapy in patients with HCV-related cirrhosis.^{10–13} In addition, the incidence of hepatocarcinogenesis is reduced in patients with SVR to IFN.¹⁷

Previous studies of patients with chronic hepatitis C indicated that HCV genotype and HCV-RNA level are the most significant factors that contribute to SVR and it is assumed that patients with genotype 1b and high viral load respond poorly to IFN therapy.⁷ In addition, IFN is less effective in patients with advanced fibrosis than those without.^{7,18} In the present study, we retrospectively reviewed the efficacy of IFN therapy and subsequent cancer prevention effect among patients with HCV-related compensated cirrhosis. Patients with both HCV

genotype 1b and high viral load were excluded because the rate of HCV elimination among these patients was extremely low and the anticancer effect of such therapy was expected to be low.

In our study, even in patients in whom fibrosis progressed to liver cirrhosis, the SVR ratio was 42.9% with genotype 1b-low HCV concentration, 25.8% with genotype 2a-high HCV concentration, 74.3% with genotype 2a-low HCV concentration, and 23.5% with genotype 2b-high HCV concentration. Thus, the therapeutic efficacy of IFN was approximately equal for liver cirrhosis and chronic hepatitis. Univariate analysis identified four parameters that significantly influenced the SVR. These

Table 4 Results of multivariate analysis for SVR to IFN for HCV-related cirrhosis

Factors	Category	Risk ratio (95% CI)	P-value
HCV-RNA level	1: low viral load	6.99 (2.72–17.9)	< 0.001
	2: high viral load	1	
Daily dose of IFN	1: ≥ 6 MU	2.62 (1.04–6.67)	0.042
	2: < 6 MU	1	

High viral load > 100 KIU/mL or > 1 Meq/mL; low viral load < 100 KIU/mL or < 1 Meq/mL.

CI, confidence interval; HCV, hepatitis C virus; IFN, interferon; SVR, sustained virological response.




Figure 1 Cumulative rate of development of hepatocellular carcinoma (HCC) in patients treated with interferon. The rates of development of HCC in sustained virological response (SVR) patients at 5 years (5.8%) and 10 years (10.3%) were significantly lower than the respective rates in non-SVR patients (25.8%, 42.5%, $P = 0.007$). —, SVR; ·····, non-SVR.

were viral load, daily dose of IFN, presence of induction therapy and choline esterase.

According to a previous study, the SVR rate was higher among patients who received long-term and high daily dose of IFN.¹³ Although the duration of IFN

therapy was marginally longer in that study compared to the present study, the results were almost similar to those reported here. In addition, as the advantage of induction therapy among patients with chronic hepatitis has been emphasized in Japan, our result indicates that induction therapy is also important for cirrhotic patients. While the total dose of IFN was also reported previously to correlate with SVR,¹⁹ no significant effect for the dose was recognized in our study. Because the design of our study was not prospective, the treatment schedule varied among patients, and therefore many biases may exist in our results. Further prospective studies are needed to confirm our findings. Multivariate analysis showed that viral load (< 100 KIU/mL) independently influenced SVR, and the SVR rate was as high as 63.2% in those patients with low viral load. As patients with genotype 1b-high viral load were not included in our study, HCV genotype was not significant. When we compared only patients with low viral load, 9/21 (42.9%) patients with genotype 1b, 26/35 patients (74.3%) with genotype 2a, and one patient with genotype 2b showed SVR. Thus, the therapeutic effect of IFN in patients with genotype 2 was significantly high ($P = 0.023$).

Table 5 Results of univariate analysis for development of hepatocellular carcinoma after IFN therapy

Factor	Category	HCC (+) <i>n</i> = 22	HCC (-) <i>n</i> = 97	<i>P</i> -value
Age	> 56 years	16 (72.7%)	50 (51.5%)	0.080
Sex	Male	16 (72.7%)	63 (64.9%)	NS
BMI	≥ 25 kg/m ²	4 (18.2%)	32 (33.0%)	NS
Albumin	> 4.0 g/dL	3 (13.6%)	49 (50.5%)	0.043
AST	> 76 IU/L	11 (50.0%)	47 (48.5%)	NS
ALT	> 100 IU/L	11 (50.0%)	44 (45.4%)	NS
Cholesterol	> 160 mg/dL	13 (59.1%)	49 (50.5%)	NS
Choline esterase	> 1.0 ΔpH	3 (14.3%)	40 (42.1%)	0.009
AFP	> 20 μg/L	9 (42.9%)	31 (32.6%)	NS
FBS	> 126 mg/mL	2 (9.5%)	13 (14.4%)	NS
Platelets	> 10×10^4 /mL	11 (50.0%)	47 (48.5%)	NS
HCV genotype	1b	16 (72.7%)	81 (83.5%)	NS
HCV-RNA level	High	10 (45.5%)	51 (52.6%)	NS
IFN efficacy	SVR	3 (14.3%)	43 (52.4%)	0.007
Total dose of IFN	≥ 400 MU	10 (45.5%)	44 (48.9%)	NS
Daily dose of IFN	> 6 MU	8 (36.4%)	48 (52.7%)	0.100
Duration of IFN	≥ 52 weeks	7 (31.9%)	22 (22.7%)	NS
Induction therapy	Yes	8 (36.4%)	50 (54.9%)	NS
Type of IFN	Alpha	15 (68.2%)	65 (67.0%)	NS
Combination with ribavirin	Yes	0 (0.0%)	10 (10.3%)	NS

High viral load ≥ 100 KIU/mL or ≥ 1 Meq/mL.

AFP, alpha fetoprotein; ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index; FBS, fasting blood sugar; HCV, hepatitis C virus; IFN, interferon; NS, not significant; SVR, sustained virological response.

Table 6 Results of multivariate analysis for development of hepatocellular carcinoma after IFN therapy

Factor	Category	Hazard ratio (95% CI)	P-value
IFN efficacy	1: SVR	0.185 (0.042–0.810)	0.025
	2: non-SVR	1	

CI, confidence interval; IFN, interferon; SVR, sustained viral response.

Our results showed that the incidence of HCC increased in a time-dependent manner after induction of IFN (16.8% at the fifth year, 29.1% at the 10th year, and 34.2% at the 15th year). Compared with the rates in Child A cirrhosis patients in our institution who had not been treated with IFN (32.5, 59.6 and 77.4%, respectively),¹ the rate of hepatocarcinogenesis after induction of IFN was half. These results suggest that hepatocarcinogenesis is significantly reduced in patients treated with IFN.

Our retrospective analysis of 123 patients with HCV-related cirrhosis who received IFN treatment showed that the outcome of IFN therapy was significantly associated with carcinogenesis. SVR was associated with a significantly reduced risk of carcinogenesis (hazard ratio: 0.185) compared with non-SVR. Three patients with SVR developed HCC; Case 1 with genotype 2a-low viral load showed SVR after 6 weeks of 3MU IFN- β therapy, but developed HCC 28 months later. Case 2 with genotype 2a-low viral load showed SVR after 6 weeks of 3MU IFN- β therapy but developed HCC 5 years later. Case 3 was also genotype 2a-low viral load who developed HCC at the end of 24 months of 6 MU IFN- α three times per week therapy. According to our previous study, the tumor-doubling time of small HCC ranges from 227 to 607 days (median, 392).²⁰ The tumor diameter at the time of diagnosis of HCC was 20 mm in Case 1, 26 mm in Case 2 and 10 mm in Case 3. Therefore, it is not clear whether hepatocarcinogenesis developed before or after eradication of HCV.

There was no relationship between the duration of IFN therapy and the rate of hepatocarcinogenesis. Likewise, the duration of IFN therapy and the total dose of IFN did not correlate with hepatocarcinogenesis. It should be noted that 38 patients had been treated with IFN prior to the present study and, thus, we cannot exclude the influence of previous IFN therapy on the development of HCC. Unfortunately, we could not collect data on the type, dose or duration of IFN for these patients because such treatment was carried out in other hospitals. To compensate for this, we analyzed those 85 patients who received IFN for the first time. In

these patients, neither the duration of IFN therapy nor the total dose of IFN was significantly associated with the development of HCC.

Our results indicated that long-term IFN therapy in non-SVR patients does not seem to improve long-term prognosis, such as the development of HCC and progression to decompensation. We reported previously that the rate of hepatocarcinogenesis was approximately equal in non-responders and untreated patients with genotype 1b-high virus load.²¹ However, the rate was lower in BR cases who were maintained on long-term IFN therapy.²¹ In three patients who developed HCC during IFN therapy in the present study, ALT normalization was noted after giving twice or thrice per week IFN- α (3MU), a therapeutic course aimed at reducing the risk of hepatocarcinogenesis. Although the duration of IFN therapy was 3.3, 4.2 and 5.3 years and serum ALT was almost below the upper limit of normal, HCC appeared in these patients during the course of IFN therapy. It is difficult to conclude from the present study that SVR is the only factor that reduced the risk of hepatocarcinogenesis. Although the result of statistical analysis was not significant in our patients, hepatocarcinogenesis was less frequent in patients with BR than in those with NR (the rate of HCC development at 5 years was 17.5% vs 29.4%, respectively, data not shown). A larger study is needed to determine the impact of BR on hepatocarcinogenesis among non-SVR patients.

In the present study, only seven of 123 patients (5.7%) discontinued IFN therapy. None developed life-threatening serious adverse effects. This outcome is probably similar to that reported in chronic hepatitis.²¹ However, careful follow up is necessary in such cirrhotic patients including liver function tests, platelet and leukocyte counts.

In summary, the SVR rate to IFN therapy was 48.0% in patients with HCV-related compensated cirrhosis except for those patients infected with genotype 1b-high virus load. The SVR was especially high (63.2%) among patients with low viral loads. The risks of hepatocarcinogenesis decreased to almost 50% in all patients treated with IFN and to only 20% among patients with SVR.

Therefore, IFN therapy is strongly recommended for patients with compensated cirrhosis with genotype 1b-low viral load or genotype 2.

REFERENCES

- 1 Kobayashi M, Ikeda K, Hosaka T *et al.* Natural history of compensated cirrhosis in the child-pugh class A compared between 490 patients with hepatitis C and 167 with B virus infections. *J Med Virol* 2006; 78: 459–65.
- 2 Ikeda K, Saitoh S, Suzuki Y *et al.* Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: a prospective observation of 2215 patients. *J Hepatol* 1998; 28: 930–8.
- 3 Davis GL, Balart LA, Schiff ER *et al.* Treatment of chronic hepatitis C with recombinant interferon alfa. A multicenter randomized, controlled trial. *N Engl J Med* 1989; 321: 1501–6.
- 4 Di Bisceglie AM, Martin P, Kassianides C *et al.* Recombinant interferon alfa therapy for chronic hepatitis C. A randomized, double-blind, placebo-controlled trial. *N Engl J Med* 1989; 321: 1506–10.
- 5 Chayama K, Saitoh S, Arase Y *et al.* Effect of interferon administration on serum hepatitis C virus RNA in patients with chronic hepatitis C. *Hepatology* 1991; 13: 1040–3.
- 6 Yoshioka K, Kakumu S, Hyashi H *et al.* Anti-hepatitis C antibodies in patients with chronic non-A, non B hepatitis: relation to disease progression and effect of interferon alpha. *Am J Gastroenterol* 1991; 86: 1495–9.
- 7 Tsubota A, Chayama K, Ikeda K *et al.* Factors predictive of response to interferon-alpha therapy in hepatitis C virus infection. *Hepatology* 1994; 19: 1088–94.
- 8 Hagiwara H, Hayashi N, Mita E *et al.* Quantitative analysis of hepatitis C virus RNA in serum during interferon alpha therapy. *Gastroenterology* 1993; 104: 877–83.
- 9 Kasahara A, Hayashi N, Mochizuki K *et al.* Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. Osaka Liver Disease Study Group. *Hepatology* 1998; 27: 1394–402.
- 10 Nishiguchi S, Kuroki T, Nakatani S *et al.* Randomised trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 1995; 346: 1051–5.
- 11 International Interferon-alpha Hepatocellular Carcinoma Study Group. Effect of interferon alpha on progression of cirrhosis to hepatocellular carcinoma: a retrospective cohort study. *Lancet* 1998; 351: 1535–9.
- 12 Hu KQ, Tong MJ. The long-term outcomes of patients with compensated hepatitis C virus-related cirrhosis and history of parenteral exposure in the United States. *Hepatology* 1999; 29: 1311–16.
- 13 Ikeda K, Saitoh S, Koboyashi M *et al.* Long-term interferon therapy for 1 year or longer reduces the hepatocellular carcinogenesis rate in patients with liver cirrhosis caused by hepatitis C virus: a pilot study. *J Gastroenterol Hepatol* 2001; 16: 406–15.
- 14 Fattovich G, Giustina G, Degos F *et al.* Morbidity and mortality in compensated cirrhosis type C: a retrospective follow-up study of 384 patients. *Gastroenterology* 1997; 112: 463–72.
- 15 Valla DC, Chevallier M, Marcellen P *et al.* Treatment of hepatitis C virus-related cirrhosis: a randomized, controlled trial of interferon alfa-2b versus no treatment. *Hepatology* 1999; 29: 1870–5.
- 16 Hashimoto M, Chayama K, Tsubota A *et al.* Typing six major hepatitis C virus genotypes by polymerase chain reaction using primers derived from nucleotide sequences of the NS5 region. *Int Hepatol Commun* 1996; 4: 263–7.
- 17 Camma C, Giunta M, Andreone P, Craxi A. Interferon and prevention of hepatocellular carcinoma in viral cirrhosis: an evidence-based approach. *J Hepatol* 2001; 34: 593–602.
- 18 Kanai K, Kako M, Okamoto H. HCV genotypes in chronic hepatitis C and response to interferon. *Lancet* 1992; 339: 1543.
- 19 Akuta N, Suzuki F, Tsubota A *et al.* Efficacy of interferon monotherapy to 394 consecutive naïve cases infected with hepatitis C virus genotype 2a in Japan. *J Hepatol* 2002; 37: 831–6.
- 20 Saitoh S, Ikeda K, Koida I *et al.* Serial hemodynamic measurements in well-differentiated hepatocellular carcinomas. *Hepatology* 1995; 21: 1530–4.
- 21 Ikeda K, Arase Y, Saitoh S *et al.* Anticarcinogenic impact of interferon on patients with chronic hepatitis C: a large-scale long-term study in a single center. *Intervirology* 2006; 49: 82–90.