

30. Wu CC, Ho YZ, Ho WL, *et al.* Preoperative transcatheter arterial chemoembolization for respectable large hepatocellular carcinoma: a reappraisal. *Br J Surg* 1995; 82: 122–6.
31. Yamasaki S, Hasegawa H, Kinoshita H, *et al.* A prospective randomized trial of preventive effect of pre-operative transcatheter arterial embolization against recurrence of hepatocellular carcinoma. *Jpn J Cancer Res* 1996; 87: 206–11.
32. Kohno H, Nagasue H, Hayashi T, *et al.* Postoperative adjuvant chemotherapy after radical hepatic resection for hepatocellular carcinoma (HCC). *Hepato-gastroenterology* 1996; 43: 1405–9.
33. Ikeda K, Arase Y, Saitoh S, *et al.* Interferon beta prevents recurrence of hepatocellular carcinoma after complete resection or ablation of the primary tumor – a prospective randomized study of hepatitis C virus-related liver cancer. *Hepatology* 2000; 32: 228–32.
34. Kubo S, Nishiguchi S, Hirohashi K, *et al.* Effects of long-term postoperative interferon-alpha therapy on intrahepatic recurrence after resection of hepatitis C virus-related hepatocellular carcinoma. *Ann Intern Med* 2001; 134: 963–7.
35. Mazzaferro V, Romito R, Sciavo M, *et al.* Prevention of hepatocellular carcinoma recurrence with alpha-interferon after liver resection in HCV cirrhosis. *Hepatology* 2006; 44: 1543–54.
36. Llovet J, Ricci V, Mazzaferro V, *et al.* Randomized phase III trial of sorafenib versus placebo in patients with advanced hepatocellular carcinoma (HCC). *J Clin Oncol* 2007; 25: 1.

Original Article

Effectiveness of combination therapy of splenectomy and long-term interferon in patients with hepatitis C virus-related cirrhosis and thrombocytopenia

Daisuke Morihara, Masahiro Kobayashi, Kenji Ikeda, Yusuke Kawamura, Hiromi Saneto, Hiromi Yatuji, Tetuya Hosaka, Hitomi Sezaki, Norio Akuta, Yoshiyuki Suzuki, Fumitaka Suzuki and Hiromitsu Kumada

Department of Hepatology, Toranomon Hospital, Tokyo, Japan

Aim: To elucidate the effectiveness of combination therapy of splenectomy and long-term interferon (IFN) on survival and hepatocarcinogenesis, we retrospectively analyzed 180 patients with hepatitis C virus (HCV)-related cirrhosis and thrombocytopenia.

Methods: Group A consisted of 121 patients who received neither splenectomy nor IFN therapy. Group B consisted of 11 patients who underwent splenectomy only. Group C consisted of 32 patients who underwent IFN therapy only. Group D consisted of 16 patients who received the combination therapy splenectomy followed by IFN therapy.

Results: The viral response in group D estimated at least 6 months after IFN therapy showed sustained viral response in four patients, biochemical response in one and no response in six. Multivariate analysis using time-dependent variables showed significant improvement of survival rate in patients on the combination therapy, but no effect on the appearance rate of hepatocarcinogenesis relative to the findings in group A.

Conclusions: In this study, the splenectomy did not directly improve the prognosis, but increased the ability for patients to undergo IFN. As a result, we considered that the combination therapy of splenectomy and long-term IFN significantly improved survival rate in patients with advanced HCV-related cirrhosis and thrombocytopenia.

Key words: cirrhosis, hypersplenism, interferon, splenectomy, thrombocytopenia

Abbreviations:

AFP, Alpha-fetoprotein; ALT, Alanine aminotransferase; AST, Aspartic aminotransferase; BR, biochemical response; CT, Computed tomography; HCC, Hepatocellular carcinoma; HCV, Hepatitis C virus; ICG R15, Indocyanine green retention rate at 15 min; IFN, Interferon; MELD score, Model for End-Stage Liver Disease score; NR, No response; PLT, platelet; SVR, Sustained virological response; TTT, Thymol turbidity test; US, Ultrasonography; ZTT, Zinc sulfate turbidity test.

INTRODUCTION

THE PRESENCE OF severe thrombocytopenia in patients with cirrhosis associated with hepatitis C viral (HCV) infection limits the use of interferon (IFN) therapy. The different treatment modalities for hepatocellular carcinoma (HCC), such as hepatic resection, radiofrequency ablation, or percutaneous ethanol injection, are also limited by low platelet (PLT) counts. In

patients with compensated cirrhosis and low model for end-stage liver disease (MELD) score, liver transplantation is not warranted and the use of antiviral therapy to slow down the progression to liver failure is not recommended. In other words, such patients are too healthy for transplantation and too thrombocytopenic to treat with antiviral agents. Splenectomy has been suggested for the treatment of secondary hypersplenism and thrombocytopenia as a means to improve PLT count.¹

If patients with HCV-related cirrhosis and thrombocytopenia could receive the benefits of splenectomy^{2,3} and IFN therapy,^{4,5} such therapy would clinically be very useful. The combination therapy of splenectomy and long-term IFN administration may improve survival rate and reduce the incidence of hepatocarcinogenesis.

Correspondence: Dr Masahiro Kobayashi, Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-8470, Japan. Email: mshkobayashi@toranomon.gr.jp
Received 2 February 2008; revised 4 November 2008; accepted 17 November 2008.

However, there are only a few reports that have examined the usefulness of this combination therapy in patients with advanced HCV-related cirrhosis and low PLT count.⁶ In this study, we retrospectively analyzed 180 patients with compensated cirrhosis and thrombocytopenia who had received the combination therapy of splenectomy and long-term IFN to determine the effects of such treatment on the survival rate and incidence of HCC.

PATIENT AND METHODS

Study population

A TOTAL OF 180 Japanese patients with cirrhosis, hypersplenism and low PLT count ($\leq 80 \times 10^3/\mu\text{L}$) were examined between 1990 and 2006. Their initial sera were positive for antibodies to HCV (anti-HCV; second-generation anti-HCV kit; ELISA, Dainabot, Tokyo, Japan), positive HCV-RNA (Amplicor HCV monitor assay version 2.0; Roche Diagnostics, Tokyo, Japan), and negative for hepatitis B surface antigen (HBsAg; radioimmunoassay, Dainabot). Anti-HCV was assayed using stored frozen sera at -80°C . They were diagnosed with liver cirrhosis between 1990 and 2006 at Toranomon Hospital, Tokyo, Japan. In addition to liver biopsy and/or peritoneoscopy, liver cirrhosis was also diagnosed utilizing clinical findings (e.g. presence of esophageal varices), and with computed tomographic (CT) or ultrasonographic (US) findings. The following protocol was applied in our hospital until 2000: Patients with a platelet count of less than $50 \times 10^3/\mu\text{L}$ are eligible for HCC surgery (such as hepatic resection, radiofrequency ablation, or percutaneous ethanol injection) provided they receive platelet transfusion. The decision to pursue splenectomy was individualized and based on the presence thrombocytopenia and/or intractable gastric varices, and discussed with the patients.

We retrospectively analyzed the effect of splenectomy on cirrhotic patients with low PLT count ($\leq 80 \times 10^3/\mu\text{L}$). Of the total 180 patients, 121 (67.2%) patients received neither antiviral therapy nor splenectomy (group A). Thirty-two (17.8%) patients received only IFN therapy (group C). The remaining 27 (15.0%) patients underwent splenectomy (11 patients underwent only splenectomy [group B] and 16 received IFN therapy after splenectomy [group D]). Splenectomy was performed for the following reasons; (i) low PLT count in 20 patients (six [54.5%] of group B and 14 [8.5%] of group D), (ii) low PLT count and part of treatment of gastric varices in three (one [9.0%] of group B and two

[12.5%] of group D), and (iii) low PLT count and refractory esophageal varices in four (four [36.4%] of group B). None of the patients required emergency splenectomy (e.g. bleeding gastric varices or other bleeding complications related to low platelet count). Our institution does not require informed consent for retrospective analysis.

Patients background and laboratory data

Table 1 summarizes the profiles and patients of groups A, B, C and D at the time of diagnosis of liver cirrhosis. Indocyanine green test was conducted in 91.2% of the patients. Patients of group D had significantly lower PLT count ($P = 0.01$) and AST ($P = 0.01$) than patients in others groups. The proportion of group A patients who regularly consumed alcohol at ≥ 80 g/day was significantly higher than other groups. Patients of group C had significantly lower TTT ($P = 0.08$) than others.

Splenectomy

Splenectomy was performed through midline or left subcostal incision depending on body habitus and previous incisions. For group B, five patients underwent splenectomy and six underwent Hassab's operation.⁷ In group D, 13 patients underwent splenectomy and three underwent Hassab's operation.

IFN treatment

Thirty-two patients received IFN therapy (group C). In group C, 21 patients received 3 million units of IFN- α (natural or recombinant) intramuscularly three times per week to maintain a low alanine aminotransferase (ALT), 11 patients received 6 million units of IFN- α to eradicate HCV. Patients of group C received IFN therapy for a median period of 0.5 years (range, 0.0–9.7 years).

Sixteen patients received the combination therapy (group D). Of these, 12 (75%) patients underwent splenectomy for the purpose of induction of antiviral therapy with IFN. The other patients (25%) had undergone splenectomy pre dating this study. In group D, 11 patients (Cases 1–4, 8, 10–13, 15–16) received 3 million units of IFN- α (natural or recombinant) intramuscularly three times per week to maintain a low ALT, 3 patients (Cases 6, 7, and 9) received 6 million units of IFN- α to eradicate HCV. For the other two patients; one (Case 5) received pegylated IFN α 2b (50 μg) monotherapy and the other patient (Case 14) received pegylated IFN α 2b (50 μg) plus ribavirin (400 mg) combination therapy to maintain low ALT (Fig. 1). Patients of group D received IFN therapy for a median period of 1.4 years (range, 0.2–12.4 years).

Table 1 Patient profiles and laboratory data at the time of diagnosis of cirrhosis

	Group A (Neither splenectomy nor IFN)	Group B (splenectomy)	Group C (IFN)	Group D (splenectomy + IFN)	P*
Demography					
No. patients	121	11	32	16	
Sex (M/F)	64/57	6/5	13/19	13/3	0.07
Age (years)†	61 (32–82)	61 (42–66)	59 (36–72)	52 (36–60)	0.41
Alcohol intake of 80 g/day or more	29	0	10	0	0.03
Diabetes mellitus	12	1	4	2	0.96
Laboratory data†					
Platelet count ($\times 10^3/\mu\text{L}$)	61 (17–80)	64 (42–75)	66 (25–80)	44 (27–78)	0.01
Prothrombin activity (%)	73 (50–101)	79 (58–94)	80 (66–100)	74 (47–100)	0.88
Albumin (g/dL)	3.5 (1.7–4.8)	3.5 (2.0–4.3)	3.4 (2.5–4.1)	3.3 (2.7–4.5)	0.64
ZTT (Kunkel)	12.3 (0.7–23.3)	10.3 (3.3–18.2)	10.8 (4.4–21.0)	12.0 (6.1–17.1)	0.29
TTT (Kunkel)	14.1 (0.4–37.2)	12.0 (4.4–16.9)	7.8 (1.2–34.0)	12.7 (2.7–34.1)	0.08
Bilirubin (mg/dL)	1.5 (0.4–7.7)	1.2 (0.7–5.3)	1.1 (0.6–2.7)	1.2 (0.8–4.4)	0.03
AST (IU/L)	64 (21–652)	83 (31–157)	75 (28–216)	60 (30–154)	0.17
ALT (IU/L)	53 (11–239)	72 (24–191)	71 (18–298)	46 (14–182)	0.01
ICG R15 (%)	38 (12–96)	41 (15–64)	32 (6–62)	32 (8–53)	0.44
Alpha-fetoprotein (ng/mL)	23 (2–909)	40 (3.9–165)	29 (5–631)	11 (4–190)	0.28

ALT, alanine aminotransferase; AST, aspartic aminotransferase; ICG R15, indocyanine green retention rate at 15 min; TTT, thymol turbidity test; ZTT, zincsulfate turbidity test.

*Kruskal-Wallis test or χ^2 -test. †Expressed by median (min, max).

The effect of IFN therapy was classified according to elimination of HCV-RNA and ALT value 6 months after the end of treatment. Sustained virological response (SVR) was defined as persistent disappearance of HCV RNA after therapy, biochemical response (BR) as normal ALT values without elimination of HCV RNA for at least 6 months after therapy, and no response (NR) as persistently elevated or transiently normalized ALT levels without loss of HCV RNA.

Follow up of patients

Patients were followed up on a monthly basis after the diagnosis of cirrhosis by monitoring hematologic, biochemical, and virologic data. Imaging studies were conducted three or more times per year in the majority of patients by using computerized tomography (CT) or ultrasonography (US). Angiography was performed only when HCC was highly suspected based on CT or US. When angiography detected a typical hypervascular nodule, it was considered a specific finding for HCC in these follow-up patients, and histological confirmation was usually not required in the majority of patients. If the angiographic study did not show any hypervascular staining in a small hepatic nodule, a fine needle biopsy was performed. In this cohort, 18 (12.2%) patients were

lost to follow up [14 patients (11.6%) from group A, two patients (18.2%) from group B, one patient (3.1%) from group C and two patients (12.5%) from group D]. The date of the last follow-up in this study was 31 March 2007, and the median observation period of studied patients was 5.9 years (range, 0.1–19.6 years).

Statistical analysis

Non-parametric procedures were used for the analysis of background characteristics of the patients, including Kruskal-Wallis and χ^2 test. Changes in laboratory tests values after splenectomy were evaluated by using Wilcoxon signed-rank test. Survival rate was calculated from the period between diagnosis of liver cirrhosis and death in each group, by using the Kaplan-Meier method.⁸ HCC appearance rate was calculated from the period between diagnosis of liver cirrhosis and appearance of HCC in each group, by again using the Kaplan-Meier method. Differences in slopes of survival and carcinogenic curves were evaluated by log-rank test. The median waiting period between diagnosis of cirrhosis and splenectomy was 1.6 months (range, 0.0–199.5 months) for groups B and C. To compensate for wait-time bias in the splenectomy groups, curves of survival and HCC appearance were also drawn from the time of diagnosis

nectomy. Leukocyte count increased about 1.6 times at 6 months after splenectomy [before splenectomy, median = 3200/mm³ (range 1800–5600); after splenectomy, 5200 (3700–9000); $P < 0.001$]. PLT count increased about 2.3 times at 6 months after splenectomy [before splenectomy, median = $47 \times 10^3/\mu\text{L}$ (range, 26–77 $\times 10^3$); after splenectomy, 110×10^3 (79–275 $\times 10^3$); $P < 0.001$]. Total bilirubin decreased about 0.6 times at 6 months after splenectomy [before splenectomy, median = 1.2 mg/dL (range, 0.6–4.4); after splenectomy, 0.7 (0.4–1.8); $P = 0.001$]. Leukocyte and PLT counts reached peak levels within a month after splenectomy and were almost stabilized at six months.

Postoperative complications following splenectomy developed in three patients; hemoperitoneum ($n = 1$), portal vein thrombosis ($n = 1$) and secondary thrombocytopenia ($n = 1$). Some patients received prophylactic anticoagulation to protect against portal vein thrombosis after splenectomy. One patient with hemoperitoneum died due to multiple organ failure, while the other patients recovered with medical treatment.

Complications of splenectomy plus IFN combination therapy

Figure 1 shows patients that underwent combination therapy (group D). During the observation period, one patient (Case 3) of group D died of liver failure caused by progression of HCC. The causes of death in three other patients were not deemed to be complications related to the combination therapy. None of the patients of group D developed serious complications (e.g. portal vein thrombosis, post-operative hemorrhage, pneumonia, sepsis) from the splenectomy. Post-operatively, none of the patients showed worsening of liver biochemical test results or developed decompensated liver disease with ascites, encephalopathy, jaundice or variceal bleeding. There were also no deaths in the immediate postoperative period. Three patients (18.8%) of group D discontinued IFN therapy for the following reasons; severe thrombocytopenia (Case 1), NSAID-induced liver injury (Case 2) and peripheral neuropathy (Case 13). In contrast, eight patients (25.8%) of group C discontinued IFN therapy. Three (37.5%) of them discontinued IFN therapy due to severe thrombocytopenia. When frequency of discontinued IFN therapy was compared with group C and D, there was no significant difference ($P = 0.73$). However, there were cases, eight in group C but 0 in group D, who required a reduction in IFN dosages during treatment as compared with the beginning of treatment ($P = 0.03$).

The splenectomy could have increased the ability for patients to undergo IFN.

Effect of IFN therapy after splenectomy

Eleven of 16 (68.8%) patients of group D had HCV genotype 1b and five (31.3%) had HCV genotype 2a (Fig. 1). The viral response was determined at least 6 months after IFN therapy; SVR was noted in four (36.4%) patients, BR in one (9.1%) and NR in six (54.5%). Three patients continue to receive IFN therapy at present. In this study, patients with SVR were all male and had genotype 2a. One of the patients with SVR received pegylated-IFN α -2b (Case 5, Fig. 1), while other patients received IFN α 2b. Meanwhile, 18 of 32 (56.3%) patients of group C had HCV genotype 1b, 12 (37.5%) had HCV genotype 2a and two (6.3%) had HCV genotype 2b. Group C had more patients with low HCV-RNA ($< 100\,000$ IU/mL) than group D (12 [37.5%] of group C and three [18.8%] of group D, $P = 0.09$). In group C, SVR was noted in 7 (21.9%) patients, BR in six (18.8%) and NR in 17 (53.1%). Two patients continue to receive IFN therapy at present.

SVR were not significantly different between group C and D ($P = 0.43$). This result might be a reason that group D had more patients with HCV genotype 1 and higher HCV-RNA than group C.

Rate of hepatocarcinogenesis

During the follow-up period of up to 17 years (median observation period of 5.9 years), HCC developed in 65 patients (36.1%): 40 (33.1%) in group A, five (45.5%) in group B, 16 (50.0%) in group C and four (25.0%) in group D. HCC appearance rates at the end of the third year were 19.9, 20.0, 25.0 and 6.3% in group A, B, C and D, 28.5, 57.3, 34.5 and 14.1% at the end of the fifth year, and 48.2, 78.7, 43.8 and 39.8% at the end of tenth year, respectively (Fig. 2). There was no significant difference in the rate of HCC appearance among the four groups (log-rank test, $P = 0.42$). In particular, the HCC appearance rate in group D was not significantly different compared with group A (log-rank test, $P = 0.50$).

In addition, the rate of carcinogenesis correlated inversely with the duration of IFN administration (Fig. 1). For group D, 9 of 14 patients were treated with IFN for ≥ 12 months. The carcinogenic rate at the end of the 5th year in the remaining patients of the same group who were treated with IFN for < 12 months (20.0%) was higher than in those treated for ≥ 12 months (9.1%). Multivariate analysis showed that the hazard ratio of carcinogenesis for patients treated with IFN for




Figure 2 Crude hepatocellular carcinoma (HCC) curves in patients of groups A, B, C and D. There was no significant difference in the HCC appearance rate among the four groups (log-rank test, $P = 0.42$).

≥ 12 months was 0.022 after adjustments for significant covariates, but was not significantly different ($P = 0.43$).

We also assessed the effects of splenectomy and long-term IFN therapy on hepatocarcinogenesis by comparing patients of group D (splenectomy + IFN administration for ≥ 12 months) with those of group A. The combination therapy reduced the hazard ratio to 0.03 (multivariate analysis with adjustments for significant covariates), though it was significant ($P = 0.83$). We also assessed compared patients of groups C and B (splenectomy alone). Administration of IFN for ≥ 12 months reduced the hazard ratio to 0.03 (multivariate analysis after adjustments for significant covariates), but was not significant ($P = 0.83$). These results suggest that the combination of splenectomy plus long-term IFN decreased the likelihood of hepatocarcinogenesis.

Effect of splenectomy and IFN combination therapy on survival

During the observation period, one of the 16 patients of group D (Case 3) died (Fig. 1). The survival rates for groups A, B, C and D were 84.2, 90.9, 87.5 and 100% at the end of the third year, 72.0, 90.9, 87.5 and 100% at the fifth year, 41.4, 36.4, 83.3 and 83.3% at the tenth year, respectively (Fig. 3). The survival rate for patients of group D was the highest compared with the other groups (log-rank test, $P = 0.002$). We also compared the effect of combination therapy on the survival rate of

patients of group A and group D. The survival rate of group D was significantly higher than of group A (log-rank test, $P = 0.004$). We also compared the effect of combination therapy on the survival rate of patients of group C and group D. The survival rate of group D was not significantly different compared with group C (log-rank test, $P = 0.29$). The combination therapy significantly improved the hazard ratio of survival to 9.69 ($P = 0.028$, multivariate analysis with adjustments for significant covariates, Table 2). These results suggest that the splenectomy simply increased the ability for patients to undergo IFN and may not directly improve patient survival.

DISCUSSION

CHRONIC HEPATITIS C virus (HCV) will continue to cause significant morbidity and mortality through to at least 2015.¹⁰ HCV infection remains a common cause of chronic liver disease and is an increasing indication for liver transplantation. Thrombocytopenia (platelet counts $< 150 \times 10^3/\mu\text{L}$) is a common complication in patients with chronic liver disease (CLD), and is reported in as many as 76% of cirrhotic patients.¹¹ The ability to increase platelet levels could significantly reduce the need for platelet transfusions and facilitate the use of IFN-based antiviral therapy and other medically indicated treatments in patients with liver disease. Current treatment options for severe




Figure 3 Survival rates for patients of groups A, B, C and D. The survival rate was significantly different for group A, B, C and D (log-rank test, $P = 0.002$). The survival rate of patients of group D was significantly higher than that of group A (log-rank test, $P = 0.004$).

thrombocytopenia include platelet transfusion, splenic artery embolization and splenectomy. We studied the usefulness of the combination therapy of splenectomy and long-term IFN in patients with advanced HCV-related cirrhosis and thrombocytopenia.

With regard to the usefulness of splenectomy, some studies reported that splenectomy improved PLT counts in cirrhotic patients with thrombocytopenia.^{2,3} Furthermore, Shimada *et al.*¹² reported that splenectomy resulted in significant falls in ammonia levels and rises in serum albumin. Thus, there is evidence that splenectomy is beneficial and results in recovery of liver function by improving of blood supply to the liver.^{6,13} In the present study, at 6 months after splenectomy, leukocyte count increased 1.6 times, PLT count increased 2.3 times, and total bilirubin decreased nearly 0.6 times,

relative to prior the procedure. Furthermore, liver function test results also improved in most patients with splenectomy.

With regard to the value of IFN therapy after splenectomy, Hayashi *et al.*⁶ reported that splenectomy in patients with HCV cirrhosis can be done safely to allow application of antiviral treatment and potentially avoid transplantation.⁶ In this study, only three of 16 (18.8%) patients discontinued IFN therapy after splenectomy. Among the three patients, IFN therapy was discontinued because of thrombocytopenia in only one (6.3%) patient. On the other hand, 13 (81.3%) of the 16 patients on combination therapy were able to complete the full course of IFN therapy, continue IFN therapy or stopped therapy due to NR. Thus, it may be said that IFN therapy is safe in most patients with advanced HCV-related cirrhosis and thrombocytopenia. Furthermore, the present results indicate that splenectomy is an effective method in patients with chronic HCV infection and hypersplenism to increase peripheral leukocyte and platelet counts so that subsequent IFN therapy can be better tolerated. In this study, regarding the reduction of IFN dosages during treatment when comparing group C and D, group D did not have any cases who a reduction in IFN dosages was necessitated by thrombocytopenia ($P = 0.03$). Hayashi *et al.*⁶ reported that five of their seven patients underwent splenectomy and then completed a full course of pegylated IFN and ribavirin

Table 2 Significance of combined therapy of survival rate in patients of advanced hepatitis C virus-related cirrhosis with low platelet count (time-dependent proportional hazard model)

Factors	Category	Hazard ratio (95% CI)	P
Combined therapy (splenectomy + IFN)	1: no	1	0.028
	2: yes	9.69 (1.28–76.9)	

IFN, interferon therapy.

treatment or stopped therapy due to NR, and that none of their patients required dose reductions or treatment discontinuation due to thrombocytopenia. In the present study, the viral response to IFN therapy was SVR in four (36.4%) patients, BR in one (9.1%) and NR in six (54.5%). SVR was not significantly different between group C and D ($P = 0.43$). This result might be a reason that group D had more patients with HCV genotype 1 and higher HCV-RNA than group C. All patients with SVR of group D had genotype 2, suggesting that SVR seems to be achievable by combination therapy in patients with HCV-related cirrhosis with genotype 2 and thrombocytopenia.

We also analyzed the effect of the combination therapy on hepatocarcinogenesis in patients with advanced HCV-related cirrhosis and low PLT count. Chen *et al.*¹⁴ reported that the 5-year tumor-free survival rate was significantly higher after hepatectomy and splenectomy than after hepatectomy alone (37 vs. 27.3%, respectively, $P = 0.003$). In contrast, Yao *et al.*¹⁵ reported that splenectomy in early stage of tumor inoculation stimulated tumor growth and metastasis in their rat model of HCC.¹⁵ In this study, the HCC appearance rate in patients who underwent splenectomy alone (group B) was not significantly different from that of the control (log-rank test, $P = 0.52$). In addition, the HCC appearance rate in patients who received the combination therapy was also not significantly different from the control (log-rank test, $P = 0.50$). We previously reported that long-term IFN therapy for 12 months or longer reduced the rate of hepatocarcinogenesis in patients with liver cirrhosis caused by HCV.⁵ Multivariate analysis of long-term follow-up showed that the combination therapy, including IFN administration for ≥ 12 months, decreased the hazard ratio of hepatocarcinogenesis to 0.03, though this was not significant ($P = 0.83$). The reason for the lack of significance might be the small population sample of this study. Yoshida *et al.*¹⁶ reported that IFN therapy significantly reduced the risk for HCC, especially among virologic and biochemical responders. That the combination therapy decreased the hazard ratio of hepatocarcinogenesis to 0.03 suggests the ability of long-term IFN to inhibit HCC, especially among non-responders.

We also examined the effects of the combination therapy on survival. In this study, multivariate analysis using time-dependent variables showed significant improvement of survival in patients who received the combination therapy (group D) compared with the control group (group A) (hazard ratio 3.40, $P = 0.017$; 95% CI 1.24–9.35). This may be considered the crucial

finding of this study. In splenectomy, Morimasa *et al.*¹⁷ reported no difference in survival rate between splenectomy and endoscopic injection sclerotherapy (EIS) for esophageal varices. Similarly, the survival rate in the splenectomy group in this study (group B) was not significantly different from the control ($P = 0.88$). Furthermore, the survival rate of group D was not significantly different compared with group C (log-rank test, $P = 0.29$). These results suggest that the splenectomy increased the ability for patients to undergo IFN and that the combination therapy of splenectomy and long-term IFN significantly improved survival rate in patients with advanced HCV-related cirrhosis and thrombocytopenia. The likely mechanism of action of the combination therapy is first improvement of leucopenia and thrombocytopenia following splenectomy, which allowed administration of IFN, and then IFN produced remission of liver fibrosis, control of necroinflammatory process, and induced suppression of the HCC growth process, consequently leading to improvement of survival rate. Moreno and Muriel¹⁸ reported that IFN resulted in remission of liver fibrosis, and that control of the necroinflammatory process can therefore induce suppression of the HCC growth process. Our results also suggested that patients with NR may need to continue the combination therapy with long-term IFN therapy.

"Pegylated IFN plus ribavirin" and "eltrombopag" are promising drugs and can be potentially used in combination therapy. Recent multicenter trials have demonstrated the superiority of pegylated IFN plus ribavirin compared to pegylated IFN alone or non-pegylated combination therapy.^{19,20} In addition, several promising novel agents that stimulate TPO and increase PLT count, such as the oral platelet growth factor eltrombopag, are currently in development for the prevention and/or treatment of thrombocytopenia.²¹ Eltrombopag may be a substitute for splenectomy or PSE. Thus, combination therapy of pegylated IFN plus ribavirin after splenectomy or eltrombopag may improve survival rate and reduce the rate of hepatocarcinogenesis.

Our study had certain limitations. In particular, in this study, four (25%) of the patients who underwent combination therapy had a history of splenectomy. A randomized control study with a larger number of cases should be conducted to confirm the effectiveness of this therapy.

In conclusion, the combination therapy of splenectomy and long-term IFN decreased the rate of hepatocarcinogenesis and significantly improved the survival rate in patients with advanced HCV-related cirrhosis and low PLT count.

ACKNOWLEDGMENTS

THE PRESENT STUDY was supported in part by research on Hepatitis from the Ministry of Health, Labour and Welfare of Japan.

REFERENCES

- 1 Coon WW. Splenectomy for thrombocytopenia due to secondary hypersplenism. *Arch Surg* 1988; 123: 369-71.
- 2 Kishi Y, Sugawara Y, Akamatsu N *et al*. Splenectomy and preemptive interferon therapy for hepatitis C patients after living-donor liver transplantation. *Clin Transpl* 2005; 19: 769-72.
- 3 Cescon M, Sugawara Y, Takayama T *et al*. Role of splenectomy in living-donor liver transplantation for adults. *Hepatogastroenterology* 2002; 49: 721-3.
- 4 Nishiguchi S, Kuroki T, Nakatani S *et al*. Randomised trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 1995; 346: 1051-5.
- 5 Ikeda K, Saitoh S, Kobayashi M *et al*. Long-term interferon therapy for 1 year or longer reduces the hepatocellular carcinogenesis rate in patients with liver cirrhosis caused by hepatitis C virus: a pilot study. *J Gastroenterol Hepatol* 2001; 16: 406-15.
- 6 Hayashi PH, Mehia C, Joachim Reimers H, Solomon HS, Bacon BR. Splenectomy for thrombocytopenia in patients with hepatitis C cirrhosis. *J Clin Gastroenterol* 2006; 40: 740-4.
- 7 Hassab MA, Younis MT, el-Kilany MS. Gastroesophageal decongestion and splenectomy in the treatment of esophageal varices secondary to bilharzial cirrhosis: anatomical and experimental studies. *Surgery* 1968; 63: 731-7.
- 8 Kaplan EL. Nonparametric estimation for incomplete observation. *J Am Stat Assoc* 1958; 53: 457-81.
- 9 Cox DR. Regression models and life tables. *J R Stat Soc* 1972; 34: 248-75.
- 10 National Institutes of Health Consensus Development Conference Statement: management of hepatitis C: 2002 - June 10-12, 2002. *Hepatology* 2002; 36: S3-20.
- 11 Giannini EG. Review article: thrombocytopenia in chronic liver disease and pharmacologic treatment options. *Aliment Pharmacol Ther* 2006; 23: 1055-65.
- 12 Shimada M, Hashizume M, Shirabe K, Takenaka K, Sugimachi K. A new surgical strategy for cirrhotic patients with hepatocellular carcinoma and hypersplenism. Performing a hepatectomy after a laparoscopic splenectomy. *Surg Endosc* 2000; 14: 127-30.
- 13 Kercher KW, Carbonell AM, Heniford BT, Matthews BD, Cunningham DM, Reindollar RW. Laparoscopic splenectomy reverses thrombocytopenia in patients with hepatitis C cirrhosis and portal hypertension. *J Gastrointest Surg* 2004; 8: 120-6.
- 14 Chen XP, Wu ZD, Huang ZY, Qiu FZ. Use of hepatectomy and splenectomy to treat hepatocellular carcinoma with cirrhotic hypersplenism. *Br J Surg* 2005; 92: 334-9.
- 15 Yao YM, Liu QG, Yang W, Zhang M, Ma QY, Pan CE. Effect of spleen on immune function of rats with liver cancer complicated by liver cirrhosis. *Hepatobiliary Pancreat Dis Int* 2003; 2: 242-6.
- 16 Yoshida H, Shiratori Y, Moriyama M *et al*. Interferon therapy reduces the risk for hepatocellular carcinoma: national surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy. *Ann Intern Med* 1999; 131: 174-81.
- 17 Tomikawa M, Hashizume M, Akahoshi T *et al*. Effects of splenectomy on liver volume and prognosis of cirrhosis in patients with esophageal varices. *J Gastroenterol Hepatol* 2002; 17: 77-80.
- 18 Moreno MG, Muriel P. Remission of liver fibrosis by interferon-alpha 2b. *Biochem Pharmacol* 1995; 50: 515-20.
- 19 Manns MP, McHutchison JG, Gordon SC *et al*. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001; 358: 958-65.
- 20 Fried MW, Shiffman ML, Reddy KR *et al*. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002; 347: 975-82.
- 21 McHutchison JG, Dusheiko G, Shiffman ML *et al*. Eltrombopag for thrombocytopenia in patients with cirrhosis associated with hepatitis C. *N Engl J Med* 2007; 357: 2227-36.

GASTROENTEROLOGY

Efficacy of entecavir treatment for lamivudine-resistant hepatitis B over 3 years: Histological improvement or entecavir resistance?

Yoshiyuki Suzuki,* Fumitaka Suzuki,* Yusuke Kawamura,* Hiromi Yatsuji,* Hitomi Sezaki,* Tetsuya Hosaka,* Norio Akuta,* Masahiro Kobayashi,* Satoshi Saitoh,* Yasuji Arase,* Kenji Ikeda,* Mariko Kobayashi,[†] Yuzo Miyakawa[‡] and Hiromitsu Kumada*

*Department of Hepatology, [†]Research Institute for Hepatology, Toranomon Hospital, and [‡]Miyakawa Memorial Research Foundation, Tokyo, Japan

Key words

entecavir, chronic hepatitis, hepatitis B virus, lamivudine, YMDD mutants.

Accepted for publication 21 November 2008.

Correspondence

Dr Yoshiyuki Suzuki, Department of Hepatology, Toranomon Hospital, 1-3-1 Kajigaya, Takatsu-ku, Kawasaki City 213-8587, Japan. Email: suzunari@interlink.or.jp

Abstract

Background and Aims: Long-term lamivudine therapy is required for patients with chronic hepatitis B, because hepatitis reappears frequently after it has withdrawn. However, hepatitis B virus (HBV) mutants resistant to lamivudine emerge frequently accompanied by breakthrough hepatitis.

Methods: Effects of entecavir were evaluated in 19 patients who had developed breakthrough hepatitis during lamivudine therapy for longer than 5 years. This study is a subgroup analysis of a previously reported study. Entecavir, in either 0.5 or 1.0 mg/day doses, was given to 10 and nine patients for 52 weeks, respectively, and then all received 1.0 mg/day entecavir for an additional 68–92 weeks.

Results: There were no differences in biochemical and virological responses in the two groups of patients with respect to the two different initial doses of entecavir. Serum levels of alanine aminotransferase were normalized in 17 (90%) patients, and hepatitis B e antigen (HBeAg) disappeared from the serum in two (14%) of the 14 patients who were HBeAg-positive before. Furthermore, a decrease in histological activity index score greater than 2 points was achieved in nine of the 11 (82%) patients in whom annual liver biopsies were performed during 3 years while they received entecavir. HBV mutants resistant to entecavir emerged in five of the 19 (26%) patients, and hepatitis flare occurred in two of them (40%).

Conclusion: Entecavir in the long term would be useful for histological improvement of breakthrough hepatitis induced by lamivudine-resistant HBV mutants in patients with chronic hepatitis B. However, the relatively high rate of entecavir resistance is a concern, and other strategies need to be considered when available.

Introduction

Worldwide, an estimated 400 million people are infected with hepatitis B virus (HBV) persistently, and some of them develop fatal liver disease, such as decompensated cirrhosis and hepatocellular carcinoma.¹ In 1995, lamivudine was introduced to the treatment of chronic hepatitis B for which interferon (IFN) had previously been the only option.^{2,3} Although lamivudine is efficient for treatment of chronic hepatitis B, drug-resistant HBV variants with mutations in the tyrosine–methionine–aspartate–aspartate (YMDD) motif occur increasingly more frequently with treatment duration, to higher than 60% within 5 years.^{4–7} Furthermore, these YMDD mutants are often accompanied by breakthrough hepatitis, and it is difficult to obtain disease control with lamivudine.

Subsequently, adefovir dipivoxil has been approved for treatment of chronic hepatitis B,^{8,9} and more recently entecavir.^{10–12} Entecavir is superior to lamivudine as the first-line treatment, and both adefovir add-on lamivudine and entecavir as switch therapy have also been employed for treatment of breakthrough.^{13,14}

The present study represent a subgroup analysis of our previously reported multicenter randomized controlled trial.¹² From a single center, biological and virological responses to entecavir were examined among 19 patients who had developed hepatitis breakthrough during long-term lamivudine therapy, with particular focus on histological responses to entecavir over 3 years and the rate of development of entecavir resistance. Because patients had been randomized to both the low (0.5 mg) and higher (1.0 mg) doses of entecavir, we were also able to compare results between these two different doses.

Methods

Patients

During 10 years from November 1995 to December 2004, 704 patients with chronic hepatitis B received 100 mg lamivudine/day and were followed for more than 5 years in the Department of Hepatology of Toranomon Hospital in metropolitan Tokyo. Lamivudine-resistant YMDD mutants developed in 274 (39%) of the patients, accompanied by breakthrough hepatitis in 176 (64% of those with mutants). Medication was changed so they received the other antivirals. The present study is a subgroup analysis of our previously reported multicenter randomized controlled trial.¹² After entecavir became available, 19 of them were switched to it and the treatment was continued for up to 3 years. None of them were infected with hepatitis C virus (HCV) or HIV type 1, or had autoimmune hepatitis. They were followed for liver function tests and serum markers of HBV infection monthly. At the start of entecavir therapy, chronic hepatitis was diagnosed in them all by liver biopsies performed under laparoscopy and/or ultrasonic imaging; cirrhosis was detected in no patients. Liver biopsies were performed annually for 3 years on 12 of the 19 (63%) patients, for evaluating the efficacy of long-term entecavir in improving histology of the liver. The study design conformed to the 1975 Declaration of Helsinki, and was approved by the ethics committee of the institution. All patients gave their informed consent to participate in this study.

Markers of HBV infection

Hepatitis B surface antigen (HBsAg) and the corresponding antibody (anti-HBs) were determined by hemagglutination (MyCell; Institute of Immunology, Tokyo, Japan), and hepatitis e antigen (HBeAg) by enzyme-linked immunosorbent assay (ELISA) (F-HBe; Sysmex, Kobe, Japan). HBV-DNA was determined by reverse transcription polymerase chain reaction (RT-PCR) with commercial kits (Amplicor, Tokyo, Japan; Roche, Tokyo, Japan), and the result was expressed in log genome equivalents (LGE)/mL with the cut-off value of 2.6 LGE/mL over a dynamic range of 2.6–7.6 LGE/mL. The six major genotypes (A–F) were determined serologically by ELISA (HBV Fenotype EIA; Institute of Immunology). The method employs the combination of epitopes on preS2-region products that is specific for each genotype.^{15,16}

Analyses for viral resistance

YMDD mutants were determined by PCR followed by restriction fragment length polymorphism after the method of Chayama *et al.*⁴ HBV mutants resistant to entecavir were examined at the baseline and sequentially while patients received entecavir. HBV-DNA was extracted from the serum and amplified by PCR, and nucleotides corresponding to amino acids 1–344 of the reverse transcriptase were sequenced directly by the dideoxy-chain method of Sanger *et al.*¹⁷

Treatment with entecavir

The 19 patients were randomized to receive two different regimens of entecavir in a double-blind study. Thus, 0.5 and 1.0 mg ente-

cavir was given daily to 10 and nine patients, respectively, for the first 52 weeks. Thereafter, patients in both groups received 1.0 mg/day entecavir, and the treatment was continued for an additional 68–92 weeks (120–144 weeks in total).

Response to entecavir

Biochemical response was defined by the normalization of serum alanine aminotransferase (ALT; < 50 IU/L in our laboratory), virological response by the disappearance of HBV-DNA from serum detectable by Amplicor (sensitivity, < 2.6 LGE/mL), and histological response by a decrease in histology activity index (HAI) score of 2 points or more. Necroinflammatory activity and fibrosis were evaluated by the METAVIR score as well.

Statistical analysis

Frequencies were compared between groups by the Mann-Whitney *U*-test and Fisher's exact test, and medians by the Wilcoxon signed rank test. Normalization in ALT levels and loss of HBV-DNA from the serum, as well as the development of entecavir-resistant HBV mutants, were compared by the method of Kaplan-Meier, and differences were evaluated by the log-rank test with use of the production limit method. *P* < 0.05 was considered significant. Analysis of data was performed with SPSS software (SPSS, Chicago, IL, USA).

Results

Comparison of baseline characteristics between patients given 0.5 and 1.0 mg entecavir daily for 52 weeks and then 1.0 mg for an additional 68–92 weeks

Table 1 compares demographic, biochemical, hematological and virological characteristics between 10 and nine patients with chronic hepatitis B who were randomized to receive 0.5 and 1.0 mg entecavir, respectively, daily for the initial 52 weeks. Thereafter, they all received 1.0 mg entecavir daily for an additional 68–92 weeks (120–144 weeks in total). There were no differences in age, sex, pretreatment ALT levels, platelet counts, frequency of HBeAg, distribution of HBV genotypes, HBV-DNA levels and types of YMDD mutants between the two groups of patients.

Normalization of ALT and loss of HBV-DNA from the serum in patients who received long-term entecavir treatment

Figure 1 depicts ALT levels in 10 and nine patients who received 0.5 and 1.0 mg entecavir daily, respectively, during the initial 52 weeks; thereafter, they all received 1.0 mg entecavir daily for an additional 68–92 weeks (120–144 weeks in total). In both groups, ALT levels increased slightly during 2–4 weeks after the start of entecavir therapy, and then decreased sharply. ALT levels were lowered within the upper limit of normal (≤ 50 IU/L) 12 and 8 weeks after the start of 0.5 and 1.0 mg entecavir daily, respectively. After then, ALT levels decreased and stayed within the

Table 1 Patients with breakthrough hepatitis induced by lamivudine-resistant hepatitis B virus (HBV) mutants who were treated with two doses of entecavir during the initial 52 weeks

	Total (<i>n</i> = 19)	Initial daily dose of entecavir	
		0.5 mg (<i>n</i> = 10)	1.0 mg (<i>n</i> = 9)
Duration of entecavir (weeks)	120–144	120–144	124–140
Age (years)	38 (29–65)	37 (29–65)	39 (30–49)
Men	17 (89%)	9 (90%)	8 (89%)
ALT (IU/L)	119 (46–1708)	111 (46–1708)	275 (49–442)
Platelets ($\times 10^3/\text{mm}^3$)	190 (93–265)	180 (93–235)	190 (108–265)
HBeAg	14 (74%)	7 (70%)	7 (78%)
Genotypes (A : B : C)	1:0:18	1:0:9	0:0:9
HBV-DNA (LGE/mL)	7.2 (5.2–8.6)	7.2 (5.2–8.6)	6.6 (5.7–8.2)
YMDD mutants (I : V : IV)	11:3:5	6:2:2	5:1:3

Median values are shown with the range in parentheses, and the ratio of HBV genotypes, as well as YIDD, YVDD and both YMDD mutants, is indicated. ALT, alanine aminotransferase; HBeAg, hepatitis e antigen; LGE, log geometric equivalents.




Figure 1 Alanine aminotransferase (ALT) levels in the 19 patients with breakthrough hepatitis induced by lamivudine-resistant hepatitis B virus mutants who received entecavir for 120 weeks. Of them, 10 patients received 0.5 mg and the remaining nine 1.0 mg entecavir daily during the initial 52 weeks (shaded), and thereafter both groups received 1.0 mg entecavir daily. The mean ALT levels (a) and the normalization of serum ALT (≤ 50 IU/L) (b) are illustrated.

normal limit among patients in both groups. Collectively in the 19 patients, the ALT level was normalized in 47% at week 12 and in 83% at week 24. Figure 1(b) compares the normalization of ALT levels between patients who received 0.5 and 1.0 mg entecavir daily during the initial 52 weeks. There were no statistical differences in the normalization of ALT levels between patients given 0.5 and 1.0 mg entecavir. Of the 14 patients positive for HBeAg at the start of entecavir, two (14%) lost HBeAg and seroconverted to anti-HBe, while HBeAg was not cleared from the serum in any of the 19 patients.

The loss of HBV-DNA from serum was compared between patients given 0.5 and 1.0 mg entecavir daily during the initial 52 weeks. A sharp decrease in HBV-DNA by more than 2 logs was achieved at 4 weeks in patients given the initial 0.5 mg entecavir daily, and at 8 weeks in those receiving the initial 1.0 mg entecavir daily. Twenty-four weeks after the start, HBV-DNA levels stabilized and stayed approximately 1 log lower in the patients with the initial 0.5 than 1.0 mg entecavir daily. The loss of HBV-DNA detectable by the quantitative method varied in patients with two different initial entecavir doses. At 24 weeks after the start of entecavir therapy, HBV-DNA became undetectable in 20% and 11%, respectively, of the patients with the initial 0.5 and 1.0 mg entecavir daily; the loss increased to 50% and 33% at 120 weeks, respectively. However, there were no significant differences in the loss of HBV-DNA between the patients receiving 0.5 and 1.0 mg entecavir daily during the initial 25 weeks.

Improvement of liver histology in the patients who were switched to entecavir after the development of breakthrough hepatitis during long-term lamivudine treatment

Of the 19 patients switched to receive entecavir, 12 (63%) underwent serial liver biopsies at the baseline and annually for 3 years while they were treated with entecavir. METAVIR scores for fibrosis stages at the start of entecavir were: F1 in six (50%) patients; F2 in three (25%); and F3 in three (25%). Activity grades were: A1 in six (50%) patients and A2 in six (50%). After they had received entecavir for 1 year, the fibrosis stage improved in two (17%), was

Table 2 Improvement in histology activity scores after entecavir during 3 years in the 12 patients who had developed breakthrough hepatitis induced by lamivudine-resistant HBV mutants

Features		Before	After	Decrement	Differences (<i>P</i> -value)
Periportal and/or bridging necrosis	Median (range)	1 (0–3)	0 (0–1)	1 (0–3)	0.003
	Mean ± SD	1.2 ± 0.9	0.1 ± 0.3	1.1 ± 0.8	
Lobular degeneration and focal necrosis	Median (range)	2 (0–3)	1 (0–1)	1 (0–2)	0.014
	Mean ± SD	2.0 ± 1.0	0.9 ± 0.3	1.0 ± 1.0	
Portal inflammation	Median (range)	1 (0–3)	1 (0–1)	1 (0–2)	0.015
	Mean ± SD	1.8 ± 1.0	0.8 ± 0.4	0.9 ± 0.9	
Fibrosis	Median (range)	2 (1–3)	1 (1–3)	0 (0–2)	0.059
	Mean ± SD	2.0 ± 1.0	1.4 ± 0.8	0.5 ± 1.1	
Total HAI score	Median (range)	6 (3–12)	3 (2–5)	3 (1–8)	0.002
	Mean ± SD	7.0 ± 2.7	3.2 ± 0.9	3.5 ± 2.4	

HAI, histology activity index; SD, standard deviation.

unchanged in nine (75%), and worsened in the remaining one (8%). The activity grade improved in nine (75%) patients and was unchanged in the remaining three (25%); it did not worsen in any patient.

One of the 12 patients could not receive liver biopsy 3 years after the start of therapy, because entecavir-resistant HBV mutants developed. Table 2 summarizes changes in HAI scores in the 11 patients who had received long-term entecavir treatment. After 3 years on entecavir therapy, improvement in HAI scores by 2 points or greater was achieved in nine (82%) of the 11 patients. Significant improvement was gained in the total HAI score, as well as scores for periportal and/or bridging necrosis, lobular degeneration/focal necrosis, and portal inflammation. Fibrosis score did not improve significantly ($P = 0.059$); it increased in two patients.

Clinical and virological courses of the representative patient are illustrated in Figure 2 and histological findings in yearly biopsies in Figure 3. The patient developed resistance to lamivudine and was switched to IFN. Hepatitis was exacerbated in him, however, and he was started on lamivudine again. IFN was given intermittently to him when ALT levels were elevated. Because he did not respond to IFN, entecavir was given to him. At that time, he had a HBV-DNA level of more than 7.6 LGE/mL and an HAI score of 8 in the liver biopsy. Soon after entecavir was started, HBV-DNA levels decreased sharply along with the normalization of ALT levels. He seroconverted from HBeAg to anti-HBe 1 year after the start of entecavir treatment. Histological improvement, increasing in parallel with the duration of entecavir treatment, was demonstrated by yearly liver biopsies in comparison with the baseline findings (Fig. 3). Necroinflammatory signs decreased remarkably along with narrowed portal areas, although the stage of fibrosis did not improve appreciably.

HBV mutants resistant to entecavir

Figure 4 illustrates the development of entecavir-resistant HBV mutants that increased in parallel with the duration of treatment. Entecavir-resistant HBV mutants developed in three of the 10 (30%) patients by 18, 84 and 120 weeks; and two of the nine (22%) patients by 132 and 148 weeks, respectively, who received 0.5 and 1.0 mg entecavir daily during the first year; thereafter, they all were given 1.0 mg entecavir daily for the next 68–92 weeks.

During the initial 130 weeks (~2.5 years), therefore, entecavir-resistant HBV mutants developed in three of the 10 (30%) patients with the initial entecavir dose of 0.5 mg daily, in remarkable contrast to no emergence of such mutants in any of the nine patients that received 1.0 mg daily.

Alanine aminotransferase levels were elevated in only two of the five (40%) patients infected with entecavir-resistant HBV mutants, however. These two patients were switched to receive adefovir in combination with lamivudine, and breakthrough hepatitis resolved in them both. All the five patients who developed entecavir-resistant HBV mutants had been infected with lamivudine-resistant YMDD mutants with M204V in the presence or absence of M204I. In outstanding contrast, entecavir-resistant HBV mutants did not develop in any of the 11 patients who had been infected with YMDD mutants with M204I alone.

No adverse effects developed in any of the 19 patients. Breakthrough hepatitis occurred in only one of the five (20%) patients in whom entecavir-resistant mutants emerged.

Discussion

We have previously reported in the Journal that entecavir suppresses serum HBV-DNA to undetectable levels and normalizes ALT levels in more than 30%, respectively, in lamivudine-resistance patients with chronic hepatitis B at 52 weeks.¹² In the present report, we have followed 19 patients from one of the participating centers for 3 years so as to establish longer-term histological efficacy and rates of viral resistance with entecavir treatment of lamivudine-resistant chronic hepatitis B.

As in the earlier report,¹² among the 19 patients described here, ALT levels were normalized in more than 90% of them 8–12 weeks after the start of entecavir until the end of treatment. Although the median HBV-DNA level dropped by 3 logs and remained low during the entecavir therapy, they became undetectable in only 20–40% of the 19 patients. In a previous report, also, the loss of detectable HBV-DNA from the serum was achieved in only 27 of the 141 (19%) patients with lamivudine-resistant HBV mutants after they had received 1.0 mg entecavir daily for 52 weeks.¹⁴ In a remarkable contrast, entecavir is much more efficient in treatment-naïve patients who had received it for 1–2 years; HBV-DNA disappeared from the serum in 67–90% of them.^{10,11,18} These differences could be attributed to some lamivudine-resistant




Figure 2 Clinical course of the representative patient. Fluctuating levels of alanine aminotransferase (ALT) and hepatitis B virus (HBV)-DNA are illustrated. Antiviral treatments as well as duration of hepatitis B e antigen (HBeAg) and anti-HBe are indicated by horizontal bars. Also given are time points when four liver biopsies were undertaken, along with histological activity index scores on the top. IFN, interferon.

HBV mutants contributing to the development of entecavir-resistance.^{13,18}

Entecavir is a cyclopentyl guanosine analog and can inhibit the polymerase of hepadnaviridae selectively by interfering with priming and reverse transcription, as well as synthesis of minus- and plus-stranded HBV-DNA species.¹⁹ In an *in vitro* expression system with HepG2 cells, entecavir exhibited an antiviral activity with EC₅₀ of 0.00375 μ M, which is 1500-fold higher than 10 μ M of lamivudine.²⁰ Dose-dependent pharmacological activity of entecavir was evident in a randomized double-blind trial.²¹ Although 0.01 mg entecavir daily decreased HBV-DNA by 2.41 logs at 22 weeks, the antiviral activity was significantly lower than 4.31 and 4.72 logs, respectively, of 0.1 and 0.5 mg daily; they were both higher than 3.36 logs by 100 mg lamivudine daily, however. Accordingly, normalization of ALT was more frequent by treatments with 0.1 and 0.5 mg entecavir daily (69% and 83%, respectively) than with 100 mg lamivudine daily (59%).

Significant decrease in HAI scores has been reported in patients with chronic hepatitis B who had received lamivudine for 1–3 years.^{22,23} Furthermore, decreases in hepatic inflammation may improve the fibrosis stage. Entecavir therapy for 52 weeks has achieved histological improvement in 55–72% of patients in phase



III clinical trials.^{10,11,14} In corroboration of these results, fibrosis stage and inflammation grade improved in the present series of patients who had received entecavir for 3 years, with a significant decrease in the HAI score (Table 2). Histological improvement would have been gained by long-term entecavir therapy, and it may further increase, should entecavir be continued further.

Long-term entecavir treatment, however, may be hampered by the development of drug-resistant mutants. Although entecavir-resistant HBV mutants rarely occur in treatment-naïve patients,¹⁸ they can emerge rather frequently in the patients infected with lamivudine-resistant HBV mutants.^{14,24} In the present study, entecavir-resistant HBV mutants developed in five of the 19 (26%) lamivudine-resistant patients during 144 weeks of treatment. The incidence was comparable with 32% in the lamivudine-resistant patients who had received entecavir for 3 years.²⁴ Only two (40%) of them developed hepatitis flares and they were switched to receive adefovir in combination with lamivudine. Entecavir-resistant HBV mutants emerging in patients with lamivudine-resistant mutants are reported to be replication-impaired and rarely induce breakthrough hepatitis.²⁵ It should be found out how entecavir-resistant HBV mutants can be managed with antiviral nucleos(t)ide analogs other than lamivudine and entecavir, or


(a) Before the start of entecavir



(b) One year after the start of entecavir



(c) Two years after the start of entecavir



(d) Three years after the start of entecavir





Figure 3 Histological changes in the representative patient during 3-year entecavir treatment (Fig. 2). With hematoxylin–eosin stain on the left, marked enlargement of portal areas is evident along with infiltration of mononuclear cells before the switch from lamivudine to entecavir (a). They decreased increasingly during the 3-year treatment with entecavir (b–d). Stage of fibrosis did not change appreciably by the staining for silver on the right.




Figure 4 Development of entecavir-resistant hepatitis B virus (HBV) mutants during the 3-year treatment. The 10 patients with the initial entecavir dose of 0.5 mg daily and the nine with that of 1.0 mg daily are compared.

combination thereof. It has been proposed that adefovir add-on lamivudine is efficacious with negligible drug resistance over 3 years.^{26,27}

Acknowledgment

This work was supported in part by grants from the Japanese Ministry of Health, Labor and Welfare.

References

- Lai CL, Ratziu V, Yuen MF, Poynard T. Viral hepatitis B. *Lancet* 2003; **362**: 2089–94.
- Dienstag JL, Schiff ER, Wright TL *et al.* Lamivudine as initial treatment for chronic hepatitis B in the United States. *N. Engl. J. Med.* 1999; **341**: 1256–63.
- Lai CL, Chien RN, Leung NW *et al.* A one-year trial of lamivudine for chronic hepatitis B. *N. Engl. J. Med.* 1998; **339**: 61–8.
- Chayama K, Suzuki Y, Kobayashi M *et al.* Emergence and takeover of YMDD motif mutant hepatitis B virus during long-term lamivudine therapy and re-takeover by wild type after cessation of therapy. *Hepatology* 1998; **27**: 1711–16.
- Honkoop P, Niesters HG, de Man RA, Osterhaus AD, Schalm SW. Lamivudine resistance in immunocompetent chronic hepatitis B. Incidence and patterns. *J. Hepatol.* 1997; **26**: 1393–5.
- Leung NW, Lai CL, Chang TT *et al.* Extended lamivudine treatment in patients with chronic hepatitis B enhances hepatitis B e antigen seroconversion rates: results after 3 years of therapy. *Hepatology* 2001; **33**: 1527–32.
- Liaw YF, Chien RN, Yeh CT, Tsai SL, Chu CM. Acute exacerbation and hepatitis B virus clearance after emergence of YMDD motif mutation during lamivudine therapy. *Hepatology* 1999; **30**: 567–72.
- Hadziyannis SJ, Tassopoulos NC, Heathcote EJ *et al.* Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N. Engl. J. Med.* 2003; **348**: 800–7.
- Marcellin P, Chang TT, Lim SG *et al.* Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *N. Engl. J. Med.* 2003; **348**: 808–16.
- Chang TT, Gish RG, de Man R *et al.* A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N. Engl. J. Med.* 2006; **354**: 1001–10.
- Lai CL, Shouval D, Lok AS *et al.* Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N. Engl. J. Med.* 2006; **354**: 1011–20.
- Suzuki F, Joji T, Yoshiaki K *et al.* Efficacy and safety of entecavir in lamivudine-refractory patients with chronic hepatitis B: Randomized controlled trial in Japanese patients. *J. Gastroenterol. Hepatol.* 2008; **23**: 1320–6.
- Perrillo R, Hann HW, Mutimer D *et al.* Adefovir dipivoxil added to ongoing lamivudine in chronic hepatitis B with YMDD mutant hepatitis B virus. *Gastroenterology* 2004; **126**: 81–90.
- Sherman M, Yurdaydin C, Sollano J *et al.* Entecavir for treatment of lamivudine-refractory, HBeAg-positive chronic hepatitis B. *Gastroenterology* 2006; **130**: 2039–49.
- Usuda S, Okamoto H, Iwanari H *et al.* Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J. Virol. Methods* 1999; **80**: 97–112.
- Usuda S, Okamoto H, Tanaka T *et al.* Differentiation of hepatitis B virus genotypes D and E by ELISA using monoclonal antibodies to epitopes on the preS2-region product. *J. Virol. Methods* 2000; **87**: 81–9.
- Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc. Natl Acad. Sci. USA* 1977; **74**: 5463–7.
- Colonna RJ, Rose R, Baldick CJ *et al.* Entecavir resistance is rare in nucleoside naive patients with hepatitis B. *Hepatology* 2006; **44**: 1656–65.
- Zoulim F. Entecavir: a new treatment option for chronic hepatitis B. *J. Clin. Virol.* 2006; **36**: 8–12.
- Lampertico P. Entecavir versus lamivudine for HBeAg positive and negative chronic hepatitis B. *J. Hepatol.* 2006; **45**: 457–60.
- Lai CL, Rosmawati M, Lao J *et al.* Entecavir is superior to lamivudine in reducing hepatitis B virus DNA in patients with chronic hepatitis B infection. *Gastroenterology* 2002; **123**: 1831–8.
- Suzuki Y, Arase Y, Ikeda K *et al.* Histological improvements after a three-year lamivudine therapy in patients with chronic hepatitis B in whom YMDD mutants did not or did develop. *Intervirology* 2003; **46**: 164–70.
- Suzuki Y, Kumada H, Ikeda K *et al.* Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J. Hepatol* 1999; **30**: 743–8.
- Papatheodoridis GV, Manolakopoulos S, Dusheiko G, Archimandritis AJ. Therapeutic strategies in the management of patients with chronic hepatitis B virus infection. *Lancet Infect. Dis.* 2008; **8**: 167–78.
- Tenney DJ, Rose RE, Baldick CJ *et al.* Two-year assessment of entecavir resistance in Lamivudine-refractory hepatitis B virus patients reveals different clinical outcomes depending on the resistance substitutions present. *Antimicrob. Agents Chemother.* 2007; **51**: 902–11.
- Lampertico P, Vigano M, Manenti E *et al.* M. Low resistance to adefovir combined with lamivudine: a 3-year study of 145 lamivudine-resistant hepatitis B patients. *Gastroenterology* 2007; **133**: 1445–51.
- Yatsuji H, Suzuki F, Sezaki H *et al.* Low risk of adefovir resistance in lamivudine-resistant chronic hepatitis B patients treated with adefovir plus lamivudine combination therapy: two-year follow-up. *J. Hepatol* 2008; **48**: 923–31.

Predictive values of amino acid sequences of the core and NS5A regions in antiviral therapy for hepatitis C: a Japanese multi-center study

Takeshi Okanoue · Yoshito Itoh · Hiroaki Hashimoto · Kohichiroh Yasui · Masahito Minami · Tetsuo Takehara · Eiji Tanaka · Morikazu Onji · Joji Toyota · Kazuaki Chayama · Kentaro Yoshioka · Namiki Izumi · Norio Akuta · Hiromitsu Kumada

Received: 31 March 2009 / Accepted: 20 April 2009 / Published online: 11 June 2009
© Springer 2009

Abstract

Background Chronic hepatitis C (CHC) genotype 1b patients with high viral load are resistant to peginterferon (PEG-IFN) and ribavirin (RBV) combination therapy, especially older and female patients.

Methods To elucidate the factors affecting early and sustained viral responses (EVR and SVR), 409 genotype 1b patients CHC with high viral loads who had received 48 weeks of PEG-IFN/RBV therapy were enrolled. The amino acid (aa) sequences of the HCV core at positions 70 and 91 and of the interferon sensitivity determining region (ISDR) were analyzed. Host factors, viral factors, and

treatment-related factors were subjected to multivariate analysis.

Results Male gender, low HCV RNA load, high platelet count, two or more aa mutations of ISDR, and wild type of core aa 70 were independent predictive factors for SVR. In patients with over 80% adherences to both PEG-IFN and RBV, male gender, mild fibrosis stage, and wild type of core aa 70 were independent predictors for SVR.

Conclusions Independent predictive factors for SVR were: no aa substitution at core aa 70, two or more aa mutations in the ISDR, low viral load, high values of platelet count, mild liver fibrosis and male gender.

T. Okanoue (✉)
Hepatology Center, Saiseikai Suita Hospital,
1-2 Kawazonocho, Suita 564-0013, Japan
e-mail: okanoue@suita.saiseikai.or.jp

T. Okanoue · Y. Itoh · H. Hashimoto · K. Yasui · M. Minami
Department of Gastroenterology and Hepatology,
Kyoto Prefectural University of Medicine,
Kawaramachi-Hirokoji Kamigyo-Ku,
Kyoto 602-8566, Japan

Y. Itoh
e-mail: yitoh@koto.kpu-m.ac.jp

H. Hashimoto
e-mail: road1820@yahoo.co.jp

K. Yasui
e-mail: yasui@koto.kpu-m.ac.jp

M. Minami
e-mail: minami@koto.kpu-m.ac.jp

T. Takehara
Department of Gastroenterology and Hepatology,
Osaka University Graduate School of Medicine,
2-2 Yamadaoka, Suita 565-0871, Japan
e-mail: takehara@gh.med.osaka-u.ac.jp

E. Tanaka
Department of Medicine, Shinshu University School
of Medicine, 3-1-1 Asahi, Matsumoto 390-8621, Japan
e-mail: etanaka@shinshu-u.ac.jp

M. Onji
Department of Gastroenterology, Ehime University,
454 Shizukawa, Tohon, Ehime 791-0295, Japan
e-mail: onjimori@m.med.ehime-u.ac.jp

J. Toyota
Department of Gastroenterology, Sapporo-Kosei General
Hospital, Kita Sanjyo, Chuo-ku, Sapporo 060-0033, Japan
e-mail: joji.toyota@ja_hokkaidoukouseiren.or.jp

K. Chayama
Department of Medicine and Molecular Science,
Graduate School of Science, Hiroshima University,
1-2-3 Kasumi, Minami-ku, Hiroshima,
Hiroshima 734-8551, Japan
e-mail: chayama@hiroshima-u.ac.jp

K. Yoshioka
Department of Hepato-Biliary-Pancreas, Fujita Health Science,
Kutsukake-cho, Toyoake 470-1192, Japan
e-mail: kyoshiok@fujita-hu.ac.jp

Keywords Chronic hepatitis C · Peginterferon and ribavirin · Core amino acid · Interferon sensitivity determining region

Abbreviations

CHC	Chronic hepatitis C
PEG-IFN	Peginterferon
RBV	Ribavirin
RVR	Rapid viral response
cEVR	Complete early viral response
LVR	Late viral response
ETR	End of treatment response
NR	Non response
SVR	Sustained viral response
ISDR	Interferon sensitivity determining region
Aa	Amino acid
ALT	Alanine aminotransferase
PLT	Platelet
HCC	Hepatocellular carcinoma

Introduction

A combination of pegylated interferon (PEG-IFN) and ribavirin (RBV) therapy for 48 weeks achieves a sustained viral response (SVR) rate of 40–50% in chronic hepatitis C (CHC) patients with a high viral load of genotype 1 [1–4]. The dose-reduction rate and the frequency of discontinuation of this treatment are high in aged patients [5]. The SVR rate of the therapy is lower in females than males, especially in older patients in Japan [6].

Around 30% of HCV carriers have serum alanine aminotransferase (ALT) levels within the upper limit of normal ranges [7, 8] and HCV carriers with persistently normal serum ALT (PNALT) and serum platelet (PLT) counts of over $15 \times 10^4/\text{mm}^3$ show low grade hepatic fibrosis and good prognosis [9]. Before treating HCV carriers, it is very important to predict non-response to PEG-IFN plus RBV therapy because of its medical cost, adverse effects, and its impact on the long term quality of life.

N. Izumi
Department of Gastroenterology and Hepatology,
Musashino Red Cross Hospital, Sakaiminamimachi,
Musashino 180-8610, Japan
e-mail: nizumi@musashino.jrc.or.jp

N. Akuta · H. Kumada
Department of Hepatology, Toranomon Hospital,
Kajigaya, Takatsu-ku, Kawasaki 213-8587, Japan

N. Akuta
e-mail: akuta-gi@umin.ac.jp

H. Kumada
e-mail: kumahiro@toranomon.gr.jp

There are many factors affecting response to IFN monotherapy and PEG-IFN/RBV therapy, including body mass index (BMI) [10, 11], steatosis [12, 13], insulin resistance [14], stage of liver fibrosis [15, 16], total cholesterol (T. Chol), triglyceride (TG), adherence to both PEG-IFN and RBV [17], race [18, 19], age [1, 2, 20], and viral factors including serum quantity of HCV RNA, HCV genotype and substitution of amino acids (aa) in the interferon sensitivity determining region (ISDR, 2209–2248) of the nonstructural protein 5A (NS5A) [21] and in the core protein [22, 23]. Early viral response is an important predictive factor in PEG-IFN/RBV therapy for CHC patients with genotype 1 and high viral loads [24–27].

The aim of this study was to elucidate the valuable predictive factors of SVR in Japanese patients with HCV genotype 1b high viral loads following 48 weeks of PEG-IFN/RBV therapy, focusing on the relationship between aa substitutions in the ISDR and at core aa 70 and 91 and early viral kinetics.

Patients and methods

Selection of patients

This retrospective study was conducted at 15 clinical sites in Japan which are part of the Study Group of Optimal Treatment of Viral Hepatitis supported by the Ministry of Health, Labor and Welfare, Japan. Eligible subjects were CHC patients, who (1) had received liver biopsy; (2) were genotype 1b with high viral load (≥ 100 KIU/ml by Cobas Amplicor Hepatitis C Virus Test, version 2.0) at the start of PEG-IFN/RBV therapy; (3) received weekly injections of PEG-IFN- α -2b (PEG-INTRON; Shering-Plough, Kenilworth, NJ) of 1.5 $\mu\text{g}/\text{kg}$ bw and oral administration of RBV (Rebetol; Shering-Plough) for 48 weeks. The amount of RBV was adjusted based on the subject's body weight; (600 mg for ≤ 60 kg bw, 800 mg for 60–80 kg bw, 1,000 mg for > 80 kg bw); (4) were examined serially for quantitative and qualitative HCV RNA; and (5) the aa sequences at positions 70 and 91 in the core region and of the ISDR in the NS5A had been determined in pretreatment sera.

Hepatitis B virus (HBV) infection, human immunodeficiency virus (HIV) infection, autoimmune hepatitis, primary biliary cirrhosis, hemochromatosis, and Wilson's disease were excluded. Histopathological diagnosis was based on the scoring system of Desmet et al. [28]. The definition of alcohol abuse included patients having a history of more than 100 kg of total ethanol intake. Complete blood counts, liver function tests, serum lipids, serum ferritin, serum fibrosis markers, fasting plasma glucose (FPG), and immune reactive insulin (IRI) were examined in most cases. Written informed consent was obtained from all

patients before treatment, and the protocol was approved by the ethics committees in each site.

Study design

Four hundred and nine patients who completed 48 weeks of treatment and were followed for more than 24 weeks after treatment were enrolled in the first study (*Study design 1*).

To elucidate the effect of aa substitution of HCV core and in the ISDR on HCV dynamics, including a rapid viral response (RVR), complete early viral response (cEVR), a late viral response (LVR) and SVR, according to gender and age (<60 years \geq 60 years), 201 of the 409 patients maintaining over 80% adherences to both PEG-IFN and RBV were enrolled in the second study (*Study design 2*).

Nucleotide sequencing of the core and NA5A gene

The nucleotide sequences encoding aa 1–191 (HCV core) and aa 2209–2248 (ISDR) were analyzed by direct sequencing as described by Akuta et al. [22, 27] and Enomoto et al. [21]. In brief, RNA was extracted from the sera and converted to cDNA and two nested rounds of polymerase chain reaction (PCR) were performed. Primers used in the PCR were as follows; (a) Nucleotide sequences of the core region: the first-round PCR was performed with CC11 (sense) and e14 (antisense) primers [22, 27], and the second-round PCR with CC9 (sense) and e14 (antisense) primers [22, 27]. (b) Nucleotide sequences of the ISDR in NS5A: the first-round PCR was performed with ISDR1 (sense) and ISDR2 (antisense) primers [21], and the second-round PCR with ISDR3 (sense) and ISDR4 (antisense) primers [21]. These sequences were compared with the consensus sequence of genotype 1b (HCV-J) [29]. Wild types virus encoded arginine and leucine at aa 70 and 91, respectively, and the aa substitutions were glutamine or histidine at aa 70 and methionine at aa 91.

Viral kinetic study

Serum HCV RNA levels were measured by PCR (Amplicor HCV RNA kit, version 2.0, Roche Diagnostics) using samples taken before treatment and at 4, 12, 24, and 48 weeks after the therapy. SVR was defined as HCV RNA negativity by qualitative analysis by PCR at 24 weeks after the treatment. RVR was defined as HCV RNA negativity at 4 weeks, cEVR as HCV RNA negativity at 12 weeks, LVR as HCV RNA negativity during 13–24 weeks and an end of treatment response (ETR) as HCV RNA negativity at the end of treatment. Patients who remained positive for HCV RNA at the end of the treatment and at 24 weeks after the therapy were defined as non-responders (NR).

Adherences to PEG-IFN and RBV

Adherences to PEG-IFN and RBV were assessed by separately calculating the actual doses of PEG-IFN and RBV received as percentages of the intended dosages. Adherences to PEG-IFN and RBV were divided into two groups; $80\% \leq$ and $<80\%$.

Statistical analysis

All data analyses were conducted using the SAS version 9.1.3 statistical analysis packages (SAS Institute, Cary, NC, USA). Individual characteristics between groups were evaluated by Mann–Whitney *U* test for numerical variables or Fisher's exact test for categorical variables. Variables exhibiting values of $p < 0.1$ in the univariate analysis were subjected to stepwise multivariate logistic regression analysis. The grade of steatosis and iron deposition in liver tissue, BMI, albumin (Alb), low density lipoprotein-cholesterol (LDL-C), homeostasis model assessment-insulin resistance (HOMA-IR), ferritin, and hyaluronic acid were excluded from multivariate logistic regression analysis because of the absence of those data in more than 10% of the patients. All p values of $p < 0.05$ by the two-tailed test were considered statistically significant.

Results

Study design 1

Baseline backgrounds, characteristics and adherences of peginterferon and ribavirin in males and females

The treatment outcome of PEG-IFN and RBV combination therapy depends on gender in Japanese patients, so in addition to aa substitutions in the ISDR in NS5A [21] or at HCV core 70 and 91 [22, 27], we compared the baseline characteristics according to gender (Table 1). Males were younger and the grade of hepatic inflammation was milder in males. The serum levels of LDL-C, PLT count, and aa substitutions of ISDR and at core 70 and 91 did not differ significantly different between males and females. The frequency of no alcohol abuse was significantly ($p < 0.0001$) higher in females than males (Some of them are not described in Table 1).

The rates of over 80% adherences to PEG-IFN and RBV were significantly lower ($p = 0.0066$, $p < 0.00001$, respectively) in females than males. Only in those above 60 years did the rate of over 80% adherence to PEG-IFN not differ significantly between males and females, but the rate of over 80% adherence to RBV was significantly lower ($p = 0.035$) in females than males (Table 1).