

Figure 2 Frequencies of core promoter, pre-core and core mutations compared between the transient hepatitis B virus infection (FH-T) and the acute self-limited hepatitis B (AHB) patients who were infected with HBV of subgenotype B1/Bj (a) or C2/Ce (b).

2.75–66.64], $P = 0.0014$) and serum HBV DNA more than 5.23 log copies/mL (OR, 5.14 [95% CI, 1.10–24.15], $P = 0.0379$) were independent risk factors for the development of FHB by transient HBV infection (Table 2). Other mutations (T1753V, T1754V, A1762T/G1764A, G1899A and A2339G) were not significantly associated with the development of FHB by transient HBV infection, however.

Baseline clinical characteristics for distinguishing between the patients with FHB by AE of ASC (FH-C) and those without FHB by AE of CHB (AE-C)

Table 3 compares baseline clinical characteristics between the 12 FH-C patients and the 12 AE-C patients who were matched for age and sex. The levels of T.bil were significantly higher in the FH-C patients (15.0 ± 7.3 vs 7.3 ± 8.8 mg/dL, $P < 0.05$), but the peak ALT and AST levels tended to be slightly higher in the FH-C patients than AE-C (887 ± 681 vs 641 ± 620 IU/L and 701 ± 451 vs 601 ± 753 IU/L, respectively). There were also no significant differences in levels of sera HBV DNA, core protein and HBcrAg between these two groups (7.44 ± 1.51 vs 6.60 ± 1.10 log copies/mL, 5.04 ± 1.45 vs 5.07 ± 1.07 log U/mL, and 6.35 ± 1.70 vs 6.29 ± 1.95 log U/mL, respectively).

HBV genotypes and enhancer II/core promoter/pre-core/core mutations between the patients with FH-C and those with AE-C

There were no significant differences in the frequencies of any HBV genotypes between the 12 FH-C patients and the 12 AE-C patients (Fig. 3a). In addition, there were also no significant differences in the frequencies

Table 2 Multivariate analysis for factors independently associated with fulminant hepatitis by transient HBV infection

Factors	Odds ratio	95% confidence interval	P-value
Total bilirubin (mg/dL)†			
<10.35	1		
≥ 10.35	7.81	1.77–34.51	0.0067
G1896A mutation			
Absent	1		
Present	13.53	2.75–66.64	0.0014
HBV DNA (log copies/mL)†			
<5.23	1		
≥ 5.23	5.14	1.10–24.15	0.0379

†Median values. HBV, hepatitis B virus.

Table 3 Baseline characteristics between patients with FH by AE of ASC (FH-C) and those without FH by AE of CHB (AE-C)

Features	FH-C (n = 12)	AE-C (n = 12)	Differences P-value
Age (years)	51.7 ± 14.7	49.9 ± 5.6	Matched
Male	10 (83%)	9 (75%)	Matched
ALT (IU/L)	887 ± 681	641 ± 620	NS
AST (IU/L)	701 ± 451	601 ± 753	NS
Total bilirubin (mg/dL)	15.0 ± 7.3	7.3 ± 8.8	<0.05
Prothrombin time (%)	25.8 ± 6.6	48.4 ± 21.5	<0.005
HBeAg positive	4 (33%)	3 (25%)	NS
Core protein (log U/mL)	5.04 ± 1.45	5.07 ± 1.07	NS
HBcrAg (log U/mL)	6.35 ± 1.70	6.29 ± 1.95	NS
HBV DNA (log copies/mL)	7.44 ± 1.51	6.60 ± 1.10	NS

AE, acute exacerbation; ALT, alanine aminotransferase; ASC, asymptomatic HBV carrier; AST, aspartate aminotransferase; CHB, chronic hepatitis B; HBcrAg, hepatitis B core related antigen; HBeAg, hepatitis B e antigen; HBV, hepatitis B virus; NS, not significant.

of any specific mutations between these two groups (Fig. 3b).

DISCUSSION

THE MAGNITUDE OF liver injuries depends on the replication level of HBV and cytotoxic immune responses of the host raised against viral epitopes in general.³¹ Various viral factors have been proposed that promote the development of FHB, represented by pre-core (G1896A) and core promoter (A1762T/G1764A) mutations.^{13–16} Impact of virological factors on the development of FHB has remained controversial, however, especially because these mutations are rarely detected in the patients from the USA and France.^{19–21} It has been argued that the development of FHB is not promoted by these mutations and is dependent on host factors including the human leukocyte antigen (HLA) environment.²²

The expression of HBeAg is terminated by G1896A mutation in the pre-core region at the translation level,³² and downregulated by the A1762T/G1764A double mutation at the transcription level.^{33,34} Lamberts *et al.* are the first to implicate a negative influence of HBeAg on the replication of HBV.³⁵ Should HBeAg suppress the replication of HBV, presumably by inhibiting the encapsidation of pre-genome,³⁵ the lack or decrease of HBeAg would enhance the reproduction of HBV. Furthermore, HBeAg acts as a tolerogen to T cells recognizing epitopes on core protein, thereby, obviating immune injury of hepatocytes.^{36,37} In the absence or decrease of HBeAg, therefore, hosts would mount vigor cytotoxic T-cell responses to core epitopes excessively

presented on hepatocytes, and develop severe liver injuries culminating in FHB.³⁸

There is a possibility that influence of viral factors such as HBV mutants with a HBeAg-negative phenotype, on the induction of FHB, may have been confounded by host factors and created disagreement. Therefore, the sheer influence of virological factors on FHB would need to be evaluated in case-control studies, as has been attempted to sort out the influence of HBV genotypes on development of cirrhosis and hepatocellular carcinoma.⁸ These backgrounds have instigated us to identify virological factors accelerating the severity of liver disease in the 50 FHB patients by transient HBV infection and the 50 AHB patients who were of the same ethnicity and matched for age as well as sex.

In this case controlled study, A1762T/G1764A, G1896A, G1899A and A2339G mutation were significantly more frequent in the patients with FH-T than AHB, providing further corroboration of previous studies;^{13–16} these mutations could enhance viral replication. Interestingly, our recent study using an *in vitro* replication model, showed that A2339G mutation in the core region enhanced viral replication and the effect of A2339G mutation may be associated with inhibition of the cleavage of the core protein by a furin-like protease, resulting in the high expression of the complete core protein.¹⁸ Such enhanced HBV would induce significant immune response, resulting in development of FHB.

In multivariate analysis, higher levels of serum HBV DNA and G1896A mutation were independent virological risk factors for the development of FHB by transient




Figure 3 Genotypes/subgenotypes (a) and mutations in core promoter, pre-core and core regions (b) between the 12 transient hepatitis B virus infection patients (FH-T) and the 12 acute self-limited hepatitis B (AHB) patients.

HBV infection (Table 2). In particular, G1896A mutation was the most important factor associated with the development of FHB. Host responses, represented by T.bil, contributed to the development of FHB as well.

As for HBV genotypes, B1/Bj alone was significantly more frequent in the FH-T patients in univariate analysis.

In the patients infected with B1/Bj, G1896A was more frequent in those with FH-T than AHB. In *in vitro* replication analysis, Ozasa *et al.*¹⁵ observed extremely high expressions of intra- and extracellular HBV DNA in culture transfected with an HBV clone of B1/Bj genotype having the G1896A mutation; a high replication would be induced by this pre-core mutation for the induction of FHB. Our clinical results stand in support of this *in vitro* analysis. Taken altogether, chances for developing severe acute or FH would be high in the patients with acute hepatitis who are infected with HBV/B1 having the pre-core mutation. By contrast, in patients infected with C2/Ce, G1896A or A1762T/G1764A, or both was much more frequent in the FH-T patients than AHB. Of note, the co-occurrence of G1896A and A1762T/G1764A mutations was invariably accompanied by either FHB or acute severe hepatitis B in this study. Hence, these pre-core and core-promoter mutations might have additive or synergetic effects for exacerbating hepatitis, when they emerge in the patients infected with C2/Ce. Such high-risk patients deserve special care and surveillance for signs and symptoms of fulminant or severe acute hepatitis B.

In the present study, serum levels of HBV DNA were significantly higher in the patients with FH-T than AHB. High serum levels of HBV DNA have been reported in patients with FHB;³⁹ they are followed by rapid decrease as the sequel of virus elimination operated by vigorous immune responses. Because of rapid and extensive elimination of HBV by the host immune system, HBV DNA in serum, in general, has decreased to low levels in patients with FHB at the presentation.⁴⁰ HBV DNA levels may be subject to the time that has elapsed from the onset of hepatitis to its measurement.³⁹ Also, serum levels of core protein (the product of the C gene) closely correlate with serum HBV DNA levels in patients with hepatitis B,²⁷ and they were compared between the FH-T patients and AHB. The core protein was determined by the newly developed CLEIA method; it is much easier and less expensive than the determination of HBV DNA. The level of core protein has turned out to be marginally higher in the FH-T patients than AHB (Table 1), and therefore might not contribute to an early diagnosis of FHB by transient infection.

Fulminant hepatitis B by AE of ASC is assumed as a different clinical condition from FHB by transient HBV infection. In this study, as there was no case-control study on virological factors associated with FHB for the patients with AE of ASC, we also attempted to identify virological factors associated with the development of FHB in the 12 FH-C and the 12 AE-C patients who were

matched for age as well as sex. Disappointingly, no differences of virological factors such as HBV genotypes and pre-core mutations, which were strongly associated with the development of FHB by transient infection, were found between the FH-C and AE-C patients (Fig. 3a,b). Furthermore, there were also no significant differences about HBeAg-positive rate and the levels of serum HBV DNA or core protein (Table 3), suggesting that several host factors may play a more important role in the development of FHB in ASC instead of virological factors. In this case-control study, however, there seems to be some problems: a small number of patients, different duration of HBV infection, different clinical stage (ASC or CHB) at the onset of AE, and HBV quasispecies complexity. Further investigations are needed to identify factors associated with FHB precipitating in asymptomatic HBV carriers.

In conclusion, virological factors associated with enhancement of viral replication seemed to be important for the development of FHB in the patients by transient HBV infection. But no virological factors were identified for differentiation of the FH-C patients from the AE-C patients. Hence, the pathogenic mechanism of FHB between transient HBV infection and AE of ASC would be different.

ACKNOWLEDGMENTS

WE WOULD LIKE to thank Dr S. Baba, Showa University Hospital, Dr Y. Koga, Kurume University School of Medicine and the other doctors for collecting serum samples in this study. We would also thank Dr N. Maki, Advanced Life Science Institute (Saitama, Japan) for measuring core protein in serum. This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare of Japan, and a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology.

REFERENCES

- 1 Fujiwara K, Mochida S, Matsui A, Nakayama N, Nagoshi S, Toda G. Fulminant hepatitis and late onset hepatic failure in Japan. *Hepatol Res* 2008; 38: 646–57.
- 2 Norder H, Hammas B, Lofdahl S, Courouce AM, Magnius LO. Comparison of the amino acid sequences of nine different serotypes of hepatitis B surface antigen and genomic classification of the corresponding hepatitis B virus strains. *J Gen Virol* 1992; 73 (Pt 5): 1201–8.
- 3 Okamoto H, Tsuda F, Sakugawa H *et al.* Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J Gen Virol* 1988; 69: 2575–83.
- 4 Stuyver L, De Gendt S, Van Geyt C *et al.* A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J Gen Virol* 2000; 81 (Pt 1): 67–74.
- 5 Arauz-Ruiz P, Norder H, Robertson BH, Magnius LO. Genotype H: a new Amerindian genotype of hepatitis B virus revealed in Central America. *J Gen Virol* 2002; 83 (Pt 8): 2059–73.
- 6 Miyakawa Y, Mizokami M. Classifying hepatitis B virus genotypes. *Intervirology* 2003; 46: 329–38.
- 7 Chu CJ, Lok AS. Clinical significance of hepatitis B virus genotypes. *Hepatology* 2002; 35: 1274–6.
- 8 Tanaka Y, Hasegawa I, Kato T *et al.* A case-control study for differences among hepatitis B virus infections of genotypes A (subtypes Aa and Ae) and D. *Hepatology* 2004; 40: 747–55.
- 9 Suguchi F, Orito E, Ichida T *et al.* Hepatitis B virus of genotype B with or without recombination with genotype C over the pre-core region plus the core gene. *J Virol* 2002; 76: 5985–92.
- 10 Huy TT, Ushijima H, Quang VX *et al.* Genotype C of hepatitis B virus can be classified into at least two subgroups. *J Gen Virol* 2004; 85 (Pt 2): 283–92.
- 11 Tanaka Y, Orito E, Yuen MF *et al.* Two subtypes (subgenotypes) of hepatitis B virus genotype C: a novel subtyping assay based on restriction fragment length polymorphism. *Hepatol Res* 2005; 33: 216–24.
- 12 Lindh M, Andersson AS, Gusdal A. Genotypes, nt 1858 variants, and geographic origin of hepatitis B virus—large-scale analysis using a new genotyping method. *J Infect Dis* 1997; 175: 1285–93.
- 13 Sato S, Suzuki K, Akahane Y *et al.* Hepatitis B virus strains with mutations in the core promoter in patients with fulminant hepatitis. *Ann Intern Med* 1995; 122: 241–8.
- 14 Omata M, Ehata T, Yokosuka O, Hosoda K, Ohto M. Mutations in the pre-core region of hepatitis B virus DNA in patients with fulminant and severe hepatitis. *N Engl J Med* 1991; 324: 1699–704.
- 15 Ozasa A, Tanaka Y, Orito E *et al.* Influence of genotypes and pre-core mutations on fulminant or chronic outcome of acute hepatitis B virus infection. *Hepatology* 2006; 44: 326–34.
- 16 Liang TJ, Hasegawa K, Rimon N, Wands JR, Ben-Porath E. A hepatitis B virus mutant associated with an epidemic of fulminant hepatitis. *N Engl J Med* 1991; 324: 1705–9.
- 17 Imamura T, Yokosuka O, Kurihara T *et al.* Distribution of hepatitis B viral genotypes and mutations in the core promoter and pre-core regions in acute forms of liver disease in patients from Chiba, Japan. *Gut* 2003; 52: 1630–7.
- 18 Sugiyama M, Tanaka Y, Kurbanov F, Nakayama N, Mochida S, Mizokami M. Influences on hepatitis B virus replication by a naturally occurring mutation in the core gene. *Virology* 2007; 365: 285–91.

- 19 Laskus T, Persing DH, Nowicki MJ, Mosley JW, Rakela J. Nucleotide sequence analysis of the pre-core region in patients with fulminant hepatitis B in the United States. *Gastroenterology* 1993; 105: 1173–8.
- 20 Liang TJ, Hasegawa K, Munoz SJ *et al.* Hepatitis B virus pre-core mutation and fulminant hepatitis in the United States. A polymerase chain reaction-based assay for the detection of specific mutation. *J Clin Invest* 1994; 93: 550–5.
- 21 Feray C, Gigou M, Samuel D, Bernuau J, Bismuth H, Brechot C. Low prevalence of pre-core mutations in hepatitis B virus DNA in fulminant hepatitis type B in France. *J Hepatol* 1993; 18: 119–22.
- 22 Karayiannis P, Alexopoulou A, Hadziyannis S *et al.* Fulminant hepatitis associated with hepatitis B virus e antigen-negative infection: importance of host factors. *Hepatology* 1995; 22: 1628–34.
- 23 Trey C, Lipworth L, Chalmers TC *et al.* Fulminant hepatic failure. Presumable contribution to halothane. *N Engl J Med* 1968; 279: 798–801.
- 24 Ng HJ, Lim LC. Fulminant hepatitis B virus reactivation with concomitant listeriosis after fludarabine and rituximab therapy: case report. *Ann Hematol* 2001; 80: 549–52.
- 25 Fujiwara K, Mochida S, Matsui A. [Prospective study for the efficiency of lamivudine for the patients with acute exacerbation of HBV carrier.] *Annual Report of Intractable Liver Disease Study Group of Japan, the Ministry of Health, Welfare and Labor* 2004. (In Japanese.)
- 26 Kimura T, Rokuhara A, Sakamoto Y *et al.* Sensitive enzyme immunoassay for hepatitis B virus core-related antigens and their correlation to virus load. *J Clin Microbiol* 2002; 40: 439–45.
- 27 Kimura T, Rokuhara A, Matsumoto A *et al.* New enzyme immunoassay for detection of hepatitis B virus core antigen (HBcAg) and relation between levels of HBcAg and HBV DNA. *J Clin Microbiol* 2003; 41: 1901–6.
- 28 Abe A, Inoue K, Tanaka T *et al.* Quantitation of hepatitis B virus genomic DNA by real-time detection PCR. *J Clin Microbiol* 1999; 37: 2899–903.
- 29 Sugauchi F, Mizokami M, Orito E *et al.* A novel variant genotype C of hepatitis B virus identified in isolates from Australian Aborigines: complete genome sequence and phylogenetic relatedness. *J Gen Virol* 2001; 82 (Pt 4): 883–92.
- 30 Shin IT, Tanaka Y, Tateno Y, Mizokami M. Development and public release of a comprehensive hepatitis virus database. *Hepatol Res* 2008; 38: 234–43.
- 31 Chisari FV, Ferrari C. Hepatitis B virus immunopathogenesis. *Annu Rev Immunol* 1995; 13: 29–60.
- 32 Carman WF, Jacyna MR, Hadziyannis S *et al.* Mutation preventing formation of hepatitis B e antigen in patients with chronic hepatitis B infection. *Lancet* 1989; 2 (8663): 588–91.
- 33 Buckwold VE, Xu Z, Chen M, Yen TS, Ou JH. Effects of a naturally occurring mutation in the hepatitis B virus basal core promoter on pre-core gene expression and viral replication. *J Virol* 1996; 70: 5845–51.
- 34 Okamoto H, Tsuda F, Akahane Y *et al.* Hepatitis B virus with mutations in the core promoter for an e antigen-negative phenotype in carriers with antibody to e antigen. *J Virol* 1994; 68: 8102–10.
- 35 Lamberts C, Nassal M, Velhagen I, Zentgraf H, Schroder CH. Pre-core-mediated inhibition of hepatitis B virus progeny DNA synthesis. *J Virol* 1993; 67: 3756–62.
- 36 Chen MT, Billaud JN, Sallberg M *et al.* A function of the hepatitis B virus pre-core protein is to regulate the immune response to the core antigen. *Proc Natl Acad Sci USA* 2004; 101: 14913–8.
- 37 Chen M, Sallberg M, Hughes J *et al.* Immune tolerance split between hepatitis B virus pre-core and core proteins. *J Virol* 2005; 79: 3016–27.
- 38 Bocharov G, Ludewig B, Bertoletti A *et al.* Underwhelming the immune response: effect of slow virus growth on CD8+ T-lymphocyte responses. *J Virol* 2004; 78: 2247–54.
- 39 Sainokami S, Abe K, Sato A *et al.* Initial load of hepatitis B virus (HBV), its changing profile, and pre-core/core promoter mutations correlate with the severity and outcome of acute HBV infection. *J Gastroenterol* 2007; 42: 241–9.
- 40 Tassopoulos NC, Papaevangelou GJ, Roumeliotou-Karayannis A, Ticehurst JR, Feinstone SM, Purcell RH. Search for hepatitis B virus DNA in sera from patients with acute type B or non-A, non-B hepatitis. *J Hepatol* 1986; 2: 410–8.

Pretreatment Prediction of Virological Response to Peginterferon Plus Ribavirin Therapy in Patients with Chronic Hepatitis C, Using Viral and Host Factors: Some Concerns

To the Editor:

We read with great interest the article published in a recent issue of *HEPATOLOGY*.¹ Shirakawa et al. enrolled 120 patients with chronic hepatitis C (CHC) infected with genotype 1 hepatitis C virus (HCV-1) and high baseline viral loads (HVL), defined by HCV RNA levels $\geq 10^5$ international units (IU)/mL measured by quantitative Cobas Amplicor assays (Roche Diagnostics Co. Ltd, Tokyo, Japan), who underwent combination therapy scheduled for 48 (85%) or 72 (15%) weeks. The authors concluded that the sequence of interferon sensitivity determining region of the HCV, the T-helper type 1 and type 2 (Th1/Th2) ratio, body weight, and neutrophil count can be useful for accurately predicting actual sustained virologic response (SVR) rate before combination therapy.

The definition of HVL ($\geq 10^5$ IU/mL) by the authors is not in accordance with the Asian Pacific Association for the Study of the Liver consensus statements ($\geq 4 \times 10^5$ IU/mL)² or other studies ($\geq 8 \times 10^5$ IU/mL).^{3,4} Whether this definition influences the results deserves further study. The rates of rapid virologic response (RVR, defined as HCV RNA negative at week 4) and SVR were 45% and 21.7%, respectively, in their study. We examined the 200 Taiwanese patients with HCV-1 CHC in our randomized trial published recently⁵ and found that 150 (75%) of these patients had serum HCV RNA $\geq 10^5$ IU/mL (HVL). The RVR and SVR rates of 150 patients were 64% and 34.7%, respectively, after receiving combination therapy with peginterferon alpha-2a plus oral ribavirin. A significantly higher SVR rate in patients treated for 48 weeks than 24 weeks (78.5%, 62 of 79 versus 35.4%, 34 of 71, $P < 0.001$) was observed. Recently, Ide et al. reported a 36% (20 of 56) SVR rate in Japanese patients with CHC and HVL treated with peginterferon alpha-2b/ribavirin for 48 weeks.⁶ Accordingly, we have noticed the SVR rate and RVR rate in Taiwanese patients with HCV-1 with HVL treated for 24 or 48 weeks were higher than Japanese patients treated for 48 or 72 weeks. The important role of the weight-based ribavirin exposure during the first 4 weeks of combination therapy on the achievement of an RVR has been highlighted previously.⁷ A cut-off point of 13 mg/kg/day of the first 4 weeks of weight-based ribavirin exposure have been reported associated with the achievement of an RVR.⁸ Patients in the study by Shirakawa et al. would have an exposure of less than 13.1 mg/kg/day ribavirin exposure by 4 weeks (starting dose: for body weight 60 kg or less; 600 mg/day, 61 kg to 80 kg; 800 mg/day, 81 kg or more; 1000 mg/day) which is lower than 17.4 mg/kg/day (starting dose: for body weight 75 kg or less; 1000 mg/day, 75 kg or more; 1200 mg/day) in Taiwanese patients infected with HCV-1. We agree that the valuable finding of pretreatment predictors for SVR rate by Shirakawa et al. should be applauded. The relative low SVR and RVR rates, even with longer duration of treatment, compared to Taiwanese patients infected with HCV-1 with HVL raise a concern of relative suboptimal ribavirin exposure, and the actual role of these pretreatment predictors have to be confirmed by further studies.

CHIA-YEN DAI^{1,3}

WAN-LONG CHUANG^{1,2}

JEE-FU HUANG^{2,4}

MING-YEN HSIEH²

MING-LUNG YU^{1,2,4}

¹Faculty of Internal Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

²Hepatobiliary Division, Department of Internal Medicine, and

³Department of Occupational Medicine, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

⁴Department of Internal Medicine, Kaohsiung Municipal Hsiao-Kang Hospital, Kaohsiung, Taiwan

References

- Shirakawa H, Matsumoto A, Joshita S, Komatsu M, Tanaka N, Umemura T, et al. Pretreatment prediction of virological response to peginterferon plus ribavirin therapy in chronic hepatitis C patients using viral and host factors. *HEPATOLOGY* 2008;48:1753-1760.
- Asian Pacific Association for the Study of the Liver (APASL) Hepatitis C Working Party, McCaughan GW, Omata M, Amarapurkar D, Bowden S, Chow WC, Chutaputti A, et al. Asian Pacific Association for the Study of the Liver consensus statements on the diagnosis, management and treatment of hepatitis C virus infection. *J Gastroenterol Hepatol* 2007;22:615-633.
- Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL Jr, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- Hadziyannis SJ, Sette H, Morgan TR, Balan V, Diago M, Marcellin P, et al. Peginterferon alfa-2a (40 kilodaltons) and ribovirin combination therapy in chronic hepatitis C: randomized study of the effect of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140:346-355.
- Yu ML, Dai CY, Huang JF, Chiu CF, Yang YH, Hou NJ, et al. Rapid virological response and treatment duration for chronic hepatitis c genotype 1 patients: a randomized trial. *HEPATOLOGY* 2008;47:1884-1893.
- Ide T, Hino T, Ogata K, Miyajima I, Kuwahara R, Kuhara K, et al. A Randomized study of extended treatment with peginterferon alpha-2b plus ribavirin based on time to HCV RNA negative-status in patients with genotype 1b chronic hepatitis C. *Am J Gastroenterol* 2009;104:70-75.
- Yu ML, Dai CY, Huang JF, Chuang WL. Rapid virological response to peginterferon and ribavirin for hepatitis C genotype 1: the role of weight-based ribavirin exposure. *HEPATOLOGY* 2008;48:1019-1020.
- Rodriguez-Torres M, Sulkowski M, Chung RT, Hamzeh FM, Jensen D. Association of pretreatment and on-treatment factors with rapid virological response in HCV genotype 1-infected patients with peginterferon alfa-2a/ribavirin [Abstract]. *HEPATOLOGY* 2007;46(Suppl):S17A.

Copyright © 2009 by the American Association for the Study of Liver Diseases. Published online in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/hep.22826

Potential conflict of interest: Nothing to report.

Reply:

First, we have to point out a mistake that Dai et al. made in their letter: the rapid virological response (RVR) and sustained virological response (SVR) rates are described inversely with respect to our data and maybe with respect to their data.

Dai et al. reported that the SVR rate of 64% in their study is apparently higher than the rate of 45% in our study. We think that this difference can be attributed to the difference in the mean ages of the patients enrolled in the two studies. The mean age in their study seems to have been around 50 years¹ and was over 10 years lower than that in our study (62 years old). Younger patients tended to show higher SVR rates in our study as well, and the SVR rate in patients who were younger than 55 years old was 61%, which is comparable to the rate reported by Dai et al.

It is possible that a higher administration dose of ribavirin during the first 4 weeks is associated with a higher RVR rate. However, we cannot comment on this because our study was not designed for that purpose.

We adopted the definition of high viral load generally used in Japan (10^5 IU/mL). Since Dai et al. suggested that a difference in the definition of high viral load influenced the results, we recalculated factors associated with SVR in 103 patients whose viral load was higher than 4×10^5 IU/mL and also in 89 patients whose viral load was higher than 8×10^5 IU/mL. The same four prediction factors as those provided by the original analysis of 120 patients whose viral load was higher than 10^5 IU/mL were again selected.

Distribution of Hepatitis B Virus Genotypes among Patients with Chronic Infection in Japan Shifting toward an Increase of Genotype A[∇]

Kentaro Matsuura,^{1,2} Yasuhito Tanaka,^{1*} Shuhei Hige,³ Gotaro Yamada,⁴ Yoshikazu Murawaki,⁵ Masafumi Komatsu,⁶ Tomoyuki Kuramitsu,⁷ Sumio Kawata,⁸ Eiji Tanaka,⁹ Namiki Izumi,¹⁰ Chiaki Okuse,¹¹ Shinichi Kakumu,¹² Takeshi Okanoue,¹³ Keisuke Hino,¹⁴ Yoichi Hiasa,¹⁵ Michio Sata,¹⁶ Tatsuji Maeshiro,¹⁷ Fuminaka Sugauchi,² Shunsuke Nojiri,² Takashi Joh,² Yuzo Miyakawa,¹⁸ and Masashi Mizokami^{1,19}

Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan¹; Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan²; Department of Internal Medicine, Hokkaido University Graduate School of Medicine, Sapporo, Japan³; Department of Basic Laboratory Sciences, Kawasaki Medical School, Kawasaki Hospital, Okayama, Japan⁴; Division of Medicine and Clinical Science, Faculty of Medicine, Tottori University, Tottori, Japan⁵; Department of Gastroenterology, Akita City Hospital, Akita, Japan⁶; Kuramitsu Clinic, Akita, Japan⁷; Department of Gastroenterology, Yamagata University School of Medicine, Yamagata, Japan⁸; Department of Medicine, Shinshu University School of Medicine, Matsumoto, Japan⁹; Division of Gastroenterology and Hepatology, Musashino Red Cross Hospital, Tokyo, Japan¹⁰; Department of Internal Medicine, Division of Gastroenterology and Hepatology, St. Marianna University School of Medicine, Kawasaki, Japan¹¹; Department of Gastroenterology, Aichi Medical University School of Medicine, Aichi, Japan¹²; Molecular Gastroenterology and Hepatology, Kyoto Prefectural University of Medicine, Kyoto, Japan¹³; Department of Gastroenterology and Hepatology, Yamaguchi University Graduate School of Medicine, Yamaguchi, Japan¹⁴; Department of Gastroenterology and Metabolism, Ehime University Graduate School of Medicine, Ehime, Japan¹⁵; Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Fukuoka, Japan¹⁶; First Department of Internal Medicine, University Hospital, Faculty of Medicine, University of the Ryukyus, Okinawa, Japan¹⁷; Miyakawa Memorial Research Foundation, Tokyo, Japan¹⁸; and Research Center for Hepatitis and Immunology, Kohnodai Hospital International Medical Center of Japan, Ichikawa, Japan¹⁹

Received 29 October 2008/Returned for modification 17 December 2008/Accepted 2 March 2009

Acute hepatitis B virus (HBV) infection has been increasing through promiscuous sexual contacts, and HBV genotype A (HBV/A) is frequent in patients with acute hepatitis B (AHB) in Japan. To compare the geographic distribution of HBV genotypes in patients with chronic hepatitis B (CHB) in Japan between 2005 and 2006 and between 2000 and 2001, with special attention to changes in the proportion of HBV/A, a cohort study was performed to survey changes in genotypes of CHB patients at 16 hospitals throughout Japan. Furthermore, we investigated the clinical characteristics of each genotype and examined the genomic characteristics of HBV/A isolates by molecular evolutionary analyses. Of the 1,271 patients, 3.5%, 14.1%, and 82.3% were infected with HBV/A, -B, and -C, respectively. In comparison with our previous survey during 2000 and 2001, HBV/A was twice as frequent (3.5% versus 1.7%; $P = 0.02$). The mean age was lower in the patients with HBV/A than in those with HBV/B or -C. Based on phylogenetic analyses of 11 full-length genomes and 29 pre-S2/S region sequences from patients, HBV/A isolates were imported from Europe and the United States, as well as the Philippines and India. They clustered with HBV/A from AHB patients and have spread throughout Japan. HBV/A has been increasing in CHB patients in Japan as a consequence of AHB spreading in the younger generation through promiscuous sexual contacts, aided by a tendency of HBV/A to induce chronic hepatitis. The spread of HBV/A infection in Japan should be prevented by universal vaccination programs.

Hepatitis B virus (HBV), a member of the *Hepadnaviridae*, is a circular, partially double-stranded DNA virus and is one of the major causes of chronic liver diseases, including chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC).

The HBV genome is composed of approximately 3,200 nucleotides. HBV is classified into eight genotypes, designated A to H, based on an intergroup divergence of 8% or more in the complete nucleotide sequence (3, 23, 26, 37). They have dis-

tinct geographical distributions and are associated with differences in clinical and virological characteristics, such as severity of liver disease and response to antiviral therapies (7, 8, 12, 13, 22, 28). Furthermore, subgenotypes have been reported for HBV/A, -B, and -C and named A1 to -3 (17, 38), B1 to -6 (31, 32, 40), and C1 to -6 (20, 31, 45). Equally, other genotypes are classified into subgenotypes. There have been increasing lines of evidence to indicate influences of HBV subgenotypes on the outcome of liver disease and the response to antiviral therapies (1, 39, 44).

In 2001, we reported the geographic distribution of HBV genotypes in Japan (27). Of the 720 Japanese patients with chronic HBV infection (CHB), 12 (1.7%) harbored HBV/A, 88 (12.2%) HBV/B, 610 (84.7%) HBV/C, 3 (0.4%) HBV/D, and 7 (1.0%) mixed genotypes. HBV/C was detected in over 94%

* Corresponding author. Mailing address: Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan. Phone: 81-52-853-8292. Fax: 81-52-842-0021. E-mail: ytanaka@med.nagoya-cu.ac.jp.

[∇] Published ahead of print on 18 March 2009.

of patients on the Japanese mainland, while HBV/B was found in 64% of those in Okinawa, the southernmost islands, and 44% of those in the Tohoku area in the northern part of the mainland.

Recently, acute HBV infection (AHB) has been increasing in Japan, predominantly through promiscuous sexual contacts. In addition, it was reported that HBV/A was more frequent in patients with acute hepatitis than in those with chronic hepatitis (29, 41, 49). Recent studies suggest that the chances for progression to chronic disease may differ among patients acutely infected with HBV of distinct genotypes (21, 25); patients infected with HBV/A run an increased risk of becoming HBV carriers. Hence, it is of utmost concern whether chronic HBV/A infection is increasing in Japan.

In the present study, we compared the geographic distribution of HBV genotypes in Japan during 2005 and 2006 with 2000 and 2001, with special attention to changes in the proportion of HBV/A. Furthermore, we investigated the clinical characteristics of each genotype and examined the genomic characteristics of HBV/A isolates by molecular evolutionary analyses.

MATERIALS AND METHODS

Patients. From September 2005 to October 2006, sera were collected from 1,370 consecutive patients with CHB at 16 representative hospitals that were liver centers in their respective regions throughout Japan for the purpose of investigating the geographic distribution of HBV genotypes in Japan. All of the patients were diagnosed after they had been followed for at least 12 months. Patients diagnosed with AHB were excluded from the study; they had a sudden onset of clinical symptoms of hepatitis, along with high-titer antibody to HBV core antigen of the immunoglobulin M class in serum. Their sera were tested for alanine aminotransferase (ALT), alkaline phosphatase (ALP), γ -glutamyl transpeptidase (γ -GTP), and hepatitis B e antigen (HBeAg), as well as antibody to HBeAg (anti-HBe) (Dinabot, Tokyo, Japan). Four clinical diagnoses were established for them. The inactive carrier state was defined by the presence of HBV surface antigen (HBsAg) with normal ALT levels over 1 year (examined at least four times at 3-month intervals) and without evidence of portal hypertension. Chronic hepatitis was defined by elevated ALT levels (>1.5 times the upper limit of normal [35 IU/liter]) persisting over 6 months (with at least three bimonthly tests). Cirrhosis was diagnosed principally by ultrasonography (coarse liver architecture, nodular liver surface, blunt liver edges, and hypersplenism), platelet counts of $<100,000/\text{cm}^3$, or a combination thereof. Histological confirmation by fine-needle biopsy of the liver was performed as required. HCC was diagnosed by ultrasonography, computerized tomography, magnetic resonance imaging, angiography, tumor biopsy, or a combination thereof.

The study protocol conformed to the 1975 declaration of Helsinki and was approved by the ethics committees of the respective institutions. Every patient or his/her next of kin gave informed consent to the purpose of the study.

Genotypes and subgenotypes of HBV. The six HBV genotypes (A to F) were determined serologically by enzyme immunoassay (EIA) using commercial kits (HBV Genotype EIA; Institutes of Immunology Co., Ltd., Tokyo, Japan). The method depends on the combination of epitopes on pre-S2 region products detected by monoclonal antibodies that were specific for each of them (46, 47). Subgenotypes of HBV/A, designated A1 and A2, were determined by direct sequencing of the pre-S2/S gene, followed by a phylogenetic analysis.

Quantification of HBV DNA and sequencing. HBV DNA levels in sera were quantitated with a commercial kit (Amplicor HBV Monitor; Roche Diagnostics, Basel, Switzerland) with a detection range from 2.6 to 7.6 log copies/ml. Nucleic acids were extracted from 100 μl of serum using the Qiaamp DNA Blood Minikit (Qiagen GmbH, Hilden, Germany). Eleven complete HBV/A genomes and 29 pre-S2/S region sequences were amplified by PCR with appropriate primer sets, as described previously (40). The amplified HBV DNA fragments were directly sequenced using the ABI Prism Big Dye kit version 3.0 (Applied Biosystems, Foster City, CA) in an ABI 3100 automated DNA sequencer (Applied Biosystems). All sequences were analyzed in both forward and reverse directions. Complete and partial HBV genome sequences were aligned using GENETYX version 11.0 (Software Development Co., Ltd., Tokyo, Japan).

TABLE 1. Characteristics of 1,271 CHB patients

Parameter	Value
Characteristic	
Male gender [no. (%)]	766 (60.3)
Age (yr; mean \pm SD)	51.4 \pm 14.0
Diagnosis	
Inactive carrier state [no. (%)]	206 (16.2)
Chronic hepatitis [no. (%)]	786 (61.8)
Cirrhosis [no. (%)]	175 (13.8)
HCC [no. (%)]	104 (8.2)
Antiviral treatment [no. (%)]	577 (45.4)
Blood tests	
Platelets ($10^4/\text{mm}^3$)	21.4 \pm 30.2
ALT (IU/liter)	59.8 \pm 103.0
ALP (IU/liter)	270.4 \pm 136.0
γ -GTP (IU/liter)	47.4 \pm 66.1
HBV markers	
HBeAg [no. (%)]	399 (31.4)
HBV DNA (median [range] [log copies/ml])	4.2 (<2.6 to >7.6)

Molecular evolutionary analysis of HBV. Reference sequences were retrieved from the DDBJ/EMBL/GenBank databases with their accession numbers for identification. To investigate the relationship between HBV isolates from patients with chronic and acute hepatitis B in Japan, HBV/A isolates (AH1 to -10) were randomly retrieved from them and sequenced in our previous study (29). Nucleotide sequences of HBV DNA were aligned by the program CLUSTAL X, and genetic distance was estimated by the six-parameter method (10) in the Hepatitis Virus Database (36). Based on these values, phylogenetic trees were constructed by the neighbor-joining method (30) with the midpoint rooting option. To confirm the reliability of the phylogenetic trees, bootstrap resampling tests were performed 1,000 times.

Statistical analysis. Categorical variables were compared between groups by the χ^2 test or Fisher's exact test and noncategorical variables by the Mann-Whitney U test. A *P* value of less than 0.05 was considered significant.

Nucleotide sequence accession numbers. The DDBJ/EMBL/GenBank accession numbers of the complete genome sequences of HBV isolates JPN_CH1 to -11 are AB453979 to AB453989.

RESULTS

Distribution of HBV genotypes among patients with CHB.

Of the 1,370 serum samples, the genotype could not be determined for 99 (7.2%) by EIA due to low HBsAg levels, leaving 1,271 for analysis in this study (Table 1). Of these, 206 (16.2%) were inactive carriers, 786 (61.8%) had chronic hepatitis, 175 (13.8%) cirrhosis, and 104 (8.2%) HCC. They had a mean age of 51.4 \pm 14.0 years and included 766 (60.3%) men. They had a median HBV DNA level of 4.2 log copies/ml, and 399 (31.4%) of them were positive for HBeAg. Antiviral treatment had been given to 577 (45.4%) of them with interferon, lamivudine, adefovir pivoxil, or entecavir.

The genotypes were HBV/A in 44 (3.5%), HBV/B in 179 (14.1%), HBV/C in 1,046 (82.2%), and HBV/D in 2 (0.2%) (Table 2). In comparison with our previous report on the distribution of genotypes in Japan in 2001 (27), HBV/A was more frequent in this study (3.5% versus 1.7%; *P* = 0.02). Of the 16 hospitals in this study, 10 overlapped with those in our previous report from 2001. In these 10 hospitals, HBV/A was more frequent in the present than in the previous survey (3.6% versus 1.7%; *P* = 0.04).

The distribution of HBV genotypes in Japan differed by

TABLE 2. Distribution of HBV Genotypes

Genotype	No. (%)	
	2005–2006 (n = 1,271)	2000–2001 ^a (n = 720)
A	44 (3.5 ^b)	12 (1.7)
B	179 (14.1)	88 (12.2)
C	1,046 (82.3)	610 (84.7)
D	2 (0.2)	3 (0.4)
Mixed	0 (0.0)	7 (1.0)

^a From Orito et al. (27).^b $P = 0.02$.

geographic location (Fig. 1). HBV/C was the most prevalent in the majority of areas. In the Tohoku area, the northern part of the Japanese mainland (Honshu), HBV/B was more prevalent than in the other areas of the Japanese mainland. In Okinawa, the southernmost islands of Japan, HBV/B was predominant. Of note, HBV/A was more frequent in the Kanto area (9.5%), the metropolitan area, and Okinawa (9.1%) than in the other areas.

Clinical differences among HBV/A, -B, and -C. Clinical backgrounds were compared among the patients infected with HBV/A, -B, and -C (Table 3). HBeAg was significantly less prevalent in the patients infected with HBV/B than in those infected with HBV/A or -C ($P < 0.01$ for each). When the positivity of HBeAg was stratified by age, HBeAg was markedly less common in patients infected with HBV/B than in those infected with HBV/A or -C who were older than 40 years of age (7/157 [4.5%] versus 4/19 [21.1%] [$P < 0.05$] or 215/755 [28.5%] [$P < 0.01$]) (Fig. 2). There were no significant differences in HBV DNA levels among patients infected with the three genotypes. As antiviral treatments might have influenced the severity of liver disease, clinical states were compared among patients infected with HBV/A, -B, and -C who did and

did not receive it; antiviral treatments did not affect the above-mentioned trends represented in Table 3 in age, diagnosis, and HBeAg, as well as ALT and HBV DNA levels (data not shown).

Additionally, we compared the distributions of age and liver diseases in patients infected with HBV/A, -B, and -C. In patients infected with HBV/C, the prevalence of cirrhosis and HCC increased in those older than 50 years of age compared to younger patients (Fig. 3), whereas in the patients infected with HBV/B, cirrhosis and HCC were rare in elderly patients. The proportion of patients younger than 40 years of age was higher in those infected with HBV/A than in those infected with HBV/B or -C (25/44 [56.8%] versus 22/179 [12.3%] or 288/1,046 [27.5%]; $P < 0.01$ for each), while cirrhosis and HCC were also found in those older than 50 years of age infected with HBV/A.

Coinfection with human immunodeficiency virus type 1 (HIV-1) was found in 6 of the 44 (13.6%) patients infected with HBV/A compared to only 3 of the 1,046 (0.3%) patients infected with HBV/C ($P < 0.0001$); it occurred in none of the 179 patients infected with HBV/B.

Phylogenetic analyses. Among the 44 HBV/A isolates, the complete genome was sequenced successfully in 11 (JPN_CH1 to -11). Seven of them were classified as HBV/A2 and four as HBV/A1. A phylogenetic tree was constructed based on the complete genome sequences of these 11 isolates, along with those from two patients with AHB and those from 40 HBV/A isolates retrieved from the database (Fig. 4). Of the seven HBV/A2 isolates, the four from patients with CHB in this study formed a cluster with the Japanese isolates retrieved from the database and two from patients with AHB. Of the other three isolates, JPN_CH5 clustered with French and U.S. isolates, JPN_CH6 with German isolates, and JPN_CH7 with




FIG. 1. Geographic distribution of HBV genotypes in patients with chronic HBV infection in Japan during 2005 and 2006.

TABLE 3. Clinical characteristics of individuals chronically infected with HBV of different genotypes

Parameter	Value for genotype:		
	A (n = 44)	B (n = 179)	C (n = 1,046)
Male gender [no. (%)]	32 (72.7)	112 (62.6)	621 (59.4)
Age (yr [mean \pm SD])	41.3 \pm 14.9 ^a	55.8 \pm 13.7 ^b	48.8 \pm 13.3
Diagnosis			
Inactive carrier state [no. (%)]	13 (29.5) ^c	63 (35.2) ^b	129 (12.3)
Chronic hepatitis [no. (%)]	26 (59)	103 (57.5)	656 (62.7)
Cirrhosis [no. (%)]	3 (6.8)	10 (5.6) ^b	162 (15.5)
HCC [no. (%)]	2 (4.5)	3 (1.7) ^b	99 (9.5)
Anti viral treatment [no. (%)]	13 (29.5) ^d	48 (26.8) ^b	516 (49.3)
Blood tests			
Platelet ($10^4/\text{mm}^3$)	23.3 \pm 21.9	25.9 \pm 35.9 ^e	20.6 \pm 29.5
ALT (IU/liter)	56.2 \pm 83.8	42.2 \pm 104.2 ^e	63.0 \pm 103.3
ALP (U/liter)	247.1 \pm 123.0	255.5 \pm 97.9	273.9 \pm 141.9
γ -GTP (U/liter)	39.6 \pm 34.6	49.3 \pm 63.4	47.5 \pm 67.6
HBV markers			
HBeAg [positive rate(%)]	15 (34.0) ^f	17 (9.5) ^b	367 (35.1)
HBV DNA (median [range]) (log copies/ml)	4.2 (<2.6-7.6)	4.1 (<2.6-7.6)	4.2 (<2.6-7.6)

^a $P < 0.01$, A versus B or C.

^b $P < 0.01$, B versus C.

^c $P < 0.01$, A versus C.

^d $P < 0.05$, A versus C.

^e $P < 0.05$, B versus C.

^f $P < 0.01$, A versus B.

Spanish and Italian isolates. All four HBV/A1 isolates in this study formed a cluster with Philippine and Indian isolates.

In addition, the pre-S2/S region sequences of a total of 29 isolates were determined, including the 11 isolates whose complete genomes were sequenced. Of these, 21 (72%) were classified as HBV/A2 and the remaining 8 as HBV/A1. A phylogenetic tree was constructed based on the pre-S2/S region sequences from the 29 isolates, along with those from 10 patients with AHB infected with HBV/A and 47 HBV/A isolates retrieved from the database (Fig. 5). The 21 HBV/A2 isolates in the present study formed a cluster with Japanese, American, and European isolates retrieved from the database and those from patients with acute hepatitis. In addition, some of them were highly homologous with each other. Likewise, HBV/A1 isolates from eight patients with chronic hepatitis in this study

were highly homologous with those from two patients with acute hepatitis and isolates from the Philippines and India. Based on the phylogenetic analyses, HBV/A isolates were imported from Europe and the United States, as well as the Philippines and India, and had infiltrated throughout Japan.

DISCUSSION

Perinatal transmission from carrier mothers to their babies has been the principal route for establishing persistent HBV infection in Asian countries (19). In Japan, passive and active immunoprophylaxis with HBV immune globulin and vaccine has been mandated for babies born to HBeAg-positive carrier mothers since 1986; this was extended to HBeAg-negative carrier mothers in 1995. As a result, HBsAg has become rare in Japanese born after 1986; it was detected in only 0.2% of first-time blood donors younger than 19 years of age in 2000 (24). However, AHB has been increasing in Japan, predominantly through promiscuous sexual contacts.

In Japan, HBV/A is detected rarely among patients with CHB but is frequent in those with acute hepatitis (14, 25, 29, 41, 43). Yotsuyanagi et al. reported the distribution of genotypes in 145 Japanese patients with AHB and found HBV/A in 27 (19%), HBV/B in 8 (5%), and HBV/C in 109 (75%) (49). HBV/A is more frequent in metropolitan areas than other areas. The majority of patients with HBV/A infection in metropolitan areas have had extramarital sexual contacts with multiple irregular partners, through which they could have contracted infection. In support of this view, among men who have sex with men (MSM) who are coinfecting with HBV and HIV-1 in Tokyo, most were infected with HBV/A (15, 35).

In Japan, AHB in adulthood becomes chronic in only ~1%




FIG. 2. Prevalence of HBeAg among patients infected with HBV of different genotypes stratified by the age.




FIG. 3. Distribution of HCC, cirrhosis, chronic hepatitis, and inactive carrier state among the 1,271 patients infected with HBV of different genotypes stratified by the age.

of cases. This is much less than the progression to chronic disease (close to 10%) in Europe and the United States, where HBV/A prevails (34). Recent studies have suggested that the chances for persistence may differ among patients acutely infected with HBV of distinct genotypes (21, 25). In particular, acute infection with HBV/A may bring about an increased risk of progression to chronic disease. Therefore, an increase of acute infection with HBV/A would result in a surge of HBV/A among patients with CHB in Japan. In actuality, in comparison with our previous results during 2000 and 2001 (27), HBV/A was twice as frequent in this study (3.5% versus 1.7%; $P = 0.02$). HBV/A has been increasing in patients with CHB in the Kanto area, where HBV/A in patients with acute hepatitis is more frequent than in the other areas. In the islands of Okinawa, also, HBV/A was found to be prevalent in this study. Of the four patients infected with HBV/A there, two were coinfecting with HIV-1. They were both MSM, and they were sus-




FIG. 4. Phylogenetic tree constructed based on the complete genome sequences of HBV/A isolates. Those from 11 patients with chronic infection in this study are shown in boldface italic (JPN_CH1 to -11), along with two isolates (JPN_AH1 and -2) from patients with acute hepatitis in Japan reported in our previous study (17). Representative isolates were retrieved from the DDBJ/EMBL/GenBank databases, including 21 HBV/Ae, 10 HBV/Aa, and 2 HBV/Ac isolates, along with 7 HBV isolates representative of the other seven genotypes. Isolates from the databases are identified by accession numbers, followed by the country of origin. The bar at the bottom spans 0.01 nucleotide substitutions per site.

pected to have been infected with HIV through sexual contacts on the Japanese mainland. It has been reported that HIV infection increases the probability that AHBs will become chronic (2, 11, 33, 48). Because they share routes of transmission and the risk for HIV-1 and HBV infections, approximately




FIG. 5. Phylogenetic tree constructed based on pre-S2/S region sequences of HBV/A isolates. Those from 29 patients with chronic infection in this study are shown in boldface italic (JPN_CH 1 to -29), along with 10 isolates (JPN_AH 1 to -10) from patients with acute hepatitis in Japan reported in our previous study (17). Representative isolates were retrieved from the DDBJ/EMBL/GenBank databases, including 28 HBV/Ae, 10 HBV/Aa, and 2 HBV/Ac isolates and 7 HBV isolates representative of the other seven genotypes. Isolates from the databases are identified by accession numbers, followed by the country of origin. The bar at the bottom spans 0.01 nucleotide substitutions per site.

90% of patients with AIDS have markers of past or ongoing HBV infection (18). Thus, HBV carriers are more frequent in the HIV-1-positive than in the HIV-1-negative population (4, 9). Among patients with HIV infection in Japan, 6.3% are HBsAg positive, in particular, 8.3% of HIV-infected MSM (16). In this study, coinfection with HIV was found in 6 of the 44 (13.6%) patients infected with HBV/A. All of them were men. Their median age was 27.7 ± 4.1 years, and five patients were positive for HBeAg. Thus, there is a possibility that HIV-1 and HBV/A coinfections are increasing among young people in Japan, and the high rate of HBeAg positivity may be influenced by immune suppression due to HIV infection.

In the phylogenetic analysis, the HBV/A2 isolates recovered in this study were homologous to those from Europe and the United States, and some of them clustered with the Japanese isolates. On the other hand, there were HBV/A1 isolates that formed a cluster with those from the Philippines and India. Furthermore, some isolates from patients with acute hepatitis who were infected with HBV/A in Japan were highly homologous to HBV/A isolates from patients with chronic hepatitis. This invites speculation that some HBV/A isolates were introduced into Japan from foreign countries, while others have already settled down there and spread from patients with chronic infection to their contacts. HBV/A would have been infiltrating throughout Japan by these two different routes.

Clinical differences among patients infected with HBV/A, -B, and -C were observed. The mean age was lower in the patients infected with HBV/A than in those infected with HBV/B or -C. As mentioned above, AHB patients infected with HBV/A have been increasing in the younger generation in Japan, and around 10% of them would have progressed to chronic infection. This is one of the reasons why the patients infected with HBV/A are younger than those infected with HBV/B or -C. Most patients infected with HBV/B were negative for HBeAg, while a high proportion of the patients infected with HBV/A and -C had it. In particular, this difference was remarkable in the patients who were older than 40 years of age. Thus, the seroconversion rate for the loss of HBeAg among younger people may be higher in infection with HBV/B than in that with HBV/A or -C. Inactive carriers were commoner in HBV/A than in HBV/C infection, as well.

These lines of evidence indicate that the activity of hepatitis is lower in HBV/B than HBV/C infection, and patients with HBV/B seroconvert from HBeAg to anti-HBe at young ages. In addition, cirrhosis and HCC were less frequent in the patients infected with HBV/B than in those infected with HBV/C. Therefore, the prognosis would be better in the patients infected with HBV/B than in those infected with HBV/C. These results are in accord with previous reports (5, 13, 28, 42). There have been few reports on the clinical features of patients with chronic hepatitis infected with HBV/A in Japan. Chu et al. have reported the distribution of HBV genotypes with reference to clinical characteristics in the United States (6). They have shown that HBV/A and HBV/C infections are accompanied by a higher frequency of HBeAg than HBV/B infection, while HBV/B is associated with a lower rate of hepatic decompensation than HBV/A and -C. In our study, inactive carriers were commoner, while cirrhosis and HCC were found less often in HBV/A than in HBV/C infection. HBeAg was more prevalent in the patients infected with HBV/A than in those

infected with HBV/B who were older than 40 years of age. Therefore, it can be said that the prognosis is better for patients infected with HBV/A than for those infected with HBV/C; it may be poorer than for those infected with HBV/B.

In conclusion, HBV/A has been increasing among CHB patients in Japan. On the basis of phylogenetic analyses, some HBV/A isolates appear to have been imported from foreign countries. They clustered with HBV/A from AHB patients and have infiltrated throughout Japan. It is very likely that acute and chronic infections with HBV/A have been increasing in Japan. Obviously, immunoprophylaxis of perinatal HBV infection, implemented since 1986 on a national basis, has been insufficient to prevent horizontal HBV/A infection diffusing among high-risk groups by transmission routes shared by HIV infection. The foreseeable spread of HBV/A infection in Japan should be prevented by universal vaccination programs extended to high-risk groups or the general population.

ACKNOWLEDGMENTS

The study was supported in part by a grant-in-aid from the Ministry of Health, Labor and Welfare of Japan and a grant-in-aid from the Ministry of Education, Culture, Sports, Science and Technology.

We thank T. Kimura and K. Sato, Institutes of Immunology Co., Ltd. (Tokyo, Japan), for determining HBV genotypes in this study and Takashi Saito, Yamagata University Hospital; Akihiro Matsumoto, Shinshu University Hospital; Yasuhiro Asahina, Musashino Red Cross Hospital; Yoshito Ito, University Hospital, Kyoto Prefectural University of Medicine; Keiko Hoshio, Tottori University Hospital; Morikazu Onji, Ehime University Hospital; Tatsuya Ide, Kurume University Hospital; and Hiroshi Sakugawa, Hospital, University of the Ryukyus, for their help throughout this work.

Kentaro Matsuura wrote the study protocol and the first draft of the manuscript and performed the experiments and statistical analysis. Yasuhiro Tanaka contributed to the experimental work and the final version of the manuscript. Shuhei Hige, Gotaro Yamada, Yoshikazu Murawaki, Masafumi Komatsu, Tomoyuki Kuramitsu, Sumio Kawata, Eiji Tanaka, Namiki Izumi, Chiaki Okuse, Shinichi Kakumu, Takeshi Okanoue, Keisuke Hino, Yoichi Hiasa, Michio Sata, and Tatsuji Maeshiro contributed to the collection of the samples and clinical data from patients and to the final version of the manuscript. Fuminaka Sugauchi, Shunsuke Nojiri, Takashi Joh, and Yuzo Miyakawa contributed to the final version of the manuscript. Masashi Mizokami had the original idea and did the planning of the study and contributed to the final version of the manuscript. All of the authors have seen and approved the final draft of the manuscript.

REFERENCES

1. Akuta, N., F. Suzuki, M. Kobayashi, A. Tsubota, Y. Suzuki, T. Hosaka, T. Someya, S. Saitoh, Y. Arase, K. Ikeda, and H. Kumada. 2003. The influence of hepatitis B virus genotype on the development of lamivudine resistance during long-term treatment. *J. Hepatol.* 38:315-321.
2. Alter, M. J. 2006. Epidemiology of viral hepatitis and HIV co-infection. *J. Hepatol.* 44:S6-S9.
3. Arauz-Ruiz, P., H. Norder, B. H. Robertson, and L. O. Magnius. 2002. Genotype H: a new Amerindian genotype of hepatitis B virus revealed in Central America. *J. Gen. Virol.* 83:2059-2073.
4. Bodsworth, N. J., D. A. Cooper, and B. Donovan. 1991. The influence of human immunodeficiency virus type 1 infection on the development of the hepatitis B virus carrier state. *J. Infect. Dis.* 163:1138-1140.
5. Chu, C. J., M. Hussain, and A. S. Lok. 2002. Hepatitis B virus genotype B is associated with earlier HBeAg seroconversion compared with hepatitis B virus genotype C. *Gastroenterology* 122:1756-1762.
6. Chu, C. J., E. B. Keeffe, S. H. Han, R. P. Perrillo, A. D. Min, C. Soldevilla-Pico, W. Carey, R. S. Brown, Jr., V. A. Luketic, N. Terrault, and A. S. Lok. 2003. Hepatitis B virus genotypes in the United States: results of a nationwide study. *Gastroenterology* 125:444-451.
7. Chu, C. J., and A. S. Lok. 2002. Clinical significance of hepatitis B virus genotypes. *Hepatology* 35:1274-1276.
8. Ding, X., M. Mizokami, G. Yao, B. Xu, E. Orito, R. Ueda, and M. Nakanishi. 2001. Hepatitis B virus genotype distribution among chronic hepatitis B virus carriers in Shanghai, China. *Intervirology* 44:43-47.

9. Fujii, T., H. Taguchi, H. Katao, S. Mori, T. Nakamura, N. Nojiri, K. Nakajima, K. Tadokoro, T. Juji, and A. Iwamoto. 1999. Seroprevalence of human herpesvirus 8 in human immunodeficiency virus 1-positive and human immunodeficiency virus 1-negative populations in Japan. *J. Med. Virol.* 57:159-162.
10. Gojobori, T., K. Ishii, and M. Nei. 1982. Estimation of average number of nucleotide substitutions when the rate of substitution varies with nucleotide. *J. Mol. Evol.* 18:414-423.
11. Hadler, S. C., F. N. Judson, P. M. O'Malley, N. L. Altman, K. Penley, S. Buchbinder, C. A. Schable, P. J. Coleman, D. N. Ostrow, and D. P. Francis. 1991. Outcome of hepatitis B virus infection in homosexual men and its relation to prior human immunodeficiency virus infection. *J. Infect. Dis.* 163:454-459.
12. Kao, J. H. 2002. Clinical relevance of hepatitis B viral genotypes: a case of *deja vu*? *J. Gastroenterol. Hepatol.* 17:113-115.
13. Kao, J. H., P. J. Chen, M. Y. Lai, and D. S. Chen. 2000. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology* 118:554-559.
14. Kobayashi, M., Y. Arase, K. Ikeda, A. Tsubota, Y. Suzuki, S. Saitoh, F. Suzuki, N. Akuta, T. Someya, M. Matsuda, J. Sato, K. Takagi, Y. Miyakawa, and H. Kumada. 2002. Viral genotypes and response to interferon in patients with acute prolonged hepatitis B virus infection of adulthood in Japan. *J. Med. Virol.* 68:522-528.
15. Koibuchi, T., A. Hitani, T. Nakamura, N. Nojiri, K. Nakajima, T. Jyuji, and A. Iwamoto. 2001. Predominance of genotype A HBV in an HBV-HIV-1 dually positive population compared with an HIV-1-negative counterpart in Japan. *J. Med. Virol.* 64:435-440.
16. Koike, K., Y. Kikuchi, M. Kato, J. Takamatsu, Y. Shintani, T. Tsutsumi, H. Fujie, H. Miyoshi, K. Moriya, and H. Yotsuyanagi. 2008. Prevalence of hepatitis B virus infection in Japanese patients with HIV. *Hepatol. Res.* 38:310-314.
17. Kurbanov, F., Y. Tanaka, K. Fujiwara, F. Sugauchi, D. Mbanya, L. Zekeng, N. Ndembu, C. Ngansop, L. Kaptue, T. Miura, E. Ido, M. Hayami, H. Ichimura, and M. Mizokami. 2005. A new subtype (subgenotype) Ac (A3) of hepatitis B virus and recombination between genotypes A and E in Cameroon. *J. Gen. Virol.* 86:2047-2056.
18. Lebovics, E., B. M. Dworkin, S. K. Heier, and W. S. Rosenthal. 1988. The hepatobiliary manifestations of human immunodeficiency virus infection. *Am. J. Gastroenterol.* 83:1-7.
19. Lok, A. S. 1992. Natural history and control of perinatally acquired hepatitis B virus infection. *Dig. Dis.* 10:46-52.
20. Lusida, M. I., V. E. Nugrahaputra, Soetjipto, R. Handajani, M. Nagano-Fujii, M. Sasayama, T. Utsumi, and H. Hotta. 2008. Novel subgenotypes of hepatitis B virus genotypes C and D in Papua, Indonesia. *J. Clin. Microbiol.* 46:2160-2166.
21. Mayerat, C., A. Mantegani, and P. C. Frei. 1999. Does hepatitis B virus (HBV) genotype influence the clinical outcome of HBV infection? *J. Viral Hepat.* 6:299-304.
22. Miyakawa, Y., and M. Mizokami. 2003. Classifying hepatitis B virus genotypes. *Intervirology* 46:329-338.
23. Norder, H., B. Hammas, S. Lofdahl, A. M. Courouce, and L. O. Magnius. 1992. Comparison of the amino acid sequences of nine different serotypes of hepatitis B surface antigen and genomic classification of the corresponding hepatitis B virus strains. *J. Gen. Virol.* 73:1201-1208.
24. Noto, H., T. Terao, S. Ryou, Y. Hirose, T. Yoshida, H. Ookubo, H. Mito, and H. Yoshizawa. 2003. Combined passive and active immunoprophylaxis for preventing perinatal transmission of the hepatitis B virus carrier state in Shizuoka, Japan during 1980-1994. *J. Gastroenterol. Hepatol.* 18:943-949.
25. Ogawa, M., K. Hasegawa, T. Naritoni, N. Torii, and N. Hayashi. 2002. Clinical features and viral sequences of various genotypes of hepatitis B virus compared among patients with acute hepatitis B. *Hepatol. Res.* 23:167-177.
26. Okamoto, H., F. Tsuda, H. Sakugawa, R. I. Sastrosoewiguno, M. Imai, Y. Miyakawa, and M. Mayumi. 1988. Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J. Gen. Virol.* 69:2575-2583.
27. Orito, E., T. Ichida, H. Sakugawa, M. Sata, N. Horiike, K. Hino, K. Okita, T. Okanoue, S. Iino, E. Tanaka, K. Suzuki, H. Watanabe, S. Hige, and M. Mizokami. 2001. Geographic distribution of hepatitis B virus (HBV) genotype in patients with chronic HBV infection in Japan. *Hepatology* 34:590-594.
28. Orito, E., M. Mizokami, H. Sakugawa, K. Michitaka, K. Ishikawa, T. Ichida, T. Okanoue, H. Yotsuyanagi, S. Iino, et al. 2001. A case-control study for clinical and molecular biological differences between hepatitis B viruses of genotypes B and C. *Hepatology* 33:218-223.
29. Ozasa, A., Y. Tanaka, E. Orito, M. Sugiyama, J. H. Kang, S. Hige, T. Kuramitsu, K. Suzuki, E. Tanaka, S. Okada, H. Tokita, Y. Asahina, K. Inoue, S. Kakumu, T. Okanoue, Y. Murawaki, K. Hino, M. Onji, H. Yatsuhashi, H. Sakugawa, Y. Miyakawa, R. Ueda, and M. Mizokami. 2006. Influence of genotypes and precore mutations on fulminant or chronic outcome of acute hepatitis B virus infection. *Hepatology* 44:326-334.
30. Saitou, N., and M. Nei. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4:406-425.
31. Sakamoto, T., Y. Tanaka, E. Orito, J. Co, J. Clavio, F. Sugauchi, K. Ito, A. Ozasa, A. Quino, R. Ueda, J. Sollano, and M. Mizokami. 2006. Novel subtypes (subgenotypes) of hepatitis B virus genotypes B and C among chronic liver disease patients in the Philippines. *J. Gen. Virol.* 87:1873-1882.
32. Sakamoto, T., Y. Tanaka, J. Simonetti, C. Osiwoy, M. L. Borresen, A. Koch, F. Kurbanov, M. Sugiyama, G. Y. Minuk, B. J. McMahon, T. Joh, and M. Mizokami. 2007. Classification of hepatitis B virus genotype B into 2 major types based on characterization of a novel subgenotype in Arctic indigenous populations. *J. Infect. Dis.* 196:1487-1492.
33. Salmon-Ceron, D., C. Lewden, P. Morlat, S. Bevilacqua, E. Jougla, F. Bonnet, L. Heripret, D. Costagliola, T. May, and G. Chene. 2005. Liver disease as a major cause of death among HIV infected patients: role of hepatitis C and B viruses and alcohol. *J. Hepatol.* 42:799-805.
34. Sherlock, S. D. J. 1997. Virus hepatitis, p. 265-392. *In* S. D. J. Sherlock (ed.), *Diseases of the liver and biliary system*, 10th ed. Blackwell Scientific Publications, London, United Kingdom.
35. Shibayama, T., G. Masuda, A. Ajisawa, K. Hiruma, F. Tsuda, T. Nishizawa, M. Takahashi, and H. Okamoto. 2005. Characterization of seven genotypes (A to E, G and H) of hepatitis B virus recovered from Japanese patients infected with human immunodeficiency virus type 1. *J. Med. Virol.* 76:24-32.
36. Shin, I. T., Y. Tanaka, Y. Tateno, and M. Mizokami. 2008. Development and public release of a comprehensive hepatitis virus database. *Hepatol. Res.* 38:234-243.
37. Stuyver, L., S. De Gendt, C. Van Geyt, F. Zoulim, M. Fried, R. F. Schinazi, and R. Rossau. 2000. A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J. Gen. Virol.* 81:67-74.
38. Sugauchi, F., H. Kumada, S. A. Acharya, S. M. Shrestha, M. T. Gamutan, M. Khan, R. G. Gish, Y. Tanaka, T. Kato, E. Orito, R. Ueda, Y. Miyakawa, and M. Mizokami. 2004. Epidemiological and sequence differences between two subtypes (Ae and Aa) of hepatitis B virus genotype A. *J. Gen. Virol.* 85: 811-820.
39. Sugauchi, F., E. Orito, T. Ichida, H. Kato, H. Sakugawa, S. Kakumu, T. Ishida, A. Chutaputti, C. L. Lai, R. G. Gish, R. Ueda, Y. Miyakawa, and M. Mizokami. 2003. Epidemiologic and virologic characteristics of hepatitis B virus genotype B having the recombination with genotype C. *Gastroenterology* 124:925-932.
40. Sugauchi, F., E. Orito, T. Ichida, H. Kato, H. Sakugawa, S. Kakumu, T. Ishida, A. Chutaputti, C. L. Lai, R. Ueda, Y. Miyakawa, and M. Mizokami. 2002. Hepatitis B virus of genotype B with or without recombination with genotype C over the precore region plus the core gene. *J. Virol.* 76:5985-5992.
41. Sugauchi, F., E. Orito, T. Ohno, Y. Tanaka, A. Ozasa, J. H. Kang, J. Toyoda, T. Kuramitsu, K. Suzuki, E. Tanaka, Y. Akahane, T. Ichida, N. Izumi, K. Inoue, H. Hoshino, S. Iino, H. Yotsuyanagi, S. Kakumu, E. Tomita, T. Okanoue, S. Nishiguchi, Y. Murawaki, K. Hino, M. Onji, H. Yatsuhashi, M. Sata, Y. Miyakawa, R. Ueda, and M. Mizokami. 2006. Spatial and chronological differences in hepatitis B virus genotypes from patients with acute hepatitis B in Japan. *Hepatol. Res.* 36:107-114.
42. Sumi, H., O. Yokosuka, N. Seki, M. Arai, F. Imazeki, T. Kurihara, T. Kanda, K. Fukai, M. Kato, and H. Saisho. 2003. Influence of hepatitis B virus genotypes on the progression of chronic type B liver disease. *Hepatology* 37:19-26.
43. Suzuki, Y., M. Kobayashi, K. Ikeda, F. Suzuki, Y. Arfase, N. Akuta, T. Hosaka, S. Saitoh, T. Someya, M. Matsuda, J. Sato, S. Watabiki, Y. Miyakawa, and H. Kumada. 2005. Persistence of acute infection with hepatitis B virus genotype A and treatment in Japan. *J. Med. Virol.* 76:33-39.
44. Tanaka, Y., I. Hasegawa, T. Kato, E. Orito, N. Hirashima, S. K. Acharya, R. G. Gish, A. Kranvis, M. C. Kew, N. Yoshihara, S. M. Shrestha, M. Khan, Y. Miyakawa, and M. Mizokami. 2004. A case-control study for differences among hepatitis B virus infections of genotypes A (subtypes Aa and Ae) and D. *Hepatology* 40:747-755.
45. Tanaka, Y., E. Orito, M. F. Yuen, M. Mukaide, F. Sugauchi, K. Ito, A. Ozasa, T. Sakamoto, F. Kurbanov, C. L. Lai, and M. Mizokami. 2005. Two subtypes (subgenotypes) of hepatitis B virus genotype C: a novel subtyping assay based on restriction fragment length polymorphism. *Hepatol. Res.* 33:216-224.
46. Usuda, S., H. Okamoto, H. Iwanari, K. Baba, F. Tsuda, Y. Miyakawa, and M. Mayumi. 1999. Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J. Virol. Methods* 80:97-112.
47. Usuda, S., H. Okamoto, T. Tanaka, K. Kidd-Ljunggren, P. V. Holland, Y. Miyakawa, and M. Mayumi. 2000. Differentiation of hepatitis B virus genotypes D and E by ELISA using monoclonal antibodies to epitopes on the preS2-region product. *J. Virol. Methods* 87:81-89.
48. Weinbaum, C. M., K. M. Sabin, and S. S. Santibanez. 2005. Hepatitis B, hepatitis C, and HIV in correctional populations: a review of epidemiology and prevention. *AIDS* 19(Suppl. 3):S41-S46.
49. Yotsuyanagi, H., C. Okuse, K. Yasuda, E. Orito, S. Nishiguchi, J. Toyoda, E. Tomita, K. Hino, K. Okita, S. Murashima, M. Sata, H. Hoshino, Y. Miyakawa, and S. Iino. 2005. Distinct geographic distributions of hepatitis B virus genotypes in patients with acute infection in Japan. *J. Med. Virol.* 77:39-46.

Predictive values of amino acid sequences of the core and NS5A regions in antiviral therapy for hepatitis C: a Japanese multi-center study

Takeshi Okanoue · Yoshito Itoh · Hiroaki Hashimoto · Kohichiroh Yasui · Masahito Minami · Tetsuo Takehara · Eiji Tanaka · Morikazu Onji · Joji Toyota · Kazuaki Chayama · Kentaro Yoshioka · Namiki Izumi · Norio Akuta · Hiromitsu Kumada

Received: 31 March 2009 / Accepted: 20 April 2009 / Published online: 11 June 2009
© Springer 2009

Abstract

Background Chronic hepatitis C (CHC) genotype 1b patients with high viral load are resistant to peginterferon (PEG-IFN) and ribavirin (RBV) combination therapy, especially older and female patients.

Methods To elucidate the factors affecting early and sustained viral responses (EVR and SVR), 409 genotype 1b patients CHC with high viral loads who had received 48 weeks of PEG-IFN/RBV therapy were enrolled. The amino acid (aa) sequences of the HCV core at positions 70 and 91 and of the interferon sensitivity determining region (ISDR) were analyzed. Host factors, viral factors, and

treatment-related factors were subjected to multivariate analysis.

Results Male gender, low HCV RNA load, high platelet count, two or more aa mutations of ISDR, and wild type of core aa 70 were independent predictive factors for SVR. In patients with over 80% adherences to both PEG-IFN and RBV, male gender, mild fibrosis stage, and wild type of core aa 70 were independent predictors for SVR.

Conclusions Independent predictive factors for SVR were: no aa substitution at core aa 70, two or more aa mutations in the ISDR, low viral load, high values of platelet count, mild liver fibrosis and male gender.

T. Okanoue (✉)
Hepatology Center, Saiseikai Suita Hospital,
1-2 Kawazonocho, Suita 564-0013, Japan
e-mail: okanoue@suita.saiseikai.or.jp

T. Okanoue · Y. Itoh · H. Hashimoto · K. Yasui · M. Minami
Department of Gastroenterology and Hepatology,
Kyoto Prefectural University of Medicine,
Kawaramachi-Hirokoji Kamigyo-Ku,
Kyoto 602-8566, Japan

Y. Itoh
e-mail: yitoh@koto.kpu-m.ac.jp

H. Hashimoto
e-mail: road1820@yahoo.co.jp

K. Yasui
e-mail: yasui@koto.kpu-m.ac.jp

M. Minami
e-mail: minami@koto.kpu-m.ac.jp

T. Takehara
Department of Gastroenterology and Hepatology,
Osaka University Graduate School of Medicine,
2-2 Yamadaoka, Suita 565-0871, Japan
e-mail: takehara@gh.med.osaka-u.ac.jp

E. Tanaka
Department of Medicine, Shinshu University School
of Medicine, 3-1-1 Asahi, Matsumoto 390-8621, Japan
e-mail: etanaka@shinshu-u.ac.jp

M. Onji
Department of Gastroenterology, Ehime University,
454 Shizukawa, Tohon, Ehime 791-0295, Japan
e-mail: onjimori@m.med.ehime-u.ac.jp

J. Toyota
Department of Gastroenterology, Sapporo-Kosei General
Hospital, Kita Sanjyo, Chuo-ku, Sapporo 060-0033, Japan
e-mail: joji.toyota@ja_hokkaidoukouseiren.or.jp

K. Chayama
Department of Medicine and Molecular Science,
Graduate School of Science, Hiroshima University,
1-2-3 Kasumi, Minami-ku, Hiroshima,
Hiroshima 734-8551, Japan
e-mail: chayama@hiroshima-u.ac.jp

K. Yoshioka
Department of Hepato-Biliary-Pancreas, Fujita Health Science,
Kutsukake-cho, Toyoake 470-1192, Japan
e-mail: kyoshiok@fujita-hu.ac.jp

Keywords Chronic hepatitis C · Peginterferon and ribavirin · Core amino acid · Interferon sensitivity determining region

Abbreviations

CHC	Chronic hepatitis C
PEG-IFN	Peginterferon
RBV	Ribavirin
RVR	Rapid viral response
cEVR	Complete early viral response
LVR	Late viral response
ETR	End of treatment response
NR	Non response
SVR	Sustained viral response
ISDR	Interferon sensitivity determining region
Aa	Amino acid
ALT	Alanine aminotransferase
PLT	Platelet
HCC	Hepatocellular carcinoma

Introduction

A combination of pegylated interferon (PEG-IFN) and ribavirin (RBV) therapy for 48 weeks achieves a sustained viral response (SVR) rate of 40–50% in chronic hepatitis C (CHC) patients with a high viral load of genotype 1 [1–4]. The dose-reduction rate and the frequency of discontinuation of this treatment are high in aged patients [5]. The SVR rate of the therapy is lower in females than males, especially in older patients in Japan [6].

Around 30% of HCV carriers have serum alanine aminotransferase (ALT) levels within the upper limit of normal ranges [7, 8] and HCV carriers with persistently normal serum ALT (PNALT) and serum platelet (PLT) counts of over $15 \times 10^4/\text{mm}^3$ show low grade hepatic fibrosis and good prognosis [9]. Before treating HCV carriers, it is very important to predict non-response to PEG-IFN plus RBV therapy because of its medical cost, adverse effects, and its impact on the long term quality of life.

N. Izumi

Department of Gastroenterology and Hepatology,
Musashino Red Cross Hospital, Sakaiminamimachi,
Musashino 180-8610, Japan
e-mail: nizumi@musashino.jrc.or.jp

N. Akuta · H. Kumada

Department of Hepatology, Toranomon Hospital,
Kajigaya, Takatsu-ku, Kawasaki 213-8587, Japan

N. Akuta

e-mail: akuta-gi@umin.ac.jp

H. Kumada

e-mail: kumahiro@toranomon.gr.jp

There are many factors affecting response to IFN monotherapy and PEG-IFN/RBV therapy, including body mass index (BMI) [10, 11], steatosis [12, 13], insulin resistance [14], stage of liver fibrosis [15, 16], total cholesterol (T. Chol), triglyceride (TG), adherence to both PEG-IFN and RBV [17], race [18, 19], age [1, 2, 20], and viral factors including serum quantity of HCV RNA, HCV genotype and substitution of amino acids (aa) in the interferon sensitivity determining region (ISDR, 2209–2248) of the nonstructural protein 5A (NS5A) [21] and in the core protein [22, 23]. Early viral response is an important predictive factor in PEG-IFN/RBV therapy for CHC patients with genotype 1 and high viral loads [24–27].

The aim of this study was to elucidate the valuable predictive factors of SVR in Japanese patients with HCV genotype 1b high viral loads following 48 weeks of PEG-IFN/RBV therapy, focusing on the relationship between aa substitutions in the ISDR and at core aa 70 and 91 and early viral kinetics.

Patients and methods

Selection of patients

This retrospective study was conducted at 15 clinical sites in Japan which are part of the Study Group of Optimal Treatment of Viral Hepatitis supported by the Ministry of Health, Labor and Welfare, Japan. Eligible subjects were CHC patients, who (1) had received liver biopsy; (2) were genotype 1b with high viral load (≥ 100 KIU/ml by Cobas Amplicor Hepatitis C Virus Test, version 2.0) at the start of PEG-IFN/RBV therapy; (3) received weekly injections of PEG-IFN- α -2b (PEG-INTRON; Shering-Plough, Kenilworth, NJ) of 1.5 $\mu\text{g}/\text{kg}$ bw and oral administration of RBV (Rebetol; Shering-Plough) for 48 weeks. The amount of RBV was adjusted based on the subject's body weight; (600 mg for ≤ 60 kg bw, 800 mg for 60–80 kg bw, 1,000 mg for > 80 kg bw); (4) were examined serially for quantitative and qualitative HCV RNA; and (5) the aa sequences at positions 70 and 91 in the core region and of the ISDR in the NS5A had been determined in pretreatment sera.

Hepatitis B virus (HBV) infection, human immunodeficiency virus (HIV) infection, autoimmune hepatitis, primary biliary cirrhosis, hemochromatosis, and Wilson's disease were excluded. Histopathological diagnosis was based on the scoring system of Desmet et al. [28]. The definition of alcohol abuse included patients having a history of more than 100 kg of total ethanol intake. Complete blood counts, liver function tests, serum lipids, serum ferritin, serum fibrosis markers, fasting plasma glucose (FPG), and immune reactive insulin (IRI) were examined in most cases. Written informed consent was obtained from all

patients before treatment, and the protocol was approved by the ethics committees in each site.

Study design

Four hundred and nine patients who completed 48 weeks of treatment and were followed for more than 24 weeks after treatment were enrolled in the first study (*Study design 1*).

To elucidate the effect of aa substitution of HCV core and in the ISDR on HCV dynamics, including a rapid viral response (RVR), complete early viral response (cEVR), a late viral response (LVR) and SVR, according to gender and age (<60 years \geq 60 years), 201 of the 409 patients maintaining over 80% adherences to both PEG-IFN and RBV were enrolled in the second study (*Study design 2*).

Nucleotide sequencing of the core and NS5A gene

The nucleotide sequences encoding aa 1–191 (HCV core) and aa 2209–2248 (ISDR) were analyzed by direct sequencing as described by Akuta et al. [22, 27] and Enomoto et al. [21]. In brief, RNA was extracted from the sera and converted to cDNA and two nested rounds of polymerase chain reaction (PCR) were performed. Primers used in the PCR were as follows; (a) Nucleotide sequences of the core region: the first-round PCR was performed with CC11 (sense) and e14 (antisense) primers [22, 27], and the second-round PCR with CC9 (sense) and e14 (antisense) primers [22, 27]. (b) Nucleotide sequences of the ISDR in NS5A: the first-round PCR was performed with ISDR1 (sense) and ISDR2 (antisense) primers [21], and the second-round PCR with ISDR3 (sense) and ISDR4 (antisense) primers [21]. These sequences were compared with the consensus sequence of genotype 1b (HCV-J) [29]. Wild types virus encoded arginine and leucine at aa 70 and 91, respectively, and the aa substitutions were glutamine or histidine at aa 70 and methionine at aa 91.

Viral kinetic study

Serum HCV RNA levels were measured by PCR (Amplicor HCV RNA kit, version 2.0, Roche Diagnostics) using samples taken before treatment and at 4, 12, 24, and 48 weeks after the therapy. SVR was defined as HCV RNA negativity by qualitative analysis by PCR at 24 weeks after the treatment. RVR was defined as HCV RNA negativity at 4 weeks, cEVR as HCV RNA negativity at 12 weeks, LVR as HCV RNA negativity during 13–24 weeks and an end of treatment response (ETR) as HCV RNA negativity at the end of treatment. Patients who remained positive for HCV RNA at the end of the treatment and at 24 weeks after the therapy were defined as non-responders (NR).

Adherences to PEG-IFN and RBV

Adherences to PEG-IFN and RBV were assessed by separately calculating the actual doses of PEG-IFN and RBV received as percentages of the intended dosages. Adherences to PEG-IFN and RBV were divided into two groups; $80\% \leq$ and $<80\%$.

Statistical analysis

All data analyses were conducted using the SAS version 9.1.3 statistical analysis packages (SAS Institute, Cary, NC, USA). Individual characteristics between groups were evaluated by Mann–Whitney *U* test for numerical variables or Fisher's exact test for categorical variables. Variables exhibiting values of $p < 0.1$ in the univariate analysis were subjected to stepwise multivariate logistic regression analysis. The grade of steatosis and iron deposition in liver tissue, BMI, albumin (Alb), low density lipoprotein-cholesterol (LDL-C), homeostasis model assessment-insulin resistance (HOMA-IR), ferritin, and hyaluronic acid were excluded from multivariate logistic regression analysis because of the absence of those data in more than 10% of the patients. All p values of $p < 0.05$ by the two-tailed test were considered statistically significant.

Results

Study design 1

Baseline backgrounds, characteristics and adherences of peginterferon and ribavirin in males and females

The treatment outcome of PEG-IFN and RBV combination therapy depends on gender in Japanese patients, so in addition to aa substitutions in the ISDR in NS5A [21] or at HCV core 70 and 91 [22, 27], we compared the baseline characteristics according to gender (Table 1). Males were younger and the grade of hepatic inflammation was milder in males. The serum levels of LDL-C, PLT count, and aa substitutions of ISDR and at core 70 and 91 did not differ significantly different between males and females. The frequency of no alcohol abuse was significantly ($p < 0.0001$) higher in females than males (Some of them are not described in Table 1).

The rates of over 80% adherences to PEG-IFN and RBV were significantly lower ($p = 0.0066$, $p < 0.00001$, respectively) in females than males. Only in those above 60 years did the rate of over 80% adherence to PEG-IFN not differ significantly between males and females, but the rate of over 80% adherence to RBV was significantly lower ($p = 0.035$) in females than males (Table 1).

Table 1 Backgrounds and characteristics of male and female patients

Factors	Gender		p value
	Male	Female	
No. of patients	256 (62.6%)	153 (37.4%)	
Age			
Median (range)	53 (18–73)	59 (23–75)	0.00001
F stage			
F0–2	206 (80.5%)	119 (77.8%)	0.592
F3–4	50 (19.5%)	34 (22.2%)	
Grade (A factor)			
A0–1	163 (63.7%)	79 (51.6%)	0.026
A2–3	93 (36.3%)	74 (48.4%)	
HCV RNA load 0 week (KIU/mL)			
Median (range)	1500 (100–5000 <)	1280 (100–5000 <)	0.384
ALT 0 week (IU/L)			
Median (range)	74.5 (16–504)	59 (19–391)	0.001
BMI			
Median (range)	23.6 (17.5–31.2)	22.1 (16.1–33.9)	0.00033
Alb (g/dL)			
Median (range)	4.0 (3.0–5.2)	3.8 (3.0–4.8)	0.011
LDL-C (mg/dL)			
Median (range)	97 (30–185)	90 (34–174)	0.612
T-Chol (mg/dL)			
Median (range)	167 (85–273)	176 (114–261)	0.0016
PLT count ($\times 10^4/\text{mm}^3$)			
Median (range)	17.0 (8.0–31.9)	16.4 (8.1–39.9)	0.350
Amino acid mutation of ISDR			
0–1	200 (78.1%)	121 (79.1%)	0.608
2 \leq	56 (21.9%)	32 (20.9%)	
Amino acid substitution of core 70			
Wild	177 (69.1%)	114 (74.5%)	0.261
Mutant	79 (30.9%)	39 (25.5%)	
Amino acid substitution of core 91			
Wild	153 (59.8%)	98 (64.1%)	0.403
Mutant	103 (40.2%)	55 (35.9%)	
PEG-IFN adherence			
<80%	41 (17.7%)	42 (30.4%)	0.0066
80% \leq	190 (82.3%)	96 (69.6%)	
Ribavirin adherence			
<80%	54 (23.6%)	73 (52.1%)	<0.00001
80% \leq	175 (76.4%)	67 (47.9%)	
Age: <60 years			
PEG adherence			
<80%	30 (17.8%)	23 (31.5%)	0.027
80% \leq	139 (82.2%)	50 (68.5%)	
Ribavirin adherence			
<80%	27 (16.2%)	31 (42.5%)	0.000029
80% \leq	140 (83.8%)	42 (57.5%)	
Age: 60 years \leq			
PEG adherence			
<80%	11 (17.7%)	19 (29.2%)	0.147
80% \leq	51 (82.3%)	46 (70.8%)	
Ribavirin adherence			
<80%	27 (43.5%)	42 (62.7%)	0.035
80% \leq	35 (56.5%)	25 (37.3%)	




Fig. 1 Relationship between time course of serum HCV RNA negativity and amino acid substitutions in the ISDR and core amino acids 70 and 91. For cases with no or only one amino acid (aa) change in the ISDR, the rates of cEVR, LVR, ETR and SVR were significantly higher in patients with wild type core aa 70 but only the rates of RVR, cEVR, and LVR were significantly higher in patients with wild type core aa 91

Amino acid substitutions

There were no significant differences in the frequency of aa substitutions in the ISDR between males and females. Core aa substitutions at positions 70 and 91 were as follows; 291 (71.1%) were wild type and 118 (28.9%) were mutant at core aa 70, and 251 (61.4%) were wild type and 158 (38.6%) were mutant at core aa 91. There were no significant differences between males and females and between patients below and above 60 years of age.

Virological responses and aa substitutions

The rate of RVR did not differ significantly between males and females. However, more male patients showed HCV RNA negativity at 12 weeks (males vs. females; 60.7 vs. 48.4%, $p = 0.018$), 24 weeks (76.8 vs. 64.2%, $p = 0.0078$) and 48 weeks (78.2 vs. 68.6%, $p = 0.049$), and the proportion of male patients in SVR was significantly higher than that of females (61.3 vs. 37.3%, $p < 0.00001$).

RVR, cEVR and SVR rates were significantly higher in patients with two or more aa mutations in the ISDR compared to patients having no or one aa substitution in that region (20 vs. 11%, $p = 0.044$; 71 vs. 52%, $p = 0.0021$; 66 vs. 49%, $p = 0.0054$, respectively). AA substitution at core position 70 resulted in significantly lower rate of cEVR, LVR, ETR, and SVR (40 vs. 63%, $p = 0.000037$; 51 vs. 81%, $p < 0.00001$; 56 vs. 83%, 41 vs. 57%; $p < 0.00001$, $p = 0.0031$, respectively). Although the patients with the wild type aa at core 91 showed significantly higher rates of RVR and cEVR, the rate of SVR was not significantly higher in those patients ($p = 0.054$).

SVR rates were 30% for patients with no or one aa substitution in the ISDR and the core aa 70 substitution, and were significantly lower compared to those with the wild type aa core 70 (Fig. 1). These findings were not confirmed in patients with no or one aa substitution in the ISDR and the core aa 91 substitution (Fig. 1).

Factors affecting SVR by univariate analysis

Univariate analysis identified nine parameters that influenced non-SVR significantly: female gender, older age, advanced staged liver fibrosis, high viral load, low serum Alb level, low PLT count, no or one aa substitution in the ISDR, aa substitution at core aa 70, and low adherence to RBV (Table 2). The frequency of steatosis and HOMA-IR were significantly ($p = 0.0057$, $p < 0.00001$, respectively) lower in patients with SVR compared with non-SVR (data not shown). However, these factors were not entered in the multivariate analysis because of the absence of the data in many cases.

Factors affecting RVR, cEVR, and SVR by multivariate logistic regression analysis

Multivariate analysis identified four parameters that influenced RVR independently: low HCV RNA load, low serum ALT level, two or more aa mutations in the ISDR and the wild type aa at core position 91 (Table 3).

Concerning cEVR, male gender, mild fibrosis stage, low HCV RNA load, two or more aa mutations in the ISDR, and the wild type aa at core positions 70 and 91 were independent predictors (Table 3).

Concerning SVR, male gender ($p < 0.0001$), low HCV RNA load ($p = 0.013$), high PLT count ($p = 0.0019$), two or more aa mutations in the ISDR ($p = 0.024$), and wild type core aa 70 ($p = 0.0045$) were found to be independent predictors (Table 3).

The predictive values of the combination of gender, PLT count, ISDR and core aa 70 are shown in Fig. 2a. In male patients having PLT of $<15 \times 10^4/\text{mm}^3$, and, no or one aa substitution in the ISDR, the SVR rate was 68% when core 70

Table 2 Univariate analysis to identify the factors of SVR

Factors	Negative of HCV RNA after 24 weeks		<i>p</i> value
	(-)	(+)	
No. of patients	214 (52.3%)	195	
Gender			
Male	157 (61.3%)	99	<0.00001
Female	57 (37.3%)	96	
Age			
Median (range)	52.5 (18–75)	58 (20–74)	<0.00001
<60 years	155 (58.1%)	112	0.0018
60 years ≤	59 (41.5%)	83	
Age: <60 years			
Male	118 (63.4%)	68	0.010
Female	37 (45.7%)	44	
Age: 60 years ≤			
Male	39 (55.7%)	31	0.0011
Female	20 (27.8%)	52	
F stage			
F0–2	190 (58.5%)	135	0.000013
F3–4	25 (29.8%)	59	
Grade (A factor)			
A0–1	138 (56.8%)	104	0.130
A2–3	81 (48.5%)	86	
HCV RNA load 0 week (KIU/mL)			
Median (range)	1300 (100–5000<)	1700 (130–5000<)	0.016
ALT 0 week (IU/L)			
Median (range)	66 (16–391)	67 (19–504)	0.892
BMI			
Median (range)	23.0 (17.3–32.4)	23.25 (16.1–33.9)	0.714
Alb (g/dL)			
Median (range)	4.0 (3.2–5.2)	3.8 (3.0–4.8)	0.0088
LDL-C (mg/dL)			
Median (range)	94.5 (31–185)	97.5 (30–182)	0.611
T-Chol (mg/dL)			
Median (range)	169.5 (85–257)	170 (103–273)	0.511
PLT count ($\times 10^4/\text{mm}^3$)			
Median (range)	18.2 (8.7–39.9)	15.1 (8.0–31.9)	<0.00001
<15	54 (36.5%)	94	<0.00001
15 ≤	160 (61.3%)	101	
Amino acid mutation of ISDR			
0–1	156 (48.6%)	165	0.0054
2 ≤	58 (65.9%)	30	
Amino acid substitution of core 70			
Wild	166 (57.0%)	125	0.0031
Mutant	48 (40.7%)	70	
Amino acid substitution of core 91			
Wild	141 (56.2%)	110	0.054
Mutant	73 (46.2%)	85	
PEG-IFN adherence			
<80%	35 (42.2%)	48	0.063
80% ≤	154 (53.8%)	132	
Ribavirin adherence			
<80%	55 (43.3%)	72	0.048
80% ≤	132 (54.5%)	110	