

Fig. 7. Chronic hepatitis B patients with hepatocytes positive for pSmad3L and negative for pSmad3C increased risk of HCC development. (A) HCC occurred subsequently among patients whose hepatocytes in chronic hepatitis B specimens were strongly positive for pSmad3L. Incidence of HCC was significantly higher in patients with abundant Smad3L phosphorylation (scores 3 to 4, solid line) in hepatocytic nuclei versus those with sparse Smad3L phosphorylation (scores 0 to 2, dotted line). (B) HCC did not occur subsequently among patients whose hepatocytes in chronic hepatitis B specimens were strongly positive for pSmad3C. HCC occurred only in patients with sparse Smad3C phosphorylation (scores 0 to 2, solid line) in hepatocytic nuclei, while no patients with abundant Smad3C phosphorylation (scores 3 to 4, dotted line) have developed HCC. Cumulative rates of HCC occurrence from chronic hepatitis B were compared between cases with high and low phosphorylation of Smad3L and Smad3C (Kaplan-Meier analysis and log-rank test). (C) HBx protein shifted hepatic TGF- β signaling from the tumor-suppressive pSmad3C pathway to the oncogenic JNK-dependent pSmad3L pathway in early stages of chronic hepatitis B. Normal hepatocytes exhibited TGF- β -dependent Smad3 phosphorylation at the C-terminal region, which is related to growth inhibition by up-regulation of p21^{WAF1}. HBx protein activates JNK, promoting the oncogenic pSmad3L signaling, which fosters cell growth by up-regulating c-Myc, in a mean time reducing tumor-suppressive pSmad3C-mediated signaling.

The general biomedical approach to HCC is shifting away from population risk assessment and empirical treatment of patients to predictive personalized medicine based on molecular classification and targeted therapy.²⁹ Better knowledge of the risk factors associated with the occurrence of HCC can improve the effectiveness of surveillance programs. Our approach has identified pSmad3L and pSmad3C as prognostic markers that may prove to be clinically useful. Such predictive markers could allow us to select patients with chronic hepatitis B who have a high or low risk of developing HCC. Although the latter group could be followed up on an annual basis, the patients with a high risk require targeted surveillance measures to allow early diagnosis of HCC.

Phosphorylation of many transcription factors is controlled by the dynamic interplay between kinases and phosphatases. In this regard, we studied the kinetics of both linker and C-terminal phosphorylation of Smad3 in parental and HBx-expressing hepatocytes in response to TGF- β (unpublished observation). In parental hepatocytes, the levels of linker and C-terminal phosphorylation peaked at 30 minutes after the start of exposure to TGF- β and then gradually declined. However, HBx-expressing hepatocytes showed constitutive phosphorylation at Smad3L during continuous exposure to TGF- β . Several lines of evidence have identified small C-terminal domain phosphatase (SCP1-3) and protein phosphatase magnesium 1A (PPM1A) as the linker and C-terminal phosphatases, respectively.^{37,38} Accordingly, SCP1-3 and PPM1A may reverse domain-specific phosphorylation in normal hepatocytes. In contrast, HBx-expressing hepatocytes may not show induction or activation of SCP1-3. Alternatively, linker phosphorylation in HBx-expressing hepatocytes might be resistant to SCP1-3.

Many researchers have been seeking key transcription factors regulating tumor-suppressive pathways that are altered in cancer. Our current model of JNK/pSmad3L signaling during HBV-related chronic liver disease suggests that specific inhibitors of the JNK/pSmad3L pathway might inhibit the progression of HCC. With respect to molecular targeting therapy for human HCC, pSmad3L and pSmad3C should be assessed as biomarkers to evaluate the benefit from specific inhibition of the JNK/pSmad3L pathway.

Acknowledgment: We thank Dr. Rik Derynck (University of California at San Francisco) and Dr. Seishi Murakami (Kanazawa University) for providing us with complementary DNAs encoding human Smad3 and HBx. We also thank Chiaki Kitano for assistance to construct ecotropic retrovirus and Natsuko Ohira for assistance with immunoblotting.

Table 5. Variables with Independent Predictive Value for HCC in Univariate and Multivariate Analyses

Characteristics	n	No. of Patients with HCC (%)	Univariate Analysis		Multivariate Analysis	
			Hazard Ratio (95% CI)	P Value	Hazard Ratio (95% CI)	P Value
pSmad3L positivity*						
Low (1 and 2)	32	1 (3)	1.00		1.00	
High (3 and 4)	28	6 (21)	3.8 (1.4-10.6)	0.01	14.8 (1.8-118.5)	0.01
pSmad3C positivity*						
High (3 and 4)	30	0 (0)	1.00		1.00	
Low (1 and 2)	30	7 (23)	2.8 (0.001-7.0)	0.03	16.4 (1.0-125.0)	0.04
Fibrotic stage†						
Low (F1 and F2)	39	4 (10)	1.00		1.00	
High (F3)	21	3 (14)	1.9 (0.7-5.4)	0.24	3.9 (0.4-38.6)	0.24
Inflammatory activity†						
Low (A0 and A1)	23	1 (4)	1.00		1.00	
High (A2 and A3)	37	6 (16)	1.8 (0.7-4.8)	0.27	0.2 (0.02-1.1)	0.06
HBV DNA (copies/mL)						
<10 ⁵	42	3 (7)	1.00		1.00	
>10 ⁵	18	4 (22)	1.9 (1.0-3.5)	0.05	2.5 (0.9-6.9)	0.08
HBeAg						
Negative	42	4 (10)	1.00		1.00	
Positive	18	3 (17)	2.1 (0.5-9.5)	0.32	9.9 (1.1-89.3)	0.03

Abbreviations: CI, confidence interval; HBeAg, hepatitis B e antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; pSmad3C, C-terminally phosphorylated Smad3; pSmad3L, linker-phosphorylated Smad3.

*Hepatocytic Smad3 phosphorylation in chronic hepatitis B specimens is scored as follows: 0, no phosphorylation; 1, <25% Smad3 phosphorylation; 2, 25% to 50% Smad3 phosphorylation; 3, 50% to 75% Smad3 phosphorylation; 4, >75% Smad3 phosphorylation.

†Necroinflammatory activity and fibrotic stage are determined histologically according to Desmet's classification.

References

- Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 2003; 362:1907-1917.
- El-Serag HB. Hepatocellular carcinoma: recent trends in the United States. *Gastroenterology* 2004;127(Suppl):275-345.
- Brechot C, Pourcel C, Louise A, Rain B, Tiollais P. Presence of integrated hepatitis B virus DNA sequences in cellular DNA of human hepatocellular carcinoma. *Nature* 1980;286:533-535.
- Kim CM, Koike K, Saito I, Miyamura T, Jay G. HBx gene of hepatitis B virus induces liver cancer in transgenic mice. *Nature* 1991;351:317-320.
- Yu DY, Moon HB, Son JK, Jeong S, Yu SL, Yoon H, et al. Incidence of hepatocellular carcinoma in transgenic mice expressing the hepatitis B virus X-protein. *J Hepatol* 1999;31:123-132.
- Koike K, Moriya K, Iino S, Yotsuyanagi H, Endo Y, Miyamura T, et al. High-level expression of hepatitis B virus HBx gene and hepatocarcinogenesis in transgenic mice. *HEPATOLOGY* 1994;19:810-819.
- Benn J, Su F, Doria M, Schneider RJ. Hepatitis B virus HBx protein induces transcription factor AP-1 by activation of extracellular signal-related and c-Jun N-terminal mitogen-activated protein kinases. *J Virol* 1996;70:4978-4985.
- Roberts AB, Sporn MB. The transforming growth factor- β s. In: Sporn MB, Roberts AB, eds. *Peptide Growth Factors and Their Receptors*. Berlin: Springer-Verlag, 1990:419-472.
- Heldin CH, Miyazono K, ten Dijke P. TGF- β signaling from cell membrane to nucleus through SMAD proteins. *Nature* 1997;390:465-471.
- Massagué J. TGF- β signal transduction. *Annu Rev Biochem* 1998;67:753-791.
- Kretschmar M, Doody J, Timokhina I, Massagué J. A mechanism of repression of TGF- β /Smad signaling by oncogenic Ras. *Genes Dev* 1999; 13:804-816.
- Mori S, Matsuzaki K, Yoshida K, Furukawa F, Tahashi Y, Yamagata H, et al. TGF- β and HGF transmit the signals through JNK-dependent Smad2/3 phosphorylation at the linker regions. *Oncogene* 2004;23:7416-7429.
- Furukawa F, Matsuzaki K, Mori S, Tahashi Y, Yoshida K, Sugano Y, et al. p38 MAPK mediates fibrogenic signal through Smad3 phosphorylation in rat myofibroblasts. *HEPATOLOGY* 2003;38:879-889.
- Matsuura I, Denisova NG, Wang G, He D, Long J, Liu F. Cyclin-dependent kinases regulate the antiproliferative function of Smads. *Nature* 2004;430:226-231.
- Yamagata H, Matsuzaki K, Mori S, Yoshida K, Tahashi Y, Furukawa F, et al. Acceleration of Smad2 and Smad3 phosphorylation via c-Jun NH(2)-terminal kinase during human colorectal carcinogenesis. *Cancer Res* 2005; 65:157-165.
- Sekimoto G, Matsuzaki K, Yoshida K, Mori S, Murata M, Seki T, et al. Reversible Smad-dependent signaling between tumor suppression and oncogenesis. *Cancer Res* 2007;67:5090-5096.
- Arany PR, Rane SG, Roberts AB. Smad3 deficiency inhibits v-ras-induced transformation by suppression of JNK MAPK signaling and increased farnesyl transferase inhibition. *Oncogene* 2008;27:2507-2512.
- Yoshida K, Matsuzaki K, Mori S, Tahashi Y, Yamagata H, Furulawa F, et al. Transforming growth factor- β and platelet-derived growth factor signal via c-Jun N-terminal kinase-dependent Smad2/3 phosphorylation in rat hepatic stellate cells after acute liver injury. *Am J Pathol* 2005;166:1029-1039.
- Block TM, Mehta AS, Fimmel CJ, Jordan R. Molecular viral oncology of hepatocellular carcinoma. *Oncogene* 2003;22:5093-5107.
- Matsuzaki K, Murata M, Yoshida K, Sekimoto G, Uemura Y, Sakaida N, et al. Chronic inflammation associated with hepatitis C viral infection perturbs hepatic TGF- β signaling, promoting cirrhosis and hepatocellular carcinoma. *HEPATOLOGY* 2007;46:48-57.
- Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Scheuer PJ. Classification of chronic hepatitis: diagnosis, grading and staging. *HEPATOLOGY* 1994;19:1513-1520.
- Cox DR. Regression models and life-tables. *J R Stat Soc (B)* 1972;34:187-220.
- Pardali K, Moustakas A. Actions of TGF- β as tumor suppressor and prometastatic factor in human cancer. *Biochim Biophys Acta* 2007;1775:21-62.

24. Chen CJ, Yang HI, Su J, Jen CL, You SL, Lu SN, et al. REVEAL-HBV study group. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006;295:65-73.
25. Feitelson MA. c-Myc overexpression in hepatocarcinogenesis. *Human Pathology* 2004;35:1299-1302.
26. Yang HI, Lu SN, Liaw YF, You SL, Sun CA, Wang LY, et al. Taiwan community-based cancer screening project group. Hepatitis B e antigen and the risk of hepatocellular carcinoma. *N Engl J Med* 2002;347:168-174.
27. Thorgeirsson SS, Lee JS, Grisham JW. Functional genomics of hepatocellular carcinoma. *HEPATOLOGY* 2006;43:145-150.
28. Theise ND, Park YN, Kojiro M. Dysplastic nodules and hepatocarcinogenesis. *Clin Liver Dis* 2002;6:497-512.
29. Thorgeirsson SS, Grisham JW. Molecular pathogenesis of human hepatocellular carcinoma. *Nat Genet* 2002;31:339-346.
30. Chisari FV, Klopchin K, Moriyama T, Pasquinelli C, Dunsford HA, Sell S, et al. Molecular pathogenesis of hepatocellular carcinoma in hepatitis B virus transgenic mice. *Cell* 1989;59:1145-1156.
31. Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, et al. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065-1067.
32. Di Bisceglie AM. Hepatitis C and hepatocellular carcinoma. *HEPATOLOGY* 1997;26 (Suppl):34S-38S.
33. Popper H, Shafritz DA, Hoofnagle JH. Relation of the hepatitis B virus carrier state to hepatocellular carcinoma. *HEPATOLOGY* 1987;7:764-772.
34. Inagaki Y, Okazaki I. Emerging insights into transforming growth factor β Smad signal in hepatic fibrogenesis. *Gut* 2007;56:284-292.
35. Kaimori A, Potter J, Kaimori JY, Wang C, Mezey E, Koteish A. Transforming growth factor- β 1 induces an epithelial-to-mesenchymal transition state in mouse hepatocytes in vitro. *J Biol Chem* 2007;282:22089-22101.
36. Weng HL, Ciuclan L, Liu Y, Hamzavi J, Godoy P, Gaitantzi H, et al. Profibrogenic transforming growth factor- β /activin receptor-like kinase 5 signaling via connective tissue growth factor expression in hepatocytes. *HEPATOLOGY* 2007;46:1257-1270.
37. Lin X, Duan X, Liang YY, Su Y, Wrighton KH, Long J, et al. PPM1A functions as a Smad phosphatase to terminate TGF β signaling. *Cell* 2006;125:915-928.
38. Wrighton KH, Willis D, Long J, Liu F, Lin X, Feng XH. Small C-terminal domain phosphatases dephosphorylate the regulatory linker regions of Smad2 and Smad3 to enhance transforming growth factor- β signaling. *J Biol Chem* 2006;281:38365-38375.

Steatosis, liver injury, and hepatocarcinogenesis in hepatitis C viral infection

KAZUHIKO KOIKE

Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

In addition to the link with development of hepatocellular carcinoma (HCC), hepatitis C virus (HCV) infection is associated with several hepatic and extrahepatic manifestations. A role of hepatic steatosis in the pathogenesis of chronic hepatitis C has been shown, implying hepatitis C as a metabolic disease. Furthermore, recent epidemiological studies have suggested a linkage between insulin resistance and chronic HCV infection. In addition to the data indicating the presence of lipid metabolism disturbance and insulin resistance in the cohort of chronic hepatitis C patients, we found evidence showing the association between these two conditions and HCV infection using mice transgenic for the HCV core gene. These mice develop HCC late in life after the phase of hepatic steatosis and insulin resistance. The nonappearance of both steatosis and HCC in HCV core gene transgenic mice that are null for the proteasome activator 28 γ implies a close relationship between lipid metabolism disturbance and hepatocarcinogenesis. Also, the core protein is shown to bind with retinoid X receptor (RXR)- α , resulting in the upregulation of some lipid metabolism enzymes, including cellular retinol binding protein II and acyl-CoA oxidase. In addition, the persistent activation of peroxisome proliferator activated receptor (PPAR)- α has recently been found in the liver of HCV core gene transgenic mice, yielding dramatic changes in lipid metabolism and hepatocyte proliferation, including HCC development. These results would provide a clue for further understanding of the role of lipid metabolism in pathogenesis of HCV infection, including liver injury and hepatocarcinogenesis.

Key words: lipid metabolism, transgenic mouse, oxidative stress, intracellular signal transduction, peroxisome proliferator activated receptor

Introduction

Worldwide, approximately 170 million people are persistently infected with hepatitis C virus (HCV), which induces a spectrum of chronic liver diseases from chronic hepatitis to cirrhosis and, eventually, to hepatocellular carcinoma (HCC).¹ HCV has been given increasing attention because of its wide and deep penetration in the community, tied with a very high incidence of HCC in persistent HCV infection. Once liver cirrhosis is established in hosts persistently infected with HCV, HCC develops at a yearly rate of approximately 7%,² resulting in the development of HCC in nearly 90% of HCV-associated cirrhotic patients in 15 years. In addition, the outstanding features in the mode of hepatocarcinogenesis in HCV infection, i.e., development of HCC in a multicentric fashion and at a very high incidence, are not common in other malignancies except for hereditary cancers such as familial polyposis of the colon. Knowledge of the mechanism underlying HCC development in persistent HCV infection, therefore, is imminently required for the prevention of HCC.

In addition to the link with development of HCC, HCV infection is associated with several hepatic and extrahepatic manifestations.³ A role of hepatic steatosis in the pathogenesis of chronic hepatitis C has been shown, implicating hepatitis C as a metabolic disease.⁴ Moreover, recent epidemiological studies have suggested a linkage between insulin resistance and chronic HCV infection.⁵ In addition to the epidemiological data indicating the presence of lipid metabolism disturbance and insulin resistance in the cohort of chronic hepatitis C patients, detailed analyses on the relationship between

Received: June 9, 2008 / Accepted: August 10, 2008
Reprint requests to: K. Koike

metabolic disorders and chronic hepatitis C have revealed evidence showing a close association between the progression of liver fibrosis and metabolic abnormalities in HCV infection.⁶ However, it is unclear yet whether a causative relationship exists between these medical conditions. Moreover, it is unclear whether such metabolic disorders contribute to hepatocarcinogenesis in HCV infection.

Possible roles of HCV in hepatocarcinogenesis

The mechanism underlying hepatocarcinogenesis in HCV infection is not yet fully understood, despite the fact that nearly 80% of patients with HCC in Japan are persistently infected with HCV.^{1,7,8} HCV infection is also common in patients with HCC in other countries, albeit to a lesser extent. These lines of evidence prompted us to seek to determine the role of HCV in hepatocarcinogenesis. Inflammation induced by HCV should be considered, of course, in a study on the hepatocarcinogenesis in hepatitis viral infection: necrosis of hepatocytes caused by chronic inflammation followed by regeneration enhances genetic aberrations in host cells, the accumulation of which culminates in HCC. This theory presupposes an indirect involvement of hepatitis viruses in HCC via hepatic inflammation. However, this context leaves us with a serious question: can inflammation alone result in the development of HCC in such a high incidence (90% in 15 years) or multicentric nature in HCV infection?

The other role of HCV would have to be weighed against an extremely rare occurrence of HCC in patients with autoimmune hepatitis in which severe inflammation in the liver persists indefinitely, even after the development of cirrhosis. This background and reasoning lead to a possible activity of viral proteins for inducing neoplasia. This possibility has been evaluated by introducing genes of HCV into hepatocytes in culture with little success. One of the difficulties in using cultured cells is the carcinogenic capacity of HCV, if any, which would be weak and would take a long time to manifest. Actually, it takes 30–40 years for HCC to develop in individuals infected with HCV. On the basis of these points of view, we started to investigate carcinogenesis in chronic hepatitis C, *in vivo*, by transgenic mouse technology.

HCV core protein has an *in vivo* oncogenic activity as revealed by animal studies

Transgenic mouse lines carrying the HCV genome were engineered by introducing the genes from the cDNA of




Fig. 1. Transgenic mouse lines carrying the hepatitis C virus (HCV) genome. Three different kinds of transgenic mouse lines, carrying the *core* gene, envelope genes, or nonstructural genes of HCV, respectively, were established under the control of the same regulatory elements. Among these mouse strains, only the transgenic mice carrying the HCV core gene develop hepatocellular carcinoma (HCC) after an early phase with hepatic steatosis in two independent lineages. The mice transgenic for the envelope genes or nonstructural genes do not develop HCC. *HCC*, hepatocellular carcinoma; *env*, envelope genes; *NS*, nonstructural genes

the HCV genome of genotype 1b.^{9,10} Established are three different kinds of transgenic mouse lines, which carry the core gene, envelope genes, or nonstructural genes, respectively, under the same transcriptional regulatory element. Among these mouse lines, only the transgenic mice carrying the core gene developed HCC in two independent lineages.¹⁰ The envelope gene transgenic mice do not develop HCC, despite high expression levels of both E1 and E2 proteins,^{11,12} and the transgenic mice carrying the entire nonstructural genes have developed no HCC (Fig. 1).

The core gene transgenic mice express the core protein of an expected size, and the level of the core protein in the liver is similar to that in chronic hepatitis C patients. Early in life, these mice develop hepatic steatosis, which is one of the histological characteristics of chronic hepatitis C, along with lymphoid follicle formation and bile duct damage.¹³ Thus, the core gene transgenic mouse model reproduces well the features of chronic hepatitis C. Of note, no pictures of significant inflammation are observed in the liver of this animal model. Late in life, these transgenic mice develop HCC. Notably, the development of steatosis and HCC has been reproduced by other HCV transgenic mouse lines, which harbor the entire HCV genome or structural genes including the core gene.^{14–16} These outcomes indicate that the core protein, *per se*, of HCV has an oncogenic potential when expressed *in vivo*.

Oxidative stress overproduction and intracellular signaling pathway activation are the major pathways in the core-induced liver pathology

It is difficult to elucidate the mechanism underlying the development of HCC, even for our simple model in which only the core protein is expressed in otherwise normal liver. There is a notable feature in the localization of the core protein in hepatocytes; while the core protein predominantly exists in the cytoplasm associated with lipid droplets, it is also present in the mitochondria and nuclei.^{10,17} On the basis of this finding, the pathways related to these two organelles, the mitochondria and nuclei, were thoroughly investigated.

One effect of the core protein is an increased production of oxidative stress in the liver. We would like to draw particular attention to the fact that the production of oxidative stress is increased in our transgenic mouse model in the absence of inflammation in the liver. This finding reflects a state of overproduction of reactive oxygen species (ROS) in the liver,¹⁸ or predisposition to it, which is staged by the HCV core protein without any intervening inflammation.^{19,20} The overproduction of oxidative stress results in the generation of deletions in mitochondrial and nuclear DNA, an indicator of genetic damage. In addition, analysis of antioxidant system revealed that some antioxidative molecules are not increased despite the overproduction of ROS in the liver of core gene transgenic mice: hemoxygenase-1 and glutathione peroxidase are not augmented whereas catalase and glutathione S-transferase levels are increased and enhanced by iron overloading (Moriya et al., manuscript in preparation). These results suggest that HCV core protein not only induces overproduction of ROS but also attenuates some of the antioxidant systems, which may explain the mechanism underlying the production of a strong oxidative stress in HCV infection compared to other forms of hepatitis.

In the absence of inflammation, thus, the core protein induces oxidative stress overproduction, which may, at least in part, contribute to hepatocarcinogenesis in HCV infection. If inflammation were added to the liver with the HCV core protein, the production of oxidative stress would be escalated to an extent that can no longer be scavenged by a physiological antagonistic system. This idea suggests that the inflammation in chronic HCV infection would have a characteristic difference in its quality from those of other types of hepatitis, such as autoimmune hepatitis. The basis for the overproduction of oxidative stress may be ascribed to the mitochondrial dysfunction.^{10,19} The dysfunction of the electron transfer system of the mitochondrion is suggested in association with the presence of the HCV core protein.²¹

Other pathways in hepatocarcinogenesis would be the alteration of the expression of cellular genes and modulation of intracellular signaling pathways. For example, tumor necrosis factor (TNF)- α and interleukin-1 β have been found to be transcriptionally activated.²² The mitogen-activated protein kinase (MAPK) cascade is also activated in the liver of the core gene transgenic mouse model. The MAPK pathway, which consists of three routes, c-Jun N-terminal kinase (JNK), p38, and extracellular signal-regulated kinase (ERK), is involved in numerous cellular events including cell proliferation. In the liver of the core gene transgenic mouse model before HCC development, only the JNK route is activated. Downstream of JNK activation, transcription factor activating protein (AP)-1 activation is markedly enhanced.^{20,21} At far downstream, both the mRNA and protein levels of cyclin D1 and CDK4 are increased. Thus, the HCV core protein modulates the intracellular signaling pathways and gives an advantage for cell proliferation to the hepatocytes. Interestingly, we found recently that a protein interacting with the core protein, proteasome activator 28 γ (PA28 γ), is indispensable for the core protein to exert its function for the development of steatosis, insulin resistance, and HCC.^{23,24}

Lipid metabolism and HCV infection

Steatosis is frequently observed in chronic hepatitis C patients and is significantly associated with increased fibrosis and progression rate of fibrosis of the liver.⁶ A comprehensive analysis of gene expression in the liver of core gene transgenic mice, in which steatosis develops from early in life, revealed that a number of genes related to lipid metabolism are significantly upregulated or downregulated (Table 1).

The composition of fatty acids that are accumulated in the liver of core gene transgenic mice is different from that in fatty liver resulting from simple obesity. Carbon-18 monounsaturated fatty acids (C18:1) such as oleic or vaccenic acids are significantly increased; this is also the case in the comparison of liver tissues from hepatitis C patients and patients with simple fatty liver due to obesity.²⁰ The mechanism of steatogenesis in hepatitis C was investigated using this mouse model. There are at least three pathways for the development of steatosis. One is the frequent presence of insulin resistance in hepatitis C patients as well as in the core gene transgenic mice, which occurs through the inhibition of tyrosine phosphorylation of insulin receptor substrate (IRS)-1.²⁵ Insulin resistance increases the peripheral release and hepatic uptake of fatty acids, resulting in an accumulation of lipid in the liver. The second pathway is the suppression of the activity of

Table 1. Cellular genes differentially expressed in hepatitis C virus (HCV) core transgenic mouse liver

	Upregulated	Downregulated
Lipid metabolism	NPC1 Catalase Very long chain acyl-CoA dehydrogenase Carboxylesterase selenoprotein P Carbonic anhydrase Adipose differentiation-related protein Bilirubin/phenol family UDP glucuronosyltransferase	Stearoyl-CoA desaturase Sterol-carrier protein X Alpha-enolase carnitine acetyltransferase Gal beta 1,4(3) GlcNAc alpha 2,3-sialyltransferase Very long chain acyl-CoA synthetase Liver transferrin 4-Hydroxyphenylpyruvate dioxygenase LAF1 transketolase s-Adenosylmethionine synthetase Apolipoprotein A-II Human guanine nucleotide regulatory protein Alpha-fetoprotein Retinol binding protein
Transcription and cell proliferation	Int-6 GCN5L1 <i>H. sapiens</i> 8.2k-Da differentiation factor USF1 Initiation factor eIF-4AI Human elongation factor-1-delta Sui1	
Inflammation	Alpha-1 protease inhibitor 3 Hemopexin	Alpha-2-macroglobulin LMW prekininogen Complement component C3 AHSG(alpha 2 HS-glycoprotein) homologue Vitronectin Epithelin 1 and 2 Murinoglobulin
Others	Microvascular endothelial differentiation gene 1 Diazepam-binding inhibitor Argininosuccinate synthetase Skeletal muscle alpha-tropomyosin Ampd3 gene DNA-binding protein	

microsomal triglyceride transfer protein (MTP) by HCV core protein²⁶; this inhibits the secretion of very low density protein (VLDL) from the liver, yielding an increase of triglycerides in the liver. The last pathway involves sterol regulatory element-binding protein (SREBP)-1c, which regulates the production of triglycerides and phospholipids. In HCV core gene transgenic mice, SREBP-1c is activated, whereas neither SREBP-2 nor SREBP-1a is upregulated.²⁷

In relation to lipid metabolism, the core protein has also been found to interact with retinoid X receptor (RXR)- α .²⁸ RXR- α is one of the nuclear receptors, which forms a homodimer or heterodimers with other nuclear receptors, including PPAR (peroxisome proliferator-activated receptor)- α , and plays a pivotal role in the regulation of the expression of genes relating to lipid metabolism, cell differentiation, and proliferation. In fact, the core protein of HCV activates genes that have an RXR- α -responsive element as well as those with a PPAR- α -responsive element, both in mice and in cultured cells.²⁸ Based on these results, we, then, examined the expression and function of PPAR- α in the liver of core gene transgenic mice.

PPAR- α activation in HCV-associated hepatocarcinogenesis

PPAR- α , one of the PPAR genes, plays a central role as a heterodimer with RXR- α in regulating fatty acid transport and catabolism. It is also known as a molecular target for lipid-lowering fibrate drugs.²⁹ On the other hand, prolonged administration of PPAR- α agonists causes HCC in rodents. Currently, there is little evidence that the low-affinity fibrate ligands are associated with human cancers, but it is possible that chronic activation of high-affinity ligands could be carcinogenic in humans.²⁹

The level of PPAR- α protein was increased in the liver of core gene transgenic mice as early as 9 months of age. PPAR- α protein is accumulated with age in the nuclei of hepatocytes together with cyclin D1 protein. However, the level of PPAR- α mRNA was not increased at any age. By pulse-chase experiment, the stability of nuclear PPAR- α was increased in the presence of the core protein. In line with the increase of PPAR- α protein, target genes of PPAR- α were activated in the liver of core gene transgenic mice; these genes include

cyclin D1, cyclin-dependent kinase (CDK)-4, acy-CoA oxidase, and peroxisome thiolase.³⁰ However, in general, the activation of PPAR- α leads to improvement but not aggravation of steatosis. Then, what is the function of PPAR- α activation that is observed in the core gene transgenic mice?

To clarify the role of PPAR- α activation in pathogenesis of steatosis and HCC, we mated a core gene transgenic mouse with a PPAR- α knockout (KO) mouse and studied the phenotype. PPAR- α KO mice have reduced expression of target genes of PPAR- α , and have mild steatosis in the liver, as expected.³¹ It was unanticipated, however, that steatosis was absent in PPAR- α -null or -heterozygous core gene transgenic mice but present in PPAR- α -intact core gene transgenic mice at the age of 9 or 24 months.³⁰ 8-Hydroxy deoxyguanosine (8-OHdG) and peroxy lipids, both of which are markers for oxidative stress, were decreased in PPAR- α KO core gene transgenic mice. Mitochondrial dysfunction in the core gene transgenic mice, which contributes to overproduction of oxidative stress,¹⁹ was also improved in PPAR- α KO core gene transgenic mice.

Finally, PPAR- α KO core gene transgenic mice did not develop HCC at the age of 24 months, whereas about one-third of PPAR- α -intact core gene transgenic mice did. It should be noted that core gene transgenic mice that are heterozygous for the PPAR- α gene also did not develop HCC.³² When clofibrate, a peroxisome proliferator, was administered for 24 months to PPAR- α -heterozygous mice, either with or without the core gene, HCC developed in a higher rate in the core gene (+) mice with greater PPAR- α activation. It should be noted that steatosis was present only in core gene (+) PPAR- α -heterozygous mice. In summary, steatosis and HCC developed in PPAR- α -intact but not in PPAR- α -heterozygous or PPAR- α -null core gene transgenic mice, indicating that not the presence but the persistent activation of PPAR- α would be important in hepatocarcinogenesis by HCV core protein. In general, PPAR- α acts to ameliorate steatosis, but with the presence of mitochondrial dysfunction, which is also provoked by the core protein, the core-activated PPAR- α may exacerbate steatosis. Persistent activation of PPAR- α with "strong" ligands such as the core protein of HCV could be carcinogenic in humans, although the low-affinity fibrate ligands are not likely associated with human cancers.

HCV core protein causes "fatty acid spiral"

Figure 2 illustrates our current hypothesis for the role of lipid metabolism in HCV-associated hepatocarcinogenesis. Immune-mediated inflammation should also play a pivotal role in hepatocarcinogenesis in HCV




Fig. 2. "Fatty acid spiral" by HCV core protein. In HCV infection, the core protein induces steatosis via several pathways, leading to "fatty acid spiral" in the presence of the mitochondrial complex 1 dysfunction and PPAR- α activation, both of which are also caused by the core protein. These intracellular alterations would contribute to hepatocarcinogenesis by inducing oxidative stress overproduction and cell-growth signal activation. In such a sense, the core protein of HCV is not a classical type oncoprotein, but rather seems to contribute to hepatocarcinogenesis by modulating intracellular metabolism and signaling. HCV, hepatitis C virus; HCC, hepatocellular carcinoma; ROS, reactive oxygen species; JNK, c-Jun N-terminal kinase; ERK, extracellular signal-regulated kinase; AP-1, activating protein-1; RXR- α , retinoid X receptor- α ; PPAR- α , peroxisome proliferator activated receptor- α ; AOX, acyl-CoA oxidase; CYP, cytochrome P450; MTP, microsomal triglyceride transfer protein; FAT, fatty acid translocase; fatty acid transport protein

infection. However, in HCV infection, the core protein induces steatosis through the aforementioned pathways, leading to "fatty acid spiral" in the presence of the mitochondrial complex 1 dysfunction and PPAR- α activation, both of which are caused by the core protein. These intracellular alterations would contribute to hepatocarcinogenesis by inducing oxidative stress overproduction and cell-growth signal activation. In such a sense, the core protein of HCV is not a classical-type oncoprotein, but rather seems to contribute to hepatocarcinogenesis by modulating intracellular metabolism and signaling.

The HCV protein may allow some steps in multistep hepatocarcinogenesis to be skipped

The results of our studies on transgenic mice have indicated a carcinogenic potential of the HCV core protein in vivo; thus, HCV would be directly involved in hepatocarcinogenesis. In research studies of carcinogenesis, the theory outlined by Kinzler and Vogelstein³³ has gained wide popularity. They have proposed that the




Fig. 3. Mechanism of HCV-associated hepatocarcinogenesis. Multiple steps are required in the induction of all cancers; it would be mandatory for hepatocarcinogenesis that genetic mutations accumulate in hepatocytes. However, in HCV infection, some of these steps may be skipped in the development of HCC in the presence of the core protein. The overall effects achieved by the expression of the core protein would be the induction of HCC, even in the absence of a complete set of genetic aberrations required for carcinogenesis. By considering such a "non-Vogelstein-type" process for the induction of HCC, a plausible explanation may be given for many unusual events happening in HCV carriers

development of colorectal cancer is induced by the accumulation of a complete set of cellular gene mutations. They have deduced that mutations in the APC gene for inactivation, those in *K-ras* for activation, and those in the *p53* gene for inactivation accumulate, which cooperate toward the development of colorectal cancer.³³ Their theory has been extended to the carcinogenesis of other cancers as well, called "Vogelstein-type" carcinogenesis (Fig. 3).

On the basis of the results we obtained for the induction of HCC by the HCV core protein, we would like to introduce a different mechanism for hepatocarcinogenesis in HCV infection. We do allow multistages in the induction of all cancers; it would be mandatory for hepatocarcinogenesis that many mutations accumulate in hepatocytes. Some of these steps, however, may be skipped in the development of HCC in HCV infection to which the core protein would contribute (see Fig. 3). The overall effect achieved by the expression of the viral protein would be the induction of HCC, even in the absence of a complete set of genetic aberrations required for carcinogenesis.

By considering such a "non-Vogelstein-type" process for the induction of HCC, a plausible explanation may be given for many unusual events happening in HCV carriers.³⁴ Now it does not seem so difficult as before to determine why HCC develops in persistent HCV infection at an outstandingly high incidence. Our theory may also give an account of the nonmetastatic and multicentric de novo occurrence characteristics of HCC, which would be the result of persistent HCV infection.

References

- Saito I, Miyamura T, Ohbayashi A, Harada H, Katayama T, Kikuchi S, et al. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990;87:6547-9.
- Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Koida I, et al. Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: a prospective observation of 2215 patients. *J Hepatol* 1998;28:930-8.
- Okuse C, Yotsuyanagi H, Koike K. Hepatitis C as a systemic disease: virus and host immunologic responses underlie hepatic and extrahepatic manifestations. *J Gastroenterol* 2007;42:857-65.
- Koike K, Moriya K. Metabolic aspects of hepatitis C: steatohepatitis distinct from NASH. *J Gastroenterol* 2005;40:329-36.
- Negro F. Insulin resistance and HCV: will new knowledge modify clinical management? *J Hepatol* 2006;45:514-9.
- Powell EE, Jonsson JR, Clouston AD. Steatosis: co-factor in other liver diseases. *Hepatology* 2005;42:5-13.
- Kiyosawa K, Sodeyama T, Tanaka E, Gibo Y, Yoshizawa K, Nakano Y, et al. Interrelationship of blood transfusion, non-A, non-B hepatitis and hepatocellular carcinoma: analysis by detection of antibody to hepatitis C virus. *Hepatology* 1990;12:671-5.
- Yotsuyanagi H, Shintani Y, Moriya K, Fujie H, Tsutsumi T, Kato T, et al. Virological analysis of non-B, non-C hepatocellular carcinoma in Japan: frequent involvement of hepatitis B virus. *J Infect Dis* 2000;181:1920-8.
- Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Ishibashi K, Matsuura Y, et al. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J Gen Virol* 1997;78:1527-31.
- Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Matsuura Y, et al. Hepatitis C virus core protein induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065-8.
- Koike K, Moriya K, Ishibashi K, Matsuura Y, Suzuki T, Saito I, et al. Expression of hepatitis C virus envelope proteins in transgenic mice. *J Gen Virol* 1995;76:3031-8.
- Koike K, Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Ishibashi K, et al. Sialadenitis resembling Sjögren's syndrome in mice transgenic for hepatitis C virus envelope genes. *Proc Natl Acad Sci USA* 1997;94:233-6.
- Bach N, Thung SN, Schaffner F. The histological features of chronic hepatitis C and autoimmune chronic hepatitis: a comparative analysis. *Hepatology* 1992;15:572-7.
- Lerat H, Honda M, Beard MR, Loesch K, Sun J, Yang Y, et al. Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352-65.
- Naas T, Ghorbani M, Alvarez-Maya I, Lapner M, Kothary R, De Repentigny Y, et al. Characterization of liver histopathology in a transgenic mouse model expressing genotype 1a hepatitis C virus core and envelope proteins 1 and 2. *J Gen Virol* 2005;86:2185-96.
- Machida K, Cheng KT, Lai CK, Jeng KS, Sung VM, Lai MM. Hepatitis C virus triggers mitochondrial permeability transition with production of reactive oxygen species, leading to DNA damage and STAT3 activation. *J Virol* 2006;80:7199-207.
- Moriya K, Fujie H, Yotsuyanagi H, Shintani Y, Tsutsumi T, Matsuura Y, et al. Subcellular localization of hepatitis C virus structural proteins expressed in transgenic liver. *Jpn J Med Sci Biol* 1997;50:169-77.
- Sasaki Y. Does oxidative stress participate in the development of hepatocellular carcinoma? *J Gastroenterol* 2006;41:1135-48.
- Moriya K, Nakagawa K, Santa T, Shintani Y, Fujie H, Miyoshi H, et al. Oxidative stress in the absence of inflammation in a mouse model for hepatitis C virus-associated hepatocellular carcinogenesis. *Cancer Res* 2001;61:4365-70.

20. Moriya K, Todoroki T, Tsutsumi T, Fujie H, Shintani Y, Miyoshi H, et al. Increase in the concentration of carbon 18 monounsaturated fatty acids in the liver with hepatitis C: analysis in transgenic mice and humans. *Biophys Biochem Res Commun* 2001;281:1207-12.
21. Okuda M, Li K, Beard MR, Showalter LA, Schole F, Lemon SM, Weinman SA. Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* 2002;122:366-75.
22. Tsutsumi T, Suzuki T, Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, et al. Intrahepatic cytokine expression and AP-1 activation in mice transgenic for hepatitis C virus core protein. *Virology* 2002;304:415-24.
23. Tsutsumi T, Suzuki T, Moriya K, Shintani Y, Fujie H, Miyoshi H, et al. Hepatitis C virus core protein activates ERK and p38 MAPK in cooperation with ethanol in transgenic mice. *Hepatology* 2003;38:820-8.
24. Miyamoto H, Moriishi K, Moriya K, Murata S, Tanaka K, Suzuki T, et al. Hepatitis C virus core protein induces insulin resistance through a PA28 γ -dependent pathway. *J Virol* 2007;81:1727-35.
25. Shintani Y, Fujie H, Miyoshi H, Tsutsumi T, Kimura S, Moriya K, et al. Hepatitis C virus and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840-8.
26. Perlemuter G, Sabile A, Letteron P, Vona G, Topilco A, Koike K, et al. Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J* 2002;16:185-94.
27. Moriishi K, Mochizuki R, Moriya K, Miyamoto H, Mori Y, Abe T, et al. Critical role of PA28 γ in hepatitis C virus-associated steatogenesis and hepatocarcinogenesis. *Proc Natl Acad Sci USA* 2007;104:1661-6.
28. Tsutsumi T, Suzuki T, Shimoike T, Moriya K, Yotsuyanagi H, Matsuura Y, et al. Interaction of hepatitis C virus core protein with retinoid X receptor- α modulates its transcriptional activity. *Hepatology* 2002;35:937-46.
29. Peters JM, Cheung C, Gonzalez FJ. Peroxisome proliferator-activated receptor-alpha and liver cancer: where do we stand? *J Mol Med* 2005;83:774-85.
30. Tanaka N, Moriya K, Kiyosawa K, Koike K, Aoyama T. Hepatitis C virus core protein induces spontaneous and persistent activation of peroxisome proliferator-activated receptor alpha in transgenic mice: implications for HCV-associated hepatocarcinogenesis. *Int J Cancer* 2008;122:124-31.
31. Akiyama TE, Sakai S, Lambert G, Nicol CJ, Matsusue K, Pimprale S, et al. Conditional disruption of the peroxisome proliferator-activated receptor gamma gene in mice results in lowered expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol efflux. *Mol Cell Biol* 2002;22:2607-19.
32. Tanaka N, Moriya K, Kiyosawa K, Koike K, Gonzalez FJ, Aoyama T. PPAR- α is essential for severe hepatic steatosis and hepatocellular carcinoma induced by HCV core protein. *J Clin Invest* 2008;118:683-94.
33. Kinzler KW, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;87:159-70.
34. Koike K. Molecular basis of hepatitis C virus-associated hepatocarcinogenesis: lessons from animal model studies. *Clin Gastroenterol Hepatol* 2005;3:S132-5.

