

virological responders (LVR) (HCV RNA-positive at 12 weeks and negative at 24 weeks after the start of treatment), because high relapse rates in LVR may indicate that treatment was not administered for a sufficient duration [Ferenci et al., 2005]. Previous studies from Europe and United States have demonstrated that LVR improves SVR rates when treatment is extended to 72 weeks, compared with standard duration of therapy, largely as a result of reducing posttreatment relapse rates [Buti et al., 2003; Berg et al., 2006; Sánchez-Tapias et al., 2006; Pearlman et al., 2007]. Thus, prolongation of therapy in LVR may improve the virological response rate. However, it is not clear at present whether prolongation of treatment improves the SVR rate of treatment-resistant Japanese patients infected with HCV/genotype 1b [Akuta et al., 2007a,b,c].

Previous studies indicated that amino acid (aa) substitutions at position 70 and/or 91 in the HCV core region of genotype 1b were predictors of poor virological response to 48-week PEG-IFN plus RBV therapy [Akuta et al., 2005, 2006, 2007a,b,c; Donlin et al., 2007], and also risk factors for hepatocarcinogenesis [Akuta et al., 2007d, 2008a]. However, it is not clear at this stage whether aa substitutions in the core region can be used before therapy to predict the outcome of 72-week regimen.

The aims of the present study in HCV genotype 1b-infected Japanese adult patients, who received PEG-IFN plus RBV, were the following: (1) To conduct a case-control study matched for sex, age, and periods from the start of treatment to the initial point of HCV RNA-negative, to compare the treatment efficacy of 72-week regimen and 48-week regimen. (2) To identify the pretreatment factors that could predict treatment efficacy of the 72-week regimen, including pretreatment aa substitutions in the core region.

PATIENTS AND METHODS

Study Population

A total of 559 HCV genotype 1b-infected Japanese adult patients were consecutively recruited into the study protocol of combination therapy with PEG-IFN α -2b plus RBV between 2001 and 2008 at Toranomon Hospital, Tokyo, Japan. They received PEG-IFN α -2b at a median dose of 1.4 μ g/kg (range, 0.7–2.1 μ g/kg) subcutaneously each week plus oral RBV at a median dose of 11.1 mg/kg (range, 3.4–16.0 mg/kg) daily. Among these, 383 patients, who could complete a total of 48 or 72 weeks of combination therapy, were enrolled in this retrospective study. The latter group consisted of 65 patients who extended combination therapy to 72-week (72-week group), and 318 patients who stopped combination therapy at the 48 weeks (48-week group). The decision to extend the combination therapy to 72 weeks was made by the patient. To compare the efficacy of the 72- and 48-week courses, all 65 patients of the 72-week group entered this study along with 130 patients of 48-week. The latter group was selected from among the 318 because they matched those

patients of the 72-week group with respect to sex, age, and periods from the start of treatment to the initial point of HCV RNA-negative (matched case-control study). The treatment efficacy was evaluated by HCV-RNA positive based on qualitative PCR analysis at the end of treatment (non-virological response; NVR), and by HCV-RNA negative based on qualitative PCR analysis at 24 weeks after the completion of therapy (SVR). Furthermore, LVR was defined as HCV RNA-positive at 12 weeks and negative at 24 weeks after the start of treatment, based on qualitative PCR analysis. All patients fulfilled the following criteria: (1) Negativity for hepatitis B surface antigen (radioimmunoassay, Dainabot, Tokyo, Japan), positivity for anti-HCV (third-generation enzyme immunoassay, Chiron Corp., Emerville, CA), and positivity for HCV RNA qualitative analysis with PCR (Amplicor, Roche Diagnostics, Mannheim, Germany). (2) Infection with HCV genotype 1b only. (3) A high viral load ($\geq 100 \times 10^3$ IU/ml) by quantitative analysis of HCV RNA with PCR (AMPLICOR GT HCV Monitor v2.0 using the 10-fold dilution method, Roche Molecular Systems Inc., Pleasanton, CA) within the preceding 2 months of enrolment. (4) No hepatocellular carcinoma. (5) Body weight > 40 kg. (6) Lack of coinfection with human immunodeficiency virus. (7) No previous treatment with antiviral or immunosuppressive agents within the preceding 3 months of enrolment. (8) None was an alcoholic; lifetime cumulative alcohol intake was <500 kg. (9) None had other forms of liver diseases, such as hemochromatosis, Wilson disease, primary biliary cirrhosis, alcoholic liver disease, or autoimmune liver disease. (10) None of the females was pregnant or a lactating mother. (11) All patients had completed a 24-week follow-up program after cessation of treatment, and SVR could be evaluated. (12) Each signed a consent form of the study protocol that had been approved by the human ethics review committee. The profile and laboratory data of 195 patients, who entered the matched case-control study, are summarized in Table I.

Laboratory Tests

Blood samples were obtained at least once every month before, during, and after treatment, and were analyzed for alanine aminotransferase (ALT) and HCV-RNA levels. The serum samples were frozen at -80°C within 4 hr of collection and thawed at the time of measurement. HCV genotype was determined by PCR using a mixed primer set derived from the nucleotide sequences of NS5 region [Chayama et al., 1993]. HCV-RNA levels were measured by quantitative PCR (AMPLICOR GT HCV Monitor v2.0 using the 10-fold dilution method, Roche Molecular Systems Inc.) at least once every month before, during, and after therapy. The dynamic range of the assay was 5.0×10^3 to 5.0×10^6 IU/ml. Samples collected during and after therapy that showed undetectable levels of HCV-RNA ($< 5.0 \times 10^3$ IU/ml) were also checked by qualitative PCR (AMPLICOR HCV v2.0, Roche Molecular Systems Inc.),

TABLE I. Patient Profile and Laboratory Data at Commencement of 48- and 72-Week Combination Therapy of Peginterferon Plus Ribavirin in Patients Infected With HCV Genotype 1b (Matched Case-Control Study)

	72-week group	48-week group	
Matching data			
Number of patients	65	130	
Sex (M/F)	28/37	57/73	Matched
Age (years)*	57 (22–70)	56 (25–68)	Matched
Periods to the initial point of HCV RNA-negative (weeks)*	17.4 (5.9–72.0)	19.7 (6.0–48.0)	Matched
Demographic data			
History of blood transfusion	18 (27.7%)	42 (32.3%)	NS
Family history of liver disease	21 (32.3%)	31 (23.8%)	NS
Body mass index (kg/m ²)*	22.6 (16.6–38.0)	22.2 (17.0–32.4)	NS
Laboratory data*			
Serum aspartate aminotransferase (IU/L)	49 (23–213)	51 (21–217)	NS
Serum alanine aminotransferase (IU/L)	64 (25–430)	68 (20–391)	NS
Serum albumin (g/dl)	3.9 (3.2–4.5)	3.8 (3.2–4.6)	NS
Gamma-glutamyl transpeptidase (IU/L)	40 (14–171)	38 (15–581)	NS
Leukocytes (/mm ³)	4,400 (2,300–8,800)	4,600 (1,200–9,400)	NS
Hemoglobin (g/dl)	14.0 (11.3–17.8)	13.9 (10.6–18.1)	NS
Platelet count ($\times 10^4$ /mm ³)	16.2 (8.2–30.7)	15.8 (6.4–31.6)	NS
ICG R15 (%)	13 (2–73)	15 (2–45)	NS
Level of viremia (KIU/ml)	2,650 (52–>5,000)	1,850 (49–>5,000)	0.013
Alfa-fetoprotein (μ g/L)	6 (2–47)	6 (2–110)	NS
Total cholesterol (mg/dl)	174 (111–276)	175 (104–274)	NS
High density lipoprotein cholesterol (mg/dl)	45 (27–86)	51 (24–78)	NS
Low density lipoprotein cholesterol (mg/dl)	104 (49–204)	107 (50–182)	NS
Triglycerides (mg/dl)	91 (35–259)	94 (35–315)	NS
Uric acid (mg/dl)	5.3 (2.6–7.7)	5.0 (2.3–8.7)	NS
Fasting blood sugar (mg/dl)	95 (79–218)	98 (76–157)	NS
Histological findings			
Stage of fibrosis (F1/F2/F3/ND)	20/12/11/1/21	44/27/22/0/37	NS
Hepatocyte steatosis (none to mild/moderate to severe/ND)	40/2/23	78/8/44	NS
Treatment			
PEG-IFN α -2b dose (μ g/kg)*	1.4 (0.8–2.1)	1.4 (0.7–1.9)	NS
Ribavirin dose (mg/kg)*	10.9 (6.6–16.0)	10.8 (3.7–14.2)	NS
Amino acid substitutions in the HCV			
Core aa 70 (arginine/glutamine (histidine)/ND)	37/23/5	11 47/6	NS
Core aa 91 (leucine/methionine/ND)	42/18/5	66/57/7	0.038
ISDR of NS5A (wild-type/mutant-type/ND)	49/5/11	99/17/14	NS

Data are number and percentages of patients, except those denoted by *, which represent the median (range) values. ND: not determined.

which has a higher sensitivity than quantitative analysis, and the results were expressed as positive or negative. The lower limit of the assay was 50 IU/ml.

Histopathological Examination of Liver Biopsies

Liver biopsy specimens were obtained percutaneously or at peritoneoscopy using a modified Vim Silverman needle with an internal diameter of 2 mm (Tohoku University style, Kakinuma Factory, Tokyo), fixed in 10% formalin, and stained with hematoxylin and eosin, Masson's trichrome, silver impregnation, and periodic acid-Schiff after diastase digestion. All specimens for examinations contained six or more portal areas. Histopathological diagnosis was confirmed by an experienced liver pathologist (H.K.) who was blinded to the clinical data. Chronic hepatitis was diagnosed based on histopathological assessment according to the scoring system of Desmet et al. [1994]. Hepatocyte steatosis was graded as either none (absent), mild (less than 1/3 of hepatocytes involved), moderate (greater than 1/3 but less than 2/3 of hepatocytes involved), or severe (greater

than 2/3 of hepatocytes involved) [D'Alessandro et al., 1991].

Detection of Amino Acid Substitutions in Core Region and NS5A Region

With the use of HCV-J (accession no. D90208) as a reference [Kato et al., 1990], the sequence of 1–191 aa in the core protein of genotype 1b was determined and then compared with the consensus sequence constructed on 50 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [Akuta et al., 2005]. The sequence of 2209–2248 aa in the NS5A of genotype 1b (IFN-sensitivity determining region [ISDR]) reported by Enomoto et al. [1995, 1996] was also determined, and the numbers of aa substitutions in ISDR were defined as wild-type (≤ 1) or mutant-type (≥ 2).

In the present study, aa substitutions of the core region and NS5A-ISDR were analyzed by direct sequencing [Enomoto et al., 1995, 1996; Akuta et al., 2005]. HCV RNA was extracted from serum samples at

the start of treatment and reverse transcribed with random primer and MMLV reverse transcriptase (Takara Syuzo, Tokyo). Nucleic acids were amplified by PCR using the following primers: (a) Nucleotide sequences of the core region: The first-round PCR was performed with CC11 (sense, 5'-GCC ATA GTG GTC TGC GGA AC-3') and e14 (antisense, 5'-GGA GCA GTC CTT CGT GAC ATG-3') primers, and the second-round PCR with CC9 (sense, 5'-GCT AGC CGA GTA GTG TT-3') and e14 (antisense) primers. (b) Nucleotide sequences of NS5A-ISDR: The first-round PCR was performed with ISDR1 (sense, 5'-ATG CCC ATG CCA GGT TCC AG-3') and ISDR2 (antisense, 5'-AGC TCC GCC AAG GCA GAA GA-3') primers, and the second-round PCR with ISDR3 (sense, 5'-ACC GGA TGT GGC AGT GCT CA-3') and ISDR4 (antisense, 5'-GTA ATC CGG GCG TGC CCA TA-3') primers ([a] hemi-nested PCR; [b] nested PCR). All samples were initially denatured at 95°C for 15 min. The 35 cycles of amplification were set as follows: denaturation for 1 min at 94°C, annealing of primers for 2 min at 55°C, and extension for 3 min at 72°C with an additional 7 min for extension. Then 1 µl of the first PCR product was transferred to the second PCR reaction. Other conditions for the second PCR were the same as the first PCR, except that the second PCR primers were used instead of the first PCR primers. The amplified PCR products were purified by the QIA quick PCR purification kit (Qiagen, Tokyo, Japan) after agarose gel electrophoresis and then used for direct sequencing. Dideoxynucleotide termination sequencing was performed with the Big Dye Deoxy Terminator Cycle Sequencing kit (Perkin-Elmer, Tokyo, Japan).

Statistical Analysis

Non-parametric tests (Mann-Whitney *U*-test, chi-squared test and Fisher's exact probability test) were used to compare the characteristics of the groups. Univariate and multivariate logistic regression analyses were used to determine those factors that significantly contributed to SVR. The odds ratios and 95% confidence intervals (95% CI) were also calculated. All *P* values less than 0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance ($P < 0.05$) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent factors. The potential pretreatment predictive factors associated with SVR included the following variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), ALT, albumin, gamma-glutamyl transpeptidase (γ GTP), leukocyte count, hemoglobin, platelets, indocyanine green retention rate at 15 min (ICG R15), level of viremia, alpha-fetoprotein, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, uric acid, fasting blood sugar, hepatocyte steatosis, stage of fibrosis, PEG-IFN dose/body weight, RBV dose/body weight, duration of treatment, and amino acid substitution in the core and ISDR of NS5A.

Statistical analyses were performed using the SPSS software (SPSS Inc., Chicago, IL).

RESULTS

Comparison of Treatment Efficacy Between 48-Week Group and 72-Week Group

Figure 1 shows comparison of the treatment efficacy between 48- and 72-week groups. SVR was achieved by 42 of 130 patients (32.3%) and 40 of 65 (61.5%) in the 48- and 72-week groups, respectively. The proportion of SVR was significantly higher in 72-week group than in the 48-week group ($P < 0.001$). Furthermore, NVR was identified in 38 of 130 patients (29.2%) and 6 of 65 (9.2%) in the 48- and 72-week groups, respectively. The proportion of NVR was significantly lower in the 72-week group than in 48-week group ($P = 0.002$).

Predictive Factors Associated With SVR in Multivariate Analysis

Univariate analysis identified 13 parameters that influenced SVR either significantly or marginally: gender (female sex; $P = 0.002$), stage of fibrosis ($F_{1,2}$; $P = 0.008$), PEG-IFN dose/body weight (≥ 1.4 µg/kg; $P = 0.001$), RBV dose/body weight (≥ 11.0 mg/kg; $P = 0.029$), platelet count ($\geq 15.0 \times 10^4/\text{mm}^3$; $P = 0.002$), level of viremia ($< 1,000$ KIU/ml; $P = 0.049$), γ GTP (< 50 IU/L; $P = 0.026$), ICG R15 ($< 15\%$; $P = 0.003$), triglycerides (< 100 mg/dl; $P = 0.038$), high-density lipoprotein cholesterol (≥ 50 mg/dl; $P = 0.018$), α -fetoprotein (< 20 µg/L; $P = 0.005$), substitution of aa 70 and 91 (Arg70 and/or Leu91; $P = 0.002$), and duration of treatment (72-week group; $P < 0.001$).

Multivariate analysis identified three independent parameters that either significantly influenced or tended to significantly influence SVR; substitution of aa 70 and 91 (Arg70 and/or Leu91; $P = 0.015$), duration of treatment (72-week group; $P = 0.014$), and high-density lipoprotein cholesterol (≥ 50 mg/dl; $P = 0.084$) (Table II).




Fig. 1. Comparison of treatment efficacy between the 48-week group and 72-week group. The proportion of patients with sustained virological response in 72-week group was significantly higher than in 48-week group ($P < 0.001$). Furthermore, the proportion of patients with non-virological response in 72-week group was significantly lower than in 48-week group ($P = 0.002$).

TABLE II. Factors Associated With Sustained Virological Response to Combination Therapy of Peginterferon Plus Ribavirin in 195 Patients Infected With HCV Genotype 1b, Identified by Multivariate Analysis

Factor	Category	Odds ratio (95% CI)	P
Substitution of aa 70 and 91	1: Gln70 (His70) and Met91	1	0.015
	2: Arg70 and/or Leu91	5.46 (1.39–21.3)	
Duration of treatment (weeks)	1: 48	1	0.014
	2: 72	3.51 (1.28–9.62)	
HDL-cholesterol (mg/dl)	1: <50	1	0.084
	2: ≥50	2.42 (0.89–6.58)	

*Only variables that achieved statistical significance ($P < 0.05$) or marginal significance ($P < 0.10$) on multivariate logistic regression are shown.

Treatment Efficacy According to Amino Acid Substitutions in Core Region

Figure 2 shows comparison of the treatment efficacy according to aa substitutions in the core region. In Gln70 (His70) and Met91, SVR was achieved by 4 of 26 patients (15.4%) and 3 of 10 (30.0%) in the 48- and 72-week groups, respectively. The proportion of SVR in 72-week group was not significantly different than in 48-week group. In Arg70 and/or Leu91, SVR was achieved by 37 of 98 patients (37.8%) and 34 of 50 (68.0%) in the 48- and 72-week groups, respectively. The proportion of SVR in 72-week group was significantly higher than in 48-week group ($P = 0.001$).

Treatment Efficacy According to Amino Acid Substitutions in NS5A-ISDR

Figure 3 shows comparison of the treatment efficacy according to aa substitutions in NS5A-ISDR. In mutant-type, SVR was achieved by 9 of 17 patients (52.9%) and 3 of 5 (60.0%) in the 48- and 72-week groups, respectively. The proportion of SVR in 72-week group was not significantly different from that in 48-week group. In wild-type, SVR was achieved by 29 of 99 patients (29.3%) and 30 of 49 (61.2%) in the 48- and 72-week groups, respectively. The proportion of SVR in 72-week group was significantly higher than that in 48-week group ($P < 0.001$).

DISCUSSION

This matched case-controlled study of PEG-IFN plus RBV for LVR infected with HCV genotype 1b, showed that treatment extension to 72 weeks seems to improve SVR rates in Japanese patients. To our knowledge, the present study is the first to report that 72-week regimen of PEG-IFN plus RBV might be also useful in Asians. Especially, the 72-week regimen significantly improved the SVR rates in LVR with Arg70 and/or Leu91 of core or wild-type of ISDR. The present study based on patients, who could complete a total of 48 or 72 weeks of combination therapy, did not show the frequencies of patients, who could not complete by side effects. Patients, who dropped out by side effects between 48 and 72 week for therapy prolonged to 72 weeks, were only 3 of 559 HCV genotype 1b-infected Japanese adult patients (data not shown), so the frequencies of side effects with 72-week regimen might be nearly equal to those with 48-week regimen. Large-scale prospective study based on the intention to treat analysis should be conducted to confirm the above finding in future.

NS5A-ISDR, reported as predictor of treatment efficacy with IFN monotherapy by Enomoto et al. [1995, 1996], is also useful as predictor of 48-week PEG-IFN plus RBV combination therapy [Murayama et al., 2007; Shirakawa et al., in press; Yen et al., 2008]. Furthermore, the present study also indicated that




Fig. 2. Comparison of treatment efficacy according to amino acid substitutions in the core region. In Gln70 (His70) and Met91, the proportion of patients with sustained virological response in 72-week group was not significantly different from that in 48-week group. However, in Arg70 and/or Leu91, the proportion of patients with sustained virological response in 72-week group was significantly higher than in 48-week group ($P = 0.001$).




Fig. 3. Comparison of treatment efficacy according to amino acid substitutions in NS5A-ISDR. In mutant-type, the proportion of patients with sustained virological response in 72-week group was not significantly different from that in 48-week group. However, in wild-type, the proportion of patients with sustained virological response in 72-week group was significantly higher than in 48-week group ($P < 0.001$).

72-week regimen of PEG-IFN plus RBV significantly improved the SVR rate in LVR with wild-type of ISDR. Unfortunately, the 72-week regimen of PEG-IFN plus RBV did not improve the SVR rate in LVR with Gln70 (His70) and Met91 of the core region. Multivariate analysis also identified Gln70 (His70) and Met91 of the core region as independent parameter that significantly influenced non-SVR. PEG-IFN plus RBV carries potential serious side effects and is costly especially when used long enough to achieve higher SVR rates. For these reasons, we need to identify those patients who do not achieve SVR, to free them of unnecessary side effects and reduce costs, preferably before the start of the combination therapy. For patients unsuitable for PEG-IFN plus RBV, including LVR with Gln70 (His70) and Met91 of the core region, low-dose intermittent IFN monotherapy might be an efficacious therapeutic regimen, because it can lead to ALT normalization and thus reduce the risk of hepatocarcinogenesis [Akuta et al., 2008b].

One limitation of this study is that LVR could not be evaluated by the COBAS AmpliPrep/COBAS TaqMan HCV Test (the lower limit of this assay; 15 IU/ml), which has a higher sensitivity than AMPLICOR HCV v2.0 (the lower limit of this assay; 50 IU/ml) [Sizmann et al., 2007]. Rapid virological response (HCV RNA-negative at 4 weeks after the start of treatment) and early virological response (HCV RNA-positive at 4 weeks and negative at 12 weeks after the start of treatment) by AMPLICOR HCV v2.0 might be diagnosed as LVR by the COBAS AmpliPrep/COBAS TaqMan HCV Test. Further studies using highly sensitive real-time PCR assay should be performed to facilitate the development of more effective therapeutic regimens in future.

We previously reported that viral factors (e.g., aa substitutions in core region) and host factors (e.g., lipid metabolic factors, sex, and AFP) might be important predictors of treatment response to 48-week PEG-IFN plus RBV in Japanese patients infected with HCV genotype 1b, in addition to treatment-related factors (e.g., RBV dose) [Akuta et al., 2005, 2006, 2007a,b,c]. The present study also identified viral (aa substitutions in the core region), host (HDL-cholesterol), and treatment-related factors (duration of treatment) that can be useful as independent and significant pretreatment predictors of SVR. Thus, substitution of aa 70 and 91 is also useful as a pretreatment predictor of 72-week regimen. Further studies that examine the structural and functional impact of aa substitutions during combination therapy should be conducted to confirm the above finding.

Another limitation of our study was that we did not examine aa substitutions in areas other than the core region and NS5A-ISDR of HCV genome, such as the interferon/ribavirin resistance determining region (IRRDR), including V3 of NS5A region, although they should be investigated in future studies [El-Shamy et al., 2008; Muñoz de Rueda et al., 2008].

We conclude that treatment efficacy of 72-week PEG-IFN plus RBV seems to be based on a dynamic tripartite

interaction of viral-, host-, and treatment-related factors. Further understanding of the complex interaction between these factors should facilitate the development of more effective therapeutic regimens.

REFERENCES

- Akuta N, Chayama K, Suzuki F, Someya T, Kobayashi M, Tsubota A, Suzuki Y, Saitoh S, Arase Y, Ikeda K, Kumada H. 2001. Risk factors of hepatitis C virus-related liver cirrhosis in young adults: Positive family history of liver disease and transporter associated with antigen processing 2 (TAP2) *0201 allele. *J Med Virol* 64:109–116.
- Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2005. Association of amino acid substitution pattern in core protein of hepatitis C virus genotype 1b high viral load and non-virological response to interferon-ribavirin combination therapy. *Intervirology* 48:372–380.
- Akuta N, Suzuki F, Sezaki H, Suzuki Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Watahiki S, Sato J, Matsuda M, Arase Y, Ikeda K, Kumada H. 2006. Predictive factors of virological non-response to interferon-ribavirin combination therapy for patients infected with hepatitis C virus of genotype 1b and high viral load. *J Med Virol* 78:83–90.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2007a. Predictive factors of early and sustained responses to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b: Amino acid substitutions in the core region and low-density lipoprotein cholesterol levels. *J Hepatol* 46:403–410.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2007b. Predictors of viral kinetics to peginterferon plus ribavirin combination therapy in Japanese patients infected with hepatitis C virus genotype 1b. *J Med Virol* 79:1686–1695.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Miyakawa Y, Kumada H. 2007c. Prediction of response to pegylated interferon and ribavirin in hepatitis C by polymorphisms in the viral core protein and very early dynamics of viremia. *Intervirology* 50:361–368.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2007d. Amino acid substitutions in the hepatitis C virus core region are the important predictor of hepatocarcinogenesis. *Hepatology* 46:1357–1364.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2008a. Substitution of amino acid 70 in the hepatitis C virus core region of genotype 1b is an important predictor of elevated alpha-fetoprotein in patients without hepatocellular carcinoma. *J Med Virol* 80:1354–1362.
- Akuta N, Suzuki F, Kawamura Y, Yatsuji H, Sezaki H, Suzuki Y, Hosaka T, Kobayashi M, Kobayashi M, Arase Y, Ikeda K, Kumada H. 2008b. Efficacy of low-dose intermittent interferon-alpha monotherapy in patients infected with hepatitis C virus genotype 1b who were predicted or failed to respond to pegylated interferon plus ribavirin combination therapy. *J Med Virol* 80:1363–1369.
- Berg T, von Wagner M, Nasser S, Sarrazin C, Heintges T, Gerlach T, Buggisch P, Goeser T, Rasenack J, Pape GR, Schmidt WE, Kallinowski B, Klinker H, Spengler U, Martus P, Alshuth U, Zeuzem S. 2006. Extended treatment duration for hepatitis C virus type 1: Comparing 48 versus 72 weeks of peginterferon-alfa-2a plus ribavirin. *Gastroenterology* 2006:1086–1097.
- Buti M, Valdés A, Sánchez-Avila F, Esteban R, Lurie Y. 2003. Extending combination therapy with peginterferon alfa-2b plus ribavirin for genotype 1 chronic hepatitis C late responders: A report of 9 cases. *Hepatology* 37:1226–1227.
- Chayama K, Tsubota A, Arase Y, Saitoh S, Koida I, Ikeda K, Matsumoto T, Kobayashi M, Iwasaki S, Koyama S, Morinaga T, Kumada H. 1993. Genotypic subtyping of hepatitis C virus. *J Gastroenterol Hepatol* 8:150–156.
- D'Alessandro AM, Kalayoglu M, Sollinger HW, Hoffmann RM, Reed A, Knechtle SJ, Pirsch JD, Hafez GR, Lorentzen D, Belzer FO. 1991. The predictive value of donor liver biopsies for the development of

- primary nonfunction after orthotopic liver transplantation. *Transplantation* 51:157–163.
- Davis GL, Balart LA, Schiff ER, Lindsay K, Bodenheimer HC, Jr., Perrillo RP, Carey W, Jacobson IM, Payne J, Dienstag JL, VanThiel DH, Tamburro C, Lefkowitz J, Albrecht J, Meschievitz C, Ortego TJ, Gibas A. 1989. Treatment of chronic hepatitis C with recombinant interferon alfa. A multicenter randomized, controlled trial. Hepatitis Interventional Group. *N Engl J Med* 321:1501–1506.
- Desmet VJ, Gerber M, Hoofnagle JH, Manna M, Scheuer PJ. 1994. Classification of chronic hepatitis: Diagnosis, grading and staging. *Hepatology* 19:1513–1520.
- Di Bisceglie AM, Martin P, Kassianides C, Lisker-Melman M, Murray L, Waggoner J, Goodman Z, Banks SM, Hoofnagle JH. 1989. Recombinant interferon alfa therapy for chronic hepatitis C. A randomized, double-blind, placebo-controlled trial. *N Engl J Med* 321:1506–1510.
- Donlin MJ, Cannon NA, Yao E, Li J, Wahed A, Taylor MW, Belle SH, Di Bisceglie AM, Aurora R, Tavis JE. 2007. Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *J Virol* 81:8211–8224.
- Dusheiko GM. 1998. The natural course of chronic hepatitis C: Implications for clinical practice. *J Viral Hepatol* 5:9–12.
- El-Shamy A, Nagano-Fujii M, Sasase N, Imoto S, Kim SR, Hotta H. 2008. Sequence variation in hepatitis C virus nonstructural protein 5A predicts clinical outcome of pegylated interferon/ribavirin combination therapy. *Hepatology* 48:38–47.
- Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Izumi N, Marumo F, Sato C. 1995. Comparison of full-length sequences of interferon sensitive and resistant hepatitis C virus 1b. Sensitivity to interferon is conferred by amino acid substitutions in the NS5A region. *J Clin Invest* 96:224–230.
- Enomoto N, Sakuma I, Asahina Y, Kurosaki M, Murakami T, Yamamoto C, Ogura Y, Izumi N, Marumo F, Sato C. 1996. Mutations in the nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 334:77–81.
- Ferenci P, Fried MW, Shiffman ML, Smith CI, Marinos G, Gonçalves FL, Jr., Häussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Chaneac M, Reddy KR. 2005. Predicting sustained virological responses in chronic hepatitis C patients treated with peginterferon alfa-2a (40 KD)/ribavirin. *J Hepatol* 43:425–433.
- Fried MW, Shiffman ML, Reddy R, Smith C, Marinos G, Gonçalves FL, Häussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Lin A, Hoffman J, Yu J. 2002. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 347:975–982.
- Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Koida I, Arase Y, Fukuda M, Chayama K, Murashima N, Kumada H. 1998. Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: A prospective observation of 2215 patients. *J Hepatol* 28:930–938.
- Kato N, Hijikata M, Ootsuyama Y, Nakagawa M, Ohkoshi S, Sugimura T, Shimotohno K. 1990. Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci USA* 87:9524–9528.
- Kenny-Walsh E. 1999. Clinical outcomes after hepatitis C infection from contaminated anti-D immune globulin. Irish Hepatology Research Group. *N Engl J Med* 340:1228–1233.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, Goodman ZD, Koury K, Ling MH, Albrecht JK. 2001. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: A randomized trial. *Lancet* 358:958–965.
- Muñoz de Rueda P, Casado J, Patón R, Quintero D, Palacios A, Gila A, Quiles R, León J, Ruiz-Extremera A, Salmerón J. 2008. Mutations in E2-PePHD, NS5A-PKRBD, NS5A-ISDR, and NS5A-V3 of hepatitis C virus genotype 1 and their relationships to pegylated interferon-ribavirin treatment responses. *J Virol* 82:6644–6653.
- Murayama M, Katano Y, Nakano I, Ishigami M, Hayashi K, Honda T, Hirooka Y, Itoh A, Goto H. 2007. A mutation in the interferon sensitivity-determining region is associated with responsiveness to interferon-ribavirin combination therapy in chronic hepatitis patients infected with a Japan-specific subtype of hepatitis C virus genotype 1B. *J Med Virol* 79:35–40.
- Niederrau C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hürter D, Nawrocki M, Kruska L, Hensel F, Petry W, Häussinger D. 1998. Progress of chronic hepatitis C: Results of a large, prospective cohort study. *Hepatology* 28:1687–1695.
- Pearlman BL, Ehleben C, Saifee S. 2007. Treatment extension to 72 weeks of peginterferon and ribavirin in hepatitis c genotype 1-infected slow responders. *Hepatology* 46:1688–1694.
- Sánchez-Tapias JM, Diago M, Escartín P, Enríquez J, Romero-Gómez M, Bárcena R, Crespo J, Andrade R, Martínez-Bauer E, Pérez R, Testillano M, Planas R, Solá R, García-Bengoechea M, García-Samaniego J, Muñoz-Sánchez M, Moreno-Otero R, TeraVic-4 Study Group. 2006. Peginterferon-alfa2a plus ribavirin for 48 versus 72 weeks in patients with detectable hepatitis C virus RNA at week 4 of treatment. *Gastroenterology* 131:451–460.
- Shirakawa H, Matsumoto A, Joshita S, Komatsu M, Tanaka N, Umemura T, Ichijo T, Yoshizawa K, Kiyosawa K, Tanaka E, the Nagano Interferon Treatment Research Group. 2008. Pretreatment prediction of virologic response to peginterferon plus ribavirin therapy in chronic hepatitis C patients using viral and host factors. *Hepatology* 48:1753–1760.
- Sizmann D, Boeck C, Boelter J, Fischer D, Miethke M, Nicolaus S, Zadak M, Babel R. 2007. Fully automated quantification of hepatitis C virus (HCV) RNA in human plasma and human serum by the COBAS® AmpliPrep/COBAS® TaqMan® System. *J Clin Virol* 38:326–333.
- Yen YH, Hung CH, Hu TH, Chen CH, Wu CM, Wang JH, Lu SN, Lee CM. 2008. Mutations in the interferon sensitivity-determining region (nonstructural 5A amino acid 2209–2248) in patients with hepatitis C-1b infection and correlating response to combined therapy of pegylated interferon and ribavirin. *Aliment Pharmacol Ther* 27:72–79.

Virological Response and Hepatocarcinogenesis in Lamivudine-Resistant Hepatitis B Virus Genotype C Patients Treated with Lamivudine plus Adefovir Dipivoxil

Norio Akuta^a Fumitaka Suzuki^a Yusuke Kawamura^a Hiromi Yatsuji^a
Hitomi Sezaki^a Yoshiyuki Suzuki^a Tetsuya Hosaka^a Masahiro Kobayashi^a
Mariko Kobayashi^b Yasuji Arase^a Kenji Ikeda^a Hiromitsu Kumada^a

^aDepartment of Hepatology and ^bLiver Research Laboratory, Toranomon Hospital, Tokyo, Japan

Key Words

Hepatitis B virus · Lamivudine · Adefovir dipivoxil · Hepatocellular carcinoma · Basic core promoter · Precore · Core region

Abstract

Aims: The long-term efficacy of adefovir dipivoxil in combination with lamivudine to chronic hepatitis B virus (HBV) infection is still unclear. **Methods:** Virological response and hepatocarcinogenesis during lamivudine + adefovir were investigated in 183 lamivudine-resistant Japanese patients with chronic genotype C-dominant HBV infection. As the predictors of virological response, an assessment of clinical parameters and a nucleotide (nt) sequence analysis of the negative regulatory element to core gene (nt 1611–2450) were performed at the start of adefovir. **Results:** The cumulative HBV-DNA non-detectable and ALT normalization rates were 93.6 and 97.6% at the end of 3 years, respectively. Multivariate analysis identified total bilirubin, AST, and nt substitutions (nt 1762, 1768, 1846, 1896, 2134, 2288, 2441) as determinants of early non-detectable HBV-DNA. The yearly incidence of hepatocellular carcinoma (HCC) during the first

3 years was 2.7%. At the diagnosis of HCC, ALT normalization, HBV-DNA non-detectable, and HBeAg-seronegative conversion rates were 75.0, 83.3, and 57.1%, respectively. Furthermore, the cumulative HBV-DNA non-detectable and ALT normalization rates were not significantly different according to the development of HCC or not. **Conclusions:** Lamivudine-resistant patients treated with lamivudine + adefovir could achieve the excellent virological response and biochemical response, but the low hepatitis activity was not enough to suppress hepatocarcinogenesis.

Copyright © 2009 S. Karger AG, Basel

Introduction

Hepatitis B virus (HBV) is a small, enveloped DNA virus known to cause chronic hepatitis and often leads to liver cirrhosis and hepatocellular carcinoma (HCC) [1, 2]. To date, interferon and five nucleoside and nucleotide analogs (lamivudine, adefovir dipivoxil, entecavir, telbivudine, and tenofovir) have been approved for the treatment of chronic HBV infection. Nucleoside and nucleotide analogues suppress HBV replication in most patients and

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2009 S. Karger AG, Basel
0300–5526/08/0516–0393\$24.50/0

Accessible online at:
www.karger.com/int

Norio Akuta, MD
Department of Hepatology, Toranomon Hospital
2-2-2 Toranomon, Minato-ku, Tokyo 105-0001 (Japan)
Tel. +81 44 877 5111, Fax +81 44 860 1623
E-Mail akuta-gi@umin.ac.jp

improve transaminase levels and liver histology [3–7]. Especially lamivudine monotherapy to naive patients for nucleoside analogues suppresses hepatocarcinogenesis [8, 9], but prolonged therapy results in the emergence of drug-resistant mutants.

Most lamivudine-resistant strains show amino acid substitutions in the YMDD (tyrosine-methionine-aspartate-aspartate) motif in the C domain of HBV polymerase [10, 11]. Both experimental and clinical studies have shown recently that adefovir and entecavir could suppress not only wild-type but also lamivudine-resistant strains and were confirmed as salvage therapy for lamivudine-refractory patients [12, 13]. Recently, Hosaka et al. [14] reported the efficacy of adefovir + lamivudine combination therapy in patients with lamivudine-resistant chronic HBV infection. However, the number of patients was limited and follow-up time was a short duration. Thus, the long-term efficacy in respect to viral response and suppression of hepatocarcinogenesis with lamivudine + adefovir is still unclear.

Virological predictors of viral response during the treatment of lamivudine + adefovir are insufficiently investigated. Negative regulatory element (NRE; nt 1611–1634), core upstream regulatory sequences (CURS; nt 1643–1742), basic core promoter (BCP; nt 1742–1849) are located mainly in the HBV X gene and play an important role in replication and hepatitis B core antigen/HBeAg formation [15–20]. Furthermore, in respect to the viral response to interferon, Erhardt et al. [21] reported that good response in HBeAg-positive patients was associated with a high number of mutations in the BCP and nt 1753–1766 as well as mutations at nt 1764, and that good response in HBeAg-negative patients correlated with a low number of mutations in the BCP and nt 1753–1766 and wild-type sequence at nt 1764. However, the significance of substitutions in NRE, CURS, BCP, precore, and core gene for viral response during the treatment of lamivudine + adefovir is still unknown.

The present study based on the long follow-up time included 183 lamivudine-resistant consecutive patients with chronic genotype C-dominant HBV infection treated with lamivudine + adefovir. The aims of the study were the following: (1) to evaluate the cumulative HBV-DNA non-detectable, alanine aminotransferase (ALT) normalization, and hepatocarcinogenesis rates during the treatment of lamivudine + adefovir, and (2) to analyze the predictive factors, including clinical parameters and a sequence analysis of the complete NRE, CURS, BCP, precore, and core gene, associated with early non-detectable HBV-DNA during the treatment of lamivudine + adefovir.

Table 1. Patient characteristics at the start of treatment with lamivudine + adefovir dipivoxil

Number	183
Male/female	150/33
Age, years ^a	47 (26–75)
Prior lamivudine therapy duration, years ^a	2.9 (0.6–10.8)
Lamivudine + adefovir treatment duration years ^a	2.2 (0.5–4.5)
HBeAg, number positive	109 (59.6%)
HBV-DNA, log copies/ml ^a	7.3 (3.3 to >7.6)
HBV genotype, number of A/B/C/D	7/7/168/1
Presence of cirrhosis	56 (30.6%)
Total bilirubin, mg/dl ^a	0.8 (0.2–6.0)
Aspartate aminotransferase, IU/l ^a	92 (18–1,413)
Alanine aminotransferase, IU/l ^a	130 (18–1,563)
γ -Glutamyl transpeptidase, IU/l ^a	58 (12–446)
Albumin, g/dl ^a	4.1 (2.3–4.7)
α -Fetoprotein, μ g/l ^a	6 (2–282)
Creatinine, mg/dl ^a	0.8 (0.4–1.3)
Platelets, $\times 10^4/\text{mm}^3$ ^a	15.0 (3.1–38.8)
Mutant type of YMDD motif (YIDD/YVDD/YIDD+YVDD)	85/42/56

^a Data are expressed as median (range).

Patients and Methods

Study Population

A total of 183 consecutive adult Japanese patients with chronic HBV infection were treated with adefovir at Toranomon Hospital, Tokyo, Japan, in addition to ongoing lamivudine treatment, for more than 24 weeks since 2002. Serum HBV-DNA and ALT levels re-increased despite the continuation of lamivudine, indicating breakthrough hepatitis, in all patients who then received adefovir along with the lamivudine. Enrolment in this study and the start of adefovir treatment were determined by the following criteria: (1) Increase in serum HBV DNA levels of ≥ 1 log copies/ml during lamivudine treatment on at least two consecutive occasions, compared with the nadir of initial antiviral efficacy. (2) Detection of mutations of the YMDD motif before the start of adefovir treatment by the PCR-based method described later and/or direct sequence analysis. (3) No history of treatment with other nucleoside analogues such as famciclovir and entecavir. The exclusion criteria were as follows: (1) patients with HCC; (2) serum creatinine levels ≥ 1.5 mg/dl; (3) patients coinfecting with hepatitis C, hepatitis delta virus, or HIV, and (4) history of other liver diseases, such as autoimmune hepatitis, alcoholic liver disease, or metabolic liver disease.

This study was conducted in accordance with the guidelines of the Declaration of Helsinki and its subsequent amendments, and informed consent was obtained from every patient. This study was approved by the Local Ethics Committee of Toranomon Hospital.

Table 1 summarizes the profiles of the patients. They included 150 men and 33 women. The median duration of treatment

with lamivudine + adefovir was 2.2 years (range 0.5–4.5). Patients received a 10-mg once-daily dose of oral adefovir, in addition to ongoing lamivudine treatment (100 mg/day). Blood samples were obtained at least once every month before, during, and after treatment with lamivudine + adefovir, and analyzed for virological markers, biochemical markers associated with liver function and renal function, and complete blood cell counts every visit. The diagnosis of cirrhosis was based on liver biopsy histology and/or on clinical criteria, including imaging studies and signs of portal hypertension. As the indicators of low hepatitis activity, non-detectable HBV-DNA level by PCR assay and normalization of ALT level were evaluated. Adverse reactions were monitored clinically by careful interview and medical examination at least once every month. Patient compliance with treatment was evaluated by questionnaire. Follow-up time represented the time from the start of the treatment with lamivudine + adefovir until the last visit.

Laboratory Tests

HBeAg, HBsAg and antibody against HBeAg (anti-HBe) were determined by commercially available radioimmunoassay systems (Abbott Japan, Tokyo, Japan). HBV DNA serum level was determined by using the Amplicor HBV monitor test (Roche Diagnostics, Tokyo, Japan). The measurement range of the assay is $10^{2.6}$ – $10^{7.6}$ copies/ml (2.6–7.6 log copies/ml). The HBV genotype was determined by enzyme-linked immunosorbent assay (ELISA) (HBV Genotype EIA, Institute of Immunology, Tokyo, Japan) based on the method of Usuda et al. [22]. Substitution at rtM204 of the YMDD motif was identified at baseline by using the Enzyme-Linked Mini-Sequence Assay with a commercial assay kit (PCR-ELMA; Genome Science, Tokyo, Japan).

Nucleotide Sequencing of Negative Regulatory Element, Core Upstream Regulatory Sequences, Basic Core Promoter, Precore, and Core Gene

The sequences of nt 1611–2450, including the complete NRE (nt 1611–1634), CURS (nt 1643–1742), BCP (nt 1742–1849), pre-core (nt 1814–1901), and core gene (nt 1901–2450), were determined by the direct sequencing method using sera at the start of adefovir treatment. Nucleotide sequences of HBV were compared with the prototype sequences of the HBV genotype C (accession No. AB033550) [23]. In the present study, the PCR genotyping could be performed in 148 patients; the remaining 35 patients could not be analyzed due to the lack of adequate serum samples obtained at the start of adefovir treatment.

HBV DNA was extracted with a Smitest EX-R&D kit (Genome Science). Nucleic acids were amplified by PCR using the following primers: (a) *Sequences of nt 1588–2130*: the single-round PCR was performed with HBVPCCPseqF01 (sense, 5'-GCT TCA CCT CTG CAC GTC GCA TG-3' [nt 1588–1610]) and HBVPCCPseqR03 (antisense, 5'-TCC AAA TTA CTT CCC ACC CAG GT-3' [nt 2130–2108]) primers. (b) *Sequences of nt 2022–2529*: the single-round PCR was performed with HBVCOREseqF01 (sense, 5'-CCT TAG AGT CTC CGG AAC ATT G-3' [nt 2022–2043]) and HBVCOREseqR02 (antisense, 5'-GCC ACT CAG GAT TAA AGA CAG G-3' [nt 2529–2508]) primers. All samples were initially denatured at 95° for 2 min. 45 cycles of amplification were set as follows: denaturation for 30 s at 94°, annealing of primers for 30 s at 60°, and extension for 30 s at 68° with an additional 7 min for extension. The amplified PCR products were purified by the QIA

Quick PCR purification kit (Qiagen, Tokyo, Japan) after agarose gel electrophoresis and then used for direct sequencing. Dideoxynucleotide termination sequencing was performed with the Big Dye Deoxy Terminator Cycle Sequencing kit (PerkinElmer, Tokyo, Japan). To avoid false-positive results, the procedures recommended by Kwok and Higuchi [24] to prevent contamination were strictly applied to these PCR assays. No false-positive results were observed in this study.

Liver Histopathological Examination

Liver biopsy specimens were obtained percutaneously or at peritoneoscopy using a modified Vim Silverman needle with an internal diameter of 2 mm (Tohoku University style, Kakinuma Factory, Tokyo, Japan), fixed in 10% formalin, and stained with hematoxylin and eosin, Masson's trichrome, silver impregnation, and periodic acid-Schiff after diastase digestion. All specimens for examination contained 6 or more portal areas. Histopathological diagnosis was made by an experienced liver pathologist (H.K.) who was blinded to the clinical data. Chronic hepatitis was diagnosed based on histopathological assessment according to the scoring system of Desmet et al. [25].

Diagnosis of Hepatocellular Carcinoma

Patients were examined for HCC by abdominal ultrasonography every 3–6 months. If HCC was suspected based on ultrasonographic results, additional procedures, such as computed tomography, magnetic resonance imaging, abdominal angiography, and ultrasonography-guided tumor biopsy if necessary, were used to confirm the diagnosis.

Statistical Analysis

The cumulative rates of non-detectable HBV-DNA and hepatocarcinogenesis were calculated using the Kaplan-Meier method and differences between the curves were tested using the log-rank test. Statistical analyses of non-detectable HBV-DNA and hepatocarcinogenesis were calculated using the period from start of treatment with lamivudine + adefovir. Stepwise Cox regression analysis was used to determine independent predictive factors that were associated with non-detectable HBV-DNA. The odds ratios and 95% confidence intervals (95% CI) were also calculated. Potential predictive factors associated with early HBV-DNA negativity included the following variables: age, sex, histological stage, HBV genotype, HBeAg, viremia level, mutant type of YMDD motif, total bilirubin, aspartate aminotransferase (AST), ALT, albumin, γ -glutamyl transpeptidase (GGT), α -fetoprotein (AFP), creatinine, platelets, nt substitutions in CURS to core gene. Each variable was transformed into categorical data consisting of two simple ordinal numbers for uni- and multivariate analyses. Variables that achieved statistical significance ($p < 0.05$) or marginal significance ($p < 0.10$) on univariate Cox proportional hazards model were tested by multivariate Cox proportional hazards model to identify significant independent factors. Statistical comparisons were performed using the SPSS software (SPSS Inc., Chicago, Ill., USA). All p values < 0.05 by the two-tailed test were considered significant.




Fig. 1. Cumulative HBV-DNA non-detectable and ALT normalization rates. Patients treated lamivudine + adefovir dipivoxil could achieve the excellent virological response (non-detectable HBV-DNA) and biochemical response (ALT normalization) as an indicator of low hepatitis activity.




Fig. 2. Cumulative hepatocarcinogenesis rates during the treatment of lamivudine + adefovir dipivoxil. The yearly incidence of HCC during the first 3 years was 2.7%.

Results

Cumulative HBV-DNA Non-Detectable and ALT Normalization Rates

The cumulative HBV-DNA non-detectable rates were 48.4, 66.8, 79.4, and 93.6 at the end of 0.5, 1, 2, and 3 years, respectively. The cumulative ALT normalization rates were 72.4, 84.0, 97.6, and 97.6 at the end of 0.5, 1, 2, and 3 years, respectively. Thus, patients treated with lamivudine + adefovir could achieve the excellent virological response (non-detectable HBV-DNA) and biochemical response (ALT normalization) as an indicator of low hepatitis activity (fig. 1).

Predictive Factors Associated with Early Non-Detectable HBV-DNA by Uni- and Multivariate Analysis

The data for the whole population sample were analyzed to determine those factors that could predict early non-detectable HBV-DNA. Univariate analysis identified 21 parameters that tended to or significantly correlated with early non-detectable HBV-DNA. These included total bilirubin ($p = 0.027$), AST ($p = 0.004$), ALT ($p = 0.072$), HBV DNA ($p < 0.001$), HBeAg ($p < 0.001$), and nt substitutions [nt 1659 ($p = 0.073$), nt 1762 ($p = 0.040$), nt 1768 ($p =$

0.084), nt 1792 ($p = 0.077$), nt 1846 ($p < 0.001$), nt 1896 ($p < 0.001$), nt 1899 ($p = 0.031$), nt 1938 ($p = 0.019$), nt 2005 ($p = 0.058$), nt 2009 ($p < 0.001$), nt 2134 ($p = 0.074$), nt 2189 ($p = 0.017$), nt 2201 ($p = 0.031$), nt 2288 ($p = 0.038$), nt 2429 ($p = 0.042$), nt 2441 ($p < 0.001$)]. These factors were entered into multivariate analysis, which then identified 9 parameters that tended to or significantly influenced early non-detectable HBV-DNA independently; total bilirubin ($p = 0.002$), aspartate aminotransferase ($p = 0.077$), and nt substitutions [nt 1762 ($p = 0.092$), nt 1768 ($p = 0.001$), nt 1846 ($p = 0.034$), nt 1896 ($p = 0.001$), nt 2134 ($p = 0.034$), nt 2288 ($p = 0.016$), nt 2441 ($p = 0.019$)] (table 2).

Cumulative Hepatocarcinogenesis Rates and the Profiles of Patients Who Developed HCC

The cumulative hepatocarcinogenesis rates were 2.2, 5.9, and 8.1% at the end of 1, 2, and 3 years, respectively (fig. 2). The yearly incidence of HCC during the first 3 years was 2.7%. Table 3 summarizes the profiles of 12 patients who developed HCC during treatment with lamivudine + adefovir. They included 9 men and 3 women. The median age at the start of adefovir was 51 years (range 35–75). The median duration from the start of lamivudine to the diagnosis of HCC was 4.9 years (range 1.9–7.5), and the median duration from the start of adefovir

Table 2. Factors associated with early non-detectable HBV-DNA during the treatment with lamivudine + adefovir dipivoxil, identified by uni- and multivariate analysis

Factor	Category	Univariate Cox proportional hazards model		Multivariate Cox proportional hazards model	
		odds ratio (95% CI)	p	odds ratio (95% CI)	p
Total bilirubin, mg/dl	1: <1.0 2: ≥1.0	1 1.503 (1.047–2.159)	0.027	1 2.055 (1.289–3.279)	0.002
Aspartate aminotransferase, IU/l	1: <80 2: ≥80	1 1.695 (1.181–2.434)	0.004	1 1.506 (0.956–2.371)	0.077
Alanine aminotransferase, IU/l	1: <100 2: ≥100	1 1.407 (0.970–2.041)	0.072	– –	– –
HBV DNA, log copies/ml	1: <7.0 2: ≥7.0	1 0.488 (0.342–0.695)	<0.001	– –	– –
HBeAg	1: negative 2: positive	1 0.428 (0.299–0.613)	<0.001	– –	– –
nt 1659	1: A 2: not A	1 2.135 (0.931–4.895)	0.073	– –	– –
nt 1762	1: A 2: not A	1 1.988 (1.032–3.829)	0.040	1 1.987 (0.893–4.421)	0.092
nt 1768	1: T 2: not T	1 1.892 (0.917–3.903)	0.084	1 5.584 (2.096–14.88)	0.001
nt 1792	1: A 2: not A	1 0.168 (0.023–1.211)	0.077	– –	– –
nt 1846	1: A 2: not A	1 2.080 (1.382–3.131)	<0.001	1 1.740 (1.043–2.902)	0.034
nt 1896	1: G 2: not G	1 2.207 (1.500–3.247)	<0.001	1 2.323 (1.430–3.775)	0.001
nt 1899	1: G 2: not G	1 1.711 (1.049–2.789)	0.031	– –	– –
nt 1938	1: T 2: not T	1 1.859 (1.107–3.124)	0.019	– –	– –
nt 2005	1: T 2: not T	1 0.661 (0.431–1.014)	0.058	– –	– –
nt 2009	1: C 2: not C	1 4.678 (2.191–9.986)	<0.001	– –	– –
nt 2134	1: C 2: not C	1 1.566 (0.957–2.561)	0.074	1 1.781 (1.044–3.038)	0.034
nt 2189	1: A 2: not A	1 1.611 (1.087–2.385)	0.017	– –	– –
nt 2201	1: T 2: not T	1 0.596 (0.373–0.953)	0.031	– –	– –
nt 2288	1: C 2: not C	1 1.518 (1.024–2.252)	0.038	1 1.733 (1.108–2.711)	0.016
nt 2429	1: C 2: not C	1 2.573 (1.033–6.408)	0.042	– –	– –
nt 2441	1: T 2: not T	1 2.815 (1.656–4.783)	<0.001	1 2.001 (1.122–3.568)	0.019

Only variables that achieved statistical significance ($p < 0.05$) or marginal significance ($p < 0.10$) on uni- and multivariate Cox proportional hazards model are shown.
95% CI = 95% confidence interval.




Fig. 3. Comparison of non-detectable HBV-DNA and ALT normalization in patients who developed HCC or not. The cumulative HBV-DNA non-detectable and ALT normalization rates were not significantly different according to the development of HCC or not. Low hepatitis activity during the treatment of lamivudine + adefovir dipivoxil was not enough to suppress hepatocarcinogenesis.

Table 3. Characteristics of 12 patients who developed HCC during treatment with lamivudine (LAM) + adefovir dipivoxil (ADV)

Case	Sex	Age years ^a	LAM to HCC years ^b	ADV to HCC years ^c	At the start of ADV			At the diagnosis of HCC		
					HBeAg	ALT IU/l	HBV-DNA log copies/ml	HBeAg	ALT IU/l	HBV-DNA log copies/ml
1	male	50	6.6	4.5	-	576	6.9	-	27	<2.6
2	male	40	4.9	3.8	-	124	6.3	-	24	<2.6
3	male	48	6.3	3.3	+	99	7.6	+	35	<2.6
4	female	58	7.4	3.3	+	214	4.4	-	13	<2.6
5	male	58	4.8	2.2	-	216	6.5	-	28	<2.6
6	male	35	7.5	1.6	+	164	7.5	-	35	<2.6
7	male	47	2.3	1.3	+	138	7.6	+	29	<2.6
8	male	50	2.5	1.1	+	272	7.6	-	72	<2.6
9	male	75	3.4	1.1	+	209	7.6	-	125	<2.6
10	female	51	1.9	0.9	-	130	5.3	-	73	<2.6
11	male	53	4.7	0.5	+	97	7.6	+	35	4.2
12	female	59	5.9	0.1	-	132	7.6	-	41	3.6

^a Age at the start of adefovir dipivoxil.

^b Duration from the start of lamivudine to the diagnosis of HCC.

^c Duration from the start of adefovir dipivoxil to the diagnosis of HCC.

to the diagnosis of HCC was 1.5 years (range 0.1–4.5). At the diagnosis of HCC, 75.0% (9/12 patients) could achieve ALT normalization, and 83.3% (10/12) could achieve HBV-DNA non-detectable. 57.1% (4/7) of HBeAg-posit-

ive at the start of adefovir could achieve HBeAg-seronegative conversion at the diagnosis of HCC. Thus, they developed HCC in spite of the excellent virological response and biochemical response.

Comparison of Non-Detectable HBV-DNA and ALT Normalization in Patients Who Developed HCC or Not

The cumulative HBV-DNA non-detectable and ALT normalization rates were not significantly different according to the development of HCC or not (fig. 3). Thus, some of the patients during treatment with lamivudine + adefovir developed HCC in spite of the early non-detectable HBV-DNA and ALT normalization, and low hepatitis activity during treatment with lamivudine + adefovir was not enough to suppress hepatocarcinogenesis.

Discussion

This is the first report that investigates virological response and hepatocarcinogenesis during the treatment of lamivudine + adefovir in lamivudine-resistant patients with chronic genotype C-dominant HBV infection. Multivariate analysis identified total bilirubin, aspartate aminotransaminase, and nt substitutions (nt 1762, 1768, 1846, 1896, 2134, 2288, 2441) as determinants of early non-detectable HBV-DNA. Erhardt et al. [21] reported that the viral response to interferon was associated with a number of mutations in the BCP and nt 1753–1766 as well as mutation at nt 1764. As determinants of early non-detectable HBV-DNA, this study did not only identify nt substitutions in BCP (nt 1762, 1768), but also identified nt substitutions in precore (nt 1846, 1896), and core (nt 2134, 2288, 2441). This discrepancy between this results and previous findings may be explained by the difference of antiviral treatment, and design of this cohort study based on the only lamivudine-resistant patients during the treatment of lamivudine + adefovir. To our knowledge, the present study is the first to report that the precore, and core gene might influence viral response during lamivudine + adefovir. One limitation of the present study based on the small number of patients was that nt substitutions in areas other than the NRE, CURS, BCP, precore, and core gene of HBV genome, could not be examined. Further prospective studies based on the large numbers of patients, that examine the clinical impact of nt substitutions during lamivudine + adefovir (e.g., virological response and hepatocarcinogenesis) and the underlying mechanisms, should be conducted to confirm the above finding.

Lampertico et al. [26] recently reported the hepatocarcinogenesis during the treatment of lamivudine + adefovir in lamivudine-resistant patients with chronic genotype D-dominant HBV infection for a long-term follow-

up period. To our knowledge, the present study is the first to report the hepatocarcinogenesis rates in patients with chronic genotype C-dominant HBV infection. Lamivudine-resistant patients treated with lamivudine + adefovir could achieve the excellent virological response and biochemical response, but the low hepatitis activity was not enough to suppress hepatocarcinogenesis. Kobayashi et al. [27] reported that the yearly incidence of HCC during the first 10 years was 3.3% in natural histories of patients with HBV genotype C-related-compensated cirrhosis without antiviral treatment, who have the higher risk for HCC development. This result showed that the yearly incidence of HCC during the first 3 years was 2.7% during the treatment of lamivudine + adefovir. Treatment of lamivudine + adefovir did not worsen natural histories of chronic HBV infection, but indicated the almost similar hepatocarcinogenesis rates in comparison to cirrhosis patients without antiviral therapy (namely, high-risk group for HCC development). Thus, lamivudine monotherapy to naive patients for nucleoside analogues without lamivudine-resistant HBV infection suppresses hepatocarcinogenesis [8, 9], but lamivudine-resistant chronic HBV patients might be also one of the high-risk groups for hepatocarcinogenesis. This study indicated the high cumulative hepatocarcinogenesis rates of 46.4% at the end of 4 years, and this reason is probably related to the small number of patients, in whom more than 4 years had elapsed since the induction of adefovir (data not shown). Further studies of a large group of patients for the longer-term follow-up period are required to clarify the true cumulative hepatocarcinogenesis rates during the treatment of lamivudine + adefovir.

Low hepatitis activity by suppression of viral replication was not enough to suppress hepatocarcinogenesis during the treatment of lamivudine + adefovir to lamivudine-refractory patients, in contrast to the suppression of hepatocarcinogenesis by lamivudine monotherapy to naive patients without lamivudine-resistant HBV infection [8, 9]. HBV DNA is often integrated into host chromosome in liver tumor tissue, possibly causing chromosomal instability [28–31]. Previous studies reported that antiviral treatment (e.g., lamivudine, adefovir, entecavir, peg-interferon) also diminished the amount of intrahepatic covalently closed circular DNA (cccDNA) as an important intermediate in the life cycle of HBV [32, 33]. However, it is possible that any residual cccDNA in the hepatocytes may still have had integrative capacity at the HBV-DNA non-detectable state during lamivudine + adefovir, and that those may induce hepatocarcinogenesis. Further investigations should be performed whether

cccDNA could influence hepatocarcinogenesis. In this study, it is regrettable that the associations of nt substitutions with the development of HCC could not be presented. This reason is related to the small number of HCC patients, who might provide misleading results (e.g., possible type II error). Further studies should be also conducted to investigate nt substitutions, which might affect the development of HCC during lamivudine + adefovir.

The hepatocarcinogenesis in many patients of this study might have started before the suppression of HBV replication under adefovir, since carcinogenesis begins several months or even years before HCC diagnosis. HCC was diagnosed from 0.1 to 1.6 years in 7 of 12 patients, so the potential beneficial effect of viral suppression might be expected to be seen after the first 2 years of adefovir therapy and only in patients who achieve HBV DNA non-detectable. Further studies should be performed to evaluate the HCC risk in patients who have remained at least 1 year in remission under adefovir to evaluate beneficial effect of viral suppression.

Previous studies reported that interferon monotherapy and lamivudine monotherapy to naive patients for nucleoside analogues without lamivudine-resistant HBV infection suppressed HBV-related hepatocarcinogenesis [8, 9, 34, 35]. In conclusion, lamivudine-resistant patients treated with lamivudine + adefovir could achieve the excellent virological response and biochemical response, but the low hepatitis activity by suppression of viral replication was not enough to suppress hepatocarcinogenesis. Further understanding including viral predictors should facilitate the development of more effective therapeutic regimens to reduce risk of hepatocarcinogenesis.

Acknowledgment

This study was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

References

- 1 Wright TL, Lau JY: Clinical aspects of hepatitis B virus infection. *Lancet* 1993;342:1340-1344.
- 2 Ganem D, Prince AM: Hepatitis B virus infection - natural history and clinical consequences. *N Engl J Med* 2004;350:1118-1129.
- 3 Nevens F, Main J, Honkoop P, Tyrrell DL, Barber J, Sullivan MT, Fevery J, De Man RA, Thomas HC: Lamivudine therapy for chronic hepatitis B: a six-month randomized dose-ranging study. *Gastroenterology* 1997;113:1258-1263.
- 4 Lai CL, Chien RN, Leung NW, Chang TT, Guan R, Tai DI, Ng KY, Wu PC, Dent JC, Barber J, Stephenson SL, Gray DF: A one-year trial of lamivudine for chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *N Engl J Med* 1998;339:61-68.
- 5 Suzuki Y, Kumada H, Ikeda K, Chayama K, Arase Y, Saitoh S, Tsubota A, Kobayashi M, Koike M, Ogawa N, Tanikawa K: Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J Hepatol* 1999;30:743-748.
- 6 Lai CL, Leung N, Teo EK, Tong M, Wong F, Hann HW, Han S, Poynard T, Myers M, Chao G, Lloyd D, Brown NA: A 1-year trial of telbivudine, lamivudine, and the combination in patients with hepatitis B e antigen-positive chronic hepatitis B. *Gastroenterology* 2005;129:528-536.
- 7 Van Bömmel F, Zöllner B, Sarrazin C, Spengler U, Hüppe D, Möller B, Feucht HH, Wiedenmann B, Berg T: Tenofovir for patients with lamivudine-resistant hepatitis B virus (HBV) infection and high HBV DNA level during adefovir therapy. *Hepatology* 2006;44:318-325.
- 8 Liaw YF, Sung JY, Chow WC, Farrell G, Lee CZ, Yuen HY, Tanwandee T, Tao QM, Shue K, Keene ON, Dixon JS, Gray DF, Sabbat J: Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 2004;351:1521-1531.
- 9 Matsumoto A, Tanaka E, Rokuhara A, Kiyosawa K, Kumada H, Omata M, Okita K, Hayashi N, Okanoue T, Iino S, Tanikawa K: Efficacy of lamivudine for preventing hepatocellular carcinoma in chronic hepatitis B: a multicenter retrospective study of 2,795 patients. *Hepatol Res* 2005;32:173-184.
- 10 Allen MI, Deslauriers M, Andrews CW, Tipples GA, Walters KA, Tyrrell DL, Brown N, Condreay LD: Identification and characterization of mutations in hepatitis B virus resistant to lamivudine. Lamivudine Clinical Investigation Group. *Hepatology* 1998;27:1670-1677.
- 11 Delaney WE 4th, Yang H, Westland CE, Das K, Arnold E, Gibbs CS, Miller MD, Xiong S: The hepatitis B virus polymerase mutation rtV173L is selected during lamivudine therapy and enhances viral replication in vitro. *J Virol* 2003;77:11833-11841.
- 12 Sherman M, Yurdaydin C, Sollano J, Silva M, Liaw YF, Cianciara J, Boron-Kaczmarek A, Martin D, Goodman Z, Colonna R, Cross A, Denisky G, Kreter B, Hinds R: Entecavir for treatment of lamivudine-refractory, HBeAg-positive chronic hepatitis B. *Gastroenterology* 2006;130:2039-2049.
- 13 Rapti I, Dimou E, Mitsoula P, Hadziyannis SJ: Adding-on versus switching to adefovir therapy in lamivudine-resistant HBeAg-negative chronic hepatitis B. *Hepatology* 2007;45:307-313.
- 14 Hosaka T, Suzuki F, Suzuki Y, Saitoh S, Kobayashi M, Someya T, Sezaki H, Akuta N, Tsubota A, Arase Y, Ikeda K, Kumada H: Adefovir dipivoxil for treatment of breakthrough hepatitis caused by lamivudine-resistant mutants of hepatitis B virus. *Intervirology* 2004;47:362-369.
- 15 Tong SP, Li JS, Vitvitski L, Trepo C: Replication capacities of natural and artificial pre-core stop codon mutants of hepatitis B virus: relevance of pregenome encapsidation signal. *Virology* 1992;191:237-245.
- 16 Okamoto H, Tsuda F, Akahane Y, Sugai Y, Yoshida M, Moriya K, Tanaka T, Miyakawa Y, Mayumi M: Hepatitis B virus with mutations in the core promoter for an e antigen-negative phenotype in carriers with antibody to e antigen. *J Virol* 1994;68:8102-8110.

- 17 Laskus T, Rakela J, Tong MJ, Nowicki MJ, Mosley JW, Persing DH: Naturally occurring hepatitis B virus mutants with deletions in the core promoter region. *J Hepatol* 1994;20:837–841.
- 18 Laskus T, Rakela J, Nowicki MJ, Persing DH: Hepatitis B virus core promoter sequence analysis in fulminant and chronic hepatitis B. *Gastroenterology* 1995;109:1618–1623.
- 19 Takahashi K, Aoyama K, Ohno N, Iwata K, Akahane Y, Baba K, Yoshizawa H, Mishiro S: The precore/core promoter mutant (T1762/A1764) of hepatitis B virus: clinical significance and an easy method for detection. *J Gen Virol* 1995;76:3159–3164.
- 20 Lindh M, Horal P, Dhillon AP, Furuta Y, Norkrans G: Hepatitis B virus carriers without precore mutations in hepatitis B e antigen-negative stage show more severe liver damage. *Hepatology* 1996;24:494–501.
- 21 Erhardt A, Reineke U, Blondin D, Gerlich WH, Adams O, Heintges T, Niederau C, Häussinger D: Mutations of the core promoter and response to interferon treatment in chronic replicative hepatitis B. *Hepatology* 2000;31:716–725.
- 22 Usuda S, Okamoto H, Iwanari H, Baba K, Tsuda F, Miyakawa Y, Mayumi M: Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J Virol Methods* 1999;80:97–112.
- 23 Okamoto H, Tsuda F, Sakugawa H, Sastrosoewignjo RI, Imai M, Miyakawa Y, Mayumi M: Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J Gen Virol* 1988;69:2575–2583.
- 24 Kwok S, Higuchi R: Avoiding false positive with PCR. *Nature* 1989;339:237–238.
- 25 Desmet VJ, Gerber M, Hoofnagle JH, Manna M, Scheuer PJ: Classification of chronic hepatitis: diagnosis, grading and staging. *Hepatology* 1994;19:1513–1520.
- 26 Lampertico P, Vigano M, Manenti E, Iavarone M, Sablon E, Colombo M: Low resistance to adefovir combined with lamivudine: a 3-year study of 145 lamivudine-resistant hepatitis B patients. *Gastroenterology* 2007;133:1445–1451.
- 27 Kobayashi M, Ikeda K, Hosaka T, Sezaki H, Someya T, Akuta N, Suzuki F, Suzuki Y, Saitoh S, Arase Y, Miyakawa Y, Kumada H: Natural history of compensated cirrhosis in the Child-Pugh class A compared between 490 patients with hepatitis C and 167 with B virus infections. *J Med Virol* 2006;78:459–465.
- 28 Bréchet C, Pourcel C, Louise A, Rain B, Tiollais P: Presence of integrated hepatitis B virus DNA sequences in cellular DNA of human hepatocellular carcinoma. *Nature* 1980;286:533–535.
- 29 Bréchet C, Hadchouel M, Scotto J, Fonck M, Potet F, Vyas GN, Tiollais P: State of hepatitis B virus DNA in hepatocytes of patients with hepatitis B surface antigen-positive and -negative liver diseases. *Proc Natl Acad Sci USA* 1981;78:3906–3910.
- 30 Shafritz DA, Shouval D, Sherman HI, Hadziyannis SJ, Kew MC: Integration of hepatitis B virus DNA into the genome of liver cells in chronic liver disease and hepatocellular carcinoma. Studies in percutaneous liver biopsies and post-mortem tissue specimens. *N Engl J Med* 1981;305:1067–1073.
- 31 Wong DK, Yuen MF, Poon RT, Yuen JC, Fung J, Lai CL: Quantification of hepatitis B virus covalently closed circular DNA in patients with hepatocellular carcinoma. *J Hepatol* 2006;45:553–559.
- 32 Wursthorn K, Lugehetmann M, Dandri M, Volz T, Buggisch P, Zollner B, Longerich T, Schirmacher P, Metzler F, Zankel M, Fischer C, Currie G, Brosgart C, Petersen J: Peginterferon- α_{2b} plus adefovir induce strong cccDNA decline and HBsAg reduction in patients with chronic hepatitis B. *Hepatology* 2006;44:675–684.
- 33 Wong DK, Yuen MF, Ngai VW, Fung J, Lai CL: One-year entecavir and lamivudine therapy results in reduction of hepatitis B virus intrahepatic covalently closed circular DNA levels. *Antivir Ther* 2006;11:909–916.
- 34 Lin SM, Sheen IS, Chien RN, Chu CM, Liaw YF: Long-term beneficial effect of interferon therapy in patients with chronic hepatitis B virus infection. *Hepatology* 1999;29:971–975.
- 35 Lin SM, Yu ML, Lee CM, Chien RN, Sheen IS, Chu CM, Liaw YF: Interferon therapy in HBeAg-positive chronic hepatitis reduces progression to cirrhosis and hepatocellular carcinoma. *J Hepatol* 2007;46:45–52.

Amino Acid Substitutions in the Hepatitis C Virus Core Region of Genotype 1b Are the Important Predictor of Severe Insulin Resistance in Patients Without Cirrhosis and Diabetes Mellitus

Norio Akuta,^{1*} Fumitaka Suzuki,¹ Miharuru Hirakawa,¹ Yusuke Kawamura,¹ Hiromi Yatsuji,¹ Hitomi Sezaki,¹ Yoshiyuki Suzuki,¹ Tetsuya Hosaka,¹ Masahiro Kobayashi,¹ Mariko Kobayashi,² Satoshi Saitoh,¹ Yasuji Arase,¹ Kenji Ikeda,¹ and Hiromitsu Kumada¹

¹Department of Hepatology, Toranomon Hospital, Tokyo, Japan

²Liver Research Laboratory, Toranomon Hospital, Tokyo, Japan

Previous studies provided a direct experimental evidence for the contribution of HCV core protein in the development of insulin resistance (IR), but the clinical impact of HCV core region on IR is still not clear. The present study evaluated the impact of Amino acid (aa) substitutions of HCV-1b core region on IR in 123 Japanese patients infected with HCV-1b without cirrhosis and diabetes mellitus, and investigated the treatment efficacy of 48-week pegylated interferon (PEG-IFN) plus ribavirin (RBV) according to HOMA-IR values. Patients with IR (HOMA-IR ≥ 2.5) and severe IR (HOMA-IR ≥ 3.5) were present in 51.2% and 27.6%, respectively. Multivariate analysis identified body mass index (≥ 25 kg/m²) and hepatocyte steatosis ($\geq 5\%$) as significant determinants of IR. Furthermore, multivariate analysis identified hepatocyte steatosis ($\geq 5\%$), aa substitutions of the core region (Gln70 (His70) and/or Met91), and age (≥ 55 years) as significant determinants of severe IR. Especially, significantly lower proportions of patients with Gln70 (His70) and/or Met91 were noted among those without severe IR (59.6%) than those with severe IR (82.4%). The rates of sustained virological response in patients with IR (50.0%) were not significantly different from those without IR (52.9%). Furthermore, the rates of non-virological response in patients with IR (28.9%) were not significantly also different from those without IR (20.6%). In conclusion, the present study indicated that substitutions of HCV-1b core region were the important predictor of severe IR in patients without cirrhosis and diabetes mellitus, but HOMA-IR values might be not useful as predictors of 48-week PEG-IFN plus RBV therapy. *J. Med. Virol.* 81:1032–1039, 2009. © 2009 Wiley-Liss, Inc.

KEY WORDS: HCV; core region; genotype; HOMA-IR; hepatocyte steatosis; cirrhosis; diabetes mellitus

INTRODUCTION

Hepatitis C virus (HCV) usually causes chronic infection that can result in chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma (HCC) [Dusheiko, 1998; Ikeda et al., 1998; Niederau et al., 1998; Kenny-Walsh, 1999; Akuta et al., 2001]. Furthermore, HCV infection also affects an increased risk of diabetes mellitus [Allison et al., 1994; Caronia et al., 1999; Mason et al., 1999; Mehta et al., 2000, 2003; Zein et al., 2000, 2005; Antonelli et al., 2005] or insulin resistance (IR) [Hickman et al., 2003; Hui et al., 2003; Lecube et al., 2004, 2006]. IR and glucose metabolism impairment are associated with liver necroinflammation [Hui et al., 2003], hepatocyte steatosis [Fartoux et al., 2005; Cammà et al., 2006; Conjeevaram et al., 2007], cirrhosis [Petrides et al., 1994], and HCC [El-Serag et al., 2001; Lai et al., 2006; Veldt et al., 2008]. Especially, in patients infected with HCV genotype 1 (HCV-1), significant fibrosis is associated with IR independent from hepatocyte steatosis [Moucari et al., 2008; Petta et al., 2008].

Previous studies reported that HCV core protein induced HCC and IR in transgenic mice, and provided

Grant sponsor: Ministry of Health, Labor and Welfare, Japan (partly supported).

*Correspondence to: Norio Akuta, MD, Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo 105-0001, Japan. E-mail: akuta-gi@umin.ac.jp

Accepted 5 February 2009

DOI 10.1002/jmv.21473

Published online in Wiley InterScience (www.interscience.wiley.com)

a direct experimental evidence for the contribution of HCV core protein in the development of HCC and IR in human HCV infection [Moriya et al., 1998; Shintani et al., 2004]. Amino acid (aa) substitutions at position 70 and/or 91 in the HCV core region of genotype 1b (HCV-1b core region) were predictors of poor virological response to 48-week pegylated interferon (PEG-IFN) plus ribavirin (RBV) combination therapy [Akuta et al., 2005, 2006, 2007a,b,c; Donlin et al., 2007; Okanou et al., 2008], and also risk factors for hepatocarcinogenesis [Akuta et al., 2007d, 2008]. Thus, previous reports supported the oncogenic potential of the HCV core region and clinically linked substitutions of aa 70 and/or 91 in HCV-1b core region to HCC [Akuta et al., 2007d, 2008], but the clinical impact of HCV-1b core region on IR is still not clear. IR develops type 2 diabetes mellitus as its major late feature, and is also associated with advanced fibrosis [Petrides et al., 1994; Petta et al., 2008]. Hence, the biological mechanisms underlying the association between HCV core region and IR are probably multifactorial, and study based on patients without diabetes mellitus and cirrhosis, that might affect IR, should be performed to investigate whether HCV core region might affect IR clinically.

Previous reports showed that IR might be predictors of poor virological response to PEG-IFN plus RBV combination therapy [D'Souza et al., 2005; Romero-Gómez et al., 2005]. Chu et al. [2008] reported that IR was a major determinant of sustained virological response (SVR) in HCV-1 patients receiving 24-week PEG-IFN plus RBV. However, to our knowledge, there is little evidence that IR affects treatment efficacy of HCV-1b patients receiving 48-week PEG-IFN plus RBV combination therapy.

The aims of the present study conducted in Japanese patients infected with HCV-1b without cirrhosis and diabetes mellitus, were the following. (1) To evaluate the HOMA-IR values of patients infected with HCV-1b. (2) To identify the impact of aa substitutions in the core region on IR in such patients, and determine the factors associated with IR, and (3) to investigate the treatment efficacy of 48-week PEG-IFN plus RBV combination therapy according to HOMA-IR values.

PATIENTS AND METHODS

Study Population

At Toranomon Hospital, Tokyo, Japan, 221 HCV-infected Japanese patients were consecutively recruited into the study protocol of the combination therapy with PEG-IFN α -2b plus RBV between December of 2001 and June of 2005. Among these, 123 patients were selected in the present retrospective study based on the following criteria. (1) Negativity for hepatitis B surface antigen (radioimmunoassay, Dainabot, Tokyo, Japan), positivity for anti-HCV (third-generation enzyme immunoassay, Chiron Corp, Emerville, CA), and positivity for HCV RNA qualitative analysis with PCR (Amplicor, Roche Diagnostics, Mannheim, Germany). (2) They were infected with HCV-1b alone. (3) HOMA-IR values

and substitutions of aa 70 and 91 in the HCV core region were determined at the commencement of treatment. (4) They were free of cirrhosis and hepatocellular carcinoma, based on biopsy examination, laboratory tests, and imaging studies at baseline. (5) None had diabetes mellitus. (6) None was an alcoholic; lifetime cumulative alcohol intake was <500 kg (mild to moderate alcohol intake). (7) All were free of coinfection with human immunodeficiency virus. (8) None had other forms of hepatitis, such as hemochromatosis, Wilson disease, primary biliary cirrhosis, alcoholic liver disease, and autoimmune liver disease. (9) Each signed a consent form of the study protocol that had been approved by the human ethics review committee. Table I summarizes the profiles and laboratory data of the 123 patients at the commencement of treatment. They included 71 males and 52 females, aged 20–70 years (median, 55 years). The treatment efficacy was evaluated by HCV-RNA positive based on qualitative PCR analysis at the end of treatment (non-virological response; NVR), and by HCV-RNA negative based on qualitative PCR analysis at 24 weeks after the completion of therapy (SVR).

Laboratory Tests

Blood samples were obtained at least once every month before, during, and after treatment, and were analyzed for alanine aminotransferase (ALT) and HCV-RNA levels. The serum samples were frozen at -80°C within 4 hr of collection and thawed at the time of measurement. HCV genotype was determined by PCR using a mixed primer set derived from the nucleotide sequences of NS5 region [Chayama et al., 1993]. HCV-RNA levels were measured by quantitative PCR (AMPLICOR GT HCV Monitor v2.0 using the 10-fold dilution method, Roche Molecular Systems, Inc.) at least once every month before, during, and after therapy. The dynamic range of the assay was 5.0×10^3 to 5.0×10^6 IU/ml. Samples collected during and after therapy that showed undetectable levels of HCV-RNA ($<5.0 \times 10^3$ IU/ml) were also checked by qualitative PCR (AMPLICOR HCV v2.0, Roche Molecular Systems, Inc.), which has a higher sensitivity than quantitative analysis, and the results were expressed as positive or negative. The lower limit of the assay was 50 IU/ml.

Histopathological Examination of Liver Biopsies

Liver biopsy specimens were obtained percutaneously or at peritoneoscopy using a modified Vim Silverman needle with an internal diameter of 2 mm (Tohoku University style, Kakinuma Factory, Tokyo, Japan). The biopsy material was fixed in 10% formalin, and stained with hematoxylin and eosin, Masson's trichrome, silver impregnation, and periodic acid-Schiff after diastase digestion. All specimens for examination contained six or more portal areas. Histopathological diagnosis was confirmed by an experienced liver pathologist (H.K.) who was blinded to the clinical data. Chronic hepatitis and liver cirrhosis were diagnosed

TABLE I. Profile and Laboratory Data of 123 Patients Infected With HCV Genotype 1b

Demographic data	
Number of patients	123
Sex (M/F)	71/52
Age (years)*	55 (20–70)
History of blood transfusion	41 (33.3%)
Family history of liver disease	37 (30.1%)
Body mass index (kg/m ²)*	23.6 (17.6–32.0)
Laboratory data*	
Serum aspartate aminotransferase (IU/L)	59 (17–266)
Serum alanine aminotransferase (IU/L)	81 (25–504)
Serum albumin (g/dl)	3.8 (3.1–4.5)
Gamma-glutamyl transpeptidase (IU/L)	50 (15–393)
Leukocytes (/mm ³)	4,800 (2,300–8,800)
Hemoglobin (g/dl)	14.4 (10.6–17.6)
Platelet count ($\times 10^4/\text{mm}^3$)	16.8 (7.5–27.7)
Indocyanine green retention rate at 15 min (%)	15 (4–41)
Serum iron ($\mu\text{g}/\text{dl}$)	138 (18–290)
Serum ferritin ($\mu\text{g}/\text{L}$)	130 (<10–711)
Creatinine clearance (ml/min)	99 (46–146)
Level of viremia (KIU/ml)	1,900 (23–>5,000)
Alpha-fetoprotein ($\mu\text{g}/\text{L}$)	6 (2–161)
Total cholesterol (mg/dl)	166 (96–294)
High-density lipoprotein cholesterol (mg/dl)	46 (10–83)
Low-density lipoprotein cholesterol (mg/dl)	101 (53–207)
Triglycerides (mg/dl)	95 (33–362)
Uric acid (mg/dl)	5.5 (2.3–9.4)
Fasting plasma glucose (mg/dl)	94 (62–120)
Fasting insulin ($\mu\text{U}/\text{ml}$)	10.5 (0.4–55.5)
HOMA-IR	2.6 (0.1–12.5)
Treatment*	
PEG-IFN α -2b dose ($\mu\text{g}/\text{kg}$)	1.4 (0.7–1.9)
Ribavirin dose (mg/kg)	11.0 (3.7–14.2)
Histological findings	
Stage of fibrosis (F1/F2/F3/ND)	53/32/20/18
Hepatocyte steatosis (<5% (Absent)/ \geq 5% (Present)/ND)	38/64/21
Amino acid substitutions in the HCV	
Core aa 70 (arginine/glutamine (histidine))	69/54
Core aa 91 (leucine/methionine)	71/52
ISDR of NS5A (wild-type/mutant-type/ND)	94/22/7

HOMA-IR, homeostasis model for assessment of insulin resistance; ND, not determined.

Data are number and percentages of patients, except those denoted by *, which represent the median (range) values.

based on histological assessment according to the scoring system of Desmet et al. [1994]. Hepatocyte steatosis was assessed as the percentage of hepatocytes containing fat droplet, and subjects were considered to have steatosis in the presence of fat droplets in \geq 5% of hepatocytes.

Diagnosis of Liver Cirrhosis, Insulin Resistance, and Diabetes Mellitus

Liver cirrhosis was diagnosed based on the presence of markedly irregular surface with nodular formation in the liver, evident on peritoneoscopy, histological assessment according to the scoring system of Desmet et al. [1994], or on computed tomography or ultrasonography. Ascites, edema, and esophageal varicosities, facilitated the diagnosis when present.

The diagnosis of type 2 diabetes was based on the revised criteria of the American Diabetes Association using a value of fasting plasma glucose of \geq 126 mg/dl on at least two occasions [American Diabetes Association, 2000]. IR was assessed by the Homeostasis Model for Assessment of Insulin Resistance (HOMA-IR) method

[Matthews et al., 1985], using the following equation: $\text{HOMA-IR} = \text{fasting plasma glucose (mg/dl)} \times \text{fasting insulin } (\mu\text{U/ml}) / 405$. HOMA-IR values of \geq 2.5, and \geq 3.5 were evaluated as IR, and severe IR, respectively.

Detection of Amino Acid Substitutions in Core Region and NS5A Region

With the use of HCV-J (accession no. D90208) as a reference [Kato et al., 1990], the sequence of 1–191 aa in the core protein of genotype 1b was determined and then compared with the consensus sequence constructed on 50 clinical samples to detect substitutions at aa 70 of arginine (Arg70) or glutamine/histidine (Gln70/His70) and aa 91 of leucine (Leu91) or methionine (Met91) [Akuta et al., 2005]. The sequence of 2209–2248 aa in the NS5A of genotype 1b (IFN-sensitivity determining region [ISDR]) reported by Enomoto et al. [1995, 1996] was also determined, and the numbers of aa substitutions in ISDR were defined as wild-type (\leq 1) or mutant-type (\geq 2).

In the present study, aa substitutions of the core region and NS5A-ISDR were analyzed by direct

sequencing [Enomoto et al., 1995, 1996; Akuta et al., 2005]. HCV RNA was extracted from serum samples at the start of treatment and reverse transcribed with random primer and MMLV reverse transcriptase (Takara Syuzo, Tokyo, Japan). Nucleic acids were amplified by PCR using the following primers: (a) Nucleotide sequences of the core region: The first-round PCR was performed with CC11 (sense, 5'-GCC ATA GTG GTC TGC GGA AC-3') and e14 (antisense, 5'-GGA GCA GTC CTT CGT GAC ATG-3') primers, and the second-round PCR with CC9 (sense, 5'-GCT AGC CGA GTA GTG TT-3') and e14 (antisense) primers. (b) Nucleotide sequences of NS5A-ISDR: The first-round PCR was performed with ISDR1 (sense, 5'-ATG CCC ATG CCA GGT TCC AG-3') and ISDR2 (antisense, 5'-AGC TCC GCC AAG GCA GAA GA-3') primers, and the second-round PCR with ISDR3 (sense, 5'-ACC GGA TGT GGC AGT GCT CA-3') and ISDR4 (antisense, 5'-GTA ATC CGG GCG TGC CCA TA-3') primers. ([a]; hemi-nested PCR. [b]; nested PCR). All samples were initially denatured at 95°C for 15 min. The 35 cycles of amplification were set as follows: denaturation for 1 min at 94°C, annealing of primers for 2 min at 55°C, and extension for 3 min at 72°C with an additional 7 min for extension. Then 1 μ l of the first PCR product was transferred to the second PCR reaction. Other conditions for the second PCR were the same as the first PCR, except that the second PCR primers were used instead of the first PCR primers. The amplified PCR products were purified by the QIA quick PCR purification kit (Qiagen, Tokyo, Japan) after agarose gel electrophoresis and then used for direct sequencing. Dideoxynucleotide termination sequencing was performed with the Big Dye Deoxy Terminator Cycle Sequencing kit (Perkin-Elmer, Tokyo, Japan).

Statistical Analysis

Non-parametric tests were used to compare variables between groups, including the Mann-Whitney *U* test, chi-squared test, and Fisher's exact probability test. Univariate and multivariate logistic regression analyses were used to determine the independent predictive factors of IR. The odds ratios and 95% confidence intervals (95%CI) were also calculated. All *P*-values <0.05 by the two-tailed test were considered significant. Variables that achieved statistical significance ($P < 0.05$) or marginal significance ($P < 0.10$) on univariate analysis were entered into multiple logistic regression analysis to identify significant independent factors. Potential predictive factors of IR included the following pretreatment variables: sex, age, history of blood transfusion, familial history of liver disease, body mass index, aspartate aminotransferase (AST), ALT, albumin, gamma-glutamyl transpeptidase (γ GTP), leukocyte count, hemoglobin, platelet, count, indocyanine green retention rate at 15 min (ICG R15), serum iron, serum ferritin, creatinine clearance, level of viremia, alfa-fetoprotein, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol,

triglycerides, uric acid, fasting plasma glucose, fasting insulin, HOMA-IR values, stage of fibrosis, hepatocyte steatosis, PEG-IFN dose/body weight, RBV dose/body weight, and aa substitution in the core and ISDR of NS5A. Statistical analyses were performed using the SPSS software (SPSS, Inc., Chicago, IL).

RESULTS

HOMA-IR Values of Patients Infected With HCV-1b Without Cirrhosis and Diabetes Mellitus

As a whole, 16.3% (20 of 123 patients), 32.5% (40 of 123), 23.6% (29 of 123), and 27.6% (34 of 123) indicated HOMA-IR values of ≤ 1.4 , 1.5–2.4, 2.5–3.4, and ≥ 3.5 , respectively (Fig. 1). Thus, patients with IR (HOMA-IR ≥ 2.5) were present in 51.2% (63 of 123), and exceeded 50%. Furthermore, patients with severe IR (HOMA-IR ≥ 3.5) were present in 27.6%. These results show that patients, infected with HCV-1b without cirrhosis and diabetes mellitus, might indicate IR frequently.

Factors Associated With Insulin Resistance in Univariate and Multivariate Analyses

The whole population sample of 123 patients were analyzed to determine factors that could be associated with IR. IR (HOMA-IR ≥ 2.5) was detected in 63 of 123 (51.2%) patients. Univariate analysis identified six parameters that tended to or significantly influenced IR. These included age (≥ 55 years, $P = 0.072$), body mass index (≥ 25 kg/m², $P < 0.001$), serum ferritin (≥ 200 μ g/L, $P = 0.071$), family history of liver disease (Absent, $P = 0.077$), hepatocyte steatosis (Present ($\geq 5\%$), $P = 0.002$), and aa substitutions of the core region (Gln70 (His70) and/or Met91, $P = 0.092$). Multivariate analysis identified two parameters that independently influenced IR, including body mass index (≥ 25 kg/m², $P = 0.001$) and hepatocyte steatosis (Present ($\geq 5\%$), $P = 0.028$).

Severe IR (HOMA-IR ≥ 3.5) was detected in 34 of 123 (27.6%) patients. Univariate analysis identified five




Fig. 1. HOMA-IR values of patients infected with HCV genotype 1b without cirrhosis and diabetes mellitus. As a whole, 16.3%, 32.5%, 23.6%, and 27.6% indicated HOMA-IR values of ≤ 1.4 , 1.5–2.4, 2.5–3.4, and ≥ 3.5 , respectively. These results show that patients, infected with HCV genotype 1b without cirrhosis and diabetes mellitus, might indicate IR frequently.

TABLE II. Factors Associated With Severe IR (HOMA-IR ≥ 3.5) in Patients Infected With HCV Genotype 1b, Identified by Multivariate Analysis

Factor	Category	Odds ratio (95% CI)	P
Hepatocyte steatosis	1: Absent (<5%)	1	0.021
	2: Present ($\geq 5\%$)	4.170 (1.235–14.08)	
Substitution of aa 70 and 91	1: Arg70 and Leu91	1	0.021
	2: Gln70 (His70) and/or Met91	3.654 (1.215–10.99)	
Age (years)	1: <55	1	0.037
	2: ≥ 55	3.015 (1.071–8.488)	

parameters that tended to or significantly influenced severe IR. These included age (≥ 55 years, $P=0.015$), body mass index (≥ 25 kg/m², $P=0.025$), hepatocyte steatosis (Present ($\geq 5\%$), $P=0.003$), triglycerides (≥ 100 mg/dl, $P=0.060$), and aa substitutions of the core region (Gln70 (His70) and/or Met91, $P=0.020$). Multivariate analysis identified three parameters that independently influenced severe IR, including hepatocyte steatosis (Present ($\geq 5\%$), $P=0.021$), aa substitutions of the core region (Gln70 (His70) and/or Met91, $P=0.021$), and age (≥ 55 years, $P=0.037$) (Table II).

aa Substitutions of Core Region and HOMA-IR Values

The entire population sample was also analyzed to determine the relationship between aa substitutions of the core region and HOMA-IR values. HOMA-IR values of 81 patients with Gln70 (His70) and/or Met91 (median; 2.9) indicated the higher levels than those of 42 patients with Arg70 and Leu91 (median; 2.3), significantly ($P=0.022$) (Fig. 2).

Furthermore, the proportions of patients with Gln70 (His70) and/or Met91 among those with HOMA-IR values of ≤ 1.4 , 1.5–2.4, 2.5–3.4, 3.5–3.9, and ≥ 4.0 were 60.0% (12 of 20 patients), 57.5% (23 of 40), 62.1% (18 of 29), 83.3% (5 of 6), and 82.1% (23 of 28), respectively




Fig. 2. aa substitutions of HCV core region and HOMA-IR values. HOMA-IR values of 81 patients with Gln70 (His70) and/or Met91 indicated the higher levels than those of 42 patients with Arg70 and Leu91, significantly ($P=0.022$).

(Fig. 3). Thus, the higher the proportion of patients with Gln70 (His70) and/or Met91, the higher HOMA-IR values, and significantly lower proportions of patients with Gln70 (His70) and/or Met91 were noted among those without severe IR (59.6% (53 of 89)) than those with severe IR (82.4% (28 of 34)) ($P=0.020$).

Treatment Efficacy of PEG-IFN Plus RBV Combination Therapy According to HOMA-IR Values

Of the 123 patients, 72 could be evaluated as 48-week regimen of PEG-IFN plus RBV combination therapy. Seventy-two patients received PEG-IFN α -2b combination therapy at a median dose of 1.5 μ g/kg (range, 0.8–1.8 μ g/kg) subcutaneously each week plus oral RBV at a median dose of 11.3 mg/kg (range, 9.7–14.2 mg/kg) daily for 48 weeks. 51.4% (37 of 72 patients) could achieve SVR, and 25.0% (18 of 72) had NVR.

In each groups with IR or without IR, SVR was achieved by 19 of 38 patients (50.0%) and 18 of 34 (52.9%), respectively. The proportions of SVR in group with IR was not significantly different from those in group without IR. Furthermore, in each groups with IR or without IR, NVR was identified in 11 of 38 patients (28.9%) and 7 of 34 (20.6%), respectively. The proportions of NVR in group with IR was not significantly different from those in group without IR.




Fig. 3. The proportions of patients with Gln70 (His70) and/or Met91 and HOMA-IR values. Higher frequencies of Gln70 (His70) and/or Met91 correlated with higher HOMA-IR values. Significantly lower proportions of patients with Gln70 (His70) and/or Met91 were noted among those without severe IR (HOMA-IR <3.5) (59.6%) than those with severe IR (HOMA-IR ≥ 3.5) (82.4%) ($P=0.020$).