

Figure 6. A working hypothesis to explain how augmented level of AREG in BMMC of RA patients may lead to hyperproliferation of synovial cells. Putative macrophages with enhanced expression of AREG precursor (ProAR) may approach to the synovial cells of the joint through blood flow, where they release AREG and activate the EGFR signaling pathway of synovial cells. Since Ets-binding site (ETS-1) of the proximal promoter of the synoviolin gene is one of the downstream targets of ERK pathway, the enhanced activation of EGFR signaling may directly activate the expression of synoviolin gene. The enhanced level of synoviolin activates the ERAD system, which may lead to hyperproliferation of synovial cells.

Overexpression of AREG has been linked to psoriasis in mice and humans.^{49,50} Psoriasis is characterized by the hyperproliferation of keratinocytes and the loss of epidermal barrier function that leads to the infiltration of inflammatory cells into the epidermis and dermis.⁵¹ AREG is also upregulated in a synoviocyte cell line derived from an RA patient in which the wild type and a dominant negative form of the orphan nuclear receptor Nurr1 were overexpressed.⁵² Interestingly, AREG overexpression in the basal epidermis of transgenic mice induces a phenotype that is associated with synovial membrane inflammation.⁴⁹ Moreover, we showed previously that AREG expression is also enhanced in the

PBMC of SLE and idiopathic thrombocytopenic purpura patients,²⁶ which suggests that AREG overexpression may also be associated with other autoimmune diseases. Notably, metalloprotease-mediated AREG shedding and the subsequent activation of EGFR appears to play a critical role in the secretion of IL-8 by the human airway epithelium-like NCI-H292 cells that is induced by tumor necrosis factor- α (TNF- α), a potent multifunctional cytokine that plays a central role in the pathogenesis of many inflammatory diseases like RA.⁵³ Since TNF- α -induced IL-8 secretion was completely inhibited by the neutralizing antibody against AREG,⁵³ this antibody could constitute a novel therapeutic tool for RA. Taken

together, we propose that enhanced expression of AREG in BMMC and PMBC may play a pivotal role in the pathogenesis of RA.

Acknowledgements: We thank the patients and healthy volunteers who participated in this study. We also thank Ms Tomoko Motoyama, Ms Kumiko Ikeue, Ms Maki Masuda, and Ms Yuki Hamada for technical assistance and Dr Patrick Hughes for critically reading the manuscript. This work was primarily supported by a grant-in-aid from the Health Science Research grant from the Ministry of Health and Welfare of Japan. This work was also supported in part by Innovation Plaza Osaka of the Japan Science and Technology Agency (JST), and by grants-in-aid for Scientific Research on Priority Areas Applied Genomics, Scientific Research (S), Exploratory Research, and the Science and Technology Incubation Program in Advanced Regions, from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Supplementary Data: Supplementary data are available online at www.dnaresearch.oxfordjournals.org.

References

- Oertelt, S., Selmi, C., Invernizzi, P., Podda, M., and Gershwin, M. E. 2005, Genes and goals: an approach to microarray analysis in autoimmunity, *Autoimmune Rev.*, **4**, 414-422.
- Glocker, M. O., Guthke, R., Kekow, J., and Thiesen, H. J. 2006, Rheumatoid arthritis, a complex multifactorial disease: on the way toward individualized medicine, *Med. Res. Rev.*, **26**, 63-87.
- van der Pouw Kraan, T. C., van Gaalen, F. A., Huizinga, T. W., Pieterman, E., Breedveld, F. C., and Verweij, C. L. 2003, Discovery of distinctive gene expression profiles in rheumatoid synovium using cDNA microarray technology: evidence for the existence of multiple pathways of tissue destruction and repair, *Genes Immun.*, **4**, 187-196.
- Ruschpler, P., Lorenz, P., Eichler, W., et al. 2003, High CXCR3 expression in synovial mast cells associated with CXCL9 and CXCL10 expression in inflammatory synovial tissues of patients with rheumatoid arthritis, *Arthritis Res. Ther.*, **5**, 241-252.
- Devauchelle, V., Marion, S., Cagnard, N., et al. 2004, DNA microarray allows molecular profiling of rheumatoid arthritis and identification of pathophysiological targets, *Genes Immun.*, **5**, 597-608.
- Cagnard, N., Letourneur, F., Essabani, A., et al. 2005, Interleukin-32, CCL2, PF4F1 and GFD10 are the only cytokine/chemokine genes differentially expressed by in vitro cultured rheumatoid and osteoarthritis fibroblast-like synoviocytes, *Eur. Cytokine Netw.*, **16**, 289-292.
- Lindberg, J., af Klint, E., Ulfgren, A. K., et al. 2006, Variability in synovial inflammation in rheumatoid arthritis investigated by microarray technology, *Arthritis Res. Ther.*, **8**, R47.
- Finis, K., Sultmann, H., Ruschhaupt, M., et al. 2006, Analysis of pigmented villonodular synovitis with genome-wide complementary DNA microarray and tissue array technology reveals insight into potential novel therapeutic approaches, *Arthritis Rheum.*, **54**, 1009-1019.
- Bovin, L. F., Rieneck, K., Workman, C., et al. 2004, Blood cell gene expression profiling in rheumatoid arthritis: discriminative genes and effect of rheumatoid factor, *Immunol. Lett.*, **93**, 217-226.
- Amano, T., Yamasaki, S., Yagishita, N., et al. 2003, Synoviolin/Hrd1, an E3 ubiquitin ligase, as a novel pathogenic factor for arthropathy, *Genes Dev.*, **17**, 2436-2449.
- Yagishita, N., Ohneda, K., Amano, T., et al. 2005, Essential role of synoviolin in embryogenesis, *J. Biol. Chem.*, **280**, 7909-7916.
- Yamasaki, S., Yagishita, N., Tsuchimochi, K., Nishioka, K., and Nakajima, T. 2005, Rheumatoid arthritis as a hyper-endoplasmic reticulum-associated degradation disease, *Arthritis Res. Ther.*, **7**, 181-186.
- Ochi, T., Hakomori, S., Adachi, M., et al. 1988, The presence of a myeloid cell population showing strong reactivity with monoclonal antibody directed to difucosyl type 2 chain in epiphyseal bone marrow adjacent to joints affected with rheumatoid arthritis (RA) and its absence in the corresponding normal and non-RA bone marrow, *J. Rheumatol.*, **15**, 1609-1615.
- Hayashida, K., Ochi, T., Fujimoto, M., et al. 1992, Bone marrow changes in adjuvant-induced and collagen-induced arthritis, *Arthritis Rheum.*, **35**, 241-245.
- Jongen-Lavrencic, M., Peeters, H. R., Wognum, A., Vreugdenhil, G., Breedveld, F. C., and Swaak, A. J. 1997, Elevated levels of inflammatory cytokines in bone marrow of patients with rheumatoid arthritis and anemia of chronic disease, *J. Rheumatol.*, **24**, 1504-1509.
- Hirohata, S., Yanagida, T., Nampei, A., et al. 2004, Enhanced generation of endothelial cells from CD34+ cells of the bone marrow in rheumatoid arthritis: possible role in synovial neovascularization, *Arthritis Rheum.*, **50**, 3888-3896.
- Nakamura-Kikuoka, S., Takahi, K., Tsuboi, H., et al. 2006, Limited VH gene usage in B-cell clones established with nurse-like cells from patients with rheumatoid arthritis, *Rheumatology*, **45**, 549-557.
- Samuels, J., Ng, Y. S., Coupillaud, C., Paget, D., and Meffre, E. 2005, Human B cell tolerance and its failure in rheumatoid arthritis, *Ann. N. Y. Acad. Sci.*, **1062**, 116-126.
- Wardemann, H., Yurasov, S., Schaefer, A., Young, J. W., Meffre, E., and Nussenzweig, M. C. 2003, Predominant autoantibody production by early human B cell precursors, *Science*, **301**, 1374-1377.
- Bugatti, S., Caporali, R., Manzo, A., Vitolo, B., Pitzalis, C., and Montecucco, C. 2005, Involvement of subchondral bone marrow in rheumatoid arthritis: lymphoid neogenesis and in situ relationship to subchondral bone marrow osteoclast recruitment, *Arthritis Rheum.*, **52**, 3448-3459.
- Jimenez-Boj, E., Redlich, K., Turk, B., et al. 2005, Interaction between synovial inflammatory tissue and bone marrow in rheumatoid arthritis, *J. Immunol.*, **175**, 2579-2588.
- Fujii, T., Tamura, K., Masai, K., Tanaka, H., Nishimune, Y., and Nojima, H. 2002, Use of stepwise

- subtraction to comprehensively isolate mouse genes whose transcription is up-regulated during spermiogenesis, *EMBO Rep.*, **3**, 367-372.
23. Arnett, F. C., Edworthy, S. M., Bloch, D. A., et al. 1988, The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis, *Arthritis Rheum.*, **31**, 315-324.
 24. Altman, R. D. 1991, Criteria for classification of clinical osteoarthritis, *J. Rheumatol. Suppl.*, **27**, 10-12.
 25. Kobori, M., Ikeda, Y., Nara, H., et al. 1998, Large scale isolation of osteoclast-specific genes by an improved method involving the preparation of a subtracted cDNA library, *Genes Cells*, **3**, 459-475.
 26. Ishii, T., Onda, H., Tanigawa, A., et al. 2006, Isolation and expression profiling of genes upregulated in the peripheral blood cells of systemic lupus erythematosus patients, *DNA Res.*, **112**, 1-11.
 27. Shoyab, M., McDonald, V. L., Bradley, J. G., and Todaro, G. J. 1988, Amphiregulin: a bifunctional growth-modulating glycoprotein produced by the phorbol 12-myristate 13-acetate-treated human breast adenocarcinoma cell line MCF-7, *Proc. Natl Acad. Sci. USA*, **85**, 6528-6532.
 28. Fischer, G. and Aumuller, T. 2003, Regulation of peptide bond cis/trans isomerization by enzyme catalysis and its implication in physiological processes, *Rev. Physiol. Biochem. Pharmacol.*, **148**, 105-150.
 29. Russell, L. and Forsdyke, D. R. 1991, A human putative lymphocyte G0/G1 switch gene containing a CpG-rich island encodes a small basic protein with the potential to be phosphorylated, *DNA Cell Biol.*, **10**, 581-591.
 30. Santiago, B., Baleux, F., Palao, G., et al. 2006, CXCL12 is displayed by rheumatoid endothelial cells through its basic amino-terminal motif on heparan sulfate proteoglycans, *Arthritis Res. Ther.*, **8**, R43.
 31. Ahn, K. S. and Aggarwal, B. B. 2005, Transcription factor NF-kappaB: a sensor for smoke and stress signals, *Ann. N. Y. Acad. Sci.*, **1056**, 218-233.
 32. Shin, H. S., Lee, H. J., Nishida, M., et al. 2003, Betacellulin and amphiregulin induce upregulation of cyclin D1 and DNA synthesis activity through differential signaling pathways in vascular smooth muscle cells, *Circ. Res.*, **93**, 302-310.
 33. Lee, A. S. 2001, The glucose-regulated proteins: stress induction and clinical applications, *Trends Biochem. Sci.*, **26**, 504-510.
 34. Lee, D. C., Sunnarborg, S. W., Hinkle, C. L., et al. 2003, TACE/ADAM17 processing of EGFR ligands indicates a role as a physiological convertase, *Ann. N. Y. Acad. Sci.*, **995**, 22-38.
 35. Curran, M. P. and Perry, C. M. 2005, Tacrolimus: in patients with rheumatoid arthritis, *Drugs.*, **65**, 993-1001.
 36. Sasakawa, T., Sasakawa, Y., Ohkubo, Y., and Mutohm, S. 2005, FK506 inhibits prostaglandin E2 production from synovial cells by suppressing peripheral blood mononuclear cells, *Int. Immunopharmacol.*, **5**, 1291-1297.
 37. Hsu, W., Rosenquist, G.L., Ansari, A. A., and Gershwin, M. E. 2005, Autoimmunity and tyrosine sulfation, *Autoimmune Rev.*, **4**, 429-435.
 38. van Lent, P. L., Figdor, C. G., Barrera, P., et al. 2003, Expression of the dendritic cell-associated C-type lectin DC-SIGN by inflammatory matrix metalloproteinase-producing macrophages in rheumatoid arthritis synovium and interaction with intercellular adhesion molecule 3-positive T cells, *Arthritis Rheum.*, **48**, 360-369.
 39. Crazzolara, R. and Bernhard, D. 2005, CXCR4 chemokine receptors, histone deacetylase inhibitors and acute lymphoblastic leukemia, *Leuk. Lymphoma.*, **46**, 1545-1551.
 40. Glabinski, A. R., O'Bryant, S., Selmaj, K., and Ransohoff, R. M. 2000, CXC chemokine receptors expression during chronic relapsing experimental autoimmune encephalomyelitis, *Ann. N. Y. Acad. Sci.*, **917**, 135-144.
 41. Wada, T., Nakashima, T., Hiroshi, N., and Penninger, J. M. 2006, RANKL-RANK signaling in osteoclastogenesis and bone disease, *Trends Mol. Med.*, **12**, 17-25.
 42. Jimi, E. and Ghosh, S. 2005, Role of nuclear factor-kappaB in the immune system and bone, *Immunol. Rev.*, **208**, 80-87.
 43. Akoh, C. C., Lee, G. C., Liaw, Y. C., Huang, T. H., and Shaw, J. F. 2004, GDSL family of serine esterases/lipases, *Prog. Lipid Res.*, **43**, 534-552.
 44. Ishizuka, M., Toyama, Y., Watanabe, H., et al. 2004, Overexpression of human acyl-CoA thioesterase upregulates peroxisome biogenesis, *Exp. Cell Res.*, **297**, 127-141.
 45. Jalanko, A., Vesa, J., Manninen, T., et al. 2005, Mice with Ppt1Deltaex4 mutation replicate the INCL phenotype and show an inflammation-associated loss of interneurons, *Neurobiol. Dis.*, **18**, 226-241.
 46. Sakaguchi, N., Takahashi, T., Hata, H., et al. 2003, Altered thymic T-cell selection due to a mutation of the ZAP-70 gene causes autoimmune arthritis in mice, *Nature*, **426**, 454-460.
 47. Tsuchimochi, K., Yagishita, N., Yamasaki, S., et al. 2005, Identification of a crucial site for synoviolin expression, *Mol. Cell Biol.*, **25**, 7344-7356.
 48. Hoffmeyer, A., Avots, A., Flory, E., Weber, C. K., Serfling, E., and Rapp, U. R. 1998, The GABP-responsive element of the interleukin-2 enhancer is regulated by JNK/SAPK-activating pathways in T lymphocytes, *J. Biol. Chem.*, **273**, 10112-10119.
 49. Cook, P. W., Brown, J. R., Cornell, K. A., and Pittelkow, M. R. 2004, Suprabasal expression of human amphiregulin in the epidermis of transgenic mice induces a severe, early-onset, psoriasis-like skin pathology: expression of amphiregulin in the basal epidermis is also associated with synovitis, *Exp. Dermatol.*, **13**, 347-356.
 50. Bhagavathula, N., Nerusu, K. C., Fisher, G. J., et al. 2005, Amphiregulin and epidermal hyperplasia: amphiregulin is required to maintain the psoriatic phenotype of human skin grafts on severe combined immunodeficient mice, *Am. J. Pathol.*, **166**, 1009-1016.
 51. Ritchlin, C. T. 2005, Pathogenesis of psoriatic arthritis, *Curr. Opin. Rheumatol.*, **17**, 406-412.
 52. Davies, M. R., Harding, C. J., Raines, S., et al. 2005, Nurrl dependent regulation of pro-inflammatory mediators in immortalised synovial fibroblasts, *J. Inflamm. (Lond)*, **2**, 15.
 53. Chokki, M., Mitsushashi, H., and Kamimura, T. 2006, Metalloprotease-dependent amphiregulin release mediates tumor necrosis factor-alpha-induced IL-8 secretion in the human airway epithelial cell line NCI-H292, *Life Sci.*, **78**, 3051-3057.

Research article

Differentiation of monocytes into multinucleated giant bone-resorbing cells: two-step differentiation induced by nurse-like cells and cytokines

Tomoko Toyosaki-Maeda*, Hiroshi Takano^{†‡}, Tetsuya Tomita[†], Yuji Tsuruta*, Miki Maeda-Tanimura*, Yasunori Shimaoka[†], Tetsu Takahashi[‡], Tsunetoshi Itoh[§], Ryuji Suzuki* and Takahiro Ochi[†]

*Department of Immunology, Shionogi Research Laboratories, Shionogi & Co Ltd, Osaka, Japan

[†]Department of Orthopedic Surgery, Osaka University Medical School, Osaka, Japan

[‡]Department of Oral Surgery, Kyushu Dental College School of Dentistry, Kokura-Kita, Kita-Kyushu, Japan

[§]Department of Immuno-Ontogeny, Tohoku University School of Medicine, Aoba-ku, Sendai, Miyagi, Japan

Correspondence: Takahiro Ochi, Department of Orthopedic Surgery, Osaka University Medical School, Yamada-Oka 2-2, Suita, Osaka 566-0871, Japan. Tel: +81 6 6879 3862; fax: +81 6 6879 3869; e-mail: ochi@ort.med.osaka-u.ac.jp

Received: 6 December 2000

Revisions requested: 9 February 2001

Revisions received: 12 June 2001

Accepted: 22 June 2001

Published: 2 August 2001

Arthritis Res 2001, **3**:306–310

© 2001 Toyosaki-Maeda *et al*, licensee BioMed Central Ltd
(Print ISSN 1465-9905; Online ISSN 1465-9913)

Abstract

Bone resorption in the joints is the characteristic finding in patients with rheumatoid arthritis (RA). Osteoclast-like cells are present in the synovial tissues and invade the bone of patients with RA. The characteristics of these cells are not completely known. In the work reported here, we generated these cells from peripheral-blood monocytes from healthy individuals. The monocytes were co-cultured with nurse-like cells from synovial tissues of patients with RA (RA-NLCs). Within 5 weeks of culture, the monocytes were activated and differentiated into mononuclear cells positive for CD14 and tartrate-resistant acid phosphatase (TRAP). These mononuclear cells then differentiated into multinucleated giant bone-resorbing cells after stimulation with IL-3, IL-5, IL-7, and/or granulocyte-macrophage-colony-stimulating factor. TRAP-positive cells with similar characteristics were found in synovial fluid from patients with RA. These results indicate that multinucleated giant bone-resorbing cells are generated from monocytes in two steps: first, RA-NLCs induce monocytes to differentiate into TRAP-positive mononuclear cells, which are then induced by cytokines to differentiate into multinucleated giant bone-resorbing cells.

Keywords: monocytes, nurse cells, osteoclasts, rheumatoid arthritis, stromal cells

Our laboratory has established that nurse-like cells (NLCs) are present in the synovial tissues and bone marrow of patients with rheumatoid arthritis (RA) [1–3]. Such cells, which were first discovered in thymus, play an important role in thymocyte maturation and differentiation [4–6]. *In vitro*, they form unique complexes with thymocytes, which initially adhere to them and then

crawl beneath them [7–9]. This phenomenon, which is unique to NLCs at various tissue sites, has been called 'pseudoemperipolesis'. NLCs from RA synovial tissue (RA-NLCs) promote survival of B cells [2,3] and maintain the growth of myeloid cells of patients with RA [1], suggesting that they contribute profoundly to pathogenesis in RA.

DMEM = Dulbecco's modified Eagle's medium; FCS = fetal calf serum; GM-CSF = granulocyte/macrophage-colony-stimulating factor; HLA = human major histocompatibility antigen; IL = interleukin; NLC = nurse-like cell; RA = rheumatoid arthritis; RANKL = receptor activator of nuclear-factor- κ B ligand; RA-NLC = nurse-like cell derived from rheumatoid arthritis synovial tissue; TRAP = tartrate-resistant acid phosphatase.

Multinucleated cells in synovial tissues have been reported to invade the bone of patients with RA [10]. The cells' expression of tartrate-resistant acid phosphatase (TRAP) and calcitonin receptor suggested that they are osteoclasts [11,12]. Although the presence of osteoclast-like cells in rheumatoid synovium is well understood, the mechanism by which they differentiate is not. In order to examine the effect of RA-NLCs on monocyte functions, we co-cultured peripheral-blood monocytes with RA-NLCs and looked for morphological and functional alterations of CD14- and TRAP-positive cells. We also found such cells in synovial fluid from patients with RA. These cells differentiated into multinucleated giant bone-resorbing cells in the presence of IL-3, IL-5, IL-7, and/or granulocyte/macrophage-colonystimulating factor (GM-CSF). In this way we defined the process by which bone-resorbing cells are generated from monocytic cells.

Materials and methods

Isolation of NLCs from RA synovial tissues

RA-NLCs were established from RA synovial tissues as previously described [1]. Briefly, synovial tissues were obtained from knee joints of five patients with RA who fulfilled American College of Rheumatology criteria for RA [13], after informed consent had been obtained. The cells were cultured in DMEM (Dulbecco's modified Eagle's medium [DMEM; Gibco BRL, Gaithersburg, MD, USA] supplemented with 10% fetal calf serum [FCS; Hyclone, Logan, UT, USA], 100 units/ml of penicillin [Gibco BRL], and 100 µg/ml of streptomycin [Gibco BRL] at 37°C in 7.5% CO₂. RA-NLCs were identified by their ability to support pseudoemperipolesis, seen *in vitro* in the migration of a T-cell lymphoma line, MOLT-17, beneath the NLCs, as previously described [3].

Isolation of mononuclear cells from RA synovial fluid

Synovial fluid was obtained from patients with RA who fulfilled the American College of Rheumatology criteria for RA [13]. The infiltrating cells were collected from the fluid by centrifugation at 1900 g and were cultured in supplemented DMEM. After 3 to 5 weeks of culture, most of the lymphocytes and granulocytes disappeared and monocyte-like cells became dominant. CD14-positive monocyte-like cells were purified from this population with a magnetic-activated cell sorter (MACS; Miltenyi Biotec GmbH, Germany) using anti-CD14 antibody conjugated to magnetic beads in accordance with the manufacturer's instructions. The purity of CD14-positive cells was analyzed using a fluorescence-activated cell sorter (FACScan™; see Supplementary material).

Isolation and culture of monocytes from peripheral blood

Peripheral-blood monocytes were collected as plastic-adherent cells, as described previously [14]. Mononuclear cells were isolated from heparinized peripheral blood from five healthy volunteers [15]. Over 97% of the adherent

cells were determined to be monocytes by morphology and CD14 expression.

Monocytes (1×10^6) were co-cultured with RA-NLCs. After 3 to 5 weeks, TRAP-positive mononuclear cells with abundant cytoplasm became dominant. They were collected by gently washing the culture with warm supplemented DMEM and their purity was confirmed cytochemically.

Formation of multinucleated giant cells by TRAP-positive mononuclear cells

The CD14-positive and TRAP-positive mononuclear cells from the synovial fluid of patients with RA were examined for expression of surface antigen and for phagocytic activity and were stimulated with various cytokines (see Supplementary material).

TRAP-positive mononuclear cells (5×10^4) were cultured in supplemented DMEM in the presence or absence of various cytokines or in conditioned medium ([15]; and see Supplementary material) for 96–120 h. In the presence of receptor activator nuclear factor κB ligand (RANKL), cultures were maintained for 14 days. At the end of the culture period, May-Grunwald-Giemsa (Wako Pure Chemical Co., Osaka, Japan) and TRAP staining (TRAP-staining kit; Sigma, St Louis, MO, USA) were conducted. The frequency of osteoclasts was evaluated from the fusion index, as previously described [16]. More than 1000 nuclei within TRAP-positive multinucleated cells (>4 nuclei/cell) were counted. The fusion index (%) was calculated according to the formula:

$$\frac{\text{total no. of nuclei within multinucleated cells} \times 100}{\text{total no. of nuclei counted}}$$

where 'multinucleated cells' are cells with >4 nuclei.

Examination of bone resorption

TRAP-positive mononuclear cells (5×10^4) were stimulated with various cytokines on a dentin slice for 7 days. In order to examine resorption areas with a scanning electron microscope, the differentiated cells were washed off the slices with distilled water. Then the slices were dehydrated, air-dried, and sputtered with gold.

Results

Morphological changes of peripheral-blood monocytes after co-culture with RA-NLCs

After peripheral-blood monocytes had been cultured with RA-NLCs for 3 to 4 weeks, we recovered TRAP-positive mononuclear cells (Fig. 1a–1c) with abundant cytoplasm and an off-center nucleus (Fig. 1a and 1b). These cells strongly expressed CD11b, CD11c, CD14, CD45, and human major histocompatibility antigen (HLA)-DR, sug-

gesting that they were of monocyte lineage (Table 1). However, they did not express CD11a, CD35, or CD68, which are expressed on freshly isolated monocytes (see Supplementary material).

Presence of TRAP-positive mononuclear cells in synovial fluid from patients with RA

We detected monocytic cells positive for CD14 and TRAP in synovial fluids of patients with RA. These cells also strongly expressed CD11b, CD11c, CD14, CD45, and HLA-DR (see Supplementary material) but not CD1a, CD1b, CD2, CD5, or CD86, which are expressed on dendritic cells derived from monocytes [17]. These results indicate that TRAP-positive monocytic cells present in synovial fluid and those induced *in vitro* in cultures with RA-NLCs are morphologically and phenotypically the same. These cells were obtained from the synovial fluid of all patients with RA, regardless of age and sex. CD14-positive monocyte-like cells accounted for 20–91% of the mononuclear cells freshly isolated from the synovial fluid of such patients (data not presented).


Differentiation of TRAP-positive mononuclear cells into multinucleated giant bone-resorbing cells in the absence of RA-NLCs

The TRAP-positive cells induced *in vitro* and those isolated from synovial fluid both differentiated into multinucleated cells after being cultured for 72 to 96 h with the conditioned medium (Fig. 1d). These multinucleated cells still possessed TRAP activity (data not shown) and formed resorption areas on dentin slices (Fig. 2), suggesting that they had bone-resorbing activity as osteoclasts.

Induction of multinucleated cells by IL-3, IL-5, IL-7 or GM-CSF

The cytokines IL-3, IL-5, IL-7, and GM-CSF induced differentiation of TRAP-positive cells induced *in vitro* or those isolated from synovial fluid into osteoclasts (Table 1). Regardless of which cytokine was used to stimulate differentiation of the osteoclasts, they were all positive for TRAP and formed resorption pits on dentin slices, suggesting that they were all identical to the cells induced by conditioned medium (data not shown). The fusion index of osteoclasts induced by a mixture of cytokines was higher than those stimulated with a single cytokine. The cytokines IL-6 and IL-8, which are produced by RA-NLCs [2], did not induce osteoclast formation. RANKL was recently reported to induce osteoclasts from human peripheral blood [18,19]; however, a mixture of macrophage-colony-stimulating factor and RANKL exhibited only weak activity for induction of osteoclasts from the TRAP-positive mononuclear cells (Table 1). Phytohemagglutinin, which was contained in the conditioned medium, did not induce differentiation. TRAP-positive cells from the synovial fluid of patients with RA have a fusion index similar to that found for the TRAP-positive cells obtained experimentally

Figure 1



Morphology of TRAP-positive mononuclear cells induced from peripheral-blood monocytes with RA-NLCs. (a) Phase-contrast micrograph of monocytes co-cultured with RA-NLCs. Mononuclear cells (arrow) are growing on the RA-NLCs. (b) Mononuclear cells collected from the culture shown in (a). May-Grunwald-Giemsa staining. (c) Detection of TRAP expressed by the mononuclear cells (TRAP-positive cells were stained red with their cytoplasm). (d) Differentiated mononuclear cells. The cells shown here are multinucleated giant cells. May-Grunwald-Giemsa staining. Scale lines = 100 μm.


from monocytes (data not shown). Induction of the osteoclasts was completely neutralized by the antibody to each cytokine (see Supplementary material).

Discussion

We have shown that the novel ability of RA-NLCs may contribute to the pathogenesis of RA by encouraging the generation of TRAP-positive mononuclear cells, which are osteoclast precursors. The TRAP-positive precursor cells have phagocytic activity and are negative for CD83, suggesting that they are different from peripheral-blood monocytes and dendritic cells [14,17].

Fujikawa *et al* reported that synovial macrophages differentiated into osteoclasts after incubation in the presence of a rat osteoblast-like cell line [20]. The fibroblasts isolated from RA synovia induced differentiation of monocytes into multinucleated cells in the presence of 1,25-dihydroxyvitamin D₃ and macrophage-colony-

Figure 2



Scanning electron micrographs of dentin slices, showing (a) resorption areas formed on dentin by TRAP-positive multinucleated giant cells derived from monocytes and stimulated with granulocyte-macrophage-colony-stimulating factor for 96 h and (b) a control slice incubated with the TRAP-positive mononuclear cells in the absence of cytokines. Scale lines = 50 μm .

stimulating factor [21]. Further study will be required to determine the identity of those monocytic cells and our cells. Fibroblastic cells in synovial fluid from patients with RA have been reported to support pseudoemperipolesis, which was considered to be the unique feature of the nurse cells, in the presence of IL-4 [22]. Shigeyama *et al* recently reported that RA synovial fluid may promote osteoclastogenesis from monocytes by expressing osteoclast differentiation factor [23]. It is likely that RA fibroblasts and RA-NLCs share several roles in the pathogenesis of RA, including activation of monocytes. However, the molecules required in our study for osteoclastogenesis from monocytes were different from those in the study of Shigeyama *et al* [23]. These findings suggest that multiple pathways for osteoclastogenesis in RA synovia may cause severe joint destruction.

There may be two steps for generation of the osteoclasts in the joints of patients with RA: first, differentiation of monocytes into TRAP-positive mononuclear cells induced and maintained by RA-NLCs, followed by cytokine-induced differentiation of these mononuclear cells into osteoclasts. The interaction between monocytes and RA-NLCs required adhesion molecules, but RANK (receptor activator of nuclear-factor- κB) and RANKL were not necessary to induce the TRAP-positive cells in preliminary studies in our laboratory (unpublished observation). The molecules required in the interaction are under investigation. In addition, we found the presence of TRAP-positive mononuclear cells which differentiated into osteoclasts in synovial fluids of patients with RA *in vitro*. Monocytes may infiltrate the affected joints and differentiate into TRAP-positive mononuclear cells under the influence of RA-NLCs. This conclusion is consistent with previous findings

Table 1

Differentiation of human TRAP-positive mononuclear cells derived from peripheral-blood monocytes into multinucleated cells after stimulation with IL-3, IL-5, IL-7, and GM-CSF

Stimulator	Concentration	Fusion index (%) ^a
None		1.8
Conditioned medium ^b	(10% v/v)	86.1
IL-1 α	(1 ng/ml)	1.1
IL-1 β	(1 ng/ml)	1.3
IL-2	(250 U/ml)	7.7
IL-3	(5 ng/ml)	64.8
IL-4	(100 U/ml)	1.3
IL-5	(1 ng/ml)	66.1
IL-6	(20 ng/ml)	5.6
IL-6 + sIL-6R	(sIL-6R: 200 ng/ml)	5.3
IL-7	(20 ng/ml)	72.4
IL-8	(20 ng/ml)	2.1
IFN- γ	(100 U/ml)	7.1
GM-CSF	(1 ng/ml)	73.8
M-CSF	(25 ng/ml)	5.8
TNF- α	(1 ng/ml)	6.9
VD ₃	(10 ⁻⁷ mol/l)	3.4
Dexamethasone	(10 ⁻⁸ mol/l)	0.8
M-CSF + IL-4	(M-CSF, 25 ng/ml; IL-4, 100 U/ml)	0.1
IL-3 + IL-7		75.4
IL-7 + GM-CSF		77.3
IL-3 + IL-7 + GM-CSF		78.7
Phytohemagglutinin	(1% v/v)	2.0
RANKL ^c	(100 ng/ml)	2.6
RANKL + M-CSF ^c	(RANKL/ODF, 100 ng/ml; M-CSF, 25 ng/ml)	38.1

^aTRAP-positive mononuclear cells were stimulated with various cytokines for 96–120 h. Fusion indices were calculated using the formula given in the text and previously [16]. Data are representative of three independent experiments using TRAP-positive cells induced from monocytes. ^bSee [15], and Supplementary material. ^cThe culture was maintained for 14 days. GM-CSF = granulocyte macrophage-colony-stimulating factor; IFN = interferon; IL = interleukin; M-CSF = macrophage-colony-stimulating factor; ODF = osteoclast differentiation factor; RANKL = receptor activator of nuclear-factor- κB ligand; sIL-6R = soluble interleukin-6 receptor; TNF = tumor necrosis factor; VD₃ = 1,25-dihydroxyvitamin D₃.

of TRAP-positive multinucleated giant cells in the synovial tissue of patients with RA [10–12]. Further studies are required to characterize these osteoclasts derived from the TRAP-positive mononuclear cells and to delineate the unique course of differentiation into bone-resorbing cells promoted by RA-NLCs.

Conclusion

In order to elucidate the role of RA-NLCs, monocytes were co-cultured with RA-NLCs. Monocytes differentiated into TRAP-positive mononuclear cells, the precursor cells of osteoclasts. Osteoclasts were generated from TRAP-positive mononuclear cells in the presence of IL-3, IL-5, IL-7, and GM-CSF. TRAP-positive cells were also present in synovial fluids of patients with RA. RA-NLCs may play a significant role in the activation of monocytes and long-term maintenance of differentiated monocytes (osteoclast precursors). The present study suggests that monocytes may differentiate into osteoclast precursor cells in the affected joints of patients with RA.

Acknowledgements

This work was supported by the Program for Promotion of Fundamental Studies in Health Science of the Organization for Drug ADR Relief, R&D Promotion and Product Review of Japan. The authors are grateful to Dr Ichiro Kurane (Department of Virology 1, National Institute of Infectious Diseases, Tokyo, Japan) for helpful discussions and critical reading of the manuscript.

References

- Tomita T, Takeuchi E, Toyosaki-Maeda T, Oku H, Kaneko M, Takano, Sugamoto H, Ohzono K, Suzuki R, Ochi T: Establishment of nurse-like stromal cells from bone marrow patients with rheumatoid arthritis: indication of characteristic bone marrow microenvironment in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 1999, 38:854-863.
- Takeuchi E, Tomita T, Toyosaki-Maeda T, Hashimoto H, Kaneko M, Takano H, Sugano K, Suzuki R, Ochi T: Establishment and characterization of nurse-cell like stromal cell lines from synovial tissues of patients with rheumatoid arthritis. *Arthritis Rheum* 1999, 42:221-228.
- Shimaoka Y, Attrep JF, Hirano T, Ishihara K, Suzuki R, Toyosaki T, Ochi T, Lipsky PE: Nurse-like cells from bone marrow and synovium of patients with rheumatoid arthritis promote survival and enhance function of human B cells. *J Clin Invest* 1998, 102:606-618.
- Pezzano M, Li Y, Philip D, Omene C, Cantey M, Saunders G, Guyden JC: Thymic nurse cell rescue of early CD4+ CD8+ thymocytes from apoptosis. *Cell Mol Biol* 1995, 41:1099-1111.
- Pezzano M, Philip D, Stephenson S, Li Y, Reid V, Maitta R, Guyden JC: Positive selection by thymic nurse cells requires IL-1 beta and is associated with an increased Bcl-2 expression. *Cell Immunol* 1996, 169:174-184.
- Iwabuchi K, Nakayama K, McCoy RL, Wang F, Nishimura T, Habu S, Murphy KM, Loh, DY: Cellular and peptide requirements for *in vitro* clonal deletion of immature thymocytes. *Proc Natl Acad Sci U S A* 1992, 89:9000-9004.
- Itoh T, Doi H, Chin S, Nishimura T, Kasahara S: Establishment of mouse thymic nurse cell clones from a spontaneous BALB/c thymic tumor. *Eur J Immunol* 1988, 18:821-824.
- Miyake K, Hasunuma Y, Yagita H, Kimoto, M: Requirement for VLA-4 and VLA-5 integrins in lymphoma cells binding to and migration beneath stromal cells in culture. *J Cell Biol* 1992, 119:653-662.
- Hiai H, Nishi Y, Miyazawa T, Matsudaira Y, Nishizuka Y: Mouse lymphoid leukemias: symbiotic complexes of neoplastic lymphocytes and their microenvironments. *J Natl Cancer Inst* 1981, 66:713-722.
- Gravallese EM, Harada Y, Wang JT, Gom AH, Thomhil TS, Goldring, SR: Identification of cell types responsible for bone resorption in rheumatoid arthritis and juvenile rheumatoid arthritis. *Am J Pathol* 1998, 152:943-951.
- Chang JS, Quinn JM, Demaziere A, Bulstrode CJ, Francis MJ, Duthie RB, Athanasou NA: Bone resorption by cells isolated from rheumatoid synovium. *Ann Rheum Dis* 1992, 51:1223-1229.
- Fujikawa Y, Shingu M, Torisu T, Itonaga I, Masumi S: Bone resorption by tartrate-resistant acid phosphatase-positive multinuclear cells isolated from rheumatoid synovium. *Br J Rheumatol* 1996, 35:213-217.
- Arnett FC, Edworthy SM, Bloch DA, Mcshane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HA, Medsger TA, Jr, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JG, Wilder RL, Hunder GG: The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988, 31:315-324.
- Akagawa KS, Takatsuka N, Nozaki Y, Komuro I, Azuma M, Ueda M, Naito M, Takahashi K: Generation of CD11b+RelB+ dendritic cells and tartrate-resistant acid phosphatase-positive osteoclast-like multinucleated giant cells from human monocytes. *Blood* 1996, 88:4029-4039.
- Toyosaki T, Tsuruta Y, Yoshioka T, Takemoto H, Suzuki R, Tomita T, Ochi T: Recognition of rheumatoid arthritis synovial antigen by CD4+, CD8- T cell clones established from rheumatoid arthritis joints. *Arthritis Rheum* 1998, 41:92-100.
- Higuchi S, Tabata N, Tajima M, Ito M, Tsurudome M, Sudo A, Uchida A, Ito Y: Induction of human osteoclast-like cells by treatment of blood monocytes with anti-fusion regulatory protein-1/CD98 monoclonal antibodies. *J Bone Miner Res* 1998, 13:44-49.
- Zhou, LJ, Tedder TF: CD14+ blood monocytes can differentiate into functionally mature CD83+ dendritic cells. *Proc Natl Acad Sci U S A* 1996, 93:2588-2592.
- Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, Mochizuki S, Tomoyasu A, Yano K, Goto M, Murakami A, Tsuda E, Morinaga T, Higashino K, Udagawa N, Takahashi N, Suda T: Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci U S A* 1998, 95:3597-3602.
- Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ: Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 1998, 93:165-178.
- Fujikawa Y, Sabokbar A, Neale S, Athanasou NA: Human osteoclast formation and bone resorption by monocytes and synovial macrophages in rheumatoid arthritis. *Ann Rheum Dis* 1996, 55:816-822.
- Takayanagi H, Oda H, Yamamoto S, Kawaguchi H, Tanaka S, Nishikawa T, Koshihara Y: A new mechanism of bone destruction in rheumatoid arthritis: synovial fibroblasts induce osteoclastogenesis. *Biochem Biophys Res Commun* 1997, 240:279-286.
- Burger JA, Zvaifler NJ, tsukada N, Firestein GS, Kipps TJ: Fibroblast-like synoviocytes support pseudoemperipolesis via a stromal cell-derived B-cell factor-1-and CD106 (VCAM-1)-dependent mechanism. *J Clin Invest* 2001, 107:305-315.
- Shigeyama Y, Pap T, Kunzler P, Simmen BR, Gay RE, Gay S: Expression of osteoclast differentiation factor in rheumatoid arthritis. *Arthritis Rheum* 2000, 43:2523-2530.

Supplementary material

Supplementary materials and methods Cell lines

Human lung fibroblasts CCD-19Lu were obtained from American Type Culture Collection (Rockville, MD, USA). Human T cell line MOLT-17 was a generous gift from Dr J Minowada (Fujisaki Cell Center, Okayama, Japan). These cell lines were cultured as recommended by the providers.

Examination of pseudoemperipolesis

Pseudoemperipolesis was measured as previously described [3]. RA synovial stromal cells (3×10^4) were incubated in supplemented DMEM in 24-well culture

plates overnight. The next day, MOLT-17 cells (1×10^6) were added to the RA-NLC culture. Stromal cells with more than three MOLT-17 cells beneath them after 6 h of incubation were considered NLCs.

Long-term maintenance of monocytes by RA-NLCs

Monocytes (2.5×10^5) were co-cultured with RA-NLCs or CCD-19Lu with or without culture inserts (Becton Dickinson, Franklin Lakes, NJ, USA) in supplemented DMEM in 48-well culture plates at 37°C in 7.5% CO₂ for up to 70 days. Half of the medium was changed once a week. The TRAP-positive mononuclear cells were collected from the culture, stained with trypan blue, and counted for viability under a microscope.

Antibodies and staining of cells

The cell-surface antigens on freshly isolated monocytes and TRAP-positive mononuclear cells were examined by staining with monoclonal antibodies specific for CD1a, CD4, CD5, CD11a, CD11b, CD11c, CD13, CD14, CD15, CD16, CD19, CD20, CD34, CD45, CD45RA, CD45RO, CD54, HLA-DR (Becton Dickinson), CD1b (Nichirei, Tokyo, Japan), CD2, CD3 (Ortho Diagnostics, Raritan, NJ, USA), CD35, CD68 (DAKO Japan, Kyoto, Japan), CD51/61, CD83, HLA-A, B and C (Pharmingen, San Diego, CA, USA), CD80, and CD86 (Ansell, Bayport, MN, USA). Antigen-expression was analyzed with a FACScan flow cytometer (Becton Dickinson). Multinucleated giant bone-resorbing cells differentiated from TRAP-positive mononuclear cells were fixed with cold acetone and stained immunohistochemically with rabbit polyclonal antibodies specific for actin (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA), carbonic anhydrase II (Rockland, Gilbertsville, PA, USA), or vitronectin receptor (Chemicon International, Inc., Temecula, CA, USA). TRAP activity in TRAP-positive mononuclear cells and multinucleated giant bone-resorbing cells was examined using a TRAP-staining kit (Sigma, St Louis, MO, USA). Neutralizing antibodies to human IL-3, IL-5, IL-7, and GM-CSF were purchased from Genzyme (Cambridge, MA, USA).

Cytokines and reagents

Conditioned media were prepared as previously reported [15]. Briefly, a mixture of peripheral-blood mononuclear cells from 10 healthy donors was stimulated with phytohemagglutinin at 37°C for 72 h. Culture supernatant fluids were collected, filtered, and used as conditioned media. IL-1 α was purchased from Immugenex (Los Angeles, CA). IL-1 β , IL-2, IL-3, IL-4, IL-5, IL-6, soluble IL-6 receptor (sIL-6R), interferon gamma (IFN- γ), granulocyte/macrophage-colony stimulating factor (GM-CSF), macrophage-colony stimulating factor, and tumor necrosis factor (TNF)- α were purchased from R&D Systems, Inc. (Minneapolis, MN, USA). IL-7 and IL-8 were purchased from Genzyme. 1,25-dihydroxyvitamin D₃ and dexamethasone were purchased from Wako Pure Chemical Co. (Osaka, Japan). Receptor

activator of nuclear-factor- κ B ligand (RANKL) was obtained from Peprotech (London, UK).

Assessment of phagocytic activity

Phagocytic activity of TRAP-positive mononuclear cells was assessed from their ingestion of heat-killed yeast. TRAP-positive mononuclear cells (1×10^6) were incubated with 2×10^7 yeast cells in phosphate-buffered saline supplemented with 10% fresh human serum, type AB, at 37°C for 45 min. The cells were washed and stained with fuchsin (Wako Pure Chemical Co.), and the cells with ingested yeast were counted under a microscope.

Inhibition of the generation of multinucleated giant cells from TRAP-positive mononuclear cells by neutralizing antibodies

Neutralizing antibodies specific for IL-3, IL-5, IL-7, and GM-CSF were used for inhibition of the generation of multinucleated giant bone-resorbing cells. Irrelevant polyclonal mouse IgG from Jackson ImmunoResearch (West Grove, PA, USA) was used as a control. The TRAP-positive mononuclear cells (5×10^4) were pre-incubated with each antibody in DMEM containing 10% FCS in microtubes at 37°C for 1 h. The cells were cultured in 4-well chamber slides, and stimulated with a cytokine for 96–120 h at 37°C in 7.5% CO₂. At the end of the culture period, the cells were stained for TRAP and the fusion index was calculated as described in the main paper.

Detection of calcitonin receptors

Calcitonin receptors on the multinucleated giant bone-resorbing cells were detected *in situ* using ¹²⁵I-human calcitonin were performed as described elsewhere [23]. TRAP-positive mononuclear cells (5×10^4) were stimulated with IL-3, IL-5, IL-7, or GM-CSF at the optimal concentrations in 4-well chamber slides (Nalge Nunc International, Rochester, NY, USA) for 96–120 h at 37°C in 7.5% CO₂. After formation of multinucleated giant bone-resorbing cells had been confirmed microscopically, the cells were incubated in 0.4 ml α MEM (Gibco BRL) with 0.1% bovine serum albumin and 0.2 mCi/ml of ¹²⁵I-labeled human calcitonin (Amersham Pharmacia Biotech, Buckinghamshire, UK) for 1 h at 22°C. Nonspecific binding was assessed on each slide in the presence of an excess amount of unlabeled human calcitonin. Then the cells were washed three times with phosphate-buffered saline solution and fixed with 2.5% formaldehyde and 2.5% glutaraldehyde in 0.1 M sodium cacodylate buffer. The slides were washed and dried as previously described [S1]. Air-dried slides were dipped in photographic emulsion (Kodak NTB3; Eastman Kodak, Rochester, NY, USA), drained, and dried for 2 h and were kept in a light-proof container with desiccant at 4°C for 10 days. The slides were developed in accordance with the manufacturer's instructions.

Supplementary Table 1

Expression of surface antigen by TRAP-positive mononuclear cells generated by co-culture with RA-NLC

Antigen	% Positive ^a		
	Monocytes from peripheral blood	TRAP-positive cells induced from monocytes	TRAP-positive cells from RA-SF
CD1a	1.24	0.11	0.24
CD1b	0.22	0.13	0.12
CD2	2.20	0.18	1.01
CD3	0.82	0.85	0.82
CD4	0.28	0.61	0.35
CD5	0.23	0.44	0.29
CD11a	LFA-1	94.72	1.16
CD11b	CR3 α chain	95.46	99.42
CD11c	CR4 α chain	99.47	99.22
CD13		99.44	96.15
CD14		95.70	99.28
CD15	Le ^x	22.81	0.49
CD16	Fc γ R III	0.72	0.24
CD19		0.33	0.13
CD20		0.36	0.11
CD34		0.54	0.18
CD35	CR1	93.49	0.50
CD45		98.73	94.28
CD45RA		0.37	0.15
RA45RO		0.65	4.18
CD51/61	VNR ^b	0.22	0.10
CD54	ICAM-1	0.21	0.30
CD68		94.59	1.06
CD80	B7/BB1	8.19	7.41
CD83		0.37	0.10
CD86	B70/FUN-1	90.01	0.55
HLA-A, -B, -C		97.44	99.68
HLA-DR		96.65	98.45

^aCells were stained with monoclonal antibody specific for various antigens, and analyzed by FACScan. Data are representative of four independent analyses. ^bVitronectin receptor. RA-SF = synovial fluid from patients with rheumatoid arthritis.

Supplementary results**Phenotypic characterization of the TRAP-positive mononuclear cells induced by NLCs**

The TRAP-positive mononuclear cells, whether induced from monocytes or collected from the synovial fluid of patients with RA, strongly expressed CD11b, CD11c, CD14, CD45, HLA-A, HLA-B, HLA-C, and HLA-DR but did not express CD1a, CD1b, CD2, CD4, CD5, CD16, CD19, CD20, or CD83. These observations suggest that

these cells belong to the monocyte/macrophage lineage (Supplementary Table 1). However, the cells did not express CD11a, CD35, or CD68, which are expressed on freshly isolated monocytes from peripheral blood (Supplementary Table 2). The TRAP-positive mononuclear cells had strong phagocytic activity against heat-killed yeast (data not presented). In addition, the cells were positive for carbonic anhydrase II, actin, and vitronectin receptor (Supplementary Fig. 1), and calcitonin receptors were also


Supplementary Table 2

Inhibition of formation of multinucleated giant cells by neutralizing antibodies specific for IL-3, IL-5, IL-7, and GM-CSF

Antibody ^b	Fusion index (%) ^a			
	IL-3 (5 ng/ml)	IL-5 (1 ng/ml)	IL-7 (20 ng/ml)	GM-CSF (1 ng/ml)
Polyclonal mouse IgG	63.8 ± 1.2	62.3 ± 3.6	66.2 ± 5.4	67.7 ± 6.0
Anti-IL-3	7.3 ± 1.6	–	–	–
Anti-IL-5	–	3.8 ± 1.8	–	–
Anti-IL-7	–	–	5.5 ± 5.8	–
Anti-GM-CSF	–	–	–	4.0 ± 1.3

^aFusion indices were calculated using the formula given in the Methods section. Each value is the mean ± SD of three independent experiments using TRAP-positive cells induced from monocytes of three individuals. ^bConcentrations of cytokines: polyclonal mouse IgG, 10 mg/ml; anti-IL-3, 10 mg/ml; anti-IL-5, 5 mg/ml; anti-IL-7, 10 mg/ml; anti-GM-CSF, 2 mg/ml. – = not examined; GM-CSF = granulocyte/macrophage-colony-stimulating factor; IL = interleukin.

Supplementary Figure 1



Immunohistochemical staining of multinucleated giant bone-resorbing cells. The cells were stained green for (a) carbonic anhydrase II and (b) vitronectin receptor. The cells were also positive for (c) actin, which showed red, ring-form staining. (d) *In situ* detection of calcitonin receptor using ¹²⁵I-human calcitonin. Black grains mark the cells expressing calcitonin receptor. The multinucleated giant bone-resorbing cells were incubated with ¹²⁵I-human calcitonin for 1 h. The cells were washed, fixed, and dried as described in the Supplementary materials and methods section. Air-dried slides were exposed at 4°C for 10 days. Scale lines = 100 μm.

detected (Supplementary Fig. 1); these four receptors are considered characteristic of osteoclasts [10,23].

Long-term maintenance of monocytes by RA-NLCs

Monocytes obtained from all five healthy individuals and co-cultured with RA-NLCs grew and differentiated into TRAP-positive mononuclear cells (Supplementary Fig. 2). We also used five RA-NLC lines, each from a different patient with RA. All five cell lines induced the generation of TRAP-positive mononuclear cells from human peripheral blood monocytes (data not presented). The generation of these TRAP-positive cells required direct contact with RA-NLCs.


Induction of osteoclasts by IL-3, IL-5, IL-7, or GM-CSF

The conditioned medium contains several cytokines. In order to determine which cytokines induce TRAP-positive mononuclear cells to differentiate into multinucleated giant bone-resorbing cells, *in vitro* induced TRAP-positive cells and those isolated from synovial fluid of patients with RA were collected and stimulated with various cytokines in the absence of RA-NLCs. In a preliminary study, a high-performance liquid chromatography fraction of the conditioned medium, which was used to induce TRAP-positive mononuclear cells to differentiate into multinucleated giant bone-resorbing cells, contained proteins with a molecular weight of approximately 20 kDa (data not shown). Therefore, we mainly examined cytokines with approximately that molecular weight. The activities of IL-3, IL-5, IL-7, and GM-CSF for inducing multinucleated giant bone-resorbing cells were completely neutralized by the antibody to the respective cytokine (Supplementary Table 2).

Supplementary reference

- S1. Nicholson GC, Moseley JM, Sexton PM, Mendelsohn FAO, Martin TJ: Abundant calcitonin receptors in isolated rat osteoclasts. Biochemical and autoradiographic characterization. *J Clin Invest* 1986, 78:355-360.

Supplementary Figure 2



Effects of culture conditions on survival of monocytes from five donors with RA-NLCs or CCD-19Lu, in the presence (separated) or absence (no indication) of culture inserts. Each culture was maintained for up to 70 days. The resulting mononuclear cells were collected, stained with trypan blue, and counted under a microscope. Each experiment was conducted in duplicate, and each point represents the mean number of viable cells in two cultures.

Rheumatoid Arthritis Synovial Stromal Cells Inhibit Apoptosis and Up-Regulate Bcl-x_L Expression by B Cells in a CD49/CD29-CD106-Dependent Mechanism¹

Kenji Hayashida,* Yasunori Shimaoka,[†] Takahiro Ochi,[†] and Peter E Lipsky^{2*}

Inflammatory sites, such as rheumatoid arthritis (RA) synovial tissue, contain large numbers of activated B cells and plasma cells. However, the mechanisms maintaining B cell viability and promoting their differentiation are not known, but interactions with stromal cells may play a role. To examine this, purified human peripheral B cells were cultured with a stromal cell line (SCL) derived from RA synovial tissue, and the effects on apoptosis and expression of Bcl-2-related proteins were analyzed. As a control, B cells were also cultured with SCL from osteoarthritis synovium or skin fibroblasts. B cells cultured with medium alone underwent spontaneous apoptosis. However, B cells cultured with RA SCL cells exhibited less apoptosis and greater viability. Although SCL from osteoarthritis synovium and skin fibroblasts also rescued B cells from apoptosis, they were less effective than RA SCL. B cell expression of Bcl-x_L was markedly increased by RA SCL in a contact-dependent manner, whereas B cell expression of Bcl-2 was unaffected. Protection of B cells from apoptosis and up-regulation of Bcl-x_L by RA SCL were both blocked by mAbs to CD106 (VCAM-1), but not CD54 (ICAM-1). Furthermore, cross-linking of CD49d/CD29 (very late Ag-4) on the surface of B cells rescued them from apoptosis and up-regulated Bcl-x_L expression. These results indicate that SCL derived from RA synovial tissue play a role in promoting B cell survival by inducing Bcl-x_L expression and blocking B cell apoptosis in a CD49d/CD29-CD106-dependent manner. *The Journal of Immunology*, 2000, 164: 1110–1116.

The synovial tissue in rheumatoid arthritis (RA)³ is characterized by infiltration with a variety of inflammatory cells, including T cells, B cells, and plasma cells (1). The lymphocyte accumulation in synovial tissue might be induced by an increased migration from peripheral blood, proliferation of these cells in situ, and/or inhibition of cell death. Increased migration into the synovium has been suggested (2, 3), whereas local proliferation appears to be minimal (4). Programmed cell death is limited in the synovium and rarely involves lymphocytes (5). Therefore, the limited programmed cell death of lymphocytes may contribute to their accumulation in the synovium. A number of mechanisms have been proposed to account for the apparently diminished apoptosis of lymphocytes, including the action of cytokines (6–17), the presence of inhibitors of apoptosis, such as soluble CD95 and Fas ligand (18, 19), and the direct impact of cell-to-cell contact (20). Among the cells that appear to be active in blocking lymphocyte apoptosis is a population of stromal cells with the characteristics of thymus and bone marrow nurse cells (21, 22). Previous studies documented that these stromal cell lines (SCL) blocked B cell apoptosis in vitro (22). The current studies

examined the mechanism by which B cells were rescued from apoptosis and demonstrated that SCL derived from RA synovium induced expression of the antiapoptotic protein, Bcl-x_L, by a mechanism that involved cell-to-cell contact mediated by CD49d/CD29-CD106 interactions.

Materials and Methods

Antibodies and reagents

Biotinylated mouse anti-human CD19 mAb and mouse anti-human CD29 mAb (4B4) were purchased from Coulter (Miami, FL). Mouse anti-human CD20 mAb conjugated with PE, goat anti-mouse IgG conjugated with FITC, mouse anti-human CD14 mAb conjugated with FITC, goat anti-mouse IgG, and goat anti-rabbit IgG conjugated with HRP were obtained from Sigma (St. Louis, MO). Mouse anti-human CD49d mAb (HP2/1) and mouse anti-human CD106 mAb (1.G11B1) were purchased from Immunotech (Miami, FL) and Serotec (Oxford, U.K.), respectively. Mouse IgG1 (MOPC) mAb, mouse anti-human IgM heavy chain (DA4.4) conjugated with biotin, mouse anti-human CD11a mAb (TS1/22), and mouse anti-human CD18 mAb (TS1/18) were prepared from hybridoma cell lines purchased from American Type Culture Collection (Manassas, VA). Mouse anti-human CD54 mAb (R6.5) was a gift from Dr. R. Rothlein (Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT). Rabbit anti-human Bcl-2 (C-21) and Bcl-x_L (L-19) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA).

Propidium iodide and streptavidin were purchased from Sigma. Triazol reagent, DNase I, and SuperScript II reverse transcriptase were obtained from Life Technologies (Frederick, MD). Taq polymerase was obtained from Promega (Madison, WI). Oligo(dT) and Ficoll/Isopaque were purchased from Pharmacia (Piscataway, NJ). DMEM with high glucose, RPMI 1640, and FBS were purchased from Life Technologies. The protein assay system, polyvinylidene difluoride membranes, and Zeta-Probe blotting membranes were purchased from Bio-Rad (Richmond, CA). ECL was purchased from Amersham (Aylesbury, U.K.).

Stromal cell lines and fibroblast lines

One RA SCL was established from synovium as previously described (22). In brief, synovial tissues of patients with RA who met American College of Rheumatology criteria (23) were obtained after obtaining informed consent and were dissociated with collagenase and trypsin. Dissociated single

*University of Texas Southwestern Medical Center, Dallas, TX 75235; and [†]Osaka University Medical School, Osaka, Japan

Received for publication April 27, 1999. Accepted for publication October 26, 1999.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported by National Institutes of Health Specialized Center of Research Grant AR-39169 and the program for Promotion of Fundamental Studies in Health Science of the Organization for Drug ADR Relief, Research and Development Promotion and Product Review of Japan.

² Address correspondence and reprint requests to Dr. Peter E. Lipsky, Department of Internal Medicine, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, TX 75235-8884. E-mail address: peter.lipsky@email.swmed.edu

³ Abbreviations used in this paper: RA, rheumatoid arthritis; SCL, stromal cell line; OA, osteoarthritis; VLA-4, very late Ag-4; FDC, follicular dendritic cells.




FIGURE 1. The SCL derived from RA synovial tissue rescue peripheral B cells from apoptosis. Peripheral B cells (1×10^5) were cultured with or without RA SCL (Sy77; 4×10^3) in 96-well microtiter plates. After various lengths of culture, B cells were harvested with gentle pipetting, and the percentages of viable cells (A) and apoptotic cells (B and C) were determined as described in *Materials and Methods*. Data are the mean \pm

cells were cultured in DMEM supplemented with 10% FBS and 10% conditioned medium, which was prepared by incubation of PBMC from 10 healthy donors in RPMI 1640 medium with 10% FBS for 48 h. The cultures were then maintained for more than 2 mo, and the SCL were cloned by limiting dilution. Thereafter, clones were maintained and replenished with fresh DMEM with 10 or 20% FBS every 3–4 days. One clone (Sy 77) was used in the present experiments. A second RA SCL (RA6/1) was established from synovial tissue in a similar manner. After three to six passages, this RA SCL contained $<2\%$ CD14⁺ cells. An SCL was also established from osteoarthritis synovium (OA5/26) using the same procedure. Skin fibroblast lines (DT1 and FB-HG) were provided by Dr. Heather Wisbey (University of Texas Southwestern Medical Center, Dallas, TX) and Dr. Herman Girschick (University of Texas Southwestern Medical Center).

B cell separation

The PBMC were isolated from heparinized blood of healthy adult volunteers by density sedimentation using Ficoll/Isopaque. CD19-positive B cells were purified using the CEPRATE streptavidin column (CellPro, Bothell, WA), following staining of the cells with subsaturating concentrations of biotinylated mouse anti-human CD19 mAb. The resultant population of B cells contained $>95\%$ CD20-positive B cells.

B cell culture

B cells were cultured in various ways. When cultured alone, B cells (1×10^6) were cultured in 1 ml of RPMI 1640 medium (RPMI 1640 with 10% FBS, 200 U/ml of penicillin G, 10 μ g/ml gentamicin, and 0.3 mg/ml L-glutamine) in 24-well culture plates. Alternatively, B cells (1×10^5) were cultured in 200 μ l of RPMI 1640 medium in 96-well flat-bottom microtiter plates. In some experiments, B cells were stimulated with anti-IgM Ab. To accomplish this, B cells were incubated with 10 μ g/ml of anti-human IgM Ab (DA4.4) conjugated with biotin for 30 min at 4°C and washed twice. Afterward, B cells (1×10^6 in wells of 24-well plates) were cultured for 1 h with streptavidin (10 μ g/ml) to cross-link surface IgM, harvested, washed, and used for experiments. When SCLs and fibroblasts were incubated with B cells, they were initially cultured in 24-well culture plates by incubating SCL (4×10^4) in 1 ml of DMEM (high glucose DMEM including 10% FBS, 200 U/ml of penicillin G, 10 μ g/ml gentamicin, and 0.3 mg/ml L-glutamine) for 1 wk. After this period of time, they became subconfluent, and B cells (1×10^6) were added in 1 ml of RPMI 1640 culture medium. After various periods of time, B cells were harvested by gentle pipetting. Contamination of harvested B cells with SCLs and fibroblasts could be determined by scatter properties using a flow cytometer and was always $<0.5\%$. Alternatively, SCLs and fibroblasts (4×10^3) were cultured in 96-well flat-bottom microtiter plates for 1 wk, and then B cells (1×10^5) were added, incubated, and harvested as described above.

Direct cell-to-cell contact between B cells and SCL was blocked with a 0.4- μ m pore size membrane (Falcon, Becton Dickinson, Franklin Lakes, NJ). For these experiments, SCL (4×10^4) were cultured as described above, and B cells (1×10^6) were added to culture contained in the membrane chamber. In the experiments examining the effect of blocking mAb, SCL (4×10^4) were cultured with 5 μ g/ml of mAb for 60 min before addition of B cells as described above. In some experiments, B cells were stimulated with anti-CD11a/CD18 and/or CD49d/CD29 mAbs. For these experiments, 5 μ g of rabbit anti-mouse IgG mAb were coated in the wells of 24-well culture plates by incubating them in Tris-HCl at pH 9.8 overnight. B cells (1×10^6) were incubated with 2 μ g of mAbs for 30 min, washed once, added to the 24-well plates precoated with rabbit anti-mouse IgG, and cultured in RPMI medium as described above.

Analysis of viable cells

The numbers of viable cells were assessed microscopically after staining with trypan blue.

Detection of apoptosis

The hypotonic propidium iodide staining method was used to detect apoptosis of cells by flow cytometry (24). Harvested B cells were resuspended in hypotonic propidium iodide reagent (0.1% sodium citrate containing 50 μ g/ml of propidium iodide and 0.1% Triton X-100). Within 60–120 min, the stained cells were analyzed by flow cytometry using the FACScan

SEM of five cultures from one of four independent experiments with similar results. *, $p < 0.05$; **, $p < 0.01$ (using Student's *t* test).

Table I. RA SCL rescues B cell apoptosis^a

		D1	D3	D6
Viability (%)				
Expt. 1 (n = 6)	Medium	92.0 ± 0.8	47.0 ± 4.1	24.7 ± 3.9
	+ RA SCL	92.3 ± 1.0	82.0 ± 3.3*	71.0 ± 2.1**
Expt. 2 (n = 5)	Medium	95.3 ± 0.2	77.0 ± 0.5	56.0 ± 2.1
	+ RA SCL	95.7 ± 0.9	89.0 ± 2.8	82.0 ± 0.6**
Expt. 3 (n = 5)	Medium	95.0 ± 0.4	56.1 ± 1.6	28.5 ± 1.1
	+ RA SCL	94.7 ± 0.9	84.9 ± 1.4**	70.5 ± 0.2**
Apoptosis (%)				
Expt. 1 (n = 6)	Medium	7.3 ± 1.3	55.0 ± 6.5	61.7 ± 5.1
	+ RA SCL	6.7 ± 1.2	18.7 ± 2.0	21.3 ± 1.9**
Expt. 2 (n = 5)	Medium	4.3 ± 0.5	25.0 ± 1.0	46.0 ± 1.9
	+ RA SCL	4.7 ± 0.2	7.7 ± 0.9**	17.7 ± 0.9**
Expt. 3 (n = 5)	Medium	5.7 ± 0.5	62.3 ± 1.4	86.5 ± 0.2
	+ RA SCL	5.1 ± 0.7	21.3 ± 1.7**	30.0 ± 0.7**

^a Peripheral blood B cells (1×10^5) were cultured with or without RA SCL (Sy77: 1×10^4) in 96-well culture plates. Percent viability was measured after trypan blue staining. Apoptosis was measured by flow cytometry after hypotonic propidium iodide staining. Data indicates mean \pm SEM from the number of replicates indicated (n). Table shows data from three of seven similar experiments.

*, $p < 0.05$; **, $p < 0.01$: statistical difference between B cells with medium alone and B cells with RA SCL using Student's *t* test.

(Becton Dickinson), and the percentage of hypodiploid cells was determined.

Western blotting


Bcl-2-related protein expression was investigated by Western blotting. B cells (2×10^6), cultured as described above, were collected and lysed with extraction buffer (PBS with 1% Triton X-100, 5 mM EDTA, 100 μ g/ml PMSF, 1 μ g/ml leupeptin, and 0.225 U/ml aprotinin). After 2 h at 4°C, debris was eliminated by centrifugation at 15,000 rpm for 10 min, and the supernatant was collected. After measurement of protein concentration with a protein assay kit (Bio-Rad), each sample was adjusted to 4 μ g/ μ l, then mixed with 2 \times sample loading buffer containing 2-ME (1.5 M) and SDS (4%), and stored at 4°C until analysis. Before analysis, the protein concentration of each sample was adjusted to 2 μ g/ μ l. Forty micrograms of protein (20 μ l of sample) was separated by 12% SDS-PAGE, transferred to polyvinylidene difluoride membranes, and blocked with 0.1% Tween and 5% skim milk overnight. The immunoblots were incubated with anti-Bcl-2 rabbit polyclonal Ab (1/50) or anti-Bcl-x rabbit polyclonal Ab (1/100) in PBS with 1% BSA for 1 h. Goat IgG F(ab')₂ anti-rabbit IgG

conjugated with HRP was used as a second Ab. All immunoblots were detected by enhanced chemiluminescence.

RNA isolation and RT-PCR

RNA was extracted from $1-2 \times 10^6$ B cells using Triazol reagent in accordance with the company's instructions. One microgram of the extracted RNA was treated with DNase I to eliminate DNA and reverse transcribed by SuperScript II reverse transcriptase at 42°C for 70 min using oligo(dT). The PCR was conducted with Taq polymerase using 1-3 μ l of cDNA (1.5 mM MgCl; annealing temperature, 56°C; 30-40 cycles). The primer pair for Bcl-2 was CAGAATCCTCTGGAACCTTGAGG (5') and CTCCAGA CATTCCGAGACC (3'), that of Bcl-x_L was GTGGAAGAGAACG GGGCTGAGG (5') and ATGTGGTGGAGCAGAGAAGG (3'), and that of G6PD was TGACCTCAGCTGCACATCC (5') and CAGTGGC CAATAAGCTCTGG (3'). The PCR products were resolved by electrophoresis on 1.5% agarose gels and identified with ethidium bromide staining. In some cases, PCR-Southern hybridization was conducted. After denaturation of the gel in 1.5 M NaCl and 0.5 M NaOH for 30 min, the PCR

FIGURE 2. The RA SCL rescue B cells from apoptosis more effectively than other cell lines. Peripheral B cells (1×10^5) were cultured with various SCLs or fibroblasts (4×10^3) in 96-well microtiter plates. After 4 days of culture, B cells were harvested with gentle pipetting, and the percentages of viable cells (A) and apoptotic cells (B) were determined as described in *Materials and Methods*. Data are the mean \pm SEM of five cultures from one of two independent experiments with similar results. Sy77, RA synovial stromal cell clone; RA6/1, RA synovial cell line; OA5/26, OA synovial cell line; DT1 and FB-HG, fibroblast cell line from human skin. *, $p < 0.05$; **, $p < 0.01$ (using Student's *t* test).






FIGURE 3. The RA SCL induce Bcl-x_L expression by B cells. *A*, After B cells (1×10^6) were cultured with RA SCL (Sy77; 4×10^4) for 1 or 3 days, expression of Bcl-2 and Bcl-x_L proteins ($40 \mu\text{g}$ of each sample) was examined with Western blotting and compared with that by fresh B cells. Percentages of viable and apoptotic cells were assessed at the same time as described in *Materials and Methods*. Representative data from one of five similar experiments with similar results are shown. *B*, After B cells (1×10^6) stimulated by cross-linking IgM Ab or control Ab were cultured with RA SCL (Sy77; 4×10^4) for 3 days, expression of Bcl-2 and Bcl-x_L proteins ($40 \mu\text{g}$ of each sample) was examined with Western blotting. Percentages of viable and apoptotic cells were assessed at the same time as described in *Materials and Methods*. Representative data from one of two experiments with similar results are shown.

products were transferred onto Zeta-Probe blotting membranes using a vacuum blotting apparatus. ³²P-labeled probes (Bcl-2, GTGACTTCCGATCAGGAAGG; Bcl-x_L, GGTATTGGTGAGTCGGATCG; GAPDH, CCTCCAGACCTGCCTGAGC) specific for PCR products were used to detect PCR products by Southern hybridization. To adjust the amount of cDNA of each sample precisely, G6PD expression was examined first using 32–35 cycles of RT-PCR to amplify 1, 2, and 3 μl of cDNA. After resolving the PCR products on agarose gels and identifying the relevant bands with ethidium bromide, the optimal amounts of cDNA were determined. Bcl-2 and Bcl-x_L expression in this amount of cDNA was examined using 32, 35, 38, and 40 cycles of PCR amplification, and the results in the linear part of the amplification curve are reported in the figures.

Results

The SCL from RA synovium can rescue B cells from apoptosis

When cultured alone, B cells spontaneously underwent apoptosis and died (Fig. 1). However, when B cells were cultured with RA SCL, apoptosis was blocked, and B cell viability was preserved. In each experiment (Table I) significant differences in viability and apoptosis between B cells cultured with medium alone and those with RA SCL were noted after 6 days of incubation. However, in some experiments (Expt. 2), but not others (Expt. 1 and 3), differences in viability were not seen on day 3. Similarly, significant differences between apoptosis noted in cultures with and without




FIGURE 4. The RA SCL rescue B cell from apoptosis more effectively than other cell lines. After B cells (1×10^6) were cultured with various cell lines (4×10^4) for 3 days, expression of Bcl-2 and Bcl-x_L proteins ($40 \mu\text{g}$ of each sample) was examined with Western blotting. Percentages of viable and apoptotic cells were assessed at the same time as described in *Materials and Methods*. Representative data from one of two experiments with similar results are shown. Sy77, RA synovial stromal cell clone; RA6/1, RA synovial cell line; OA5/26, OA synovial cell line; DT1 and FB-HG, fibroblast lines from human skin.

SCL were not always noted on day 3. To determine whether RA SCL were unique in their ability to promote the viability of B cells, the activities of a variety of other cell lines were tested (Fig. 2). All the cell lines rescued B cells from apoptosis to varying degrees, but the viability-promoting activity of RA SCL was significantly better than those of the other cell lines.

Bcl-X is up-regulated by B cells cultured with RA SCL

To determine whether rescue from apoptosis reflected up-regulation of antiapoptotic proteins, expression of Bcl-2 related proteins was investigated before severe apoptosis (Fig. 3A). B cell expression of Bcl-2 protein was unchanged as a result of culture with or without SCL. In contrast, Bcl-x_L protein was not detected in fresh B cells, but was markedly up-regulated within 1 day of culture with SCL, but not when B cells were cultured alone. Increased B cell viability and decreased apoptosis were again observed upon coculture with SCL on day 3, but the changes were less marked because of the short length of the incubation. RA-SCL also rescued anti-IgM-stimulated B cells from apoptosis (Fig. 3B). Moreover,




FIGURE 5. Cell contact is required to induce Bcl-x_L and rescue B cells from apoptosis. B cells (1×10^6) and SCL (4×10^4) were cultured together or separated by a $0.4\text{-}\mu\text{m}$ pore size membrane. Where indicated B cells (1×10^6) were added to the bottom chamber with SCL to examine the influence of soluble factor induced after mixed culture. After a 3-day culture, B cells were analyzed for expression of Bcl-2 and Bcl-x_L protein and were assessed for viability and apoptosis. Representative data from one of four similar experiments with similar results are shown.




FIGURE 6. Roles of CD54 (ICAM-1) and CD106 (VCAM-1) in SCL-mediated up-regulation of Bcl-x_L and inhibition of B cell apoptosis. B cells (1×10^6) were cultured with or without SCL (4×10^4) for 2 days in the presence or the absence of mAbs to CD54 and/or CD106. Afterward, B cells were analyzed for Bcl-2 and Bcl-x_L expression and for viability and apoptosis. Representative data from one of three experiments with similar results are shown.

RA-SCL up-regulated Bcl-x_L, but not Bcl-2, expression by anti-IgM-stimulated B cells. Finally, the impact of coculture with other cell lines on B cell expression of Bcl-x_L was examined (Fig. 4). Bcl-x_L expression was more markedly up-regulated by coculture with RA-SCL. These results show that RA SCL can induce Bcl-x_L protein expression in resting and stimulated B cells more effectively than in other cell lines tested and, in addition, can rescue B cells from apoptosis more effectively.

Direct contact is necessary for RA SCL-mediated rescue of B cell from apoptosis and induction of Bcl-x_L in B cells

As shown in Fig. 5, both up-regulation of Bcl-x_L and rescue of B cells from apoptosis were inhibited when direct contact between RA SCL and B cells was blocked by interposition of a 0.4- μ m pore size membrane. Importantly, B cells in the upper chamber could not be rescued from apoptosis by soluble factors even when they were generated by RA SCL cultured with B cells in the lower chamber. It should be noted that the effect of the membrane separation appeared to be somewhat greater for up-regulation of Bcl-x_L than for rescue from apoptosis, in that there was minimal up-regulation of Bcl-x_L when the cells were separated, but there was some rescue from apoptosis. These results indicate that direct cell-to-cell contact between B cells and SCL cells is importantly involved in both up-regulation of Bcl-x_L by B cells and rescue from apoptosis, with the former more dependent than the latter.

Interactions mediated by CD106 (VCAM-1) and CD49d/CD29 (VLA-4) play a significant role in the rescue of B cells from apoptosis and induction of Bcl-x_L

The RA SCL expresses both CD54 and CD106 (22). Blocking CD54-mediated interactions with an mAb that blocks binding of both CD11a/CD18 and CD11b/CD18, had no effect on the ability of SCL to up-regulate B cell Bcl-x_L and rescue B cells from apoptosis (Fig. 6). By contrast, an mAb to CD106 inhibited both SCL-induced up-regulation of Bcl-x_L and rescue of B cells from apoptosis, but not completely.

Cross-linking CD49/CD29 rescues B cells from apoptosis and induces Bcl-x mRNA and protein

Cross-linking CD49d/CD29 with mAb up-regulated Bcl-x_L mRNA and protein in the absence of RA SCL and also rescued B




FIGURE 7. Stimulation by VLA-4 can induce Bcl-x_L protein and mRNA in B cells. B cells (1×10^6) were cultured for 3 days with mAbs to CD11a/CD18 and/or CD29/CD49d in 24-well culture plates coated with rabbit anti-mouse IgG. Afterward, the expression of Bcl-2 and Bcl-x_L protein was assessed by Western blotting, and the percentages of viable and apoptotic cells were measured as described in *Materials and Methods*, and mRNA for Bcl-2, Bcl-x_L, and G6PD were assessed by RT-PCR as described in *Materials and Methods*. The results of Southern blotting are shown in Expt. 1, and the expression of PCR products by ethidium bromide staining is shown in Expt. 2. One to three micrograms of cDNA was used for each amplification. The number of PCR cycles was modified to ensure that the PCR products obtained were from the linear phase of amplification. The cycle number of each PCR is shown in the figure. Representative data from two of five experiments with similar results are shown.

cells from apoptosis. Cross-linking CD11a/CD18 modestly rescued B cells from apoptosis, but had no impact on expression of Bcl-x_L (Fig. 7). Because of the smaller amount of Bcl-x_L induced by cross-linking CD49d/CD29, it was only detected with a more prolonged exposure. In this circumstance, some Bcl-x_L expression could also be detected in B cells cultured with medium alone (Expt. 1). It should be noted that mRNA and protein expressions were regulated in a qualitatively similar manner, although the quantitative effects were markedly different, with a much greater impact on mRNA levels noted. This suggests that there may be translational or post-translational regulation of Bcl-x_L expression in these cells that is controlled differently from the regulation of mRNA levels.

Discussion

The results of this study indicate that SCL derived from RA synovial tissue can induce Bcl-x_L and rescue peripheral blood B cells from apoptosis, and this ability of RA SCL is greater than that of OA SCL and skin fibroblasts. The data also indicate that a direct physical interaction between RA SCL and B cells mediated by the

coreceptor pair VCAM-1 and VLA-4 provides signals that induce Bcl- x_L and plays a role in the rescue of B cells from programmed cell death. A similar phenomenon in the rheumatoid synovium may contribute to the local accumulation and activation of B cells at this inflammatory site.

It has become apparent that maintenance of the viability of resting B cells is a dynamic process, involving the active countering of proapoptotic mechanisms. One of the major mechanisms to maintain the viability of lymphocytes involves the activity of the Bcl-2 family of molecules that inhibit programmed cell death following mitochondrial disruption (25). Previous results had indicated that a population of SCL isolated from bone marrow or rheumatoid synovium could prevent apoptosis of resting B cells and also stimulate their differentiation into Ab-forming cells (22). These SCL constitutively expressed VCAM-1 (22). The current studies demonstrate that engagement of B cell VLA-4 by VCAM-1 expressed on SCL up-regulates Bcl- x_L , an antiapoptotic Bcl-2 family member, and protects resting B cells from programmed cell death. These results are the first demonstration that engagement of VLA-4 on B cells can induce Bcl- x_L and also the first mechanistic explanation of the means by which interaction with SCL can protect B cells from apoptosis.

In RA synovial tissue, minimal apoptosis of lymphocytes has been noted despite intensive infiltration (5). Previous reports have suggested that soluble factors produced by RA synovial fibroblasts might promote the viability of synovial T cells (26). However, the previous studies clearly showed that the viability of resting B cells could be maintained not by fibroblast cell lines, but, rather, by a specific population of synovial SCL (22). Moreover, the maintenance of B cell viability could not be explained by soluble factors produced by SCL. In the current studies, soluble factors produced by RA SCL could maintain B cell viability modestly, although the major contribution of RA SCL to the rescue of B cells from apoptosis involved direct cell-to-cell contact mediated by VCAM-1.

The mechanism, by which B cells are rescued from apoptosis, has been extensively investigated in secondary lymphoid tissue (20, 27–29). In germinal centers, B cell expression of Bcl- x_L is increased (20, 30). Various combinations of surface molecules and cytokines have been reported to induce Bcl- x_L in B cells or B cell lines. In peripheral B cells, CD40 ligand (CD154) and IL-13 can induce Bcl- x_L (31). CD40 ligand and IgM cross-linking can also induce Bcl- x_L on tonsil B cells and/or WEHI-231 cells (27, 32), whereas the combination of the polyclonal activator, SAC, and IL-10 can also induce Bcl- x_L on tonsil B cells (29). Finally, anti-IgM stimulation induced Bcl- x_L protein and rescued murine A20 cells from apoptosis (33). Besides interactions with Ag and activated T cells, direct contact with follicular dendritic cells (FDC) could contribute to B cell survival. In this regard, FDC and SCL have certain similarities, including the constitutive expression of the adhesion molecules, ICAM-1 and VCAM-1, and their involvement in spontaneous interactions with B cells (22). Because SCL do not express CD40 ligand (22), the possibility that adhesion molecules play a role in maintaining the viability of resting B cells was considered. The results of the mAb blocking experiments indicated that direct cell contact with SCL through VCAM-1-VLA-4 interactions is important in rescuing resting B cells from apoptosis. In view of the central importance of VCAM-1-VLA-4 interactions between B cells and FDC (34–36), a similar process could be involved in up-regulating Bcl- x_L and protecting against apoptosis in the germinal center. We have previously noted the functional and phenotypic similarities between SCL and FDC. However, there are differences between these cell types, including the observation that SCLs do not express CD21 and CD35 on their surface.

It should be noted that the finding that VCAM-1-mediated interactions could not block B cell apoptosis completely suggested that other mechanisms could be involved. SCL are known to produce IL-6, IL-8, GM-CSF, G-CSF, and hyaluronic acid and might also express surface molecules such as CD157 that could contribute to B cell survival (22, 37). Despite the possible contributions of these other molecules, it is clear that VCAM-1-VLA-4 interactions play a central role in the SCL-dependent rescue of resting B cells from apoptosis.

Bcl- x_L expression is largely regulated at the level of gene transcription (25). In this regard, it has been reported that leukemia inhibitory factor induced Bcl- x_L via the gp130 and STAT1 signaling pathway in cardiac myocytes (38). In addition, an erythropoietin-stimulated Jak2 kinase-initiated signal pathway has been reported to inhibit radiation-induced apoptosis and induce Bcl- x_L in the DA3 murine myeloid cell line (39). However, there were no previous reports of a relationship between Bcl- x_L expression and VLA-4 stimulation. VLA-4 stimulation has been reported to result in phosphorylation of focal adhesion kinase, which can then activate phosphatidylinositol 3-kinase and the generation of PIP3 (40–42). Subsequently, Akt (activated form serine/threonine kinase) can be phosphorylated and activated, and provide a cell survival signal (43). Besides this putative pathway of activation, the promoter region of human and mouse Bcl- x_L contains several transcription binding sites, including sites for Ets-1, AP-4, NF-E2, Lyf-1, AP-1, Oct-1, GATA-1, and Evi-1, which are candidates to be involved in up-regulation of transcription (44). Current studies are focusing on the mechanism of Bcl- x_L up-regulation following VLA-4 engagement.

Synovial stromal cells in RA have many potential functions. These cells induce B cell activation and Ig production and also secrete many cytokines and chemokines related to inflammation (22, 37, 45–48). Additionally, they can protect naive B cells from apoptosis by the VLA-4-VCAM-1-dependent mechanism, as described here. These findings suggest that synovial stromal cells may play a central role in propagating the inflammatory response characteristic of RA, especially the accumulation of B cells and their activation to produce Ig locally within the synovium.

Acknowledgments

We thank Dr. Amrie C. Grammer for her careful review of the manuscript.

References

- Harris, E. D. 1990. Rheumatoid arthritis: pathology and implications for therapy. *N. Engl. J. Med.* 322:1277.
- McInnes, I. B., J. Al-Mughales, M. Field, B. P. Leung, F. P. Huang, R. Dixon, R. D. Sturrock, P. C. Wilkinson, and F. Y. Liew. 1996. The role of interleukin-15 in T-cell migration and activation in rheumatoid arthritis. *Nat. Med.* 2:175.
- Iannone, F., V. M. Corrigan, G. H. Kingsley, and G. S. Panayi. 1994. Evidence for the continuous recruitment and activation of T cells into the joints of patients with rheumatoid arthritis. *Eur. J. Immunol.* 24:2706.
- Qu, Z., C. H. Garcia, L. M. O'Rourke, S. R. Planck, M. Kohli, and J. T. Rosenbaum. 1994. Local proliferation of fibroblast-like synoviocytes contributes to synovial hyperplasia. *Arthritis Rheum.* 37:212.
- Firestein, G. S., M. Yeo, and N. J. Zvaifler. 1995. Apoptosis in rheumatoid arthritis synovium. *J. Clin. Invest.* 96:1631.
- Gomez, J., A. Gonzalez, C. Martinez-A, and A. Rebollo. 1998. IL-2-induced cellular events. *Crit. Rev. Immunol.* 18:185.
- Lichtenstein, A., Y. Tu, C. Fady, R. Vesio, and J. Berenson. 1995. Interleukin-6 inhibits apoptosis of malignant plasma cells. *Cell. Immunol.* 162:248.
- Schwarze, M. M., and R. G. Hawley. 1995. Prevention of myeloma cell apoptosis by ectopic bcl-2 expression or interleukin 6-mediated up-regulation of bcl- x_L . *Cancer Res.* 55:2262.
- Armant, M., G. Delespesse, and M. Sarfati. 1995. IL-2 and IL-7 but not IL-12 protect natural killer cells from death by apoptosis and up-regulate bcl-2 expression. *Immunology* 85:331.
- Vella, A. T., S. Dow, T. A. Potter, J. Kappler, and P. Marrack. 1998. Cytokine-induced survival of activated T cells in vitro and in vivo. *Proc. Natl. Acad. Sci. USA* 95:3810.

11. Yonish-Rouach, E., D. Resnitzky, J. Lotem, L. Sachs, A. Kimchi, and M. Oren. 1991. Wild-type p53 induces apoptosis of myeloid leukaemic cells that is inhibited by interleukin-6. *Nature* 352:345.
12. Teague, T. K., P. Marrack, J. W. Kappler, and A. T. Vella. 1997. IL-6 rescues resting mouse T cells from apoptosis. *J. Immunol.* 158:5791.
13. Cohen, S. B., J. B. Crawley, M. C. Kahan, M. Feldmann, and B. M. Foxwell. 1997. Interleukin-10 rescues T cells from apoptotic cell death: association with an upregulation of Bcl-2. *Immunology* 92:1.
14. Poe, J. C., D. H. Wagner, Jr., R. W. Miller, R. D. Stout, and J. Suttles. 1997. IL-4 and IL-10 modulation of CD40-mediated signaling of monocyte IL-1 β synthesis and rescue from apoptosis. *J. Immunol.* 159:846.
15. Meazza, R., S. Basso, A. Gaggero, D. Detotero, L. Trentin, R. Pereno, B. Azzarone, and S. Ferrini. 1998. Interleukin (IL)-15 induces survival and proliferation of the growth factor-dependent acute myeloid leukemia M-07e through the IL-2 receptor β/γ . *Int. J. Cancer* 78:189.
16. Dooms, H., M. Desmedt, S. Vaucaeneghem, P. Rottiers, V. Goossens, W. Fiers, and J. Grooten. 1998. Quiescence-inducing and antiapoptotic activities of IL-15 enhance secondary CD4⁺ T cell responsiveness to antigen. *J. Immunol.* 161:2141.
17. Adachi, S., M. Kubota, Y. W. Lin, A. Okuda, K. Matsubara, Y. Wakazono, H. Hirota, K. Kuwakado, and Y. Akiyama. 1994. In vivo administration of granulocyte colony-stimulating factor promotes neutrophil survival in vitro. *Eur. J. Haematol.* 53:129.
18. Suda, T., H. Hashimoto, M. Tanaka, T. Ochi, and S. Nagata. 1997. Membrane Fas ligand kills human peripheral blood T lymphocytes, and soluble Fas ligand blocks the killing. *J. Exp. Med.* 186:2045.
19. Hashimoto, H., M. Tanaka, T. Suda, T. Tomita, K. Hayashida, E. Takeuchi, M. Kaneko, H. Takano, S. Nagata, and T. Ochi. 1998. Soluble Fas ligand in the joints of patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum.* 41:657.
20. Koopman, G., R. M. J. Keehnen, B. Lindhout, W. Newman, Y. Shimizu, G. A. van Seventer, C. de Groot, and S. T. Pals. 1994. Adhesion through the LFA-1 (CD11a/CD18)-ICAM-1 (CD54) and the VLA-4 (CD49d)-VCAM-1 (CD106) pathways prevents apoptosis of germinal center B cells. *J. Immunol.* 152:3760.
21. Pezzano, M., Y. Li, D. Philip, C. Omene, M. Cantey, G. Saunders, and J. C. Guayden. 1995. Thymic nurse cell rescue of early CD4⁺ CD8⁺ thymocytes from apoptosis. *Cell. Mol. Biol.* 41:1099.
22. Shimaoka, Y., J. F. Attrep, T. Hirano, K. Ishihara, R. Suzuki, T. Toyosaki, T. Ochi, and P. E. Lipsky. 1998. Nurse-like cells from bone marrow and synovium of patients with rheumatoid arthritis promote survival and enhance function of human B cells. *J. Clin. Invest.* 102:606.
23. Arnett, F. C., S. M. Edworthy, D. A. Bloch, D. J. McShane, J. F. Fries, N. S. Cooper, L. A. Healey, S.R. Kaplan, M.H. Liang, H.S. Luthra, et al. 1988. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum.* 31:315.
24. Nicoletti, I., G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi. 1991. A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J. Immunol. Methods* 139:271.
25. Adams, J. M., and S. Cory. 1998. Apoptosis: the Bcl-2 protein family: arbiters of cell survival. *Science* 281:1322.
26. Salmon, M., D. Scheel-Toellner, A. P. Huissoon, D. Pilling, N. Shamsadeen, H. Hyde, A. D. D'Angeac, P. A. Bacon, P. Emery, and N. Akbar. 1997. Inhibition of T cell apoptosis in rheumatoid synovium. *J. Clin. Invest.* 99:439.
27. Tuscano, J. M., K. M. Druey, A. Riva, J. Pena, C. B. Thompson, and J. H. Kehrl. 1996. Bcl-x rather than Bcl-2 mediates CD40-dependent centrocyte survival in the germinal center. *Blood* 88:1359.
28. Zhang, X., L. Li, J. Choe, S. Krajewski, J. C. Reed, C. Thompson, and Y. S. Choi. 1996. Up-regulation of Bcl-XL expression protects CD40-activated human B cells from Fas-mediated apoptosis. *Cell. Immunol.* 173:149.
29. Li, L., S. Krajewski, J. C. Reed, and Y. S. Choi. 1997. The apoptosis and proliferation of SAC-activated B cells by IL-10 are associated with changes in Bcl-2, Bcl-x_L, and Mcl-1 expression. *Cell. Immunol.* 178:33.
30. Krajewski, S., M. Krajewska, A. Shabaik, H. G. Wang, S. Irie, L. Fong, and J. C. Reed. 1994. Immunohistochemical analysis of in vivo patterns of Bcl-x expression. *Cancer Res.* 54:5501.
31. Lomo, J., H. K. Blomhoff, S. E. Jacobsen, S. Krajewski, J. C. Reed, and E. B. Smeland. 1997. Interleukin-13 in combination with CD40 ligand potently inhibits apoptosis in human B lymphocytes: upregulation of Bcl-x_L and Mcl-1. *Blood* 89:4415.
32. Choi, M. S. K., L. H. Boise, A. R. Gottschalk, J. Quintans, C. B. Thompson, and G. G. B. Klaus. 1995. The role of bcl-x_L in CD40-mediated rescue from anti-u-induced apoptosis in WEHI-231 B lymphoma cells. *Eur. J. Immunol.* 25:1352.
33. Bras, A., C. Martínez-A, and E. Baixeras. 1997. B cell receptor cross-linking prevents Fas-induced cell death by inactivating the IL-1 β -converting enzyme protease and regulation Bcl-2/Bcl-x expression. *J. Immunol.* 159:3168.
34. Clark, E. A. K. H. Grabstein, A. M. Gown, M. Skelly, T. Kaisho, T. Hirano, and G. L. Shu. 1995. Activation of B lymphocyte maturation by a human follicular dendritic cell line, FDC-1. *J. Immunol.* 155:545.
35. Freedman, A. S., J. M. Munro, G. E. Rice, M. P. Bevilacqua, C. Morimoto, B. W. McIntyre, K. Rhyngaert, J. S. Pober, and L. M. Nadler. 1990. Adhesion of human B cells to germinal centers in vitro involves VLA-4 and INCAM-110. *Science* 249:1031.
36. Koopman, G., H. K. Parmentier, H. J. Schuurman, W. Newman, C. J. L. M. Meijer, and S. T. Pals. 1991. Adhesion of human B cells to follicular dendritic cells involves both the lymphocyte function associated antigen 1/intercellular adhesion molecule 1 and very late antigen 4/vascular adhesion molecule 1 pathways. *J. Exp. Med.* 173:1297.
37. Takeuchi, E., T. Tomita, T. Toyosaki-Maeda, M. Kaneko, H. Takano, H. Hashimoto, K. Sugamoto, R. Suzuki, and T. Ochi. 1999. Establishment and characterization of nurse cell-like stromal cell lines from synovial tissues of patients with rheumatoid arthritis. *Arthritis Rheum.* 42:221.
38. Fujio, Y., K. Kunisada, H. Hirota, K. Yamauchi-Takahira, and T. Kishimoto. 1997. Signals through gp130 upregulate bcl-x gene expression via STAT1-binding cis-element in cardiac myocytes. *J. Clin. Invest.* 99:2898.
39. Quelle, F. W., J. Wang, J. Feng, D. Wang, J. L. Cleveland, J. N. Ihle, and G. P. Zambetti. 1998. Cytokine rescue of p53-dependent apoptosis and cell cycle arrest is mediated by distinct Jak kinase signaling pathways. *Gene Dev.* 12:1099.
40. Sato, T., K. Tachibana, Y. Nojima, N. D'Avirro, and C. Morimoto. 1995. Role of the VLA-4 molecule in T cell costimulation; Identification of the tyrosine phosphorylation pattern induced by the ligation of VLA-4. *J. Immunol.* 155:2938.
41. Maguire, J. E., K. M. Danahay, L. C. Burkly, and G. A. van Seventer. 1995. T cell receptor- and $\beta 1$ integrin-mediated signals synergize to induce tyrosine phosphorylation of focal adhesion kinase (pp125FAK) in human T cells. *J. Exp. Med.* 182:2079.
42. Chen, H. C., and J. L. Guan. 1994. Association of focal adhesion kinase with its potential substrate phosphatidylinositol 3-kinase. *Proc. Natl. Acad. Sci. USA* 91:10148.
43. Kennedy, S. G., A. J. Wagner, S. D. Conzen, J. Jordan, A. Bellacosa, P. N. Tsichlis, and N. Hay. 1997. The PI3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes Dev.* 11:701.
44. Grillot, D. A. M., M. Gouzale-Gracia, D. Ekhterae, L. Duan, N. Inohara, S. Ohta, M. Seldin, and G. Nunez. 1997. Genomic organization, promoter region analysis, and chromosome localization of the mouse bcl-X gene. *J. Immunol.* 158:4750.
45. Dechanet, J., P. Merville, I. Durand, J. Bancheau, and P. Miossec. 1995. The ability of synoviocytes to support terminal differentiation of activated B cells may explain plasma cell accumulation in rheumatoid synovium. *J. Clin. Invest.* 95:456.
46. Robinson, E., E. C. Keystone, T. J. Schall, N. Gillett, and B. N. Fish. 1995. Chemokine expression in rheumatoid arthritis (RA): evidence of RANTES and macrophage inflammatory protein (MIP)-1 β production by synovial T cells. *Clin. Exp. Immunol.* 101:398.
47. Hosaka, S., T. Akahoshi, C. Wada, and H. Kondo. 1994. Expression of the chemokine superfamily in rheumatoid arthritis. *Clin. Exp. Immunol.* 97:451.
48. Rathanaswami, P., M. Hachicha, M. Sadick, T. J. Schall, and S. R. McColl. 1993. Expression of the cytokine RANTES in rheumatoid synovial fibroblasts: differential regulation of RANTES and interleukin-8 genes by inflammatory cytokines. *J. Biol. Chem.* 268:5834.

Nurse-like Cells from Bone Marrow and Synovium of Patients with Rheumatoid Arthritis Promote Survival and Enhance Function of Human B Cells

Yasunori Shimaoka,* Jeanne F. Attrep,[†] Toshio Hirano,[§] Katsuhiko Ishihara,[§] Ryuji Suzuki,[‡] Tomoko Toyosaki,[‡] Takahiro Ochi,^{*} and Peter E. Lipsky[†]

*Department of Orthopedic Surgery, Osaka University Medical School, 2-2, Yamada-oka, Suita, Osaka 565, Japan; †Harold C. Simmons Arthritis Research Center, Department of Internal Medicine, University of Texas Southwestern Medical Center at Dallas, Dallas, Texas 75235; ‡Division of Molecular Oncology, Department of Oncology, Biomedical Research Center, Osaka University Medical School, 2-2, Yamada-oka, Suita, Osaka 565, Japan; §Department of Immunology, Shionogi Research Laboratories, 5-12-4, Sagisu, Fukushima-ku, Osaka 553, Japan

Abstract

Thymic nurse cells are known to interact with T cells and play a role in their functional maturation. However, the role of nurse cells in B cell maturation and differentiation is less well established, especially at extralymphoid sites. To address this issue, nurse-like cell clones from bone marrow and synovial tissue of patients with RA (RA-NLC) were established and characterized. RA-NLC constitutively expressed CD29, CD49c, CD54 (ICAM-1), CD106 (VCAM-1), CD157 (BST-1), and class I MHC molecules, and secreted IL-6, IL-7, IL-8, granulocyte-macrophage colony-stimulating factor (GM-CSF) and granulocyte colony-stimulating factor (G-CSF). Bone marrow-derived and synovial RA-NLC differed in that the former secreted IL-7 and expressed a greater density of CD157 constitutively and after stimulation with IFN γ , whereas the latter secreted G-CSF and more IL-6. Stimulation of both bone marrow and synovial RA-NLC induced expression of CD40 and class II MHC, but not CD154 (CD40L) or CD35. RA-NLC rescued peripheral B cells from spontaneous apoptosis and promoted survival of B cells for > 4 wk. B cell survival was blocked by antibodies to CD106 or CD157. RA-NLC also increased Ig production from B cells. After long-term culture (4–6 wk) with RA-NLC, but not alone or with fibroblasts, outgrowth of B cells was observed. All B cell lines derived from these cultures had been transformed by EBV, although the RA-NLC themselves were not infected with EBV. Precursor frequency analysis indicated that ~ 1 in 12,500 peripheral B cells could give rise to these EBV-transformed B cell lines upon coculture with RA-NLC. These results indicate that RA-NLC from bone marrow and synovium have the capacity to rescue B cells from spontaneous apoptosis, facilitate Ig production, and promote the outgrowth of EBV-transformed B lymphoblastoid cells. These findings suggest that RA-NLC may play a role in the local and systemic hyperre-

activity of B cells characteristic of rheumatoid arthritis. (*J. Clin. Invest.* 1998. 102:606–618.) Key words: RA • B cells • apoptosis • stromal cells

Introduction

Nurse cells were first recognized in cell suspensions from dissociated thymus (1, 2). Thymic nurse cells form unique complexes with bone marrow-derived T cell precursors, and play an important role in thymocyte maturation and differentiation, eventually leading to the generation of positively (3, 4) or negatively (5) selected T cells that subsequently migrate to the T cell-dependent areas of secondary lymphoid organs. After initial adhesion to thymic nurse cells, thymocytes crawl beneath them in vitro (6, 7). This phenomenon, known as pseudoemperipolesis, is peculiar to nurse cells (8), and has been used to identify nurse-like cells (NLC)¹ at various tissue sites. Previous studies have characterized NLC clones from human dermal tissue (9), and nurse-like stromal cell lines have been established from bone marrow of RA patients (10; Takeuchi, E., T. Tomita, T. Toyosaki-Maeda, H. Hashimoto, M. Kaneko, H. Takano, K. Sugamoto, R. Suzuki, and T. Ochi, manuscript submitted for publication). Moreover, NLC have been established from the synovium of RA patients (11; Tomita, T. E. Takeuchi, T. Toyosaki-Maeda, H. Oku, M. Kaneko, H. Takano, K. Sugamoto, K. Ohzono, R. Suzuki, and T. Ochi, manuscript submitted for publication). A common feature of all NLC was their capacity to support pseudoemperipolesis, similarly to thymic nurse cells. Thus, nurse cells may play an important role in extralymphoid immune responsiveness, as well as promoting maturation of T cells in the thymus.

In addition to their association with T cells, nurse cells have also been noted to interact physically with B cell lymphoma lines (9). This finding suggests that NLC may also promote B cell function, but the role of NLC in B cell maturation and differentiation has not been delineated. Furthermore, the functional potential of extrathymic nurse cells is not known.

RA is characterized by chronic inflammation with infiltration of a variety of inflammatory cells, including those of myeloid origin as well as T and B lymphocytes into the affected synovium. One feature of rheumatoid inflammation is local B cell activation, resulting in the production of large amounts of

Address correspondence to Peter E. Lipsky, M.D., Department of Internal Medicine, University of Texas Southwestern Medical Center at Dallas, 5323 Harry Hines Blvd., Dallas, TX 75235-8884.

Received for publication 19 February 1998 and accepted in revised form 28 May 1998.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.
0021-9738/98/08/0595/13 \$2.00

Volume 102, Number 3, August 1998, 606–618

<http://www.jci.org>

1. *Abbreviations used in this paper:* BST-1, bone marrow stromal antigen-1; DT1, human dermal fibroblast cell line; FDC, follicular dendritic cells; G-CSF, granulocyte colony-stimulating factor; GM-CSF, granulocyte-macrophage colony-stimulating factor; MFI, mean fluorescence intensity; NLC, nurse-like cells; PE, phycoerythrin; RA-NLC, rheumatoid arthritis NLC; VCAM, vascular cell adhesion molecule.