

タミン値の測定, 日本睡眠学会第 34 回定期
学術集会, 大阪 2009 年 10 月

(ユ) 佐藤伸一, 神林崇, 近藤英明, 松渕浪
子, 星野恭一, 清水徹男: REM-NREM 睡眠
時と同様に観察されたウレタン麻酔マウスに
おける呼吸数急上昇とそれに同期した周期
的脳波振幅変化, 日本睡眠学会第 34 回定
期学術集会, 大阪 2009 年 10 月

(ヨ) 徳永純, 佐藤伸一, 神林崇, 清水徹男:
ピエゾ素子センサーを用いた非侵襲的計測
による新生ラット心拍数および自律神経系の
発達過程の解明, 日本睡眠学会第 34 回定
期学術集会, 大阪 2009 年 10 月

(ラ) 清水徹男: 【シンポジウム】GP と専門医
の住み分けをめぐる, 日本睡眠学会第 34

回定期学術集会, 大阪 2009 年 10 月

(リ) 神林崇, 本多真, 吉田祥, 清水徹男:
【シンポジウム】髄液オレキシンと血液 HLA の
補助診断としての適応と限界, 日本睡眠学会
第 34 回定期学術集会, 大阪 2009 年 10 月

(ル) 岩城忍, 清水徹男, 三島和夫: 【シンポ
ジウム】うつ病と不眠の疫学的関連性, 日本
睡眠学会第 34 回定期学術集会, 大阪 2009
年 10 月

(レ) 清水徹男: 【教育講演】うつ病と睡眠障
害, 第 105 回日本精神神経学会学術総会,
神戸 2009 年 8 月

H. 知的財産権の出願・登録状況

なし

Ⅲ. 研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Enomoto M, Tsutsui T, Higashino S, Otaga M, Higuchi S, Aritake S, Hida A, Tamura M, Matsuura M, <u>Kaneita Y</u> , Takahashi K, <u>Mishima K</u>	Sleep-related Problems and Use of Hypnotics in Inpatients of Acute Hospital Wards.	General Hospital Psychiatry	32	276-283	2010
Soshi T, Kuriyama K, Aritake S, Enomoto M, Hida A, Tamura M, Kim Y, <u>Mishima K</u>	Sleep deprivation influences diurnal variation of human time perception with prefrontal activity change: a functional near-infrared spectroscopy study.	PLoS One	5	e8395	2010
Aritake-Okada S, <u>Kaneita Y</u> , <u>Uchiyama M</u> , <u>Mishima K</u> , Ohida T	Non-Pharmacological Self-Management of Sleep Among the Japanese General Population.	Journal of Clinical Sleep Medicine	5	464-469	2009
Yamahara M, Noguchi T, Okawa M, <u>Yamada N</u>	The relationship between subjective sleep disturbance and complexity of 24-hour activity utilizing fractal theory in psychiatric inpatients.	Sleep and Biological Rhythms	7(1)	11-16	2009
Mitsunari Habukawa , <u>Naohisa Uchimura</u> ,Masaharu Maeda, Nozomu Kotorii ,Hisao Maeda	Sleep Findings in Young Adult Patients with Posttraumatic Stress Disorder.	Biol Psychiatry	62	1179-1182	2007
Ozone M, Yagi T, Itoh H, Tamura Y, <u>Inoue Y</u> , et al	Effects of Zolpidem on Cyclic Alternating Pattern, an Objective Marker of Sleep Instability, in Japanese Patients with Psychophysiological Insomnia: A Randomized crossover Comparative Study with Placebo.	Pharmacopsychiatry	41(3)	106-114	2008.05
Oka Y, Suzuki S, <u>Inoue Y</u>	Bedtime Activities, Sleep Environment, and Sleep/Wake Patterns of Japanese Elementary School Children.	Behavioral Sleep Medicine	6(4)	220-233	2008.10
Léger D, Poursain B, Neuberger D, <u>Uchiyama M</u>	An international survey of sleeping problems in the general population.	Curr Med Res Opin	24	307-17	2008

IV. 研究成果の刊行物・別刷

Sleep-related problems and use of hypnotics in inpatients of acute hospital wards

Minori Enomoto, B.Sc.^{a,c}, Takako Tsutsui, Ph.D.^b, Sadanori Higashino, Ph.D.^b, Masaaki Otaga, M.S.W.^b, Shigekazu Higuchi, Ph.D.^a, Sayaka Aritake, Ph.D.^a, Akiko Hida, Ph.D.^a, Miyuki Tamura, Ph.D.^a, Masato Matsuura, M.D., Ph.D.^c, Yoshitaka Kaneita, M.D., Ph.D.^d, Kiyohisa Takahashi, M.D., Ph.D.^e, Kazuo Mishima, M.D., Ph.D.^{a,*}

^aDepartment of Psychophysiology, National Institute of Mental Health, National Center of Neurology and Psychiatry, Tokyo 187-8553, Japan

^bDepartment of Social Services, National Institute of Public Health, Ministry of Health, Labour and Welfare, Saitama 351-0197, Japan

^cSection of Biofunctional informatics, Graduate School of Health Care Sciences, Tokyo Medical and Dental University, Tokyo 113-8519, Japan

^dDivision of Public Health, Department of Social Medicine, Nihon University School of Medicine, Tokyo 173-8610, Japan

^eAino University, Osaka 567-0012, Japan

Received 6 November 2009; accepted 26 January 2010

Abstract

Objective: Although sleep disorders are highly prevalent among patients with physical disorders, only limited information is available about the actual status of sleep-related problems in inpatients of acute hospital wards. We conducted a multicenter cross-sectional observational survey investigating the prevalence of sleep disorders and use of hypnotic-sedative drugs among inpatients of acute wards in 44 general hospitals in Japan.

Method: Questionnaire-, actigraph- and observation-based sleep evaluations were simultaneously performed in 557 adult inpatients [mean age 72.8±12.8 (S.D.) years] of acute wards during a one-month period in July 2007.

Results: Of the 421 patients with data available, 22.3% had at least one of the following sleep disorders: sleep apnea syndrome, restless legs syndrome, periodic limb movement disorder and nocturnal behavior disorder. Similarly, 62.7% had insomnia, 6.9% had severe daytime sleepiness and 12.8% had other sleep-related symptoms. Only 13.8% were free of any sleep-related problem. Although 33.7% of insomnia patients were taking hypnotic-sedative drugs, 65.2% of them complained of residual insomnia symptoms.

Conclusion: The findings obtained in this study have revealed the remarkably high prevalence of sleep-related problems experienced by inpatients of acute hospital wards in Japan. Proper diagnosis of sleep disorders should be made among patients with physical disorders.

© 2010 Elsevier Inc. All rights reserved.

Keywords: Sleep disorders; Insomnia; Acute hospital wards; Physical illness; Hypnotic use

1. Introduction

Sleep disorders, including insomnia, sleep apnea syndrome (SAS), restless legs syndrome (RLS) and periodic limb movement disorder (PLMD), are highly prevalent and particularly common in elderly patients with physical disorders. Sleep disorders reduce patients' quality of life (QOL) by causing symptoms such as daytime sleepiness and cognitive impairment and may also exacerbate underlying disorders by inhibiting respiratory, cardiovascular and

* Corresponding author. Tel.: +81 42 346 2014; fax: +81 42 346 2072.

E-mail addresses: minori@ncnp.go.jp (M. Enomoto), tsutsui@niph.go.jp (T. Tsutsui), sadanori@u-shizuoka-ken.ac.jp (S. Higashino), otaga@niph.go.jp (M. Otaga), shige@ncnp.go.jp (S. Higuchi), sayaca@ncnp.go.jp (S. Aritake), hida@ncnp.go.jp (A. Hida), tamura@ncnp.go.jp (M. Tamura), matsu.mtec@tmd.ac.jp (M. Matsuura), kaneita@med.nihon-u.ac.jp (Y. Kaneita), ktaka@ncnp.go.jp (K. Takahashi), mishima@ncnp.go.jp (K. Mishima).

metabolic functions. In one study of older patients in a skilled-care geriatric hospital in Japan, the presence of insomnia was associated with a higher risk of mortality during the 2-year follow-up period [1].

The prevalence of these sleep disorders increases with age [2], and the high incidence of physical disorders among the elderly population is a contributing factor. Previous epidemiologic studies have revealed that the prevalence of insomnia among the general population is 10.2–48.0% [3–6], and insomnia frequently occurs in association with chronic pain disorders, respiratory diseases and neurological diseases [7]. SAS, RLS or PLMD also frequently coexists with various physical diseases including hypertension [8], ischemic heart disease [9,10], chronic kidney failure [11], iron-deficiency anemia [12] and neurological diseases such as Parkinson's disease [13]. It is also noteworthy that medications used for the treatment of sleep disorders may worsen physical disorders; for example, most standard hypnotics benzodiazepines cause *sleep apnea* by reducing the muscle tone of the upper respiratory tract during sleep [14].

The fact that physical and sleep disorders can coexist at a high frequency should always be taken into account when making an accurate diagnosis and developing a treatment strategy that provides a favorable risk-benefit balance. Nevertheless, we currently have only limited information about the actual status of sleep-related problems experienced by inpatients of acute hospital wards. Thus, the objectives of the present study were to investigate the breakdown and prevalence of sleep disorders and use of hypnotic-sedative drugs in acute ward inpatients and to identify problems in the clinical practice of sleep medicine.

2. Methods

2.1. Subjects and method

Study subjects who were 20 years of age or more were randomly selected from among the inpatients of acute hospital wards, excluding psychiatric and tuberculosis wards, of 44 general hospitals in Japan. The patients'

identities were coded at each hospital ward, and then patients were randomly sampled. The investigation was carried out among 557 subjects [316 males, 241 females; mean age, 72.8±12.8 (S.D.) years; range 22–96 years] who had provided informed consent or whose family member had provided informed consent, simultaneously at all hospitals during a period of 1 month in July 2007. Each patient's primary disorder was classified according to the International Classification of Diseases and Related Health Problems Version 10 (*ICD-10*) (Table 1). The ethics committee at each research site approved the present study.

2.2. Investigation methods

The investigation was conducted over 2 days for each patient to check his or her sleep condition and details of treatment. The investigation consisted of subjective sleep evaluation using a self-administered questionnaire (Table 2), objective sleep evaluation by actigraphy, observational sleep evaluation by nursing staff and a survey of medication use as recorded in the medical records.

The questionnaire was designed to identify the presence of insomnia, SAS, RLS, PLMD, nocturnal behavior disorder (NBD), daytime sleepiness and nocturnal sleep-related symptoms. In the questionnaire, Q1–Q6 were completed by the patients, and Q7 and Q8 were completed by medical staff. Although NBD can be further divided into nocturnal delirium, REM sleep behavior disorder, behavioral and psychological symptoms of dementia and other symptoms, these disorders were not distinguished in view of the primary objective of the present study and technical restrictions.

For objective sleep evaluation, subjects were asked to wear an actigraph [Lifecorder PLUS (LC), Suzuken, Nagoya, Japan] [15] on their waist for two consecutive days for continuous recording of the intensity of activity. Total sleep time (TST; the sum of all sleep time during time in bed), total wake time (TWT; the sum of all wake time during time in bed) and sleep efficiency (SE; the percentage of TST relative to time in bed) were then calculated from the LC data. Time in bed (TIB) was defined as the time during

Table 1
Illness identified in enrolled patients

System organ/disease class	Total 557 (100%)	SAS, RLS, PLMD and NBD 94 (100%)	Insomnia			Good Sleep 63 (100%)
			Improved 31 (100%)	Untreated 175 (100%)	Not-Improved 58 (100%)	
Diseases of the circulatory system	140 (25.1)	20 (21.3)	7 (22.6)	44 (25.1)	9 (15.5)	15 (23.8)
Neoplasms	127 (22.8)	19 (20.2)	5 (16.1)	47 (26.9)	26 (44.8)	8 (12.7)
Diseases of the respiratory system	68 (12.2)	11 (11.7)	3 (9.7)	17 (9.7)	8 (13.8)	9 (14.3)
Diseases of the digestive system	62 (11.1)	13 (13.8)	2 (6.5)	21 (12.0)	7 (12.1)	8 (12.7)
Diseases of the nervous system	45 (8.1)	11 (11.7)	2 (6.5)	9 (5.1)	3 (5.2)	5 (7.9)
Diseases of the genitourinary system	16 (2.9)	4 (4.3)	1 (3.2)	5 (2.9)	1 (1.7)	3 (4.9)
Diseases of the musculoskeletal system and connective tissue	14 (2.5)	2 (2.1)	1 (3.2)	7 (4.0)	0 (0.0)	3 (4.9)
Certain infectious and parasitic diseases	8 (1.4)	0 (0.0)	1 (3.2)	3 (1.7)	1 (1.7)	0 (0.0)
Other diseases	77 (13.8)	14 (14.9)	9 (29.0)	22 (12.6)	3 (5.2)	12 (19.0)

SAS; sleep apnea syndrome, RLS; restless legs syndrome, PLMD; periodic limb movement disorder, NBD; nocturnal behavior disorder.

Table 2

Question items and percentages of respondents in the analyzed 421 inpatients

Items	1)	2)	3)	4)
Q1. How long did it take from light off until you went to sleep? 1) less than 15 minutes 2) 15-29 minutes 3) 30-59 minutes 4) more than 60 minutes	50.4	20.4	14.5	14.7
Q2. How many times did you awake during last night? 1) none 2) 1-2 times 3) 3-4 times 4) more than 5 times	21.4	35.9	24.9	17.8
Q3. What time did you get up this morning (h:min)?	22.6*	77.4		
Q4. Did you get up in the morning unrefreshed or nonrestored? 1) good 2) fair 3) insufficient 4) poor	38.7	37.1	19.2	5.0
Q5. Do you have daytime sleepiness?*** 1) none 2) some 3) moderate 4) severe	22.8	22.8	47.5	6.9
Q6. Did you experience any of the following symptoms during last night (completed by a patient)				
Q6-a creeping sensation or restless discomfort in the limbs	1) yes 2) no	5.9	94.1	
Q6-b legs or arms jerk	1) yes 2) no	2.4	97.6	
Q6-c hot flash	1) yes 2) no	4.8	95.2	
Q6-d night sweat	1) yes 2) no	6.9	93.1	
Q6-e palpitation	1) yes 2) no	1.2	98.8	
Q6-f anxiety or panic	1) yes 2) no	1.0	99.0	
Q6-g sleep paralysis	1) yes 2) no	0.0	100.0	
Q6-h nightmare	1) yes 2) no	3.1	96.9	
Q7. Did the patient experience any of the following symptoms during last night (completed by nursing staffs)				
Q7-a loud snoring, or apnea lasting for 10 seconds or longer	1) yes 2) no	10.0	90.0	
Q7-b periodic legs or arms jerk	1) yes 2) no	2.1	97.9	
Q7-c sleep-talking, delirium or abnormal behaviors such as wandering	1) yes 2) no	6.9	93.1	
Q8. Whether or not the patient took any hypnotic-sedative drug(s) for treatment of insomnia within the past one week and the name of the drug(s) if any (completed by nursing staffs) 1) yes 2) no Name of drugs []	27.6	72.4		

* Patients who woke up 30 minutes or earlier than the desired time without falling asleep again (Q3).

** answered at 2 pm.

which patients were supposed to be in bed as specified by each hospital ward, and specifically the time from “lights out” to the time at which patients were expected to wake. Mean TIB was approximately between 9 p.m. and 6 a.m. Observations by the nursing staffs on each of the wards confirmed that the patients were in bed during TIB on the evenings of the study.

For the observational sleep evaluations, several nursing staffs alternated in order to record continuously the subjects' sleep states. Opening and closing of eyes, breathing, movement and any unusual behavior of the subjects were observed and recorded at a distance so as to not disturb the subjects.

2.3. Differential diagnosis of sleep disorders

The diagnostic flow for the patients included in the investigation is shown in Fig. 1. Some of the preselected subjects ($n=136$) were either excluded from data analyses or could not participate due to reasons such as sudden change in physical condition such as fever, severe dementia, consciousness disturbance due to organic brain damages, need for emergency examination, hospital transfer or discharge or

due to missing data on their amount of physical activity. As a result, a total of 421 patients comprised the analysis population [228 males, 193 females; mean age, 72.5 ± 12.6 (S.D.) years; range 22-96 years]. The number of respondents for each question item is shown in Table 2.

Patients were initially examined for the presence of SAS (positive answer to Q7-a), RLS (positive answer to Q6-a), PLMD (positive answer to Q6-b or Q7-b) or NBD (positive answer to Q7-c). Those who were NOT diagnosed with SAS, RLS, PLMD or NBD were subsequently examined for the presence of insomnia. Patients were judged as having insomnia when the subjective sleep investigation indicated the presence of any one of the following:

- i. Disturbances of initiating sleep (DIS): Q1, the answer indicates 30 min or more.
- ii. Disturbances of maintaining sleep (DMS): Q2, the answer indicates three times or more.
- iii. Early morning awakening (EMA): Q3, the answer indicates wake time 30 minutes or earlier than the desired time without falling asleep again.
- iv. Non-restorative sleep (NRS): Q4, the answer indicates insufficient or poor sleep.




Fig. 1. Diagnostic flow of the subjects in this study. See text and Table 2 for explanation.

The subjects were also divided into the following four groups according to the presence or absence of insomnia and use or not of hypnotic-sedative drugs for insomnia treatment: the good sleep (GS) group consisting of those without insomnia and taking no medication, the improved (I) group consisting of those without insomnia and taking medication (s), the untreated (UT) group consisting of those with insomnia but taking no medication, and the not-improved (NI) group consisting of those with insomnia and taking medication(s). Of these groups, the I, UT and NI groups were grouped together and defined as the insomnia group (Fig. 1).

2.4. Daytime sleepiness

The 421 patients were examined for the presence or absence of daytime sleepiness according to the following criteria: Q5, the answer indicates the presence of moderate or severe sleepiness.

2.5. Sleep-related symptoms

The 421 patients were examined for the presence or absence of other sleep-related symptoms, such as hot flashes in the foot or body (Q6-c), night sweats (Q6-d), palpitations (Q6-e), anxiety and panic (Q6-f), sleep paralysis (Q6-g) and nightmares (Q6-h).

2.6. Statistical analysis

One-way analysis of variance followed by Tukey's multiple comparison tests was used to identify significant differences in sleep parameters (TST, TWT and SE) among

the insomnia group and GS group. Sleep parameters were also compared between each sleep disorder group and the GS group using a two-tailed Student's *t* test. Analysis values are expressed as mean±S.D. Multiple logistic regression analysis was carried out to calculate the odds ratio (OR) and 95% confidence interval (CI) for assessing the association of primary disorders, sleep disorders and use of hypnotic-sedative drugs with severe sleepiness. Presence of severe sleepiness was used as the dependent variable, and primary disorders, sleep disorders and use of hypnotic-sedative drugs were used as independent variables. We performed multiple logistic regression analyses to control for all sociodemographic (sex and age) and other factors. Statistical significance was set at $P < .05$. All analyses were made using SPSS 11.5 for Windows.

3. Results

3.1. Prevalence of sleep disorders

The breakdown of the diagnoses of sleep disorders is shown in Fig. 1. Of the 421 inpatients, 42 (10.0%, M/F=29/13) had SAS, 25 (5.9%, 14/11) had RLS, 17 (4.0%, 11/6) had PLMD and 29 (6.9%, 19/10) had NBD. A total of 94 (22.3%) had at least one of the four sleep disorders. Seventeen patients had two sleep disorders concurrently.

Of the 421 inpatients, 58 (13.8%, NI) and 175 (41.6%, UT) complained of insomnia symptoms. A total of 264 (62.7%), including the NI, UT and I (31, 7.4%) groups were given a diagnosis of insomnia. The most common insomnia

Table 3

Comparison of objective sleep parameters determined by LC in the insomnia and good sleep patients

	SAS n=42	P	RLS n=25	P	PLMD n=17	P	NBD n=29	P	Insomnia				Good sleep n=63		
									Untreated n=175	P	Improved n=31	P		Not-improved n=58	
TST (min)	367.6±119.2	.06	331.9±117.7	0	354.6±111.5	.01	359.8±126.1	.04	369.2±102.5	.04	400.7±118.4	n.s	399.7±91.0	n.s	409.4±102.4
TWT (min)	172.4±119.2	.05	208.1±117.7	0	185.4±111.5	.01	180.2±126.1	.04	170.3±102.3	.03	139.4±118.4	n.s	140.3±91.0	n.s	129.3±103.3
SE (%)	68.1±22.1	.05	61.5±21.8	0	65.7±20.6	.01	66.6±23.4	.04	68.4±19.0	.03	74.2±21.9	n.s	74.0±16.9	n.s	76.1±19.1

Value are expressed as mean±S.D..

P value vs. Good sleep group.

n.s; not significant.

symptom was DMS (60.1%), followed by DIS (41.2%), EMA (33.9%) and NRS (31.8%). Only 63 (15.0%) were free of the above-mentioned sleep disorders and were assigned to the GS group.

3.2. Objective sleep parameters

Sleep parameters in each sleep disorder group are summarized in Table 3. There were significant differences in TST [$F(3,323)=3.24$, $P=.022$], TWT [$F(3,323)=3.28$, $P=.021$] and SE [$F(3,323)=3.31$, $P=.020$] among the insomnia group and GS group. TST ($P=.039$) was significantly shorter and TWT ($P=.033$) and SE ($P=.032$) were significantly longer in the NI group than in the GS group. Patients with RLS ($P<.01$) and NBD ($P<.05$) also presented a significantly shorter TST, significantly longer TWT and significantly lower SE than those in the GS group. A similar tendency was observed for patients with SAS or PLMD ($P<.06$). On the other hand, we found no significant differences in the sleep parameters between the medicated group (the I or NI group) and the GS group, regardless of whether or not any subjective improvement was observed.

3.3. Daytime sleepiness

Of the 421 inpatients, 229 (54.4%) experienced moderate to severe sleepiness and 29 (6.9%) experienced severe sleepiness. Severe sleepiness was commonly observed in those with sleep disorders; it was most commonly observed in patients with multiple sleep disorders (27.8%, 5/18), followed by those with PLMD (18.2%, 2/11), SAS (17.9%, 5/28) and NBD (17.7%, 3/17). Multiple logistic regression analysis revealed that SAS (adjusted OR=3.78, 95% CI, 1.24–11.53, $P<.05$) and PLMD (adjusted OR=5.93, 95% CI, 1.50–23.4, $P<.05$) showed a significantly positive association with the presence of severe sleepiness.

3.4. Other sleep-related symptoms

Of the 421 inpatients, 19 (4.5%, M/F=7/12) had hot flashes, 29 (6.9%, 13/16) had night sweats, 5 (1.2%, 1/4) had palpitations, 4 (1.0%, 2/2) had anxiety or panic and 13 (3.1%, 7/6) had nightmares. None of the patients experienced sleep paralysis.

3.5. Prevalence of use of hypnotic-sedative drugs

Of the 421 inpatients, 116 (27.6%) were taking some kind of hypnotic-sedative drug for the treatment of insomnia symptoms. The breakdown of the prescribed drugs was as follows: benzodiazepine hypnotics including zolpidem and zopiclone accounted for 73.2% (26.1% for ultrashort-acting, 30.6% for short-acting and 16.5% for intermediate-acting), benzodiazepine anxiolytic accounted for 5.8%, antipsychotics accounted for 15.6% and other drugs accounted for 5.2% of all prescribed drugs. In the insomnia group, those receiving medication therapy for insomnia only accounted for 33.7% (the I+NI group). Two thirds of the patients receiving medication therapy (65.2%, corresponding to the NI group) complained of persistent insomnia symptoms. In addition, 36.0% of RLS patients, 29.4% of PLMD patients, 26.2% of SAS patients and 17.2% of NBD patients were taking at least one of the above hypnotic-sedative drugs.

4. Discussion

This is the first multicenter study investigating the prevalence of sleep disorders in inpatients of acute wards in general hospitals. Sleep disorders are extremely common disorders among community residents, and are even more so among patients with underlying physical diseases as in the subjects of the present study. Insomnia, as well as other sleep disorders, while frequently thought to be transitory or secondary to a physical disease, can become prolonged without appropriate treatment in the early stages. Furthermore, chronic sleep disorders can exacerbate lifestyle-related diseases such as hypertension and diabetes, and increase the risk of psychiatric symptoms such as depression and anxiety, not to cause subjective distress [16,17]. Many sleep disorders go undetected and are not appropriately treated in clinical practice. Therefore, this study was conducted to alert practitioners of sleep disorders to this situation, by shedding more light on their current status in general medical practice.

In the present study, we investigated the prevalence of sleep disorders and the use of hypnotic-sedative drugs in 421 inpatients with mean age of 72.5 years by questionnaire-, actigraph- and observation-based sleep evaluations, and have revealed a high prevalence of diverse types of sleep disorders

in the study population. SAS, RLS, PLMD, NBD and insomnia, in particular, were highly prevalent (10.0, 5.9, 4.0, 6.9 and 62.7%, respectively). The inpatients also suffered from various sleep-related symptoms (1.0–6.9%, except for sleep paralysis), which are common conditions with physical disorders and which could cause disrupted sleep [18–21]. In fact, the patients with these sleep disorders also showed poor sleep parameters recorded by actigraphy, which objectively indicates that they have poor-quality sleep during the night. Consequently, of the 421 patients, only 13.8% were free of any type of sleep disorder diagnosed, severe daytime sleepiness or sleep-related symptoms, revealing that sleep-related problems are very common clinical problems among inpatients of acute hospital wards.

Due to restrictions on the disclosure of personal information, the only information available regarding the underlying diseases of the patients was the names of the primary diseases according to the major classification of the *ICD-10*. We were thus unable to analyze respective medical conditions that are commonly associated with these sleep disorders, such as chronic pain, cardiovascular diseases, chronic renal failure, hemodialysis and iron deficiency anemia.

The prevalence of SAS and RLS is generally high in elderly people and patients with physical disorders. However, even though the mean age of our patients was high (72.5 years) and they had physical disorders in the exacerbation phase, contrary to our expectations, the prevalence of SAS and RLS was not higher in the study population than in community dwellers of previous studies. For example, the prevalence of SAS in middle-aged to elderly people has been shown to be 9–10% in males and 4–10% in females [22,23], which is comparable to that in the present study population (10% in the entire population, 12.7% in males, 6.7% in females). In the present study, patients were defined as having SAS if they reported loud snoring or apnea lasting for 10 seconds or more, because loud snoring is the most prominent symptom of upper airway resistance syndrome, which is included in the category of SAS [7,24]. Nevertheless, the prevalence of SAS patients including those who snored loudly in the present study was similar to that in the general population. Similarly, a large-scale survey which employed a self-administered questionnaire and used a definition of RLS similar to that in the present study has reported that the prevalence of RLS among Japanese people aged 70 years or more is 4.1% (3.4% in males, 4.6% in females), which is practically identical to that in the present study (5.9% in total, 6.1% in males, 5.7% in females) [25]. Furthermore, the frequency of NBD was as low as 6.9%, despite the occurrence rate of delirium per admission varying between 11 and 42% [26]. The low NBD frequency of the present study compared to that of all previous studies is thought to be because patients with severe physical conditions or with organic brain damages were excluded from the analyses.

In many of the epidemiologic studies on the prevalence of sleep disorders, sleep evaluation is performed during a period of one week to one month. The fact that sleep evaluation in this study was performed on a single night might have held down the prevalence of sleep disorders. However, since the physical status of the inpatients of acute hospital wards can change in a very short period of time and their sleep condition is also subject to change, we assumed that the results obtained from a long investigation period would not properly reflect the actual status of their sleep-related problems. Extension of the duration for determining the presence or absence of sleep disorders may result in a dramatic increase in the prevalence of the sleep disorders in inpatients of acute hospital wards.

Patients with physical disorders, especially with advanced age, are generally vulnerable to insomnia [27–29]. We have found that approximately two thirds (62.7%) of the representative patients in acute wards in Japan are suffering from insomnia. It was confirmed not only from the subjective complaints of patients but also from the objective sleep evaluation that the quality of sleep for patients with insomnia receiving no treatment or who had other sleep disorders was significantly lower than that for patients in the GS group (Table 3). A survey among 1500 community dwellers aged 55–84 years in the United States has demonstrated that the quality of sleep decreases in proportion to an increase in the number of physical disorders suffered [27]. Several studies have also reported a high prevalence (34–69%) of insomnia in outpatients of primary care clinics or regular inpatients with acute or chronic physical disorders [30–33]. The findings of the present study for acute ward inpatients are consistent with those obtained in the previous studies in spite of shorter-term sleep evaluation.

In many cases of sleep disorders, daytime sleepiness often occurs to compensate for low-quality sleep during the night. In the present study, 47.5% of the patients experienced mild or severe sleepiness and 6.9% experienced severe sleepiness, which was particularly high in those with multiple sleep disorders, including SAS, RLS, PLMD and NBD. The results of multiple logistic regression analysis indicated that severe sleepiness is significantly associated with SAS and PLMD, and not with an underlying disease or type of hypnotic-sedative drug.

Only one-third (33.7%) of the patients with insomnia included in the present investigation received treatment for insomnia symptoms. In addition, two-thirds (65.2%) of the patients receiving medication therapy complained of residual insomnia symptoms. The relatively low frequency of patients prescribed hypnotic-sedative drugs in the present study, which is very similar to that reported in the Meissner's study [30], suggests the possibility that physicians are not fully aware of the presence of insomnia in their patients.

The prescribed drugs mainly consisted of benzodiazepine hypnotics including intermediate-acting agents and antipsychotics. Caution should always be exercised when

using these hypnotic-sedative drugs in inpatients with physical disorders, especially in elderly patients. This is because elderly patients present a poor risk-benefit balance for hypnotic-sedative drugs due to such reasons as decreased drug metabolizing capacity, increased drug sensitivity, risk of fall and fracture or suppressed mental function, and worsening of underlying diseases induced by medication [34–37].

Moreover, administered hypnotic-sedative drugs may be ineffective or even worsen underlying diseases unless sleep disorders are properly diagnosed. In fact, 23.8% of the patients with SAS were prescribed hypnotic-sedative drugs including benzodiazepines and 36.0% of the patients with RLS were taking hypnotic-sedative drugs other than clonazepam. These results suggest that medications that are not necessarily appropriate for treatment of individual patients' sleep disorders are often selected in actual clinical practice, possibly causing a reduction in the patients' ADL and QOL.

Several limitations should be noted when interpreting the results of the present study. First, as elderly patients aged 65 years or more accounted for a large portion (76.0%) of the 421 inpatients, it is speculated that the high prevalence of sleep-related problems observed in the patients of the present investigation were associated with not only sleep disorders attributable to physical disorders but also age-related changes in sleep property.

Second, one-fourth (24.4%) of the initially enrolled 557 patients were excluded. Patients who were unable to answer questions on the day of the survey because of a change in their physical condition (e.g. fever, consciousness disturbance or need for emergency examination) or those patients with missing data due to interruptions in LC data collection were excluded. Some of these excluded patients might have developed some type of sleep disorder during their stay in hospital.

Third, insomnia defined in the present study is different from insomnia that meets the general criteria of the International Classification of Sleep Disorders, second edition (ICSD-2) [7], because we did not consider the presence or absence of "daytime impairment related to the nighttime sleep difficulty". This investigation item was not included in the present study because it was difficult to determine whether the patients' diverse psychosomatic symptoms observed during the daytime were attributable to insomnia or physical disorders.

Fourth, the questionnaire employed in the present study has not been validated. A set number of items taken from the original were configured so as to reduce the burden on inpatients who were in poor physical condition. Therefore, the questionnaire can only suggest the possibility of certain disorders such as SAS, PLMD and RLS; it does not predict the presence of these disorders with high accuracy. However, the frequency of sleep disorders and the percentage of patients exhibiting symptoms of insomnia found in the present study closely resemble the data of several other

studies. This is thought to be indirect evidence that, to a certain degree, the survey items work effectively to detect patients suffering from sleep disorders.

Fifth, the sleep/wake scoring algorithm used for the LC data in the present study has been validated for a sample of healthy young subjects [15], but not for elderly subjects with physical disorders, as in the present study's sample. However, as the results demonstrate, meaningful differences were detected in the sleep parameters calculated with this algorithm for total sleep time, total wake time, and efficiency of sleep between the UT group with insomnia and the GS group. Given this, the clinical application of the LC and sleep/wake scoring algorithm for the subjects of the present study can be considered a sound approach to a certain degree.

5. Conclusion

In the present study, which initially involved 557 inpatients who had been admitted to acute hospital wards in 44 general hospitals, we have revealed an extremely high prevalence of sleep disorders using subjective and objective sleep evaluation scales, and have also indicated several problems in the current practice of sleep medicine. Proper diagnosis of sleep disorders should be made while being aware of the high prevalence of sleep disorders among elderly patients with physical disorders, and a treatment strategy that provides a favorable risk-benefit balance must be developed.

Acknowledgments

This study was supported by a Grant-in-Aid for Cooperative Research from the Ministry of Health, Labor, and Welfare of Japan (H19-kokoro-ippan-013, H20-tyoju-ippan-001).

References

- [1] Manabe K, Matsui T, Yamaya M, Sato-Nakagawa T, Okamura N, Arai H, et al. Sleep patterns and mortality among elderly patients in a geriatric hospital. *Gerontology* 2000;46:318–22.
- [2] Wolkove N, Elkholy O, Baltzan M, Palayew M. Sleep and aging: 1. Sleep disorders commonly found in older people. *CMAJ* 2007;176: 1299–304.
- [3] Bixler EO, Kales A, Soldatos CR, Kales JD, Healey S. Prevalence of sleep disorders in the Los Angeles metropolitan area. *Am J Psychiatry* 1979;136:1257–62.
- [4] Ford DE, Kamerow DB. Epidemiologic study of sleep disturbances and psychiatric disorders. An opportunity for prevention. *JAMA* 1989; 262:1479–84.
- [5] Kim K, Uchiyama M, Okawa M, Liu X, Ogihara R. An epidemiological study of insomnia among the Japanese general population. *Sleep* 2000;23:41–7.
- [6] Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev* 2002;6:97–111.

- [7] American Academy of Sleep Medicine. International classification of sleep disorders :Diagnostic and coding manual. 2nd ed. Westchester: Illinois; 2005.
- [8] Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000;342:1378–84.
- [9] Javaheri S, Parker TJ, Liming JD, Corbett WS, Nishiyama H, Wexler L, et al. Sleep apnea in 81 ambulatory male patients with stable heart failure. Types and their prevalences, consequences, and presentations. *Circulation* 1998;97:2154–9.
- [10] Oldenburg O, Lamp B, Faber L, Teschler H, Horstkotte D, Topfer V. Sleep-disordered breathing in patients with symptomatic heart failure: a contemporary study of prevalence in and characteristics of 700 patients. *Eur J Heart Fail* 2007;9:251–7.
- [11] Hui DS, Wong TY, Li TS, Ko FW, Choy DK, Szeto CC, et al. Prevalence of sleep disturbances in Chinese patients with end stage renal failure on maintenance hemodialysis. *Med Sci Monit* 2002;8:CR331–6.
- [12] Allen RP, Earley CJ. Restless legs syndrome: a review of clinical and pathophysiologic features. *J Clin Neurophysiol* 2001;18:128–47.
- [13] Nomura T, Inoue Y, Miyake M, Yasui K, Nakashima K. Prevalence and clinical characteristics of restless legs syndrome in Japanese patients with Parkinson's disease. *Mov Disord* 2006;21:380–4.
- [14] Hanly P, Powles P. Hypnotics should never be used in patients with sleep apnea. *J Psychosom Res* 1993;37(Suppl 1):59–65.
- [15] Enomoto M, Eudo T, Suenaga K, Miura N, Nakano Y, Kohtoh S, et al. Newly developed waist actigraphy and its sleep/wake scoring algorithm. *Sleep Biol Rhythms* 2009;7:17–22.
- [16] Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med* 2002;165:1217–39.
- [17] Sivertsen B, Krokstad S, Overland S, Mykletun A. The epidemiology of insomnia: associations with physical and mental health. The HUNT-2 study. *J Psychosom Res* 2009;67:109–16.
- [18] Ulloa EW, Salup R, Patterson SG, Jacobsen PB. Relationship between hot flashes and distress in men receiving androgen deprivation therapy for prostate cancer. *Psychooncology* 2009;18:598–605.
- [19] Woodward S, Freedman RR. The thermoregulatory effects of menopausal hot flashes on sleep. *Sleep* 1994;17:497–501.
- [20] Stein MB, Chartier M, Walker JR. Sleep in nondepressed patients with panic disorder: I. Systematic assessment of subjective sleep quality and sleep disturbance. *Sleep* 1993;16:724–6.
- [21] Partinen M. Epidemiology of sleep disorders. In: Kryger M, Roth T, Dement W, editors. *Principals and practice of sleep medicine*. 2nd ed. Philadelphia: WB Saunders; 1994. p. 437–52.
- [22] Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 1993;328:1230–5.
- [23] Kim J, In K, You S, Kang K, Shim J, Lee S, et al. Prevalence of sleep-disordered breathing in middle-aged Korean men and women. *Am J Respir Crit Care Med* 2004;170:1108–13.
- [24] The Report of an American Academy of Sleep Medicine Task Force. Sleep-related breathing disorders in adults: recommendations for syndrome definition and measurement techniques in clinical research. *Sleep* 1999;22:667–89.
- [25] Enomoto M, Li L, Aritake S, Nagase Y, Kaji T, Tagaya K, et al. Restless legs syndrome and its correlation with other sleep problems in the general population of Japan. *Sleep Biol Rhythms* 2006;4:153–9.
- [26] Siddiqi N, House AO, Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age Ageing* 2006;35:350–64.
- [27] Foley D, Ancoli-Israel S, Britz P, Walsh J. Sleep disturbances and chronic disease in older adults: results of the 2003 National Sleep Foundation Sleep in America Survey. *J Psychosom Res* 2004;56:497–502.
- [28] Stewart R, Besset A, Bebbington P, Brugha T, Lindesay J, Jenkins R, et al. Insomnia comorbidity and impact and hypnotic use by age group in a national survey population aged 16 to 74 years. *Sleep* 2006;29:1391–7.
- [29] Taylor DJ, Mallory LJ, Lichstein KL, Durrence HH, Riedel BW, Bush AJ. Comorbidity of chronic insomnia with medical problems. *Sleep* 2007;30:213–8.
- [30] Meissner HH, Riemer A, Santiago SM, Stein M, Goldman MD, Williams AJ. Failure of physician documentation of sleep complaints in hospitalized patients. *West J Med* 1998;169:146–9.
- [31] Shochat T, Umphress J, Israel AG, Ancoli-Israel S. Insomnia in primary care patients. *Sleep* 1999;22(Suppl 2):S359–65.
- [32] Terzano MG, Parrino L, Cirignotta F, Ferini-Strambi L, Gigli G, Rudelli G, et al. Studio Morfeo: insomnia in primary care, a survey conducted on the Italian population. *Sleep Med* 2004;5:67–75.
- [33] Aikens JE, Rouse ME. Help-seeking for insomnia among adult patients in primary care. *J Am Board Fam Pract* 2005;18:257–61.
- [34] Neutel CI, Perry S, Maxwell C. Medication use and risk of falls. *Pharmacoepidemiol Drug Saf* 2002;11:97–104.
- [35] Mahoney JE, Webb MJ, Gray SL. Zolpidem prescribing and adverse drug reactions in hospitalized general medicine patients at a Veterans Affairs hospital. *Am J Geriatr Pharmacother* 2004;2:66–74.
- [36] Glass J, Lancotot KL, Hermann N, Sproule BA, Busto UE. Sedative hypnotics in older people with insomnia: meta-analysis of risks and benefits. *BMJ* 2005;331:1169.
- [37] Chang CM, Wu EC, Chang S, Lin KM. Benzodiazepine and risk of hip fractures in older people: a nested case-control study in Taiwan. *Am J Geriatr Psychiatry* 2008;16:686–92.

Sleep Deprivation Influences Diurnal Variation of Human Time Perception with Prefrontal Activity Change: A Functional Near-Infrared Spectroscopy Study

Takahiro Soshi^{1,2}, Kenichi Kuriyama^{1,2*}, Sayaka Aritake¹, Minori Enomoto¹, Akiko Hida¹, Miyuki Tamura¹, Yoshiharu Kim², Kazuo Mishima¹

¹ Department of Psychophysiology, National Center of Neurology and Psychiatry, National Institute of Mental Health, Tokyo, Japan, ² Department of Adult Mental Health, National Center of Neurology and Psychiatry, National Institute of Mental Health, Tokyo, Japan

Abstract

Human short-time perception shows diurnal variation. In general, short-time perception fluctuates in parallel with circadian clock parameters, while diurnal variation seems to be modulated by sleep deprivation per se. Functional imaging studies have reported that short-time perception recruits a neural network that includes subcortical structures, as well as cortical areas involving the prefrontal cortex (PFC). It has also been reported that the PFC is vulnerable to sleep deprivation, which has an influence on various cognitive functions. The present study is aimed at elucidating the influence of PFC vulnerability to sleep deprivation on short-time perception, using the optical imaging technique of functional near-infrared spectroscopy. Eighteen participants performed 10-s time production tasks before (at 21:00) and after (at 09:00) experimental nights both in sleep-controlled and sleep-deprived conditions in a 4-day laboratory-based crossover study. Compared to the sleep-controlled condition, one-night sleep deprivation induced a significant reduction in the produced time simultaneous with an increased hemodynamic response in the left PFC at 09:00. These results suggest that activation of the left PFC, which possibly reflects functional compensation under a sleep-deprived condition, is associated with alteration of short-time perception.

Citation: Soshi T, Kuriyama K, Aritake S, Enomoto M, Hida A, et al. (2010) Sleep Deprivation Influences Diurnal Variation of Human Time Perception with Prefrontal Activity Change: A Functional Near-Infrared Spectroscopy Study. *PLoS ONE* 5(1): e8395. doi:10.1371/journal.pone.00808395

Editor: Alex O. Holcombe, University of Sydney, Australia

Received: May 20, 2009; **Accepted:** November 30, 2009; **Published:** January 1, 2010

Copyright: © 2010 Soshi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by Special Coordination Funds of the Ministry of Education, Culture, Sports, and Technology of Japan and by a Grant-in-Aid for Cooperative Research from the Ministry of Health, Labour, and Welfare of Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: kenichik@ncnp.go.jp

Introduction

Temporal perception is fundamental to environmental adaptation. Developing time management skills enables us not only to avoid life-threatening situations, but also to gain rewards and establish motor and cognitive skills. Higher organisms have at least two endogenous clock systems [1,2]. One of these is a circadian pacemaker located in the suprachiasmatic nucleus of the hypothalamus [3,4] which is driven by a self-sustaining oscillator with a period of about 24 h and provides the time of day as the hour hand of the clock [5]. Another is a stopwatch-like system which perceives brief temporal intervals as the minute hand of the clock [1,6].

Most studies in human time perception refer to the issue that time perception is sensitive to the length of duration perceived, where the perception of a longer interval with a range of several minutes to hours (long-time perception) [7] shows different properties from the perception of a shorter interval with a range of seconds or within a minute (short-time perception), as well as from the perception of a sub-second interval [8–10]. Short-time perception robustly reflects the rate of the stopwatch-like system [8,11,12], being shortened or prolonged when the system “speeds up” or “slows down”, respectively [13]. When the stopwatch-like

system speeds up, the duration of events may be felt as more extended than the real interval: That may equal to “time expansion”, which enables us to deal with more things or to perform more rapidly.

Neural correlates of the stopwatch-like system are suspected to include subcortical (cerebellum and basal ganglia) structures in addition to the right prefrontal cortex (PFC) [8,9,14,15]. A three-stage model of temporal processing proposed by Pouthas et al. [9] comprises a first clock stage in which an endogenous clock system measures primitive timing intervals, a second storage stage in which measuring time is stored in short-term working memory and a third decision stage in which temporal judgment is finally made in reference to short- and long-term temporal memories. A similar model of temporal processing has been proposed by other researchers [16–18], and consensus has almost been reached that the cerebellum and the basal ganglia may contribute to a basic timing process that corresponds to the first stage of the three-stage model [19,20]. The right PFC, on the other hand, is implicated in making judgments with working memory or attentional processes contributing to temporal processing [9,21–23], although some researchers regard right PFC activity as constituting temporal processing per se [24,25].

Biological rhythm studies have reported that short-time perception driven by the stopwatch-like system is not independent of the influence of the circadian pacemaker; short-time perception correlates with circadian markers such as core body temperature and melatonin, showing diurnal variation in consequence [7,13,26–28]. On the other hand, after sleep deprivation, there is less diurnal variation in short-time perception dissociating with endogenous circadian markers [27,29,30]; the biological stopwatch is “sped up” as a result [13].

The present study aims at investigating neural responsibility for attenuation of the diurnal variation in short-time perception after sleep deprivation, utilizing the representative experimental paradigm of the time production (TP) method. The hypothesis is that an attenuated variation of short-time perception is associated with an alteration in PFC activity as a consequence of sleep deprivation. With regard to neural vulnerability to sleep deprivation, the PFC shows complicated hemodynamic or metabolic patterns during cognitive performance after sleep deprivation [31–33]. In this state, the PFC shows hypoactivity during working memory or arithmetic tasks [34,35] but hyperactivity during verbal learning or attention-loaded tasks, as well as verbal working memory tasks [36–38]. Although hemodynamic or metabolic responses to sleep deprivation on various cognitive tasks may be influenced by the task difficulty and weight of contribution from PFC activity [37], the neural vulnerability of the PFC to sleep deprivation must have consistent repercussions on cognitive performance. Taken together, the vulnerability of the PFC to sleep deprivation presumably influences short-time perception.

We utilize the brain imaging method of functional near-infrared spectroscopy (fNIRS) for the experimental purpose. fNIRS is a noninvasive optical imaging technique to measure changes of cerebral blood flow and volume through fluctuations in local glucose and oxygen coupled with neural electric activity [39,40]. It is well suited to sleep deprivation studies because it neither produces high noise levels on scanning nor requires that participants severely restrict their body movement, and thus is unlikely to seriously influence the sleep-deprived condition. It is also suitable for monitoring the participants' condition while they are performing tasks.

Results

Sleep Data

Nocturnal sleep duration on the interval day between the first and the second experiments for two different sleep schedules [sleep-controlled (SC) - sleep-deprived (SD), and SD - SC] was not significantly different [SC-SD (8.01 ± 0.96 h) vs. SD-SC (8.70 ± 1.72 h); $t(1,16) = 1.102$, $p = 0.287$]. Sleep duration between the first and the second TP sessions during the participants' stay in the laboratory also did not differ significantly between the two sleep schedules [SC-SD (6.40 ± 0.18 h) vs. SD-SC (6.63 ± 0.11 h); $t(1,16) = 1.145$, $p = 0.269$].

Time Production Data

Produced time in the SC condition was 11.16 ± 0.27 s on day 1 and 11.60 ± 0.33 s on day 2, and in the SD condition was 11.14 ± 0.24 s on day 1 and 10.94 ± 0.35 s on day 2 (Fig. 1).

A significant interaction of condition \times day was obtained [$F(1,16) = 9.888$, $p = 0.006$]. Pair-wise comparisons of two levels within the factor of day in each condition revealed that the produced time on day 2 was significantly longer than that on day 1 in the SC condition [$F(1,17) = 4.575$, $p = 0.047$]. In the SD condition, on the other hand, the produced time on day 2 was not significantly different from that on day 1 [$F(1,17) = 0.753$,




Figure 1. Mean interval variation in SC and SD conditions (n = 18). Black filled triangles with error bars (left: day 1, right: day 2) and the linking line indicate mean produced intervals and the trend of fluctuation in the SC condition, respectively. Gray filled circles with error bars (left: day 1, right: day 2) and the linking line show mean produced intervals and the trend of variation in the SD condition, respectively. Solid curves connecting two markers indicate significant differences ($p < 0.05$). doi:10.1371/journal.pone.0008395.g001

$p = 0.398$]. Produced time on day 2 in the SD condition, relative to that in the SC condition, was significantly attenuated [$F(1,17) = 6.123$, $p = 0.024$].

There was no significant interaction of the between-subject factor of schedule [condition \times schedule: $F(1,16) = 0.254$, $p = 0.621$; condition \times day \times schedule: $F(1,16) = 0.755$, $p = 0.398$]. These findings taken together with those for sleep data indicate that time production performance was not influenced by sleep schedule.

fNIRS Data

We conducted statistical analyses for fNIRS data utilizing mean values of changes in oxygenated hemoglobin (oxy-Hb) concentration. The overall ANOVA for the midline area did not show any significant difference [condition: $F(1,13) = 0.322$, $p = 0.580$; condition \times day: $F(1,13) = 1.715$, $p = 0.213$; condition \times channel: $F(3,39) = 0.458$, $p = 0.713$; condition \times day \times channel: $F(3,39) = 1.777$, $p = 0.167$]. The overall ANOVA for the lateral site showed only a significant interaction for condition \times day \times hemisphere [$F(1,13) = 5.373$, $p = 0.037$], suggesting that PFC activity in the SD condition, compared with that in the SC condition, was more enhanced in the left hemisphere on day 2 (Fig. 2B). Subsequent planned ANOVAs confirmed that there was a significant difference between conditions on day 2 [condition \times hemisphere: $F(1,26) = 9.049$, $p = 0.006$], but not on day 1 [condition \times hemisphere: $F(1,26) = 0.024$, $p = 0.879$]. Additional ANOVAs showed that the more enhanced PFC activation in the SD condition on day 2 appeared in the left hemisphere [$F(1,52) = 6.142$, $p = 0.017$], but not in the right hemisphere [$F(1,52) = 0.225$, $p = 0.637$].

Enhanced oxy-Hb concentration changes on day 2 in the SD condition, compared with those in the SC condition, were observed in the left anterior PFC (LAPFC) region of interest (ROI) including chs. 17, 21 and 22 (see Fig. 3A). Comparing the mean change in concentration in the SD condition with that in the SC condition, a significant effect was observed in the overall ROI [$t(1,13) = 2.810$, $p = 0.015$ ($p < 0.0375$: adjusted α level under the




Figure 2. Time course of grand averaged oxygenated hemoglobin (oxy-Hb) concentration changes during 10-s time production trial in the left anterior prefrontal (LAPFC) ($n = 14$). Comparison of oxy-Hb concentration changes between the SC (green line) and the SD (red line) conditions on (A) day 1 and (B) day 2 in the LAPFC (chs.17, 21, 22). Scores of 0, 11 and 27 below the waveform box indicate the onset of the trial, the end of the trial, and the finish of the post-trial rest, respectively. Oxy-Hb changes are given as z-scores. Any significant differences in oxy-Hb concentration changes between the SC and the SD conditions in the right hemisphere except for in the left hemisphere are not observed on day 2. Marked differences in oxy-Hb concentration changes between conditions are observed in the LAPFC on day 2. doi:10.1371/journal.pone.0008395.g002

control of the false discovery rate; FDR]). Each channel in the ROI also showed a significant difference [ch.22: $t(1,13) = 3.066$, $p = 0.009$ ($p < 0.0125$: FDR); ch.17: $t(1,13) = 2.872$, $p = 0.013$ ($p < 0.025$: FDR)] or trend [ch.21: $t(1,13) = 2.007$, $p = 0.066$ ($p < 0.05$: FDR)]. A significant positive correlation was observed between produced time and change in oxy-Hb concentration in the overall ROI on day 2 in the SD condition [$r = 0.535$, $p = 0.049$; Fig. 3B].

Discussion

Sleep is believed to be a neural state during which both consolidation of memories and homeostatic preservations are taking place [41–44]. Sleep deprivation, even for the course of an extended active period of the day, eliminates these effects, and possibly results in a deterioration in cognitive activity [13,30,45].

The present study demonstrated that sleep deprivation modulates short-time perception behavior, as was suggested by earlier studies [27,30]. It has been shown previously that a short-time perception profile exhibits diurnal variation, reaching a peak (the longest produced time) around 09:00 and a nadir (the shortest produced time) around 21:00 with a regular sleep-wake cycle under experimental conditions [13]. Here, we measured produced time around both these expected peak and nadir periods in order to obtain approximate peak-to-peak amplitude of the diurnal variation of short-time perception. As a result, we found that the produced time at 09:00 was significantly attenuated after sleep deprivation compared with that after physiological sleep. It was unlikely that the alteration in properties of time perception after sleep deprivation was caused by circadian phase shifting (different peak times) of short-time perception rhythms between the SC and SD sessions because all participants were exposed to similar time cues such as dim light (< 100 lux) and prohibited from physical exercise throughout the experiments, and the produced time levels around 21:00 were in fact identical between the two sessions. This seems to indicate that sleep deprivation per se modulated short-time perception. Given that produced time represents the rate of

the biological stopwatch-like system [1], the attenuation of produced time observed on day 2 in the SD condition might be associated with the stopwatch-like system that runs faster [13].

A functional correlation of increased activation of the LAPFC after sleep deprivation with short-time perception behavior was observed in the present study, although unlike in previous studies, we failed to detect any significant changes in right PFC activity [8,9,14,15]. This discrepancy in the localized PFC activation findings between the present and previous studies is possibly attributable to experimental settings, tasks used and imaging modalities [9,14,24,46]. It has also been argued that increased activation of the PFC after sleep deprivation is associated with neural compensation for cognitive function [36,38,47], and the LAPFC activity observed in the present study should also be argued in the context of neural compensation: the LAPFC activity likely contributed to the third decision stage of temporal judgment in the three-stage model [9], with adaptive neural activity in the SD condition, although the produced time on day 2 in the SD condition was different from that in the SC condition.

It remains open as to whether or not a subcortical network including the cerebellum and the basal ganglia, assumed to be associated with timing processing, contributes to attenuating the diurnal fluctuation of short-time perception. Sustained wakefulness affects dopaminergic neuronal activity related with cognitive function as well as temporal function [48–50]. If alteration of subcortical activities might be caused by sleep deprivation, attenuation of short-time perception probably reflects subcortical vulnerability, and the change in PFC activity is possibly a by-product. Further studies are warranted to elucidate the processes involved.

As has been reported, short-time perception is modulated by affective states [51], an intense fever [52,53] and psychotic disorders [54,55]. Diurnal variation of short-time perception associated with the rest-activity cycle is expedient for adapting in daily life, similarly to the vegetative functions in blood circulation [56], aspiration [57], assimilation [58] or incretion [59,60]. These functions robustly synchronize to diurnal variation oscillated by a




Figure 3. Change in hemodynamic response and functional correlation in prefrontal cortex (PFC) activity after sleep deprivation. (A) Topographic image representing increased activation in the PFC on day 2 in the SD condition ($n=14$). The image is constructed by using subtracted values between the SC and the SD conditions (SD minus SC). The area of dotted white circles emphasizes the left anterior PFC (LAPFC) including chs. 17, 21 and 22 for region of interest (ROI) analysis. (B) Correlation between normalized produced intervals and oxy-Hb concentration changes in the LAPFC ROI on day 2 in the SD condition ($n=14$). Functional connectivity in changes of short-time perception with LAPFC activity under sustained wakefulness is suggested by a correlation coefficient (r) of 0.535 with statistical significance ($p < 0.05$).

doi:10.1371/journal.pone.0008395.g003

circadian pacemaker, and simultaneously prepare for desynchronization from diurnal variation under stressful conditions. Temporarily collapsing the diurnal variation of short-time perception may be an important function for surviving a crisis, such as in a situation with an acute emergency. Time perception in humans, as well as in other organisms, should be fundamentally synchronized to the physical state for constant adaptation to daily lifestyle regularity. However, once we are placed under stress, time perception must desynchronize from regular physical homeostasis and be shortened enabling time expansion and assumedly allowing us to adopt suitable strategies for coping with a stressful environment by doing or thinking of things more rapidly. Serious consideration of an adaptive nature of the human PFC function [61,62] suggests that the PFC might play a switch-like role in

short-time perception as a situation demands, to meet the demands for adaptation.

Materials and Methods

Participants

Eighteen healthy right-handed males (mean age 22.4 yr; range 20–28 yr) participated in laboratory-based experiments in a 4-day protocol. A psychiatrist conducted a medical examination to confirm that none of the participants had a history of neurological or psychiatric disorder or abused psychoactive drugs. Participants were asked to avoid intake of caffeine, nicotine and alcohol for one week prior to and during the experiment. For one week before the experiment, they were also required to keep a regular sleep-wake habit in which they slept from around 00:00 and awoke at around 08:00, as confirmed by a sleep log and an activity recorder (Actiwatch-L, Mini-Mitter Co., Inc. Bend, OR). The mean habitual sleep duration was 6.5 ± 0.1 h, and habitual sleep onset and wake times were $00:34 \pm 00:13$ and $07:22 \pm 00:10$, respectively. All procedures for the study were carried out in accordance with the principles outlined in the Declaration of Helsinki. The experimental protocol was approved by the ethics committee of the National Center of Neurology and Psychiatry, Japan. All participants gave written informed consent to take part in the study.

Experimental Protocol

All participants performed a 10-s TP task in the SC and SD conditions, scheduled in random order with a 1-day interval (Fig. 4). On the first day (day 1) in both the SC and SD conditions, participants arrived at the laboratory at about 18:00. fNIRS probe and optodes were attached to the surface of the scalp approximately one hour before the start of the TP session (at 20:00). The session, which lasted approximately 15 min, generally started around 21:00–22:00. After the end of the session, in the SC condition, the participants rested without sleep and exercise until 00:00 and then stayed in bed under complete darkness (<0.1 lux) until 08:00 on the second day (day 2). In contrast, in the SD condition, participants were asked to stay awake quietly under room light (100 lux) until 08:00 the next morning. During the nighttime period of enforced wakefulness, researchers monitored the participants' status via a video monitoring system. On day 2, the TP session started again around 09:00–10:00. All the experiments were performed at the time isolation facility of the National Center of Neurology and Psychiatry. The ambient temperature and humidity in the time isolation facility were maintained at $24.0 \pm 0.5^\circ\text{C}$ and $60.0 \pm 5\%$, respectively, and temporal information was not provided to participants throughout the experimental schedule.

Time Production Task

TP tasks were arranged in an event-related design for detecting the hemodynamic response for a single trial. It has been confirmed that output of the task shows diurnal variation correlated with core body temperature or melatonin under a constant condition [7,13,26,28]. It shortens from morning into night, and is prolonged again from night to the next morning [7,13,26,29]. On the basis of our previous findings [26,27], TP sessions in the present study were conducted between 21:00–22:00 on day 1 and 09:00–10:00 on day 2, corresponding to the expected nadir and the peak periods of the diurnal variation of TP in the present participants with a regular sleep-wake cycle, respectively.

Each TP session consisted of 15 trials. The inter-trial interval was about 30 s. Participants were asked to produce a 10-s interval




Figure 4. Experimental protocol. Time production sessions (TPS) were conducted twice at around 21:00 on day 1 and about 09:00 on day 2 in the sleep-controlled (SC) and the sleep-deprived (SD) settings. The two conditions were randomly scheduled with a 1-day interval. Each TPS included 15 trials, each of which participants started and ended with their own button presses. Functional near-infrared spectroscopy (fNIRS) data were recorded continuously through the sessions.
doi:10.1371/journal.pone.0008395.g004

and to begin and end each trial by pressing a key button [13]. Duration from the first to the second button presses was counted as the produced time. Participants were given no feedback about accuracy in the trials.

fNIRS Data Acquisition

Regional hemodynamic changes in brain tissue were monitored throughout the TP sessions by a continuous wave-type fNIRS system (OMM3000; Shimadzu Co., Tokyo, Japan), which outputs near-infrared light at three wavelengths (780, 805 and 830 nm). All transmitted intensities of the three wavelengths were recorded every 130 ms at 22 channels in order to estimate concentration changes in oxy-Hb, deoxygenated hemoglobin and total hemoglobin, on the basis of the modified Beer-Lambert equation as a function of light absorbance of Hb and pathlength. A 3×5 optode probe was utilized, in which light detectors and emitters were alternately positioned at an equal distance of 3 cm. Based on the international 10–20 system [63], the lower central edge of each probe was suited above Fpz, along the reference curve of T3 - Fpz - T4. The 22 channels (see Fig. 5) probably covered the middle and superior PFC regions (BA9, 46, 10) [64].

fNIRS Data Analysis

Oxy-Hb data was chosen to examine event-related response in the PFC since it is an optimal index for changes of regional cerebral blood flow [65]. Data from 4 participants who were recruited to the experiment in the early stage were excluded from fNIRS analyses because they were monitored with a distinct optode probe. We applied a high-pass filter to raw data, re-sampled at 10 Hz, using the low-cutoff frequency of 0.05 Hz. Smoothing was performed by the moving average method (a boxcar filter) with a sliding time

window of 1.1 s. Data were converted into z-scores by the normalization process $[(X_i - M) / SD]$, where X_i is the value at each time point, M is the mean of all points and SD is the standard deviation of all points [66]. Concentration changes time-locked to trial onset were extracted from 5 s before to 27 s after the onset, covering a mean produced time of about 11 s and a mean rest interval of about 16 s. A total of 15 epochs were obtained for each experimental day (day 1 or day 2) in each condition (SC or SD).




Figure 5. Approximate positions of 22 channels in fNIRS superimposed on the 3D head model. The 22 measuring channels were produced by optodes placed equidistantly over the prefrontal cortex (PFC) area. The lower line of the 3×5 optode probe was positioned along the reference curve linking T3, T4 and Fpz.
doi:10.1371/journal.pone.0008395.g005

Before individual averaging, baselines were corrected with mean z-scores of 5 s before trial onset. Grand averaged concentration changes in the LAPFC region, based on statistical analyses, were superimposed (Fig. 2), and subtracted values were utilized for reconstructing a topographic image (Fig. 3A).

Statistical Analysis

To examine whether the order of the sleep conditions [SC-SD ($n = 7$) vs. SD-SC ($n = 11$)] influenced outcomes of time production performance, we compared nocturnal sleep duration in both sleep schedules during the 1-day interval between the first and the second experiments and the sleep duration between the first and the second TP sessions during the participants' stay in the laboratory. We also analyzed the time production data of 18 participants using three-way ANOVA including two within-subject factors of condition (SC and SD) and day (day 1 and day 2), and a between-subject factor of schedule (SC-SD and SD-SC). If significant interactions were obtained, follow-up ANOVAs were conducted.

fNIRS data of 14 participants were tested separately for the lateral and the midline sites by overall ANOVA including within-subject factors of condition, day, hemisphere (left, right) and channel [9 chs: left, 1, 2, 5, 6, 10, 14, 15, 19, 20; right (the corresponding order): 4, 3, 9, 8, 13, 18, 17, 22, 21] for the lateral site, or factors of condition, day and channel (4 chs: 7, 11, 12, 16) for the midline area, utilizing mean z-scores of all temporal points (0–27 s). When the overall ANOVA reached statistical significance of the interaction coefficients, planned ANOVAs were subsequently performed with pooling error terms across the entire analyses, for the purpose to examine our assumption that sleep deprivation modulates the PFC activities on the time production trial. If degrees of freedom exceeded one degree, Greenhouse-Geisser correction was applied on the basis of the sphericity test. Depending on the topographic image reconstructed with subtract-

ed values of changes in oxy-Hb concentration (SD minus SC; Fig. 3A), and a confirmed methodology for fNIRS probe placement [64], we defined the LAPFC, including chs. 17, 21 and 22, which corresponds to the medial dorsal part of the PFC (Brodmann's area 10) [64], as the ROI, and further examined the influence of sleep-deprivation on neural response in the LAPFC. Paired *t*-tests were performed for the ROI and the three constituent channels, with α levels adjusted by Benjamini and Hochberg's method to control the FDR [67]: $H_{(i)}$ denotes the null hypothesis based on the corresponding *p*-value $P_{(i)}$, $i = 1, 2, \dots, m$. $P_{(i)}$ is increasingly ordered such as $P_{(1)} < P_{(2)} < \dots < P_{(m)}$. The FDR control procedure starts with $i = m$, and continues to compare $P_{(i)}$ with $\alpha(i/m)$ until $P_{(i)}$ satisfy the constraint $P_{(i)} \leq \alpha(i/m)$. The relation between produced times standardized within each participant and oxy-Hb concentration changes in the ROI on day 2 in the SD condition was tested with Pearson's product-moment correlation coefficient to examine functional connectivity between short TP and neural response of the LAPFC.

Outcomes of the tests for produced time and fNIRS data were considered significant at level α if $p < 0.05$. Results are shown as mean and standard errors (SEs). Statistical analysis was performed by SPSS 12.0 J (SPSS Japan Inc., Tokyo).

Acknowledgments

We are grateful to S. Higuchi, H. Suzuki and M. Abe for valuable comments and discussions during the preparation of the experiment and the paper.

Author Contributions

Conceived and designed the experiments: TS KK SA ME KM. Performed the experiments: TS KK SA ME AH MT KM. Analyzed the data: TS KK KM. Wrote the paper: TS KK KM. Contributed to discussion: TS, KK, YK, KM. Supervised the entire project: KM.

References

- Morell V (1996) Setting a biological stopwatch. *Science* 271: 905–906.
- Ruan GX, Allen GC, Yamazaki S, McMahon DG (2008) An autonomous circadian clock in the inner mouse retina regulated by dopamine and GABA. *PLoS Biol* 6: 2249–2265.
- Inoue ST, Kawamura H (1979) Persistence of circadian rhythmicity in a mammalian hypothalamic 'island' containing the suprachiasmatic nucleus. *Proc Natl Acad Sci U S A* 76: 5962–5966.
- Moore RY, Silver R (1998) Suprachiasmatic nucleus organization. *Chronobiol Int* 15: 475–487.
- Moore-Ede MC, Czeisler CA, Richardson GS (1983) Circadian time-keeping in health and disease. Part 1. Basic properties of circadian pacemakers. *N Eng J Med* 309: 469–476.
- Gibbon J, Church RM (1981) Time left: linear versus logarithmic subjective time. *J Exp Psychol Anim Behav Process* 7: 87–107.
- Aschoff J (1998) Human perception of short and long time intervals: its correlation with body temperature and the duration of wake time. *J Biol Rhythms* 13: 437–442.
- Lewis PA, Miall RC (2003) Brain activation patterns during measurement of sub- and supra-second intervals. *Neuropsychologia* 41: 1583–1592.
- Pouthas V, Geroge N, Poline JB, Pfeuty M, VandeMoortele PF, et al. (2005) Neural network involved in time perception: an fMRI study comparing long and short interval estimation. *Hum Brain Mapp* 25: 433–441.
- Rammsayer TH (2006) Effects of pharmacologically induced changes in NMDA receptor activity on human timing and sensorimotor performance. *Brain Res* 1073–1074: 407–416.
- Mattel MS, Meck WH (2004) Cortico-striatal circuits and interval timing: coincidence detection of oscillatory processes. *Cogn Brain Res* 58: 119–124.
- Rammsayer T, Ulrich R (2005) No evidence for qualitative differences in the processing of short and long temporal intervals. *Acta Psychol* 120: 141–171.
- Kuriyama K, Uchiyama M, Suzuki H, Tagaya H, Ozaki A, et al. (2005) Diurnal fluctuation of time perception under 30-h sustained wakefulness. *Neurosci Res* 53: 123–128.
- Maquet P, Lejeune H, Pouthas V, Bonnet M, Casini L, et al. (1996) Brain activation induced by estimation of duration: a PET study. *Neuroimage* 3: 119–126.
- Rao S, Mayer AR, Harrington DL (2001) The evolution of brain activation during temporal processing. *Nat Neurosci* 4: 317–323.
- Treisman M (1963) Temporal discrimination and the indifference interval implications for a model of the 'internal clock.' *Psychol Monogr* 77 (13): 1–31.
- Ivry RB, Spencer RM (2004) The neural representation of time. *Curr Opin Neurobiol* 14: 225–32.
- Zakay D, Block RA (2004) Prospective and retrospective duration judgments: an executive-control perspective. *Acta Neurobiol Exp (Wars)* 64: 319–28.
- Casini L, Ivry RB (1999) Effects of divided attention on temporal processing in patients with lesions of the cerebellum or frontal lobe. *Neuropsychology* 13: 10–21.
- Lewis PA, Miall RC (2002) Brain activity during non-automatic motor production of discrete multi-second intervals. *Neuroreport* 13: 1731–1735.
- Harrington DL, Haaland KY, Knight RT (1998) Cortical networks underlying mechanisms of time perception. *J Neurosci* 18: 1085–1095.
- Macar F, Lejeune H, Bonnet M, Ferrara A, Pouthas V, et al. (2002) Activation of the supplementary motor area and of attentional networks during temporal processing. *Exp Brain Res* 142: 475–485.
- Livesey AC, Wall MB, Smith AT (2007) Time perception: manipulation of task difficulty dissociates clock functions from other cognitive demands. *Neuropsychologia* 45: 321–331.
- Smith A, Lidzba K, Taylor E, Rubia K (2003) A right hemispheric frontocerebellar network for time discrimination of several hundreds of milliseconds. *Neuroimage* 20: 344–350.
- Rubia K, Smith A (2004) The neural correlates of cognitive time management: a review. *Acta Neurobiol Exp* 64: 329–340.
- Ashoff J, Daan S (1997) Human time perception in temporal isolation: effects of illumination intensity. *Chronobiol Int* 14: 585–596.
- Miró E, Cano MC, Espinosa-Fernández L, Buela-Casal G (2003) Time estimation during prolonged sleep deprivation and its relation to activation measures. *Hum Factors* 45: 148–159.
- Kuriyama K, Uchiyama M, Suzuki H, Tagaya H, Ozaki A, et al. (2003) Circadian fluctuation of time perception in healthy human subjects. *Neurosci Res* 46: 23–31.
- Pöppel E, Giedke H (1970) Diurnal variation of time perception. *Psychol Forsch* 34: 182–198.

30. Aschoff J (1988) Circadian timing. *Ann N Y Acad Sci* 423: 442–468.
31. Horne JA (1993) Human sleep, sleep loss and behavior: implication for the prefrontal cortex and psychiatric disorder. *B J Psychiatry* 162: 413–419.
32. Drummond SPA, Brown GG (2001) The effects of total sleep deprivation on cerebral responses to cognitive performance. *Neuropsychopharmacology* 25 (5 suppl): S68–73.
33. Durmer JS, Dinges DF (2005) Neurocognitive consequence of sleep deprivation. *Semin Neurol* 25: 117–129.
34. Drummond SPA, Brown GG, Stricker JL, Buxton RB, Wong EC, et al. (2000b) Sleep deprivation-induced reduction in cortical functional response to serial subtraction. *Neuroreport* 10: 3745–3748.
35. Mu Q, Nahas Z, Johnson KA, Yamanaka K, Mishory A, et al. (2005) Decreased cortical activation response to verbal working memory following sleep deprivation. *Sleep* 28: 55–67.
36. Drummond SPA, Brown GG, Gillin JC, Stricker JL, Wong EC, et al. (2000a) Altered brain response to verbal learning following sleep deprivation. *Nature* 403: 655–657.
37. Drummond SPA, Gillin JC, Brown GG (2001) Increased cerebral response during a divided attention task following sleep deprivation. *J Sleep Res* 10: 85–92.
38. Chee MW, Choo WC (2004) Functional imaging of working memory after 24 hr of total sleep deprivation. *J Neurosci* 24: 4560–4567.
39. Hoshi Y, Tamura M (1993) Dynamic multichannel near-infrared optical imaging of human brain activity. *J Appl Physiol* 75: 1842–1846.
40. Villringer A, Planck J, Hock C, Schleinkofer L, Dirnagl U (1993) Near infrared spectroscopy (NIRS): a new tool to study hemodynamic changes during activation of brain function in human adults. *Neurosci Lett* 154: 101–104.
41. Tononi G, Cirelli C (2006) Sleep function and synaptic homeostasis. *Sleep Med Rev* 10: 49–62.
42. Walker MP, Stickgold R (2004) Sleep-dependent learning and memory consolidation. *Neuron* 44: 121–133.
43. Kuriyama K, Stickgold R, Walker SP (2004) Sleep-dependent learning and motor skill complexity. *Learn Mem* 11: 705–713.
44. Kuriyama K, Mishima K, Suzuki H, Aritake S, Uchiyama M (2008) Sleep accelerates the improvement in working memory performance. *J Neurosci* 28: 10145–10150.
45. Aschoff J (1985) On the perception of time during prolonged temporal isolation. *Human Neurobiol* 4: 41–52.
46. Coull JT, Nobre AC (2008) Dissociating explicit timing from temporal expectation with fMRI. *Curr Opin Neurobiol* 18: 137–144.
47. Drummond SPA, Meloy MJ, Yanagi MA, Orff HJ, Brown GG (2005) Compensatory recruitment after sleep deprivation and the relationship with performance. *Psychiatry Res* 140: 211–223.
48. Rammesayer TH, Vogel WH (1992) Pharmacologic properties of the internal clock underlying time perception in humans. *Neuropsychobiology* 26 (1–2): 71–80.
49. Meck WH, Penny TB, Pouthas V (2008) Cortico-striatal representation of time in animals and humans. *Curr Opin Neurobiol* 18: 145–152.
50. Volkow ND, Wang GJ, Telang F, Fowler JS, Logan J, et al. (2008) Sleep deprivation decreases binding of [¹¹C]Raclopride to dopamine D₂/D₃ receptors in the human brain. *J Neurosci* 28: 8454–8461.
51. Angrilli A, Cherubini P, Pavese A, Mangfredini S (1997) The influence of affective factors on time perception. *Percept Psychophys* 59: 972–982.
52. Hoagland H (1933) The physiological control of judgments of duration: evidence for a chemical clock. *J Gen Psychol* 9: 267–287.
53. Kleber RJ, Lhamon WT, Goldstone S (1963) Hyperthermia, hyperthyroidism, and time judgment. *J Comp Physiol Psychol* 56: 362–365.
54. Tysk L (1984) Time perception and affective disorders. *Percept Mot Skills* 58: 455–464.
55. Bschor T, Ising M, Bauer M, Lewitzka U, Skerstuepeit M, et al. (2004) Time experience and time judgment in major depression, mania and healthy subjects: a controlled study of 93 subjects. *Acta Psychiatr Scand* 109: 222–229.
56. Kraft M, Martin RJ (1995) Chronobiology and chronotherapy in medicine. *Dis Mon* 41: 501–75.
57. Jindal SK, Aggarwal AN, Gupta D (2002) Diurnal variability of peak expiratory flow. *J Asthma* 39: 363–73.
58. Davidson AJ, Castañón-Cervantes O, Stephan FK (2004) Daily oscillations in liver function: diurnal vs circadian rhythmicity. *Liver Int* 24: 179–86.
59. Boyar RM, Witkin M, Carruth A, Ramsey J (1979) Circadian cortisol secretory rhythms in Cushing's disease. *J Clin Endocrinol Metab* 48: 760–5.
60. Fevre M, Segel T, Marks JF, Boyar RM (1978) LH and melatonin secretion patterns in pubertal boys. *J Clin Endocrinol Metab* 47: 1383–6.
61. Duncan J (2001) An adaptive coding model of neural function in prefrontal cortex. *Nat Rev Neurosci* 2: 820–829.
62. Miller EK, Cohen JD (2001) An integrative theory of prefrontal cortex function. *Annu Rev Neurosci* 24: 167–202.
63. Jasper HH (1958) The ten-twenty electrode system of international federation. *Electroencephalogr Clin Neurophysiol* 86: 7–14.
64. Okamoto M, Dan H, Sakamoto K, Takeo K, Shimizu K, et al. (2004) Three-dimensional probabilistic anatomical cranio-cerebral correlation via the international 10–20 system oriented for transcranial functional brain mapping. *Neuroimage* 21: 99–111.
65. Hoshi Y, Kobayashi N, Tamura M (2001) Interpretation of near-infrared spectroscopy signals: a study with a newly developed perfused rat brain model. *J Appl Physiol* 90: 1657–1662.
66. Matsuda G, Hiraki K (2006) Sustained decrease in oxygenated hemoglobin during video games in the dorsal prefrontal cortex: a NIRS study of children. *Neuroimage* 29: 706–11.
67. Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Statist Soc B* 57: 289–300.

SCIENTIFIC INVESTIGATIONS

Non-Pharmacological Self-Management of Sleep Among the Japanese General Population

Sayaka Aritake-Okada, Ph.D.¹; Yoshitaka Kaneita, M.D., Ph.D.²; Makoto Uchiyama, M.D., Ph.D.³; Kazuo Mishima, M.D., Ph.D.¹; Takashi Ohida, M.D., Ph.D.²

¹Department of Psychophysiology, National Institute of Mental Health, National Center of Neurology and Psychiatry, Ogawa-higashi, Kodaira, Tokyo, Japan; ²Division of Public Health, Department of Social Medicine, Nihon University School of Medicine, Ohyaguchikami-machi, Itabashi-ku, Tokyo, Japan; ³Division of Psychiatry, Department of Psychiatry, Nihon University School of Medicine, Ohyaguchikami-machi, Itabashi-ku, Tokyo, Japan

Study Objectives: The present study was conducted to clarify the prevalence of non-pharmacological self-management (nPSM) practices for obtaining good sleep and to identify favorable nPSM practices that could be applied for reducing excessive daytime sleepiness (EDS). We analyzed epidemiological data for an authentic representative sample of the Japanese population.

Methods: Data obtained from 24,686 adults via a self-administered questionnaire completed in the Active Survey of Health and Welfare 2000 were used for analyses. The prevalence of individual nPSM practices was calculated by gender. Subsequently, the associations between such practices and EDS were examined using logistic regression analyses.

Results: "Having a bath" was the most prevalent nPSM practice for both men (59.0%) and women (64.4%), followed by "maintaining a regular schedule" (men: 49.0%, women: 58.6%), "reading or listening to music" (men: 43.4%, women: 49.4%), "snacking on food and/or bev-

verages" (men: 36.1%, women: 27.9%), and "exercising" (men: 26.2%, women: 29.4%). The prevalence of "maintaining a regular schedule" increased with age. Multiple logistic regression analyses revealed that having a bath and maintaining a regular schedule had negative associations with EDS, whereas snacking on food and/or beverages had a positive association.

Conclusions: Having a bath and maintaining a regular schedule were identified as favorable nPSM practices for reducing EDS, whereas snacking on food and/or beverages was considered to be an unfavorable nPSM practice.

Keywords: Self-management, sleep, Japan, epidemiology, excessive daytime sleepiness, prevalence

Citation: Aritake-Okada S; Kaneita Y; Uchiyama M; Mishima K; Ohida T. Non-pharmacological self-management of sleep among the Japanese general population. *J Clin Sleep Med* 2009;5(5):464-469.

Sleep disturbance is known to be associated with the onset of mental disorders such as depression. It is also well known that sleep disturbance is an eventual risk factor for various somatic disorders such as diabetes mellitus, obesity, and cardiovascular disease.¹⁻⁵ In addition, excessive daytime sleepiness (EDS) resulting from sleep disturbance may lead to industrial and traffic accidents.⁶⁻⁸ Thus, in developed countries, employing pertinent measures to prevent sleep disturbance is widely recognized as an important issue in promoting industrial hygiene and public health.

People in general use various non-pharmacological self-management (nPSM) strategies to obtain good sleep. However, most previous studies have focused on so-called pharmacological management practices such as the use of alcohol or hypnot-

ic medications, and many epidemiological findings regarding such practices have been reported. For example, a study in the US reported that the prevalence of using hypnotic medications to improve the quality and quantity of sleep ranged from 10% to 18%, while that of consuming alcohol ranged from 10% to 13%, with the use of hypnotic medications being more prevalent among women and the consumption of alcoholic beverages more prevalent among men.⁹ A study in Japan obtained similar findings with respect to the use of hypnotics (women: 5.9%, men: 4.3%),¹⁰ and to the larger proportion of men consuming alcoholic beverages to induce sleep one or more times a week compared to women (48.3% and 18.3%, respectively).¹⁰ Although physiological data associated with nPSM practices such as exercising, having a bath, reading, or snacking on food and/or beverages have been reported, few findings of epidemiological studies are available. In a survey of self-management practices employed by Americans to obtain sleep, Ancoli-Israel et al. found that the prevalence of exercising was higher among non-insomniacs than among insomniacs.⁹ Morin et al. conducted a similar study of Americans and reported that the prevalence of reading was the highest, followed by listening to music.¹¹ However, no epidemiological study of nPSM practices for obtaining good sleep has been conducted in any Asian

Submitted for publication November, 2008

Submitted in final revised form April, 2009

Accepted for publication May, 2009

Address correspondence to: Yoshitaka Kaneita, M.D., Ph.D., Division of Public Health, Department of Social Medicine, Nihon University School of Medicine, Ohyaguchikami-machi, Itabashi-ku, Tokyo 173-8610, Japan; Tel: +81-3-3972-8111, Ext: 2272; Fax: +81-3-3972-5878; E-mail: kaneita@med.nihon-u.ac.jp