

- Nakayama H, Ishimaru F, Avitah N, Sezaki N, Fujii N, Nakase K *et al.* (1999). Decreases in Ikaros activity correlate with blast crisis in patients with chronic myelogenous leukemia. *Cancer Res* **59**: 3931–3934.
- Rosenbaum H, Harris AW, Bath ML, McNeill J, Webb E, Adams JM *et al.* (1990). An E mu-v-abl transgene elicits plasmacytomas in concert with an activated myc gene. *EMBOJ* **9**: 897–905.
- Scheijen B, Jonkers J, Acton D, Berns A. (1997). Characterization of pal-1, a common proviral insertion site in murine leukemia virus-induced lymphomas of c-myc and Pim-1 transgenic mice. *J Virol* **71**: 9–16.
- Seidel MG, Look AT. (2001). E2A-HLF usurps control of evolutionarily conserved survival pathways. *Oncogene* **20**: 5718–5725.
- Shinto Y, Morimoto M, Katsumata M, Uchida A, Aozasa K, Okamoto M *et al.* (1995). Moloney murine leukemia virus infection accelerates lymphomagenesis in E mu-bcl-2 transgenic mice. *Oncogene* **11**: 1729–1736.
- Smith KS, Rhee JW, Cleary ML. (2002). Transformation of bone marrow B-cell progenitors by E2a-Hlf requires coexpression of Bcl-2. *Mol Cell Biol* **22**: 7678–7687.
- Smith KS, Rhee JW, Naumovski L, Cleary ML. (1999). Disrupted differentiation and oncogenic transformation of lymphoid progenitors in E2A-HLF transgenic mice. *Mol Cell Biol* **19**: 4443–4451.
- Wang J, Iwasaki H, Krivtsov A, Febbo PG, Thorner AR, Ernst P *et al.* (2005). Conditional MLL-CBP targets GMP and models therapy-related myeloproliferative disease. *EMBOJ* **24**: 368–381.
- Warming S, Liu P, Suzuki T, Akagi K, Lindtner S, Pavlakis GN *et al.* (2003). Evi3, a common retroviral integration site in murine B-cell lymphoma, encodes an EBF2-related Krüppel-like zinc finger protein. *Blood* **101**: 1934–1940.
- Wolff L, Garin MT, Koller R, Bies J, Liao W, Malumbres M *et al.* (2003a). Hypermethylation of the Ink4b locus in murine myeloid leukemia and increased susceptibility to leukemia in p15(Ink4b)-deficient mice. *Oncogene* **22**: 9265–9274.
- Wolff L, Koller R, Hu X, Anver MR. (2003b). A Moloney murine leukemia virus-based retrovirus with 4070A long terminal repeat sequences induces a high incidence of myeloid as well as lymphoid neoplasms. *J Virol* **77**: 4965–4971.
- Yamashita N, Osato M, Huang L, Yanagida M, Kogan SC, Iwasaki M *et al.* (2005). Haploinsufficiency of Runx1/AML1 promotes myeloid features and leukaemogenesis in BXH2 mice. *Br J Haematol* **131**: 495–507.
- Yoshihara T, Inaba T, Shapiro LH, Kato JY, Look AT. (1995). E2A-HLF-mediated cell transformation requires both the trans-activation domains of E2A and the leucine zipper dimerization domain of HLF. *Mol Cell Biol* **15**: 3247–3255.
- Zhuang Y, Soriano P, Weintraub H. (1994). The helix-loop-helix gene E2A is required for B cell formation. *Cell* **79**: 875–884.
- Zörnig M, Schmidt T, Karsunky H, Grzeschiczek A, Möröy T. (1996). Zinc finger protein GFI-1 cooperates with myc and pim-1 in T-cell lymphomagenesis by reducing the requirements for IL-2. *Oncogene* **12**: 1789–1801.

Supplementary Information accompanies the paper on the Oncogene website (<http://www.nature.com/onc>)

