

- 635 (73) Paller AS, Syder AJ, Chan YM, Yu QC, Hutton E, Tadini G, et al. Genetic and clinical
636 mosaicism in a type of epidermal nevus. *N Engl J Med* 1994;331(21):1408-15.
- 637 (74) Tsubota A, Akiyama M, Sakai K, Goto M, Nomura Y, Ando S, et al. Keratin 1 gene
638 mutation detected in epidermal nevus with epidermolytic hyperkeratosis. *J Invest
639 Dermatol* 2007;127(6):1371-4.
- 640 (75) De Laurenzi V, Rogers GR, Hamrock DJ, Marekov LN, Steinert PM, Compton JG, et al.
641 Sjogren-Larsson syndrome is caused by mutations in the fatty aldehyde dehydrogenase
642 gene. *Nat Genet* 1996;12(1):52-7.
- 643 (76) Sjogren T, Larsson T. Oligophrenia in combination with congenital ichthyosis and spastic
644 disorders: a clinical and genetic study. *Acta Psychiatr Neurol Scand* 1957;32(suppl.113),1-
645 112.
- 646 (77) Refsum S, Salomonsen L, Skatvedt M. Heredopathia atactica polyneuritiformis in
647 children. *J Pediatr* 1949;35(3),335-343.
- 648 (78) Reed WB, Stone VM, Boder E, Ziprkowski L. Hereditary syndromes with auditory and
649 dermatological manifestations. *Arch Dermatol* 1967;95(5):456-61.
- 650 (79) Chanarin I, Patel A, Slavin G, Wills EJ, Andrews TM, Stewart G. Neutral-lipid storage
651 disease: a new disorder of lipid metabolism. *Br Med J* 1975;1(5957):553-5.
- 652 (80) Dorfman ML, Herskho C, Eisenberg S, Sagher F. Ichthyosiform dermatosis with systemic
653 lipidosis. *Arch Dermatol* 1974;110(2):261-6.

- 654 (81) MacLeod JMH. Three cases of 'ichthyosis follicularis' associated with baldness. Brit J
655 Derm. 1909;21:165-189.
- 656 (82) Hamm H, Meinecke P, Traupe H. Further delineation of the ichthyosis follicularis,
657 atrichia, and photophobia syndrome. Eur J Pediatr 1991;150(9):627-9.
- 658 (83) Happle R. X-linked dominant chondrodysplasia punctata. Review of literature and report
659 of a case. Hum Genet 1979;53(1):65-73.
- 660 (84) Braverman N, Lin P, Moebius FF, Obie C, Moser A, Glossmann H, et al. Mutations in the
661 gene encoding 3 beta-hydroxysteroid-delta 8, delta 7-isomerase cause X-linked dominant
662 Conradi-Hunermann syndrome. Nat Genet 1999;22(3):291-4.
- 663 (85) Castano SE, Segurado RA, Guerra TA, Simon de las HR, Lopez-Rios F, Coll Rosell MJ.
664 Ichthyosis: the skin manifestation of multiple sulfatase deficiency. Pediatr Dermatol
665 1997;14(5):369-72.
- 666 (86) Dierks T, Schmidt B, Borissenko LV, Peng J, Preusser A, Mariappan M, et al. Multiple
667 sulfatase deficiency is caused by mutations in the gene encoding the human C(alpha)-
668 formylglycine generating enzyme. Cell 2003;113(4):435-44.
- 669 (87) Happle R, Kuster W. Ichthyosis variegata: a new name for a neglected disease. J Am Acad
670 Dermatol 1997;36(3 Pt 1):500.
- 671 (88) Brusasco A, Tadini G, Cambiaghi S, Ermacora E, Grimalt R, Caputo R. A case of
672 congenital reticular ichthyosiform erythroderma--ichthyosis 'en confettis'. Dermatology
673 1994;188(1):40-5.

- 674 (89) Bygum A, Westermark P, Brandrup F. Ichthyosis prematurity syndrome: a well-defined
675 congenital ichthyosis subtype. J Am Acad Dermatol 2008;59(5 Suppl):S71-S74.
- 676 (90) Klar J, Schweiger M, Zimmerman R, Zechner R, Li H, Torma H, et al. Mutations in the
677 fatty acid transport protein 4 gene cause the ichthyosis prematurity syndrome. Am J Hum
678 Genet 2009;85(2):248-53.
- 679 (91) Phadnis SV, Griffin DR, Eady RA, Rodeck CH, Chitty LS. Prenatal diagnosis and
680 management strategies in a family with a rare type of congenital ichthyosis. Ultrasound
681 Obstet Gynecol 2007;30(6):908-10.
- 682 (92) Faghri S, Tamura D, Kraemer KH, DiGiovanna JJ. Trichothiodystrophy: a systematic
683 review of 112 published cases characterises a wide spectrum of clinical manifestations. J
684 Med Genet 2008;45(10):609-21.
- 685 (93) Kraemer KH, Patronas NJ, Schiffmann R, Brooks BP, Tamura D, DiGiovanna JJ.
686 Xeroderma pigmentosum, trichothiodystrophy and Cockayne syndrome: a complex
687 genotype-phenotype relationship. Neuroscience 2007;145(4):1388-96.
- 688 (94) Morice-Picard F, Cario-André M, Rezvani H, Sarasin A, Lacombe D, Taieb A. New
689 clinico-genetic classification of trichothiodystrophy. Am J Med Genet 2009; 149A:2020-
690 20.
- 691 (95) Camisa C, Hessel A, Rossana C, Parks A. Autosomal dominant keratoderma,
692 ichthyosiform dermatosis and elevated serum beta-glucuronidase. Dermatologica
693 1988;177(6):341-7.

- 694 (96) Korge BP, Ishida-Yamamoto A, Punter C, Dopping-Hepenstal PJ, Iizuka H, Stephenson A,
695 et al. Loricrin mutation in Vohwinkel's keratoderma is unique to the variant with
696 ichthyosis. *J Invest Dermatol* 1997;109(4):604-10.
- 697 (97) Maestrini E, Monaco AP, McGrath JA, Ishida-Yamamoto A, Camisa C, Hovnanian A, et
698 al. A molecular defect in loricrin, the major component of the cornified cell envelope,
699 underlies Vohwinkel's syndrome. *Nat Genet* 1996;13(1):70-7.
- 700 (98) Sprecher E, Ishida-Yamamoto A, Mizrahi-Koren M, Rapaport D, Goldsher D, Indelman
701 M, et al. A mutation in SNAP29, coding for a SNARE protein involved in intracellular
702 trafficking, causes a novel neurocutaneous syndrome characterized by cerebral dysgenesis,
703 neuropathy, ichthyosis, and palmoplantar keratoderma. *Am J Hum Genet* 2005;77(2):242-
704 51.
- 705 (99) Gissen P, Johnson CA, Morgan NV, Stapelbroek JM, Forshew T, Cooper WN, et al.
706 Mutations in VPS33B, encoding a regulator of SNARE-dependent membrane fusion,
707 cause arthrogryposis-renal dysfunction-cholestasis (ARC) syndrome. *Nat Genet*
708 2004;36(4):400-4.
- 709 (100) Nezelof C, Martin G, Pruvost J. [A fatal syndrome associating a micromelic dwarfism, an
710 ichtyosiform skin disorder and a severe combined immunologic deficiency. Report of a
711 case and survey of the literature (author's transl)]. *Ann Pediatr (Paris)* 1979;26(5):309-14.
- 712 (101) Jang JY, Kim KM, Kim GH, Yu E, Lee JJ, Park YS, et al. Clinical characteristics and
713 VPS33B mutations in patients with ARC syndrome. *J Pediatr Gastroenterol Nutr*
714 2009;48(3):348-54.

- 715 (102) Montpetit A, Cote S, Burstein E, Drouin C, Lapointe L, Boudreau M, et al. Disruption of
716 AP1S1, causing a novel neurocutaneous syndrome, perturbs development of the skin and
717 spinal cord. Proc Natl Acad Sci USA 2009;4(12):e1000296.
- 718 (103) Feldmeyer L, Huber M, Fellmann F, Beckmann JS, Frenk E, Hohl D. Confirmation of the
719 origin of NISCH syndrome. Hum Mutat 2006;27(5):408-10.
- 720 (104) Hadj-Rabia S, Baala L, Vabres P, Hamel-Teillac D, Jacquemin E, Fabre M, et al. Claudin-1
721 gene mutations in neonatal sclerosing cholangitis associated with ichthyosis: a tight
722 junction disease. Gastroenterology 2004;127(5):1386-90.
- 723 (105) Baala L, Hadj-Rabia S, Hamel-Teillac D, Hadchouel M, Prost C, Leal SM, et al.
724 Homozygosity mapping of a locus for a novel syndromic ichthyosis to chromosome 3q27-
725 q28. J Invest Dermatol 2002;119(1):70-6.
- 726 (106) Basel-Vanagaite L, Attia R, Ishida-Yamamoto A, Rainshtein L, Ben AD, Lurie R, et al.
727 Autosomal recessive ichthyosis with hypotrichosis caused by a mutation in ST14,
728 encoding type II transmembrane serine protease matriptase. Am J Hum Genet
729 2007;80(3):467-77.
- 730 (107) Alef T, Torres S, Hausser I, Metze D, Tursen U, Lestringant GG, et al. Ichthyosis,
731 follicular atrophoderma, and hypotrichosis caused by mutations in ST14 is associated with
732 impaired profilaggrin processing. J Invest Dermatol 2009;129(4):862-9.
- 733 (108) Lestringant GG, Kuster W, Frossard PM, Happel R. Congenital ichthyosis, follicular
734 atrophoderma, hypotrichosis, and hypohidrosis: a new genodermatosis? Am J Med Genet
735 1998;75(2):186-9.

- 736 (109) Pujol RM, Moreno A, Alomar A, De Moragas JM. Congenital ichthyosiform dermatosis
737 with linear keratotic flexural papules and sclerosing palmoplantar keratoderma. Arch
738 Dermatol 1989;125(1):103-6.
- 739 (110) Dahlqvist J, Klar J, Pigg M, van Steensel MA, Pujol RM, Vahlquist A, et al. The keratosis
740 linearis with ichthyosis congenita and sclerosing keratoderma KLICK syndrome gene
741 maps to 1 Mb chromosome 13q. 7th Annual European Society for Dermatological
742 Research Meeting, Zurich, Switzerland, Sept 2007:P523.
- 743 (111) Gottron H. Congenital angelegte symmetrische progressive erythrokeratodermie.
744 Zentralbl Haut Geschlechtskrankh 1922;4:493-4.
- 745 (112) Darier MJ. Erythro-kératodermie verruqueuse en nappes, symétrique et progressive. Bull
746 Soc Fr Dermatol Syphiligr 1911;2:252-264.
- 747 (113) Mendes da Costa S. Erythro- et keratodermia variabilis in a mother and daughter. Acta
748 Derm Venereol 1925;6:255-261.
- 749 (114) Richard G, Brown N, Rouan F, Van der Schroeff JG, Bijlsma E, Eichenfield LF, et al.
750 Genetic heterogeneity in erythrokeratodermia variabilis: novel mutations in the connexin
751 gene GJB4 (Cx30.3) and genotype-phenotype correlations. J Invest Dermatol
752 2003;120(4):601-9.
- 753 (115) Richard G, Smith LE, Bailey RA, Itin P, Hohl D, Epstein EH, Jr., et al. Mutations in the
754 human connexin gene GJB3 cause erythrokeratodermia variabilis. Nat Genet
755 1998;20(4):366-9.

- 756 (116) Macari F, Landau M, Cousin P, Mevorah B, Brenner S, Panizzon R, et al. Mutation in the
757 gene for connexin 30.3 in a family with erythrokeratoderma variabilis. Am J Hum Genet
758 2000;67(5):1296-301.
- 759 (117) van Steensel MA, Oranje AP, Van der Schroeff JG, Wagner A, van GM. The missense
760 mutation G12D in connexin30.3 can cause both erythrokeratoderma variabilis of Mendes
761 da Costa and progressive symmetric erythrokeratoderma of Gottron. Am J Med Genet A
762 2009;149A(4):657-61.
- 763 (118) Burns FS. A case of generalized congenital keratoderma with unusual involvement of the
764 eyes, ears, and nasal and buccous membranes. J Cutan Dis 1915;33:255-260.
- 765 (119) Skinner BA, Greist MC, Norins AL. The keratitis, ichthyosis, and deafness (KID)
766 syndrome. Arch Dermatol 1981;117(5):285-9.
- 767 (120) Gulzow J, Anton-Lamprecht I. [Ichthyosis hystrix gravior typus Rheydt: an otologic-
768 dermatologic syndrome (author's transl)]. Laryngol Rhinol Otol (Stuttg) 1977;56(11):949-
769 55.
- 770 (121) Richard G, Rouan F, Willoughby CE, Brown N, Chung P, Ryynanen M, et al. Missense
771 mutations in GJB2 encoding connexin-26 cause the ectodermal dysplasia keratitis-
772 ichthyosis-deafness syndrome. Am J Hum Genet 2002;70(5):1341-8.
- 773 (122) van Steensel MA, Steijlen PM, Bladergroen RS, Hoefsloot EH, van Ravenswaaij-Arts
774 CM, van GM. A phenotype resembling the Clouston syndrome with deafness is associated
775 with a novel missense GJB2 mutation. J Invest Dermatol 2004;123(2):291-3.

- 776 (123) Netherton EW. A unique case of trichorrhexis nodosa; bamboo hairs. AMA Arch Derm
777 1958;78(4):483-7.
- 778 (124) Levy SB, Goldsmith LA. The peeling skin syndrome. J Am Acad Dermatol 1982;7(5):606-
779 13.
- 780 (125) Chavanas S, Bodemer C, Rochat A, Hamel-Teillac D, Ali M, Irvine AD, et al. Mutations in
781 SPINK5, encoding a serine protease inhibitor, cause Netherton syndrome. Nat Genet
782 2000;25(2):141-2.
- 783 (126) Komatsu N, Suga Y, Saijoh K, Liu AC, Khan S, Mizuno Y, et al. Elevated human tissue
784 kallikrein levels in the stratum corneum and serum of peeling skin syndrome-type B
785 patients suggests an over-desquamation of corneocytes. J Invest Dermatol
786 2006;126(10):2338-42.
- 787 (127) Vohwinkel KH. Keratoma hereditarium mutilans. Arch Derm Syph 1929;158:354-364.
- 788 (128) Maestrini E, Korge BP, Ocana-Sierra J, Calzolari E, Cambiaghi S, Scudder PM, et al. A
789 missense mutation in connexin26, D66H, causes mutilating keratoderma with
790 sensorineural deafness (Vohwinkel's syndrome) in three unrelated families. Hum Mol
791 Genet 1999;8(7):1237-43.
- 792 (129) Stulli, L. Di una variata cutanea. Lettera al direttore dell'Antologia. Estratti dall'Antologia
793 di Firenze 1826;71-72,1-3.
- 794 (130) Fischer J, Bouadjar B, Heilig R, Huber M, Lefevre C, Jobard F, et al. Mutations in the
795 gene encoding SLURP-1 in Mal de Meleda. Hum Mol Genet 2001;10(8):875-80.

- 796 (131) Papillon M, Lefèvre P. Deux cas de kératodermie palmaire et plantaire symétrique
797 familiale (maladie de Meleda) chez le frère et la soeur. Coexistence dans les deux cas
798 d'altérations dentaires graves. Bull Soc Fr Dermatol Syphiligr 1924;31:82-87.
- 799 (132) Toomes C, James J, Wood AJ, Wu CL, McCormick D, Lench N, et al. Loss-of-function
800 mutations in the cathepsin C gene result in periodontal disease and palmoplantar keratosis.
801 Nat Genet 1999;23(4):421-4.
- 802 (133) Coulombe PA, Hutton ME, Letai A, Hebert A, Paller AS, Fuchs E. Point mutations in
803 human keratin 14 genes of epidermolysis bullosa simplex patients: genetic and functional
804 analyses. Cell 1991;66(6):1301-11.
- 805 (134) Lane EB, Rugg EL, Navsaria H, Leigh IM, Heagerty AH, Ishida-Yamamoto A, et al. A
806 mutation in the conserved helix termination peptide of keratin 5 in hereditary skin
807 blistering. Nature 1992;356(6366):244-6.
- 808 (135) Fine JD, Eady RA, Bauer EA, Bauer JW, Bruckner-Tuderman L, Heagerty A, et al. The
809 classification of inherited epidermolysis bullosa (EB): Report of the Third International
810 Consensus Meeting on Diagnosis and Classification of EB. J Am Acad Dermatol
811 2008;58(6):931-50.
- 812 (136) Haenssle HA, Finkenrath A, Hausser I, Oji V, Traupe H, Hennies HC, et al. Effective
813 treatment of severe thermoregulation by oral retinoids in a patient with recessive
814 congenital lamellar ichthyosis. Clin Exp Dermatol 2008;33(5):578-81.

- 815 (137) DiGiovanna JJ, Priolo M, Itin P. Approach towards a new classification for ectodermal
816 dysplasias: integration of the clinical and molecular knowledge. Am J Med Genet A
817 2009;149A(9):2068-70.
- 818 (138) Salinas CF, Jorgenson RJ, Wright JT, DiGiovanna JJ, Fete MD. 2008 International
819 Conference on Ectodermal Dysplasias Classification: Conference report. Am J Med Genet
820 A 2009;149A(9):1958-69.
- 821 (139) Plantin P, Gavanou J, Jouan N, Leroy JP, Guillet G. [Collodion skin: a misdiagnosed but
822 frequent clinical aspect of anhidrotic ectodermal dysplasia during the neonatal period].
823 Ann Dermatol Venereol 1992;119(11):821-3.
- 824 (140) Thomas C, Suranyi E, Pride H, Tyler W. A child with hypohidrotic ectodermal dysplasia
825 with features of a collodion membrane. Pediatr Dermatol 2006;23(3):251-4.
- 826 (141) Navarro CL, De Sandre-Giovannoli A, Bernard R, Boccaccio I, Boyer A, Genevieve D, et
827 al. Lamin A and ZMPSTE24 (FACE-1) defects cause nuclear disorganization and identify
828 restrictive dermopathy as a lethal neonatal laminopathy. Hum Mol Genet
829 2004;13(20):2493-503.
- 830 (142) Lowry RB, Machin GA, Morgan K, Mayock D, Marx L. Congenital contractures, edema,
831 hyperkeratosis, and intrauterine growth retardation: a fatal syndrome in Hutterite and
832 Mennonite kindreds. Am J Med Genet 1985;22(3):531-43.
- 833 (143) Antoine T. Ein Fall von allgemeiner, angeborener Haut-atrophie. Monatsschr Geburtsh
834 Gynaekol 1929;81:276-83.

- 835 (144) Manning MA, Cunniff CM, Colby CE, El-Sayed YY, Hoyme HE. Neu-Laxova syndrome:
836 detailed prenatal diagnostic and post-mortem findings and literature review. Am J Med
837 Genet A 2004;125A(3):240-9.
- 838 (145) Happle R, Koch H, Lenz W. The CHILD syndrome. Congenital hemidysplasia with
839 ichthyosiform erythroderma and limb defects. Eur J Pediatr 1980;134(1):27-33.
- 840 (146) Konig A, Happle R, Bornholdt D, Engel H, Grzeschik KH. Mutations in the NSDHL gene,
841 encoding a 3beta-hydroxysteroid dehydrogenase, cause CHILD syndrome. Am J Med
842 Genet 2000;90(4):339-46.
- 843 (147) Happle R, Matthiass HH, Macher E. Sex-linked chondrodysplasia punctata? Clin Genet
844 1977;11(1):73-6.
- 845 (148) Darier J. Psorospermose folliculaire végétante. Ann Dermatol Syphiligr (Paris) 1889;10,
846 597-612.
- 847 (149) White J. A case of keratosis (ichthyosis) follicularis. J Cutan Dis 1889;7,201-209.
- 848 (150) Hailey H, Hailey H. Familial benign chronic pemphigus. Arch Dermatol Syph
849 1939;39,679-685.
- 850 (151) Sakuntabhai A, Ruiz-Perez V, Carter S, Jacobsen N, Burge S, Monk S, et al. Mutations in
851 ATP2A2, encoding a Ca²⁺ pump, cause Darier disease. Nat Genet 1999;21(3):271-7.
- 852 (152) Hu Z, Bonifas JM, Beech J, Bench G, Shigihara T, Ogawa H, et al. Mutations in ATP2C1,
853 encoding a calcium pump, cause Hailey-Hailey disease. Nat Genet 2000;24(1):61-5.

- 854 (153) Madison KC. Barrier function of the skin: "la raison d'etre" of the epidermis. J Invest
855 Dermatol 2003;121(2):231-41.
- 856 (154) Attenborough D. Life on Earth. Boston: Little, Brown; 1980.
- 857 (155) Blank IH. Further observations on factors which influence the water content of the stratum
858 corneum. J Invest Dermatol 1953;21(4):259-71.
- 859 (156) Winsor T, Burge GE. Differential roles of layers of human epigastric skin on diffusion rate
860 of water. Arch Intern Med 1944;74,428-436.
- 861 (157) Elias PM. Epidermal lipids, barrier function, and desquamation. J Invest Dermatol
862 1983;80 Suppl:44s-9s.
- 863 (158) Williams ML. The ichthyoses--pathogenesis and prenatal diagnosis: a review of recent
864 advances. Pediatr Dermatol 1983;1(1):1-24.
- 865 (159) Demerjian M, Crumrine DA, Milstone LM, Williams ML, Elias PM. Barrier dysfunction
866 and pathogenesis of neutral lipid storage disease with ichthyosis (Chanarin-Dorfman
867 syndrome). J Invest Dermatol 2006;126(9):2032-8.
- 868 (160) Elias PM, Schmuth M, Uchida Y, Rice RH, Behne M, Crumrine D, et al. Basis for the
869 permeability barrier abnormality in lamellar ichthyosis. Exp Dermatol 2002;11(3):248-56.
- 870 (161) Elias PM, Crumrine D, Rassner U, Hachem JP, Menon GK, Man W, et al. Basis for
871 abnormal desquamation and permeability barrier dysfunction in RXLI. J Invest Dermatol
872 2004;122(2):314-9.

- 873 (162) Hachem JP, Houben E, Crumrine D, Man MQ, Schurer N, Roelandt T, et al. Serine
874 protease signaling of epidermal permeability barrier homeostasis. *J Invest Dermatol*
875 2006;126(9):2074-86.
- 876 (163) Holleran WM, Ginns EI, Menon GK, Grundmann JU, Fartasch M, McKinney CE, et al.
877 Consequences of beta-glucocerebrosidase deficiency in epidermis. Ultrastructure and
878 permeability barrier alterations in Gaucher disease. *J Clin Invest* 1994;93(4):1756-64.
- 879 (164) Schmuth M, Fluhr JW, Crumrine DC, Uchida Y, Hachem JP, Behne M, et al. Structural
880 and functional consequences of loricrin mutations in human loricrin keratoderma
881 (Vohwinkel syndrome with ichthyosis). *J Invest Dermatol* 2004;122(4):909-22.
- 882 (165) Descargues P, Deraison C, Bonnart C, Kreft M, Kishibe M, Ishida-Yamamoto A, et al.
883 Spink5-deficient mice mimic Netherton syndrome through degradation of desmoglein 1 by
884 epidermal protease hyperactivity. *Nat Genet* 2005;37(1):56-65.
- 885 (166) Yanagi T, Akiyama M, Nishihara H, Sakai K, Nishie W, Tanaka S, et al. Harlequin
886 ichthyosis model mouse reveals alveolar collapse and severe fetal skin barrier defects.
887 *Hum Mol Genet* 2008;17(19):3075-83.
- 888 (167) Matsuki M, Yamashita F, Ishida-Yamamoto A, Yamada K, Kinoshita C, Fushiki S, et al.
889 Defective stratum corneum and early neonatal death in mice lacking the gene for
890 transglutaminase 1 (keratinocyte transglutaminase). *Proc Natl Acad Sci USA*
891 1998;95(3):1044-9.
- 892 (168) Epp N, Furstenberger G, Muller K, de JS, Leitges M, Hausser I, et al. 12R-lipoxygenase
893 deficiency disrupts epidermal barrier function. *J Cell Biol* 2007;177(1):173-82.

- 894 (169) Furuse M, Hata M, Furuse K, Yoshida Y, Haratake A, Sugitani Y, et al. Claudin-based tight
895 junctions are crucial for the mammalian epidermal barrier: a lesson from claudin-1-
896 deficient mice. *J Cell Biol* 2002;156(6):1099-111.
- 897 (170) Feingold KR. The regulation of epidermal lipid synthesis by permeability barrier
898 requirements. *Crit Rev Ther Drug Carrier Syst* 1991;8(3):193-210.
- 899 (171) Ghadially R, Brown BE, Sequeira-Martin SM, Feingold KR, Elias PM. The aged
900 epidermal permeability barrier. Structural, functional, and lipid biochemical abnormalities
901 in humans and a senescent murine model. *J Clin Invest* 1995;95(5):2281-90.
- 902 (172) Williams ML, Elias PM. From basket weave to barrier. Unifying concepts for the
903 pathogenesis of the disorders of cornification. *Arch Dermatol* 1993;129(5):626-9.
- 904 (173) Juanes SD, Epp N, Latzko S, Neumann M, Furstenberger G, Hausser I, et al. Development
905 of an Ichthyosiform Phenotype in Alox12b-Deficient Mouse Skin Transplants. *J Invest*
906 *Dermatol* 2009;129(6):1429-36.
- 907 (174) Ballabio A, Parenti G, Carrozzo R, Sebastio G, Andria G, Buckle V, et al. Isolation and
908 characterization of a steroid sulfatase cDNA clone: genomic deletions in patients with X-
909 chromosome-linked ichthyosis. *Proc Natl Acad Sci USA* 1987;84(13):4519-23.
- 910 (175) Chipev CC, Korge BP, Markova N, Bale SJ, DiGiovanna JJ, Compton JG, et al. A leucine-
911 ---proline mutation in the H1 subdomain of keratin 1 causes epidermolytic hyperkeratosis.
912 *Cell* 1992;70(5):821-8.

- 913 (176) Compton JG, DiGiovanna JJ, Santucci SK, Kearns KS, Amos CI, Abangan DL, et al.
914 Linkage of epidermolytic hyperkeratosis to the type II keratin gene cluster on chromosome
915 12q. *Nat Genet* 1992;1(4):301-5.
- 916 (177) Grzeschik KH, Bornholdt D, Oeffner F, Konig A, del Carmen BM, Enders H, et al.
917 Deficiency of PORCN, a regulator of Wnt signaling, is associated with focal dermal
918 hypoplasia. *Nat Genet* 2007;39(7):833-5.
- 919 (178) Jansen GA, Ofman R, Ferdinandusse S, Ijlst L, Muijsers AO, Skjeldal OH, et al. Refsum
920 disease is caused by mutations in the phytanoyl-CoA hydroxylase gene. *Nat Genet*
921 1997;17(2):190-3.
- 922 (179) Jansen GA, Waterham HR, Wanders RJ. Molecular basis of Refsum disease: sequence
923 variations in phytanoyl-CoA hydroxylase (PHYH) and the PTS2 receptor (PEX7). *Hum*
924 *Mutat* 2004;23(3):209-18.
- 925 (180) Oeffner F, Fischer G, Happel R, Konig A, Betz RC, Bornholdt D, et al. IFAP syndrome is
926 caused by deficiency in MBTPS2, an intramembrane zinc metalloprotease essential for
927 cholesterol homeostasis and ER stress response. *Am J Hum Genet* 2009;84(4):459-67.
- 928 (181) Rothnagel JA, Dominey AM, Dempsey LD, Longley MA, Greenhalgh DA, Gagne TA, et
929 al. Mutations in the rod domains of keratins 1 and 10 in epidermolytic hyperkeratosis.
930 *Science* 1992;257(5073):1128-30.
- 931 (182) Rothnagel JA, Traupe H, Wojcik S, Huber M, Hohl D, Pittelkow MR, et al. Mutations in
932 the rod domain of keratin 2e in patients with ichthyosis bullosa of Siemens. *Nat Genet*
933 1994;7(4):485-90.

- 934 (183) Smith FJ, Irvine AD, Terron-Kwiatkowski A, Sandilands A, Campbell LE, Zhao Y, et al.
935 Loss-of-function mutations in the gene encoding filaggrin cause ichthyosis vulgaris. *Nat*
936 *Genet* 2006;38(3):337-42.
- 937 (184) Stefanini M, Lagomarsini P, Giliani S, Nardo T, Botta E, Peserico A, et al. Genetic
938 heterogeneity of the excision repair defect associated with trichothiodystrophy.
939 *Carcinogenesis* 1993;14(6):1101-5.
- 940 (185) Takayama K, Salazar EP, Broughton BC, Lehmann AR, Sarasin A, Thompson LH, et al.
941 Defects in the DNA repair and transcription gene ERCC2(XPD) in trichothiodystrophy.
942 *Am J Hum Genet* 1996;58(2):263-70.
- 943 (186) Tsuji S, Choudary PV, Martin BM, Stubblefield BK, Mayor JA, Barranger JA, et al. A
944 mutation in the human glucocerebrosidase gene in neuronopathic Gaucher's disease. *N*
945 *Engl J Med* 1987;316(10):570-5.
- 946 (187) Mizrahi-Koren M, Shemer S, Morgan M, Indelman M, Khamaysi Z, Petronius D, et al.
947 Homozygosity mapping as a screening tool for the molecular diagnosis of hereditary skin
948 diseases in consanguineous populations. *J Am Acad Dermatol* 2006;55(3):393-401.
- 949 (188) Lugassy J, Hennies HC, Indelman M, Khamaysi Z, Bergman R, Sprecher E. Rapid
950 detection of homozygous mutations in congenital recessive ichthyosis. *Arch Dermatol Res*
951 2008;300(2):81-5.
- 952 (189) Roop D. Defects in the barrier. *Science* 1995;267(5197):474-5.

- 953 (190) Bitoun E, Bodemer C, Amiel J, de PY, Stoll C, Calvas P, et al. Prenatal diagnosis of a
954 lethal form of Netherton syndrome by SPINK5 mutation analysis. *Prenat Diagn*
955 2002;22(2):121-6.
- 956 (191) Muller FB, Hausser I, Berg D, Casper C, Maiwald R, Jung A, et al. Genetic analysis of a
957 severe case of Netherton syndrome and application for prenatal testing. *Br J Dermatol*
958 2002;146(3):495-9.
- 959 (192) Sprecher E, Chavanas S, DiGiovanna JJ, Amin S, Nielsen K, Prendiville JS, et al. The
960 spectrum of pathogenic mutations in SPINK5 in 19 families with Netherton syndrome:
961 implications for mutation detection and first case of prenatal diagnosis. *J Invest Dermatol*
962 2001;117(2):179-87.
- 963 (193) Rothnagel JA, Longley MA, Holder RA, Kuster W, Roop DR. Prenatal diagnosis of
964 epidermolytic hyperkeratosis by direct gene sequencing. *J Invest Dermatol*
965 1994;102(1):13-6.
- 966 (194) Rothnagel JA, Lin MT, Longley MA, Holder RA, Hazen PG, Levy ML, et al. Prenatal
967 diagnosis for keratin mutations to exclude transmission of epidermolytic hyperkeratosis.
968 *Prenat Diagn* 1998;18(8):826-30.
- 969 (195) Tsuji-Abe Y, Akiyama M, Nakamura H, Takizawa Y, Sawamura D, Matsunaga K, et al.
970 DNA-based prenatal exclusion of bullous congenital ichthyosiform erythroderma at the
971 early stage, 10 to 11 weeks' of pregnancy, in two consequent siblings. *J Am Acad*
972 *Dermatol* 2004;51(6):1008-11.

- 973 (196) Sillen A, Holmgren G, Wadelius C. First prenatal diagnosis by mutation analysis in a
974 family with Sjogren-Larsson syndrome. *Prenat Diagn* 1997;17(12):1147-9.
- 975 (197) Yanagi T, Akiyama M, Sakai K, Nagasaki A, Ozawa N, Kosaki R, et al. DNA-based
976 prenatal exclusion of harlequin ichthyosis. *J Am Acad Dermatol* 2008;58(4):653-6.
- 977 (198) Akiyama M, Titeux M, Sakai K, McMillan JR, Tonasso L, Calvas P, et al. DNA-based
978 prenatal diagnosis of harlequin ichthyosis and characterization of ABCA12 mutation
979 consequences. *J Invest Dermatol* 2007;127(3):568-73.
- 980 (199) Anton-Lamprecht I. The Skin. In: Papdimitriou JM, Henderon DW, Sagnolo DV, eds.
981 Diagnostic ultrastructure of non-neoplastic diseases. Diagnostic ultrastructure of non-
982 neoplastic diseases. Edinburgh: Churchill-Livingstone; 1992. p. 459-550.
- 983 (200) Anton-Lamprecht I. Ultrastructural identification of basic abnormalities as clues to genetic
984 disorders of the epidermis. *J Invest Dermatol* 1994;103(5 Suppl):6S-12S.
- 985 (201) Anton-Lamprecht I, Schnyder UW. Ultrastructural distinction of autosomal dominant
986 ichthyosis vulgaris and X-linked recessive ichthyosis. *Clin Genet* 1976;10(4):245-7.
- 987 (202) Oji V, Seller N, Sandilands A, Gruber R, Gerss J, Huffmeier U, et al. Ichthyosis vulgaris:
988 novel FLG mutations in the German population and high presence of CD1a+ cells in the
989 epidermis of the atopic subgroup. *Br J Dermatol* 2009; 160(4):771-81.
- 990 (203) Dale BA, Holbrook KA, Fleckman P, Kimball JR, Brumbaugh S, Sybert VP.
991 Heterogeneity in harlequin ichthyosis, an inborn error of epidermal keratinization: variable

- 992 morphology and structural protein expression and a defect in lamellar granules. *J Invest*
993 *Dermatol* 1990;94(1):6-18.
- 994 (204) Akiyama M, Sakai K, Sato T, McMillan JR, Goto M, Sawamura D, et al. Compound
995 heterozygous ABCA12 mutations including a novel nonsense mutation underlie harlequin
996 ichthyosis. *Dermatology* 2007;215(2):155-9.
- 997 (205) Ishida-Yamamoto A. Loricrin keratoderma: a novel disease entity characterized by nuclear
998 accumulation of mutant loricrin. *J Dermatol Sci* 2003;31(1):3-8.
- 999 (206) Arnold ML, Anton-Lamprecht I, Melz-Rothfuss B, Hartschuh W. Ichthyosis congenita
1000 type III. Clinical and ultrastructural characteristics and distinction within the
1001 heterogeneous ichthyosis congenita group. *Arch Dermatol Res* 1988;280(5):268-78.
- 1002 (207) Brusasco A, Gelmetti C, Tadini G, Caputo R. Ichthyosis congenita type IV: a new case
1003 resembling diffuse cutaneous mastocytosis. *Br J Dermatol* 1997;136(3):377-9.
- 1004 (208) Niemi KM, Kanerva L, Kuokkanen K. Recessive ichthyosis congenita type II. *Arch*
1005 *Dermatol Res* 1991;283(4):211-8.
- 1006 (209) Niemi KM, Kuokkanen K, Kanerva L, Ignatius J. Recessive ichthyosis congenita type IV.
1007 *Am J Dermatopathol* 1993;15(3):224-8.
- 1008 (210) Niemi KM, Kanerva L, Kuokkanen K, Ignatius J. Clinical, light and electron microscopic
1009 features of recessive congenital ichthyosis type I. *Br J Dermatol* 1994;130(5):626-33.

- 1010 (211) Pigg M, Gedde-Dahl T, Jr., Cox D, Hausser I, Anton-Lamprecht I, Dahl N. Strong founder
1011 effect for a transglutaminase 1 gene mutation in lamellar ichthyosis and congenital
1012 ichthyosiform erythroderma from Norway. Eur J Hum Genet 1998;6(6):589-96.
- 1013 (212) Descargues P, Deraison C, Prost C, Fraitag S, Mazereeuw-Hautier J, D'Alessio M, et al.
1014 Corneodesmosomal cadherins are preferential targets of stratum corneum trypsin- and
1015 chymotrypsin-like hyperactivity in Netherton syndrome. J Invest Dermatol
1016 2006;126(7):1622-32.
- 1017 (213) Wells RS, Kerr CB. The histology of ichthyosis. J Invest Dermatol 1966;46(6):530-535.
- 1018 (214) Fleckman P, Brumbaugh S. Absence of the granular layer and keratohyalin define a
1019 morphologically distinct subset of individuals with ichthyosis vulgaris. Exp Dermatol
1020 2002;11(4):327-36.
- 1021 (215) Bergman R, Khamaysi Z, Sprecher E. A unique pattern of dyskeratosis characterizes
1022 epidermolytic hyperkeratosis and epidermolytic palmoplantar keratoderma. Am J
1023 Dermatopathol 2008;30(2):101-5.
- 1024 (216) Ross R, DiGiovanna JJ, Capaldi L, Argenyi Z, Fleckman P, Robinson-Bostom L.
1025 Histopathologic characterization of epidermolytic hyperkeratosis: a systematic review of
1026 histology from the National Registry for Ichthyosis and Related Skin Disorders. J Am
1027 Acad Dermatol 2008;59(1):86-90.
- 1028 (217) Sperling LC, DiGiovanna JJ. "Curly" wood and tiger tails: an explanation for light and
1029 dark banding with polarization in trichothiodystrophy. Arch Dermatol 2003;139(9):1189-
1030 92.