

both have more than 700 beds, and are teaching hospitals. Targets of audits were consecutive patients who received RBC or platelet transfusions during hospitalization and were discharged from the hospitals between July 2006 and September 2006. These patients were identified from the database, as well as from lists of patients transfused at the hospitals. Five of the authors (M.S, Y.I, T.S, H.S and K.Y) participated in the chart review. Through reviews of medical records, reviewers identified conditions (clinical course, findings of physical examinations and laboratory tests) in which a blood product was used. From the information collected, we attempted to determine the appropriateness of blood product use. Appropriateness was judged based on the 'Guidelines for blood transfusions' developed by the ad-hoc group of the Japanese Ministry of Health, Labor, and Welfare [17]. The Japanese guidelines were developed based on international guidelines and other national guidelines [2,18–21], and its recommendations are substantially similar to those of other guidelines. We also checked the accuracy of administrative claim data on timing, amount and type of blood product used. Some patients (e.g., those with haematopoietic diseases or chronic anaemia) received more than one blood transfusion during their hospitalization. For these cases, we reviewed the first episode of transfusion and judged the appropriateness of the transfusion decision. Differences in judgment were discussed among reviewers and one external expert until a final agreement on appropriateness was reached. The authors fed back results of the audits to chiefs of transfusion committees of each hospital, and obtained their consent.

Through audits at the two hospitals, we identified diagnoses, surgical procedures and comorbidities underlying the use of RBC and platelet transfusions. Using the ICD-10 code and surgical procedure classification code from the Ministry of Health and Welfare in Japan, 2006), we surveyed the frequency of these risk factors in the database, as well as proportions of transfused patients among patients with each risk factor.

Logistic regression model to predict proportions of transfused patients at hospitals

Using DPC data of 587 045 cases provided by 73 acute-care hospitals, we developed a multiple logistic regression model to predict proportions of transfused patients at each hospital. In this model, the dependent variable, the occurrence of blood transfusion during hospitalization, was defined as a binary variable. Each case received at least one blood transfusion during hospitalization was classified as '1; Used'. Each case that did not receive any blood transfusions during hospitalization was classified as '0; Not used'. Independent variables included risk factors, which either were identified through chart reviews at the two hospitals or

were reported in previous studies to be associated with the risk of transfusion [14–16,22]. Performance of the logistic regression model was assessed by calculating the area under the receiver operating characteristic (ROC) curve. The area under a ROC curve relating relative proportions of correctly and incorrectly classified predictions over a wide and continuous range of threshold levels can measure discrimination capacity of the prediction rule [23,24]. Using the regression model, we calculated expected proportions of patients transfused at each hospital. By comparing observed (O) and expected (E) values, we calculated O/E ratios.

Case-mix adjusted model to predict total units of blood products transfused from the distribution of diagnostic groups at hospitals

The second model utilized the distribution of diagnostic groups to predict the total units of RBCs or platelets transfused at each hospital. In this model, we calculated the mean units of blood products used per case for each diagnostic group (defined by the first 10 digits, representing disease code + surgical code), and used these mean values as expected values for RBC or platelet use in that diagnostic group. All cases in each group were used to calculate means. Mean units used per case in the i th group can be represented by $Q(i)$, the number of cases belonging to that group can be represented by $M(i)$, and the total number of diagnostic groups can be represented by W . Thus, the expected total units of blood product used in the hospital can be described by the following formula:

$$\sum_{i=1}^W \{Q(i) \times N(i)\}$$

From the distribution of diagnostic groups in each hospital, we estimated total units of RBCs or platelets used at the hospital. By comparing expected and observed values, we calculated O/E ratios for total units transfused at each hospital. Predictive ability of this model was assessed by the correlation coefficient, r^2 . Moreover, we compared results of the retrospective audit with O/E ratios. To assess case-mix adjusted blood product use, expected total RBC or platelet use was compared with observed values to obtain observed to expected ratios (O/E ratios) at each hospital. We used SPSS (Version 11, SPSS Inc.) for all statistical analyses. **3**

Results

Risk factors of blood transfusion

Table 1 shows characteristics of the hospitals in Japan that were studied, as well as the frequency and amount of RBC products and concentrated platelets transfused at the

Table 1 Characteristics of 73 studied hospitals and the distribution of blood products use in each hospital

Factor	Range	Percentile		
		25%	50%	75%
No of bed	43–1106	239	350	530
No of hospitalization (month)	93–2468	361	559	953
Mean length of stay (days)	9.6–41.0	13.2	15.2	18.2
Hospital mortality rate (%)	0.1–8.8%	3.4	4.5	5.2
No of patients with RBC transfusion (100 beds/month)	0.8–22.6	5.2	7.5	10.1
No of patients with PLT transfusion (100 beds/month)	0–6.1	0.65	1.2	2.4
All RBC units transfused (Units/100 beds/month)	2.7–228.6	33.3	50.2	73.7
All PLT units transfused (Units/100 beds/month)	0–499.3	19.1	45.7	120.0

RBC, red blood cell; PLT, platelet.

hospitals. Of the 587 045 cases targeted, 33 155 (5.7%) patients received RBC transfusions, and 7065 (1.2%) patients received platelet transfusions. Frequency of blood transfusion and number of units transfused per patient-bed varied greatly between hospitals. Based on retrospective audits at two hospitals (Hospitals A and B), Table 2

summarizes underlying conditions for use of RBC and platelet transfusions, as well as appropriateness of blood transfusions at each hospital.

From the retrospective audit, we identified patient characteristics, diagnoses and surgical procedures potentially associated with risk of blood transfusion. Table 3 shows the distribution of these potential risk factors in 587 045 cases in the database, as well as the probability of receiving ≥ 1 transfusion during hospitalization of the population with each risk factor. Cases with the risk were identified using the disease classification (ICD-10) code and surgery classification code (from the fee schedule system for medical reimbursement by the Ministry of Health and Welfare, Japan). Factors with the greatest risk for RBC transfusion included cardiovascular surgeries, haematopoietic malignancies, aplastic anaemia and myelodysplastic syndrome (MDS). Factors with the greatest risk for platelet transfusion included blood diseases such as acute leukaemia, aplastic anaemia and MDS, as well as cardiovascular surgery.

Logistic regression models predicting proportions of patients with transfusions

Table 4 shows risk factors for RBC and platelet transfusions and coefficients and odds ratios (ORs: 95% confidence intervals) of multiple logistic regression analyses. Greatest

Underlying condition	Hospital A		Hospital B					
	<i>n</i>	%	Appropriate transfusion	%	<i>n</i>	%	Appropriate transfusion	%
<i>RBC transfusion</i>								
Acute bleeding	52	20	43	83	43	16	33	77
Perioperative								
Cardiovascular surgery	39	15	24	62	35	13	25	71
Others	55	21	30	55	50	19	33	66
Chronic anaemia								
Haematopoietic malignancies/disorders	35	14	26	74	67	25	60	90
Others	52	20	40	77	47	18	42	89
Chronic bleeding								
All	24	9	23	96	24	9	23	96
All	257	100	186	72	266	100	216	81
<i>Platelet transfusion</i>								
Haematopoietic malignancies/disorders								
Cancer	30	38	4	13	51	59	40	78
<i>Surgery</i>								
Cardiovascular surgery								
Others	23	29	4	17	12	14	10	83
Others	5	6	3	60	4	5	0	0
Others	10	13	3	30	12	14	2	17
All	79	100	23	29	87	100	56	64

GI indicates gastrointestinal; MDS, Myelodysplastic syndrome; RBC, red blood cell.

Table 2 Underlying conditions of transfusion and proportions of appropriate transfusion

Table 3 Prevalence of risk factors of transfusion and proportions of patients transfused during hospitalization

Factor	n	Prevalence		Proportion of patients with transfusion	
		(1000 admissions)	(%)	RBC transfusion (%)	PLT transfusion (%)
Age					
20-64	259588	442.2	3.5	1.0	
65-79	220247	375.2	6.5	1.5	
80+	107210	182.6	9.0	1.1	
Blood disorders					
Acute leukaemia	1986	3.4	59	60	
Chronic leukaemia	476	0.8	28	16	
Malignant lymphoma	1625	2.8	36	17	
Multiple myeloma	6032	10.3	14	11	
Aplastic anaemia	940	1.6	58	34	
MDS	1476	2.5	63	37	
DIC	4372	7.4	43	26	
Cancer*					
Without chemotherapy	93103	158.6	8.3	0.8	
With chemotherapy	42088	71.7	4.7	1.0	
Surgery					
Cardiovascular surgery					
Without CPB	2205	3.8	50	18	
With CPB	2629	4.5	73	43	
Hip fracture surgery					
Severe trauma	7457	12.7	28	0.9	
Obstetric bleeding	823	1.4	10	1.5	
GI bleeding	19808	33.7	34	2.5	
Chronic renal failure	24691	42.1	17	2.1	

GI, gastrointestinal; DIC, disseminated intravascular coagulation; CPB, cardiopulmonary bypass; MDS, Myelodysplastic syndrome; RBC, red blood cell.

risk factors for RBC transfusion were cardiovascular surgery with cardiopulmonary bypass (CPB) (OR = 106.9), followed by acute leukaemia (OR = 46.9), MDS (OR = 39.2), cardiovascular surgery without CPB (OR = 35.0), gastrointestinal haemorrhage (OR = 13.2), and hip fracture surgery (OR = 11.0). Risk of RBC transfusion increased with patient age.

Greatest risk factors for platelet transfusion were acute leukaemia (OR = 310) and cardiovascular surgery with CPB (OR = 185), followed by MDS (OR = 78.9), cardiovascular surgery without CPB (OR = 46.5), disseminated intravascular coagulation (DIC) (OR = 43.2), multiple myeloma (OR = 39.5) and aplastic anaemia (OR = 33.6). No association was observed between patient age and risk of platelet transfusion.

These logistic regression models showed good prediction performances. Areas under ROC curves for proportions of patients with RBC transfusions and platelet transfusions were 0.83 and 0.90, respectively.

Case-mix adjusted models predicting total units of blood products used

Figure 1 shows expected vs. observed use of RBCs and platelets at each hospital. The goodness-of-fit of this model, assessed using χ^2 for linear regression, was 0.88 for RBC use and 0.57 for platelet use. Therefore, use of blood products at each hospital was successfully predicted by the case-mix adjusted model.

Table 5 compares results of the medical chart review (percentage of appropriateness) and O/E ratios calculated using the two prediction models. In general, proportions of appropriate transfusions were higher for RBC than for platelet transfusions. Also, Hospital B showed better performance in terms of appropriate transfusions than did Hospital A. Both models successfully assessed the proportion of appropriate transfusions.

Discussion

Due to critical situations and complex clinical conditions underlying the use of blood products, it is often difficult for physicians to make appropriate decisions about blood transfusions. Surveys of blood product usage for specific diagnoses have demonstrated wide variation in comparable patients at different institutions [8,26-29]. Some studies have shown that a system approach is effective for improving transfusion practices. System approaches include implementation of institutional guidelines, education, prospective audits and approval of transfusion orders [12]. Of these approaches, 'clinical audits' with feedback can effectively decrease the amount of blood product used [28-35]. However, these effects on blood product use are often temporary, and continuous efforts are required for maintenance of good transfusion practices [30,36]. Moreover, audits require considerable labour and cost. Therefore, it is difficult to implement audits for routine and continuous evaluations. If another method were available for routine evaluations, then it would contribute greatly to comparison of transfusion performance between hospitals and resultant improvement in transfusion practices at hospitals.

We proposed methods which utilizing administrative data to evaluate hospital-wide use of blood products in Japan. Few studies have employed administrative data to perform such comparisons. Comparison of blood usage across hospitals usually is difficult given that risks of transfusion differ between patients and between hospitals. On the contrary, some studies have demonstrated that a patient classification

Factor	RBC			PLT		
	Regression coefficient	Odds ratio	95% CI	Regression coefficient	Odds ratio	95% CI
Age*						
65-74	0.40	1.5	1.4-1.5	0.11	1.1	1.0-1.2
80+	0.75	2.1	2.1-2.2	-0.06	0.9	0.9-1.0
Female	-0.12	0.89	0.86-0.91	0.06	1.1	1.0-1.1
Haematopoietic disorders						
Acute leukaemia	3.85	46.9	42.4-51.8	5.74	310	278-346
Chronic leukaemia	2.33	10.3	8.1-12.9	3.26	26.2	19.2-35.6
Aplastic anaemia	3.12	22.7	19.4-26.6	3.52	33.6	27.3-41.4
Multiple myeloma	2.84	17.2	15.4-19.2	3.68	39.5	33.8-46.1
Malignant lymphoma	1.51	4.5	4.1-4.9	3.10	22.2	20.0-24.7
MDS	3.67	39.2	34.9-44.1	4.37	78.9	68.6-90.7
DIC	2.44	11.4	10.6-12.3	3.77	43.2	39.3-47.5
Obstetric bleeding	1.82	6.2	4.9-7.8	0.99	2.7	1.4-5.0
Severe trauma	2.65	14.1	12-16.6	2.08	8.0	5.9-11
Cardiovascular surgery						
Without CPB	3.56	35.0	32.0-38.3	3.84	46.5	40.8-52.9
With CPB	4.67	106.9	97.7-117	5.22	185	169-203
Malignant tumour						
Without chemotherapy	0.93	2.5	2.5-2.6	0.32	1.4	1.3-1.5
With chemotherapy	0.66	1.9	1.8-2	0.81	2.2	2.0-2.5
GI bleeding	2.58	13.2	12.8-13.7			
Hip fracture surgery	2.40	11.0	10.4-11.6			
Chronic renal failure	1.59	4.90	4.7-5.1			
Liver cirrhosis				1.25	3.5	3.0-4.0
Constant	-4.10			-5.78		

Table 4 Logistic regression models for predicting transfusion of red blood cell and platelet

GI, gastrointestinal; DIC, disseminated intravascular coagulation; CPB, cardiopulmonary bypass; MDS, Myelodysplastic syndrome; RBC, red blood cell 95% CI, 95% confidence intervals for odds ratios.

system such as DRG can provide a means for inter-institutional benchmarking and cost comparison [14-16,37]. DRG is widely used for healthcare payments and analyses of hospital activities. The basic concept underlying the employment of DRG for reimbursement is that 'treatments for similar patients consume a similar degree of medical resources'. Therefore, patients within the same DRG are considered to have similar resource (e.g., blood product) utilization.

By the use of a patient classification system and identification of risk factors of transfusion, we can properly compare blood product use between different healthcare providers. Furthermore, such a method should be useful for identifying providers with extremely high levels of blood product use. If a simple risk adjustment model could be developed by analyzing large healthcare databases representing information obtainable with little labour and at low

COLOUR




Fig. 1 The multiple linear regression model to predict total unit use of RBCs and platelets transfused at each hospital

Table 5 Results of the medical chart review (percentage of appropriateness) and O/E ratios calculated using the two prediction models

Patient	Hospital	Red blood cell			Platelet		
		% of appropriateness	O/E (Model 1)	O/E (Model 2)	% of appropriateness	O/E (Model 1)	O/E (Model 2)
Overall	A	72	1.09 (1.02–1.18)	1.11 (1.05–1.16)	35	1.54 (1.34–1.81)	1.60 (1.41–1.83)
	B	81	0.93 (0.89–0.93)	0.78 (0.75–0.81)	64	1.13 (1.03–1.25)	0.78 (0.75–0.89)
Haematopoietic malignancy	A	71	1.35 (1.11–1.72)	1.61 (1.40–1.90)	25	1.50 (1.19–2.00)	2.30 (1.91–2.89)
	B	91	1.09 (0.97–1.25)	0.90 (0.82–0.99)	78	1.17 (1.02–1.36)	0.78 (0.75–0.89)
Cardiovascular surgery	A	62	1.25 (1.08–1.48)	1.73 (1.52–2.01)	17	1.76 (1.35–2.53)	1.23 (1.04–1.52)
	B	71	0.96 (0.88–1.05)	0.86 (0.79–0.93)	83	0.99 (0.85–1.18)	0.95 (0.86–1.06)

cost, it would be possible to evaluate the frequency of blood product use by various healthcare providers. It should be noted, however, that such a method is not designed to replace the detailed analyses conducted by auditing committees. Instead of pinpoint evaluations on blood product utilization at the patient level, the purpose of our method is to provide continuous surveillance and routine general evaluations on a larger scale that allows for multi-institutional comparisons with a satisfactory level of accuracy.

We selected two indicators to evaluate hospital-wide use of blood products in Japan: (i) proportion of patients that received a blood transfusion and (ii) total amount (number of units) of blood products used. These are typical indicators used for assessment of hospital-level blood usage [9,16,25,26]. We developed two regression models to calculate risk-adjusted blood use. A logistic regression model was used to predict the percentage of transfused patients in each hospital. A multiple linear regression model was used to predict hospital-wide total units of RBCs or platelets transfused.

With a limited number of variables, the logistic regression model could effectively predict proportions of patients that received a blood transfusion. Our predictive model consisted of 19 variables that were easily collected from healthcare data in Japan. In the database of approximately 587 000 cases used for the study, the amount of blood products used for patients with any of these risk factors represented 80–90% of the total amount of blood products used. The multiple linear regression model used distribution of diagnostic groups to predict total unit use of RBCs and platelets transfused at each hospital. Although both models showed good prediction abilities, the logistic regression model better predicted RBC use than platelet use. Also, the multiple linear regression model better predicted RBC use than platelet use (Fig. 1).

The following issues should be noted when evaluating blood product use with O/E ratios. First, the mean value of the group is used as the reference value (expected value) when performing evaluations with O/E ratios, thus the

evaluation is relative. For instance, if blood product use in the entire group is excessive, there is a possibility that the O/E ratio will be <1 in hospitals with high blood product use. Second, when O/E ratios are low, it is difficult to distinguish whether this is a result of appropriate blood product use, or under use. However, with blood transfusions, over use and misuse of blood product have been more of a problem than under use. Thus, when use is higher than average, there is a high likelihood that there has been inappropriate blood product use. On the contrary, when use is lower than average, there is a possibility that the healthcare provider is transfusing the bare minimum required.

Actually, O/E ratios calculated by use of the two models were very relevant to proportions of appropriate blood use (Table 5). Larger O/E ratios were associated with a smaller proportion of appropriate transfusions as judged by medical chart reviews. No particular difference was seen between the O/E ratio for proportion of patients receiving transfusions and that for the total amount (number of units) of RBC use. Both methods were considered to have successfully evaluated appropriate blood use.

Based on these findings, we conclude that the assessment of blood product use employing O/E ratios can be used, not only as an index for valid and appropriate transfusions but also as an index for blood product use that takes patient risk into consideration. Additionally, our research strongly indicates that valid comparisons may be made across hospitals in Japan.

Evaluation of blood product use at the hospital level is important in several ways. Wide variation in blood product use exists among hospitals. By comparing blood product use in different hospitals, risk-adjusted assessment of blood product use has the potential to contribute towards appropriate use of blood products [7,10,25]. Because of the labour and cost involved in gathering data on blood product use, however, almost no previous attempts had been made to collect and analyse data from many hospitals in Japan. DPC data are advantageous, as it gathers into a unified format the clinical information and data on

treatment procedures for all hospitalized patients. Therefore, it is possible to use a shared assessment standard to compare conditions for blood product use between hospitals, and to engage in discussion about clinical standards.

Acknowledgments

The authors are grateful to Dr Yasunori Ueda (Kurashiki Central Hospital, Kurashiki) for his comments on our article. We also thank to Dr Osamu Yonekawa (Seirei Hamamatsu General Hospital) for his supports to this study.

4.5 References

- 1 American Red Cross: Practice guidelines for blood transfusion: a compilation from recent peer-reviewed literature. May 2002. (available at http://www.redcrossblood.org/portal/SC/pdf/hospitals/pgbt_print.pdf)
- 2 The Task Force on Blood Component Therapy: Practice guidelines for blood component therapy. *Anesthesiology* 1996; 85:1219–1220
- 3 Baele PL, Muylle L, Noens L, Gulliksson H, Brands A, Isbister J, Van der Linden P, Hübner R, Berneman Z, Lamy M, Ferrant A, Lambermont M, Sondag D: Guidelines for the transfusion of red cells. *Acta Clin Belg* 2008; 63:301–312
- 4 O'Shaughnessy DF, Atterbury C, Bolton Maggs P, Murphy M, Thomas D, Yates S, Williamson LM, British Committee for Standards in Haematology, Blood Transfusion Task Force: Guidelines for the use of fresh-frozen plasma, cryoprecipitate and cryosupernatant. *Br J Haematol* 2004; 126:11–28
- 5 Blood Transfusion Task Force: Guidelines for the use of platelet transfusions. British Committee for Standards in Haematology. *Br J Haematol* 2003; 122:10–23
- 6 Simon TL, Alverson DC, AuBuchon J, Cooper ES, DeChristopher PJ, Glenn GC, Gould SA, Harrison CR, Milam JD, Moise KJ Jr, Rodwig FR Jr, Sherman LA, Shulman IA, Stehling L: Practice parameter for the use of red blood cell transfusions: developed by the Red Blood Cell Administration Practice Guideline Development Task Force of the College of American Pathologists. *Arch Pathol Lab Med* 1998; 122:130–138
- 7 Gombotz H, Rehak PH, Shander A, Hofmann A: Blood use in elective surgery: the Austrian benchmark study. *Transfusion* 2007; 47:1468–1480
- 8 Ozier Y, Pessione F, Samain E, Courtois F, French Study Group on Blood Transfusion in Liver Transplantation: Institutional variability in transfusion practice for liver transplantation. *Anesth Analg* 2003; 97:671–679
- 9 Mathoulin-Pélissier S, Salmi LR, Verret C, Demoures B: Blood transfusion in a random sample of hospitals in France. *Transfusion* 2000; 40:1140–1146
- 10 Surgenor DM, Wallace EL, Churchill WH, Hao S, Hale WB, Schnitzer J: Utility of DRG and ICD-9-CM classification codes for the study of transfusion issues. Transfusions in patients with digestive diseases. *Transfusion* 1989; 29:761–767
- 11 Cook SS, Epps J: Transfusion practice in central Virginia. *Transfusion* 1991; 31:355–360
- 12 Tinmouth A, Macdougall L, Fergusson D, Amin M, Graham ID, Hebert PC, Wilson K: Reducing the amount of blood transfused: a systematic review of behavioral interventions to change physicians' transfusion practices. *Arch Intern Med* 2005; 165:845–852
- 13 Slonim AD, Joseph JG, Turenne WM, Sharangpani A, Luban NL: Blood transfusions in children: a multi-institutional analysis of practices and complications. *Transfusion* 2008; 48:73–80
- 14 Syrjälä MT, Kytöniemi I, Mikkolainen K, Ranimo J, Lauharanta J: Transfusion practice in Helsinki University Central Hospital: an analysis of diagnosis-related groups (DRG). *Transfus Med* 2001; 11:423–431
- 15 Jefferies LC, Sachais BS, Young DS: Blood transfusion costs by diagnosis-related groups in 60 university hospitals in 1995. *Transfusion* 2001; 41:522–529
- 16 Lim YA, Lee WG, Cho SR, Hyun BH: A study of blood usage by diagnoses in Korean university hospital. *Vox Sang* 2004; 86:54–61
- 17 Japanese Ministry of Health, Labor, and Welfare. Practice guideline for the use of blood components. 2005. (available at http://www.jrc.or.jp/vcms_lf/iyaku_hin_benefit_guideline_sisin_090805.pdf)
- 18 American College of Surgeons: Advanced Trauma Life Support Course Manual. American College of Surgeons 1997; 103–112.
- 19 Lundsgaard-Hansen P: Component therapy of surgical hemorrhage: red cell concentrates, colloids and crystalloids. *Bibl Haematol* 1980; 46:147–69
- 20 British Committee for Standards in Haematology, Blood Transfusion Task Force: Guidelines for the use of platelet transfusions. *Br J Haematol* 2003; 122:10–23
- 21 Schiffer CA, Anderson KC, Bennett CL, Bernstein S, Elting LS, Goldsmith M, Goldstein M, Hume H, McCullough JJ, McIntyre RE, Powell BL, Rainey JM, Rowley SD, Rebullia P, Troner MB, Wagnon AH: Platelet transfusion for patients with cancer: clinical practice guidelines of the American Society of Clinical Oncology. American Society of Clinical Oncology. *J Clin Oncol* 2001; 19:1519–1538
- 22 Corwin HL, Gettinger A, Pearl RG, Fink MP, Levy MM, Abraham E, MacIntyre NR, Shabot MM, Duh MS, Shapiro MJ: The CRIT Study: Anemia and blood transfusion in the critically ill – current clinical practice in the United States. *Crit Care Med* 2004; 32:39–52
- 23 Zweig MH, Campbell G: Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993; 39:561–577
- 24 Pearce J, Ferrier S: Evaluating the predictive performance of habitat models developed using logistic regression. *Ecol Modell* 2000; 133:225–245
- 25 Sanguis Study Group: Use of blood products for elective surgery in 43 European hospitals. The Sanguis Study Group. *Transfus Med* 1994; 4:251–268
- 26 Snyder-Ramos SA, Mohnle P, Weng Y, Bottiger BW, Kulier A, Levin J, Mangano DT: The ongoing variability in blood transfusion practices in cardiac surgery. *Transfusion* 2008; 48:1284–1299
- 27 Stover EP, Siegel LC, Parks R, Levin J, Body SC, Maddi R, D'Ambra MN, Mangano DT, Spiess BD: Variability in

- transfusion practice for coronary artery bypass surgery persists despite national consensus guidelines: a 24-institution study. Institutions of the Multicenter Study of Perioperative Ischemia Research Group. *Anesthesiology* 1998; 88:327-333
- 28 Morrison JC, Sumrall DD, Chevalier SP *et al.*: The effect of provider education on blood utilization practices. *Am J Obstet Gynecol* 1993; 169:1240-1245
- 29 Muller U, Exadaktylos A, Roeder C *et al.*: Effect of a flow chart on use of blood transfusions in primary hip and knee replacement: prospective and after study. *BMJ* 2004; 328:934-938
- 30 Pentti J, Syrjala M, Pettila V: Computerized quality assurance of decisions to transfuse blood components to critically ill patients. *Acta Anaesthesiol Scand* 2003; 47:973-978
- 31 Rehm JP, Otto PS, West WW *et al.*: Hospital-wide educational program decreases red blood cell transfusions. *J Surg Res* 1998; 75:183-186
- 32 Rosen NR, Bates LH, Herod G: Transfusion therapy: improved patient care and resource utilization. *Transfusion* 1993; 33:341-347
- 33 Shanberge JN: Reduction of fresh-frozen plasma use through a daily survey and education program. *Transfusion* 1987; 27:226-227
- 34 Solomon RR, Clifford JS, Gutman SI: The use of laboratory intervention to stem the flow of fresh-frozen plasma. *Am J Clin Pathol* 1988; 89:518-521
- 35 Soumerai SB, Salem-Schatz S, Avorn J *et al.*: A controlled trial of educational outreach to improve blood transfusion practice. *J Am Med Assoc* 1999; 270:961-966
- 36 Tobin SN, Campbell DA, Boyce NW: Durability of response to a targeted intervention to modify clinician transfusion practices in a major teaching hospital. *Med J Aust* 2001; 174:445-448
- 37 Rashiq S, Shah M, Chow AK, O'Connor PJ, Finegan BA: Predicting allogenic blood transfusion use in total joint arthroplasty. *Anesth Analg* 2004; 99:1239-1244

DPCに基づく包括支払い制度導入後の 乳癌治療への影響と変化

京都大学大学院医学研究科医療経済学分野¹⁾
大隈和英¹⁾、今中雄一¹⁾、福田治久¹⁾、関本美穂¹⁾、
猪飼 宏¹⁾、濱田啓義¹⁾

The 109th Annual Congress of Japan Surgical Society
第109回日本外科学会定期学術集会

研究の主旨

- 我が国の乳癌は、死亡・罹患率ともに増加傾向にあり、年間死亡数9,885人(2003年)、罹患数は36,139人(1999年)であり、大腸がんを抜き女性の悪性腫瘍の第1位を占める。日本人女性の平均寿命の延長もあり、乳癌治療の重要性は益々高まっている。
- 乳癌治療は、手術治療を主体に、薬物治療、放射線治療、緩和医療に至るまで多彩であり、集学的治療の代表的なものである。その多様性の一方、「診療のばらつき」の存在を検証することは、EBMの見地から重要と思われる。
- DPC制度下の乳癌患者の治療の変化について診療情報を様々な角度から比較評価して、診療のトレンドとそれに伴う医療の質について検証する。


研究の方法

QIP

Quality Indicator/Improvement Project

- プロジェクト: QIP (Quality Indicator/Improvement Project) は、DPCデータを利用して診療のプロセス・アウトカムや経済性を反映する客観的な数値指標を開発・測定し、その情報を定期的に参加病院にフィードバックして、医療の質の向上に寄与する目的としている。前身は1995年から開始され、学内の倫理委員会の承認下、国内120以上の有力病院の参加を以って、厚労省の科研費により運営されている。
- 対象規模: QIP参加44病院のDPCデータ 10,659件
- 対象疾患: MDC09「乳房の疾患」のDPCコード090010
ICD-10「乳癌(C50\$)」または「乳房の上皮内癌(D50\$)」
- 除外疾患: 以下は解析から除外した。
 1. (乳房の)「皮膚のその他の悪性新生物、体幹の皮膚(C445)」
 2. 「Paget 病()」
- データ期間: 2004年4月～2008年2月

サンプルの内容



1) 疫学的解析

～DPCデータの妥当性評価として『全国乳がん患者登録調査報告』（第36号 2005年次症例）との比較～


1) 疫学

1. 性差

	全症例	乳癌登録
男性	27	74
(%)	0.33	0.5
女性	8,223	15,816
(%)	99.7	99.5
合計	8,250	15,890

2. 年齢

	手術症例	全症例	癌登録	%手術症例	%全症例	%癌登録
≤19	0	0	4	0	0	0.0
20-29	11	61	83	0.4	0.7	0.5
30-39	162	520	987	6.3	6.3	6.3
40-49	625	1819	3535	24.3	22.0	22.4
50-59	674	2353	4436	26.2	28.5	28.1
60-69	563	1969	3652	21.9	23.9	23.1
70-79	363	1116	2186	14.1	13.5	13.8
80-89	156	366	849	6.1	4.4	5.3
90≥	17	46	78	0.7	0.6	0.3
合計	2571	8250	15816	100	100	100
			不明 6			不明 0.0



	手術症例	全症例	癌登録
度数	2,571	8,250	15,816
最小値	24.2	20.2	
最大値	96.4	102.3	
平均値	58.2	58.0	57.7
中央値	57.5	57.8	57.0
最頻値	51.4	50.1	57.0
標準偏差	13.3	12.8	13.0

我々のDPCデータの性差・年齢階層は『乳癌学会の全国乳がん患者登録調査報告 2005年版』と非常に近似する。



1) 疫学

T因子

	手術症例	全TNM	乳癌登録	%手術症例	%全TNM	%癌登録
T0	2	13	227	0.1	0.3	1.4
T1	1212	1700	6185	47.1	38.5	39.1
T2	845	1625	5730	32.9	36.8	36.3
T3	107	267	679	4.2	6.0	4.3
T4	147	514	812	5.7	11.7	5.1
TIS	258	284	1345	10.0	6.6	8.5
合計	2571	4403	15816	100	100	100
			不明 838			不明 5.3

N因子

	手術症例	全TNM	乳癌登録	%手術症例	%全TNM	%癌登録
N0	1890	2563	12378	73.5	58.2	78.3
N1	581	1407	2707	22.6	32.0	17.1
N2	83	292	479	3.2	6.6	3
N3	17	141	147	0.7	3.2	0.9
合計	2571	4403	15816	100	100	100
			不明 838			不明 0.7





1) 疫学

M因子

	手術症例	全TNM	乳癌登録	%手術症例	%全TNM	%癌登録
M0	2516	3952	15228	97.9	87.8	96.3
M1	55	451	357	2.1	12.2	2.3
合計	2571	4403	15816	100	100	100
			不明231			不明 1.5

病期分類

	手術症例	全TNM	乳癌登録	%手術症例	%全TNM	%癌登録
0	258	284	1280	10.0	6.5	8.1
I	1049	1332	5568	40.8	30.3	35.2
IIA	657	1083	4384	25.6	24.6	27.7
IIB	342	699	1751	13.3	15.9	11.1
IIIA	86	239	495	3.3	5.4	3.1
IIIB	112	223	602	4.4	5.1	3.8
IIIC	12	92	91	0.5	2.1	0.6
IV	55	451	357	2.1	10.2	2.3
合計	2571	4403	15816	100	100	100
			不明1288			不明 8.1



2) 治療



1. 手術術式

2. 術後補助療法

- 化学療法の実施状況
- 放射線療法の実施状況
- リハビリテーションの実施状況

2) 治療 手術術式

我々のDPCデータにおける手術術式は『乳癌学会の全国乳がん患者登録調査報告2005年版』と非常に近似する。




	乳房温存療法	全乳房切除	胸筋温存乳房切除	胸筋合併乳房切除術	両側手術	その他	なし	不明
□ QIP施行率	55.6	7.9	32.2	0.9	1.1	2.5		
■ 乳がん登録での施行率	53.2	5.9	39.3	0.4		1	0.1	0

2) 治療 術後補助療法


化学療法の実施状況

	手術症例	全TNM	全症例	%手術症例	%全TNM	%全症例
なし	2201	2517	4492	85.6	57.1	54.4
経口	274	391	676	10.7	8.9	8.2
経静脈 or 動注	90	1472	3012	3.5	33.4	36.5
その他	6	23	70	0.23	0.52	0.85
合計	2571	4403	8250	100	100	100




放射線療法の実施状況

	%手術症例	%全TNM	%全症例
なし	42.0	43.2	34.7
あり	1.1	2.9	2.1
記載なし	56.9	53.9	63.2
合計	100	100	100



リハビリテーションの実施状況

	%手術症例	%全TNM	%全症例
なし	17.4	30.8	26.2
あり	25.6	15.3	10.6
記載なし	56.9	53.9	36.8
合計	100	100	100




3) クロス集計結果

1. 病期別解析
2. 年齢別解析
3. 年次別解析
4. 施設別解析


3) クロス集計結果

1. 病期別解析

術式<全TNM症例>



術式<手術症例>




3) クロス集計結果

1. 病期別

乳房温存術施行率(=温存率)

○両側、その他の術式を除外


病期	BCS	Mx	合計件数	温存率%
0	169	85	254	66.5
I	755	271	1026	73.6
IIA	319	313	632	50.5
IIB	138	188	326	42.3
IIIA	19	62	81	23.5
IIIB	20	82	102	19.6
IIIC	1	8	9	11.1
IV	8	42	50	16.0
合計	1429	1051	2480	57.6



3)クロス集計結果 2.年齢別


温存率

年齢階層	BCS	Mx	合計	温存率(%)
20-29	10	1	11	90.9
30-39	95	58	153	62.1
40-49	374	228	602	62.1
50-59	390	257	647	60.3
60-69	325	224	549	59.2
70-79	168	181	349	48.1
80-89	59	94	153	38.6
90+	8	8	16	50.0
	1429	1051	2480	57.6




リンパ節郭清率

年齢階層	郭清なし	郭清あり	合計	施行率(%)
20-29	1	10	11	90.9
30-39	28	125	153	81.7
40-49	133	469	602	77.9
50-59	126	521	647	80.5
60-69	131	418	549	76.1
70-79	96	253	349	72.9
80-89	61	92	153	60.1
90+	14	2	16	87.5
	590	1890	2480	76.2




3)クロス集計結果 2.年齢別

化学療法<手術症例>



化学療法<全症例>




手術症例での化学療法の施行率は、年齢との相関はないが、80歳以上では点滴での化学療法は施行されていない。


全症例では年齢が上昇に伴い、点滴化学療法の施行率は有意に減少した。80歳以上での化学療法は経口投与が主体となっている。

3) クロス集計結果 3.年次別

病期<手術症例>




病期<全症例>




3) クロス集計結果 3.年次別

併存症<手術症例>




併存症<全症例>



3) クロス集計結果 3.年次別


温存率




	BCS	Mx	件数	施行率 %
2004	148	134	282	52.5
2005	350	247	597	58.6
2006	457	340	797	57.3
2007	441	311	752	58.6
2008	33	19	52	63.5
合計	1429	1051	2480	57.6

3) クロス集計結果 3.年次別

在院日数<手術症例>




在院日数<全症例>




3)クロス集計結果 4.施設別

温存率<全病期>




温存率< I・II期>




温存率に影響を与えると思われる、専門医の勤務状況や地域性などにも、今後の検討を加えたい。

3)クロス集計結果 4.施設別

在院日数<手術症例>




在院日数<全TNM症例>



日帰り手術を施行している施設は7件あった。病院別在院日数において(手術や治療の合併症)有害事象または再入院については未解析である。

3) クロス集計結果 4. 施設別

医療費(中央値) <手術症例>



施設	包括	出来高	在院日数	件数
	46092	65359	34	9
	69561.33	47895	26.5	4
	40215.2	46299.5	20.3333	32
	38659.67	45227	17.3333	9
	43256	37166	17	3
	51518.75	46381	16.6667	8
	41558.67	41551	16.6	13
	40017	50300	14.7778	47
	31927	38679.5	14	4
	35075.5	40812	13	34
	31880.5	39408	12.8889	15
	34888.5	40264.5	12.25	18
	30233	38303	11.8	59
	23967	27456	11	1
	30309.38	42325	10.9638	403
	30624.33	41499	10.6667	51
	28186	39324	10	11
	29117.33	48128	9.66667	7
	25775.6	39233	9.13636	57
	26290.14	41627	9	36
	23248	36764.5	7.42424	188
	20486.63	38284	6.82222	99
	20854.5	36225	6.63158	86
	19234.53	38161	6.29412	105
x	20581.5	38273.5	7.2	90
合計	28240.6	40423	9.71324	1389

結果 まとめ

1. 我々のDPCデータは、学会の癌登録のデータと良く近似しており、限定的ながら解析データとしての妥当性はある。
2. 年齢別では、高齢化に伴い標準治療から低侵襲の治療が選択される傾向がある。70歳以上での乳房温存術と化学療法施行率の低下、および80歳以上でのリンパ節郭清の施行率の低下が観察された。
3. 年次別では、病期および併存症の分布には経時的変化を認め、手術症例での在院日数の短縮を認めた。また入院における補助療法の施行率が低下した。
4. 病期別では、在院日数は病期の進行に伴い延長した。
5. 入院手術症例での医療費中央値の平均は、包括部分282,406円、出来高部分404,230円であった。在院日数の延長、病期進行度と併存症の増加に伴い医療費は増加したが、経時的な変化は認めなかった。