

1. Intravenous human albumin
2. Intravenous furosemide
3. Intravenous hetastarch
4. Intravenous dobutamine
5. Intravenous dopamine
6. Do not know

W) Hypotension (80/40 mmHg) due to acute myocardial infarction

1. Intravenous human albumin
2. Intravenous hetastarch
3. Intravenous dobutamine
4. Intravenous dopamine
5. Intravenous furosemide
6. Do not know

X) Hypoxemia (SpO₂ 88%) with drowsiness due to CO₂ narcosis in a patient with acute aggravation of COPD

1. Low flow oxygen (through nasal cannula)
2. High flow oxygen (by reservoir oxygen mask)
3. Oral diazepam
4. Intravenous theophylline
5. Intravenous midazolam
6. Do not know

Y) Respiratory suppression due to morphine HCl

1. Intravenous flumazenil
2. Intravenous naloxone
3. Intravenous doxapram
4. Intravenous midazolam
5. Intravenous pentazocine
6. Do not know

Z) Acute aortic dissection (Stanford type B) with hypertension and tachycardia

1. Sublingual nifedipine
2. Intravenous nicardipine
3. Intravenous propranolol
4. Intravenous dobutamine
5. Intravenous dopamine
6. Do not know

(2) Assume that your working schedule is flexible so that you can accommodate emergency or nighttime outpatient care. For each of the situations below, to what extent will you be able to manage the patient? Please select only 1 item that you think is the closest to the real situation.

A) Primary emergency in daytime (undiagnosed, acute patient who visits without notice)

1. I can manage almost all patients.
2. I can manage more than half of the patients.
3. I can manage some of the patients.
4. I can manage few of the patients.

B) Secondary emergency in daytime (undiagnosed, acute patient who arrives by ambulance)

1. I can manage almost all patients.
2. I can manage more than half of the patients.
3. I can manage some of the patients.
4. I can manage few of the patients.

C) Primary emergency at nighttime (undiagnosed, acute patient who visits without notice)

1. I can manage almost all patients.
2. I can manage more than half of the patients.
3. I can manage some of the patients.
4. I can manage few of the patients.

D) Secondary emergency at nighttime (undiagnosed, acute patient who arrives by ambulance)

1. I can manage almost all patients.
2. I can manage more than half of the patients.
3. I can manage some of the patients.
4. I can manage few of the patients.

5. 研究班員名簿

研究代表者：徳田 安春（筑波大学）

研究分担者：

1. 福井 次矢（聖路加国際病院）
2. 後藤 英司（横浜市立大学）
3. 大滝 純司（東京医科大学）
4. 小俣 富美雄（聖路加国際病院）
5. 高橋 理（聖路加国際病院）
6. 大出 幸子（聖路加国際病院）

研究協力者

石田 也寸志（聖路加国際病院）

副島 久美子（聖路加国際病院）

シャピロ 美奈（聖路加国際病院）

ジョシュア・ジェイコブス（米国ハワイ大学）

