

CAS No.	ER score	AR score						
25122-46-7	-0.43	-0.5	508-96-3	0.92	-1.09	8056-51-7	0.18	0.82
25122-47-8	-1.09	-0.93	51092-19-4	0.16	0.57	807-38-5	1.12	0.93
25122-49-0	-0.67	-0.24	514-61-4	-0.11	1.02	84693-92-5	1.08	0.52
25122-56-9	-0.89	-0.88	53-39-4	0.5	1.25	855-19-6	0.01	-0.62
25122-57-0	-0.56	-0.89	53716-43-1	1.27	2.21 *	855-22-1	-0.14	-0.32
2529-45-5	0.25	0.01	54024-22-5	0.31	0.07	855-55-0	0.3	-0.27
2571-22-4	-0.97	-1.23	54048-10-1	0.26	0.63	88158-43-4	1.56	1.54 *
2868-02-2	0.67	1.27	54063-32-0	-0.29	-0.13	88158-44-5	0.83	2.37 *
2881-21-2	0.62	1.21	55056-58-1	-0.34	0.35	89231-65-2	-0.71	-0.75
3000-39-3	-0.79	-5.45	5534-09-8	-2.24	-0.42	899-39-8	0.35	0.63
3091-89-2	0.66	0.11	5593-20-4	-0.68	-0.96	92817-12-4	0.77	-0.02
31002-79-6	1.14	-0.06 *	5611-51-8	-0.16	1.07 *	92901-21-8	0.91	1.06 *
312-93-6	-0.27	0.21	56933-60-9	0.3	1.15	92901-23-0	1.56	1.54 *
33564-31-7	-0.84	-0.46	57781-14-3	-1.06	-0.8	92901-24-1	0.83	2.37 *
33755-46-3	-0.73	1.51	58096-47-2	-0.62	-0.95	92901-25-2	0.91	1.06 *
33784-05-3	0.35	0.97	58-19-5	0.88	1.14	92901-26-3	1.09	-0.15 *
3385-03-3	0.57	0.11	58-20-8	-0.41	-1.44	95440-71-4	-0.51	-0.65
338-95-4	0.77	0.95	58652-20-3	0.43	0.51	95796-50-2	-0.49	-0.36
338-98-7	0.4	0.05	5949-44-0	1.16	0.31	95796-51-3	-0.94	-0.96
3484-46-6	-1.49	-2.79	595-57-3	-0.95	-0.53	965-66-2	1.07	0.9
3597-38-4	0.6	0.39	601-63-8	0.9	-0.9	965-90-2	0.31	0.36
37270-71-6	-0.13	0.61	64272-13-5	-0.86	-0.16	35942-13-3	-1.08	-1.65
3801-06-7	0.37	-0.14	64272-18-0	-0.59	1.11	52183-73-0	-2.28	-3.83
38196-44-0	-0.46	-0.29 *	65-06-5	0.02	1.12	630-64-8	-6.94	
38196-45-1	-0.48	-0.48	6533-00-2	0.18	0.82	75384-50-0	-1.21	-1.42
3836-23-5	-0.95	-0.24	65466-50-4	-0.08	-0.36	103834-29-3	-1.16	-1.94
3863-59-0	-0.63	-0.28	65980-97-4	-0.18	0.74	11018-89-6	-1.87	
387-79-1	0.99	1.04	6677-98-1	-0.22	0.08	1108-68-5	1.57	-0.59
39791-38-3	0.06	0.68	6677-99-2	-0.16	0.44	14110-73-7	-0.09	-0.06
39845-47-1	1.25	0.22	6678-00-8	-0.02	-0.66	14930-96-2	0.86	-2.99
4050-05-9	0.25	0.83	67-73-2	0.24	-0.07	18542-37-5	-1.39	-1.65
4075-12-1	0.05	0.5	67-78-7	-0.6	0.82	18695-02-8	-1.45	
41020-66-0	0.33	0.52	68374-16-3	0.86	0.12	21256-71-3		-0.46 *
41020-69-3	-0.23	0.44	6987-59-3	0.66	0.72	3175-90-4	-0.2	-1.48
432-60-0	0.18	-0.32	72-63-9	0.34	0.76	3175-91-5	-0.64	-2.9
4351-48-8	-0.05	0.09	7359-79-7	0.1	-0.52	3175-92-6	-0.47	-2.38
4351-59-1	-0.16	-0.32	75883-07-7	0.29	0.85	34206-60-5	0.77	-2.25
472-56-0	0.84	-0.71	79-60-7	1.03	0.22	35413-85-5	-1.07	-2.92
472-57-1	0.84	-0.71	79-64-1	0.97	-0.02	38274-01-0	-0.74	-1.6
4968-09-6	0.7	-0.39	797-58-0	0.03	0.76	38274-02-1	-0.51	-0.93
50394-27-9	0.06	0.5	8015-29-0	-0.13	0.61	4029-68-9	-1.03	-0.56
50629-82-8	0.64	1.36	8015-30-3	-0.13	0.61	40788-89-4	-0.72	-2.25

\* : ARスコア計算に2pnuモデルを使用した化合物(その他は1t63モデルを使用)  
空欄は計算値が得られなかった化合物

CAS No.	ER score	AR score			
4099-30-3	2.05	0.1	1231-34-1	0.71	-1.22
41451-75-6		-0.65 *	1235-78-5	0.19	-1.69
470-42-8	0.98	-0.31	15520-11-3	-1.22	-2.56 *
472-26-4	0.73	-0.07	1900-69-2	-1.35	-1.8
472-95-7	-0.72	-2.25	2223-71-4	-0.87	-2.54
472-96-8	-0.72	-2.25	24238-95-7	-0.25	-1.26
516-37-0		-1.66	29065-33-6	-0.63	-1.63
5352-63-6	-0.27		29080-98-6	-0.63	-1.63
564-92-1	-0.72	-2.25	3007-19-0	-0.64	-1.83
58581-03-6	-1.32	-2.19	31207-65-5	-0.94	-1.09
68845-01-2	-0.11	-1.66	3738-00-9	0.25	-1.34
72777-88-9	-0.17	-1.39	3787-85-7	-0.52	-1.81
87818-61-9	-0.36	-0.89	3950-05-8	-0.38	-0.81
87819-60-1	-0.36	-0.89	41199-19-3	-0.39	-1.7
7299-40-3	-1.36	-2.88	46729-07-1	-0.77	-2.46
13720-13-3	-0.78	-1.96	468-84-8	0.3	-0.39
13720-37-1	-0.79	-1.76	473-03-0	0.64	-1.14
473-16-5	0.02	-1.03	5441-56-5	-1.13	-1.4
20748-87-2	-1.6		57345-19-4	0.42	-1.13
7425-14-1	-1.73		62406-83-1	-0.52	-1.92
68310-54-3	-1.24		63449-90-1	-0.34	-2.06
16397-75-4	-2.22		67874-73-1	-1.56	-2.02
68956-51-4	-2.15		67874-74-2	-0.87	-2.17
7434-89-1	-1.85		68039-42-9	-1.42	-1.94
15840-96-7	-1.57	-2.11	68797-74-0	-1.09	-2.19
25085-98-7	-0.14	-1.37			
510-78-1	0.82	-0.33			
62256-00-2	-1	-1.42			
69178-35-4	-3.8	-0.3 *			
69178-39-8	-2.68	-0.66			
74113-77-2	0.4				
1449-09-8	1.2	-0.35			
21238-33-5	0.78				
469-38-5	2.13	-0.92			
469-39-6	1.87	0.03			
511-61-5	0.95	-1.91			
10118-56-6	0.12	-2.46			
10411-92-4	-1.35	-1.8			
1153-34-0	0.3	-1.15			
1153-35-1	0.29	-0.66			
1227-93-6	0.34	-1.15			
1231-33-0	0.14	-0.92			

\* : ARスコア計算に2pnuモデルを使用した化合物(その他は1t63モデルを使用)  
空欄は計算値が得られなかった化合物

## Ⅲ. 研究成果の刊行に関する一覧表

## 雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Mariko Shirota, Takayuki Seki, Kazumi Tago, Hiroyasu Katoh, Hideki Marumo, Mami Furuya, Tomoko Shinoda and Hiroshi Ono.	Screening of toxicological properties of 4-methylbenzoic acid by oral administration to rats.	J Toxicol Sci.	33	431-445	2008
Suzuki A, Igarashi K, Aisaki KI, Kanno J, Saga Y.,	NANOS2 interacts with the CCR4-NOT deadenylation complex and leads to suppression of specific RNAs.	Proc Natl Acad Sci U S A.	107	3594-3599	2010
Matsunaga N, Kanno J, Hamada C, Yoshimura I.	An experimental design for judging synergism on consideration to endocrine disruptor animal experiments.	Environmetrics	20	1-13	2009
Woo GH, Takahashi M, Inoue K, Fujimoto H, Igarashi K, Kanno J, Hirose M, Nishikawa A, Shibutani M.,	Cellular distributions of molecules with altered expression specific to thyroid proliferative lesions developing in a rat thyroid carcinogenesis model.	Cancer Sci.	100	617-625	2009
Myers JP, vom Saal FS, Akingbemi BT, Arizono K, Belcher S, Colborn T, Chahoud I, Crain DA, Farabollini F, Guillette LJ Jr, Hassold T, Ho SM, Hunt PA, Iguchi T, Jobling S, Kanno J, Laufer H, Marcus M, McLachlan JA, Nadal A, Oehlmann J, Olea N, Palanza P, Parmigiani S, Rubin BS, Schoenfelder G, Sonnenschein C, Soto AM, Talsness CE, Taylor JA, Vandenberg LN, Vandenbergh JG, Vogel S, Watson CS, Welshons WV, Zoeller RT.	Why public health agencies cannot depend on good laboratory practices as a criterion for selecting data: the case of bisphenol A.	Environ Health Perspect.	117	309-315	2009
Ishimaru N, Takagi A, Kohashi M, Yamada A, Arakaki R, Kanno J, Hayashi Y.	Neonatal exposure to low-dose 2,3,7,8-tetrachlorodibenzo-p-dioxin causes autoimmunity due to the disruption of T cell tolerance.	J Immunol.	182	6576-6586	2009
Upham BL, Park JS, Babica P,	Structure-activity-dependent regulation of cell communication by perfluorinated fatty	Environ Health Perspect.	117	545-551	2009

Sovadinova I, Rummel AM, Trosko JE, Hirose A, Hasegawa R, <u>Kanno J</u> , Sai K.	acids using in vivo and in vitro model systems.				
Kanno J	Overview: "Children's toxicology", a renovating study field of irreversible "early exposure-delayed effects".	J Toxicol Sci.	34	SP199-SP200	2009
Sekine H, Mimura J, Oshima M, Okawa H, <u>Kanno J</u> , Igarashi K, Gonzalez FJ, Ikuta T, Kawajiri K, Fujii-Kuriyama Y.	Hypersensitivity of aryl hydrocarbon receptor-deficient mice to lipopolysaccharide-induced septic shock.	Mol Cell Biol.	29	6391-6400	2009
Saegusa Y, Woo GH, Fujimoto H, Inoue K, Takahashi M, Hirose M, Igarashi K, <u>Kanno J</u> , Mitsumori K, Nishikawa A, Shibutani M.	Gene Expression Profiling and Cellular Distribution of Molecules with Altered Expression in the Hippocampal CA1 Region after Developmental Exposure to Anti-Thyroid Agents in Rats.	J Vet Med Sci.	72	187-195	2010
Vom Saal FS, Akingbemi BT, Belcher SM, Birnbaum LS, Crain DA, Eriksen M, Farabollini F, Guillette LJ Jr, Hauser R, Heindel JJ, Ho SM, Hunt PA, Iguchi T, Jobling S, <u>Kanno J</u> , Keri RA, Knudsen KE, Laufer H, LeBlanc GA, Marcus M, McLachlan JA, Myers JP, Nadal A, Newbold RR, Olea N, Prins GS, Richter CA, Rubin BS, Sonnenschein C, Soto AM, Talsness CE, Vandenberg JG, Vandenberg LN, Walser-Kuntz DR, Watson CS, Welshons WV, Wetherill Y, Zoeller RT.	Chapel Hill bisphenol A expert panel consensus statement: integration of mechanisms, effects in animals and potential to impact human health at current levels of exposure.	Reprod Toxicol.	24	131-138	2007
Kato Y, Ikushiro S, Takiguchi R, Haraguchi K, Koga N, Uchida S, Sakaki T, Yamada S, <u>Kanno J</u> , Degawa M.	A novel mechanism for polychlorinated biphenyl-induced decrease in serum thyroxine level in rats.	Drug Metab Disposition	35	1949-1955	2007
Wetherill YB, Akingbemi BT, <u>Kanno J</u> , McLachlan JA, Nadal A,	In vitro molecular mechanisms of bisphenol A action.	Reprod Toxicol	24	178-198	2007

Sonnenschein C, Watson CS, Zoeller RT, Belcher SM.					
K. Miyagawa, M. Narita, M. Narita, H. Akama, <u>T. Suzuki</u>	Memory impairment associated with a dysfunction of the hippocampal cholinergic system induced by prenatal and neonatal exposures to bisphenol-A.	Neurosci Lett.	418	236- 241	2007
M. Narita, K. Miyagawa, K. Mizuo, T. Yoshida, <u>T. Suzuki</u>	Changes in central dopaminergic systems and morphine reward by prenatal and neonatal exposure to bisphenol-A in mice: evidence for the importance of exposure period.	Addict Biol.	12	167- 172	2007
Kazuya Miyagawa, Minoru Narita, Michiko Narita, Keiichi Niikura, Hisahiko Akama, Yuri Tsurukawa, and <u>Tsutomu Suzuki</u>	Changes in central dopaminergic systems with the expression of Shh or GDNF in mice perinatally exposed to bisphenol A.	Jpn. J. Neuropsychopharmacol	27	69-75	2007
Kenichi Kobayashi, Katsumi Ohtani, Hisayo Kubota, <u>Muneyuki Miyagawa</u>	Dietary exposure to low doses of bisphenol A: effects on reproduction and development in two generations of C57BL/6 J mice.	Congenital Anomalies			2010 In press
石丸直澄、林 良夫	ダイオキシンによる免疫異常	臨床免疫・アレルギー科	51	60-65	2009
Kinouchi N, Ohsawa Y, Ishimaru N, Ohuchi H, Sunada Y, <u>Hayashi Y</u> , Tanimoto Y, Moriyama K, Noji S.	Atelocollagen-mediated local and systemic application of myostatin-targeting siRNA increase skeletal muscle mass.	Gene Therapy		1-5	2008
Kohashi M, Ishimaru N, Arakaki R, <u>Hayashi Y.</u>	Effective treatment with oral administration of rebamipide in a mouse model of Sjögren's syndrome.	Arthritis Rheumatism	58	389- 400	2008
Hikosaka Y, Nitta T, Ohigashi I, Yano K, Ishimaru N, <u>Hayashi Y</u> , Matsumoto M, Matsuo K, Penninger JM, Takayanagi H, Yokota Y, Yamada H, Yoshikai Y, Inoue J, Akiyama T, Takahama Y.	The cytokine RANKL produced by positively selected thymocytes fosters medullary thymic epithelial cells that express autoimmune regulator.	Immunity	29	438- 450	2008
Ishimaru N, Arakaki R, Yoshida S, Yamada A, Noji S, <u>Hayashi Y.</u>	Expression of the retinoblastoma protein RbAp48 in exocrine glands leads to Sjögren's syndrome-like autoimmune exocrinopathy.	J Exp Med.	205	2915- 2927	2008
Ishimaru N, Yamada A, Kohashi M, Arakaki R, Takahashi T, Izumi K, <u>Hayashi Y.</u>	Development of inflammatory bowel disease in Long-Evans Cinnamon rats based on CD4+CD25+Foxp3+ regulatory T cell dysfunction.	J Immunology	180	6997- 7008	2008
Izawa T, Ishimaru N, Moriyama K, Kohashi M, Arakaki R, <u>Hayashi Y.</u>	Crosstalk between RANKL and Fas signaling in dendritic cells controls immune tolerance.	Blood	110	242- 250	2007
石丸直澄、岸本英	T細胞レセプターシグナルとNF-kB	臨床免疫・アレルギー科	48	29-35	2007

博、林 良夫		ルギー科			
Ohtani N, Imamura Y, Yamakoshi K, Hirota F, Nakayama R, Kubo Y, Ishimaru N, Takahashi A, Hirao A, Shimizu T, Mann DJ, Saya H, <u>Hayashi Y</u> , Arase S, Matsumoto M, Kazuki N, Hara E.	Visualizing the dynamics of p21 <sup>(Waf1/Cip1)</sup> cyclin-dependent kinase inhibitor expression in living animals.	PNAS	104	15034-15039	2007
Omori T, Idehara K, Kojima H, Sozu T, Arima K, Goto H, Hanada T, Ikarashi Y, Inoda T, Kanazawa Y, Kosaka T, Maki E, Morimoto T, Shinoda S, Shinoda N, <u>Takeyoshi M</u> , Tanaka M, Uratani M, Usami M, Yamanaka A, Yoneda T, Yoshimura I, Yuasa A.	Interlaboratory validation of the modified murine local lymph node assay based on adenosine triphosphate measurement.	J Pharmacol Toxicol Methods	58	11-26	2008
<u>Takeyoshi M</u> , Iida K, Suzuki K, Yamazaki S.	Skin sensitization potency of isoeugenol and its dimers evaluated by a non-radioisotopic modification of the local lymph node assay and guinea pig maximization test.	J Appl Toxicol.	28	530-534	2008
Basketter D, Cockshott A, Corsini E, Gerberick GF, Idehara K, Kimber I, Van Loveren H, Matheson J, Mehling A, Omori T, Rovida C, Sozu T, <u>Takeyoshi M</u> , Casati S.	An evaluation of performance standards and non-radioactive endpoints for the local lymph node assay. The report and recommendations of ECVAM Workshop 65.	ATLA	36	243-257	2008
Tarama R, Kato H, Ishikawa Y, Miyaura H, <u>Takeyoshi M</u> , Iwata H.	Gene expression changes induced by type IV allergy-inducible chemicals in dendritic cells.	J Vet Med Sci.	70	673-680	2008
<u>Nagao T.</u> , Yoshimura S.	Early embryonic losses in mice induced by diethylstilbestrol.	<i>Cong. Anom.</i>	49	269-273	2009
Ryo OHTA, Mariko SHIROTA, Yukiko KANAZAWA, Tomoko SHINDO, Mami FURUYA, Takayuki SEKI, Hiroshi ONO, Kohichi, KOJIMA, Sayaka ASAI, Gen WATANABE, Kazuyoshi TAYA	Effects of Transmaternal Exposure to Gnistein in Hatano High- and Low- Avoidance Rats,	Exp Anim.	58	471-479	2009
太田 亮、宮原 敬、	内分泌攪乱性確定試験としてのラットー	秦野研究所年	30	17-24	2007

又吉 健、大向英夫、小野 宏	生涯試験の試み	報			
菅野純、相崎健一、五十嵐勝秀、北嶋聡、中津則之、児玉幸夫、高木篤也	トキシコゲノミクスの新展開 Percellome プロジェクトによる 2,3,7,8-TCDD—2,3,7,8-TCDF 比較	細胞工学	26	1391-1396	2007
菅野純、北嶋聡、相崎健一、五十嵐勝秀、中津則之、高木篤也、小川幸男、児玉幸夫	Percellome Project による毒性トランスクリプトミクスの新しい試み	細胞工学	26	71-77	2007
Fujimoto N., Suzuki T., Ohta S., Kitamura S	Identification of rat prostatic secreted proteins using mass spectrometric analysis and androgen-dependent mRNA expression.	<i>J. Andrology</i>	30	669-678	2009
S Kitamura, S Shinohara, E Iwase, K Sugihara, N Uramaru, H Shigematsu, N Fujimoto, S Ohta	Affinity for thyroid hormone and estrogen receptors of hydroxylated polybrominated diphenyl ethers	J Health Sci.	54	607-614	2008
S Kitamura, K Sugihara, S Sanoh, N Fujimoto, S Ohta	Metabolic activation of proestrogens in the environment by cytochrome P450 system	J Health Sci.	54	343-355	2008
Suzuki, T., Fujimoto, N., Kitamura, S., Ohta, S.	Effects of environmental antiandrogenic chemicals on expression of androgen-responsive genes in rat prostate lobes.	J Health Sci.	53	401-405	2007
Suzuki, T., Fujimoto, N., Kitamura, S., Ohta, S.	Quantitative determination of lobe specificity of mRNA expression of androgen-dependent genes in the rat prostate gland.	Endocrine. J.	54	123-132	2007
Fujimoto, N., Nakajima, O., Kitamura, S., Ohta, S.	<i>In vivo</i> function of the 5' flanking region of mouse estrogen receptor beta gene.	J Steroid Biochem Mol Biol.	105	57-62	2007
Saegusa Y, Woo GH, Fujimoto H, Inoue K, Takahashi M, Hirose M, Igarashi K, Kanno J, Mitsumori K, Nishikawa A, Shibutani M.	Gene Expression Profiling and Cellular Distribution of Molecules with Altered Expression in the Hippocampal CA1 Region after Developmental Exposure to Anti-Thyroid Agents in Rats.	J Vet Med Sci.	72	187-195	2010
Tanemura K, Igarashi K, Matsugami TR, Aisaki K, Kitajima S, Kanno J.	Intrauterine environment-genome interaction and children's development (2): Brain structure impairment and behavioral disturbance induced in male mice offspring by a single intraperitoneal administration of domoic acid (DA) to their dams.	J Toxicol Sci.	34	SP279-SP286	2009
Iida K, Mimura J, Itoh K, Ohyama C, Fujii-Kuriyama Y, Shimazui T, Akazawa H, Yamamoto M.	Suppression of AhR signaling pathway is associated with the downregulation of UDP-glucuronosyltransferases during BBN-induced urinary bladder carcinogenesis in mice.	<i>J Biochem.</i>	47	353-360	2010

Skine H, Mimura J, Oshima M, Kanno J, Igarashi K, Gonzalez FJ, Ikuta T, Kawajiri K, <u>Fujii-Kuriyama Y.</u>	Hypersensitivity of aryl hydrocarbon receptor- deficient mice to lipopolysaccharide-induced septic shock.	<i>Mol Cell Bio.l</i>	29	6391-6400	2009
Kimura A, Naka T, Nakahama T, Chinen I, Masuda K, Nohara K, <u>Fujii-Kuriyama Y,</u> Kishimoto T.	Aryl hydrocarbon receptor in combination with Stat 1 regulates LPS-induced inflammatory responses.	<i>J Exp Med.</i>	206	2027-2035	2009
Kawajiri K, Kobayashi Y, Ohtaka F, Ikuta T, Matsushima Y, Mimura J, Pettersson S, Pollenz RS, Sakaki T, Hirokawa T, Akiyama T, Kurosumi M, Poellinger L, Kato S, <u>Fujii-Kuriyama Y.</u>	Aryl hydrocarbon receptor suppresses intestinal carcinogenesis in Apc <sup>min/+</sup> mice with natural ligands.	<i>Proc Natl Acad Sci USA</i>	106	13481-13486	2009
Funatake CJ, Ao K, Suzuki T, Murai H, Yamamoto M, <u>Fujii-Kuriyama Y,</u> Kerkvliet NI, Nohara K.	Expression of constitutively-active aryl hydrocarbon receptor in T-cell enhances the down-regulation of CD62L, but does not alter expression of CD25 or suppress the allogeneic CTL response.	<i>J Immuno-toxicology</i>	6	194-203	2009
Nohara K, Suzuki T, Ao K, Murai H, Miyamoto Y, Inoue K, Pan X, Motohashi H, <u>Fujii-Kuriyama Y,</u> Yamamoto M, Tohyama C.	Constitutively active aryl hydrocarbon receptor expressed in T cells increases immunization- induced IFN-gamma production in mice but does not suppress T <sub>H</sub> 2-cytokine production or antibody production.	<i>Int Immunol.</i>	21	769-777	2009
Nakata A, Urano D, <u>Fujii-Kuriyama Y,</u> Mizuno H, Tago K, Itoh.	G-protein signaling negatively regulates the stability of aryl hydrocarbon receptor.	<i>EMBO Rep</i>	10	622-628	2009
Oshima M, Mimura J, Sekine H, Okawa H, <u>Fujii-Kuriyama Y.</u>	SUMO modification regulates the transcriptional repressor function of aryl hydrocarbon receptor repressor.	<i>J Biol Chem.</i>	284	11017-11026	2009
Shimada T, Hiramatsu N, Hayakawa K, Takahashi S, Kasai A, Tagawa Y, Mukai M, Yao J, <u>Fujii-Kuriyama Y,</u> Kitamura M.	Dual suppression of adipogenesis by cigarette smoke through activation of the aryl hydrocarbon receptor and induction of endoplasmic reticulum stress.	<i>Am J Physiol Endocrinol Metab.</i>	296	E721-E730	2009
Togo Ikuta, Takeshi Namiki, <u>Yoshiaki Fujii-Kuriyama,</u> Kaname Kawajiri	AhR protein trafficking and function in the skin	Biochemical Pharmacology	77	588-596	2009
Fumiaki Ohtake, <u>Yoshiaki Fujii-Kuriyama,</u> Shigeaki Kato	AhR acts as an E3 ubiquitin ligase to modulate steroid receptor functions	Biochemical Pharmacology	77	474-484	2009
Kimura A, Naka T,	Aryl hydrocarbon receptor regulates Stat1	PNAS	105	9721-	2008



Nohara K, <u>Fujii-Kuriyama Y</u> , Kishimoto T.	activation and participates in the development of Th17 cells.			9726	
Hirabayashi Y, Yoon BI, Li GX, <u>Fujii-Kuriyama Y</u> , Kaneko T, Kanno J, Inoue T.	Benzene-induced hematopoietic toxicity transmitted by AhR in wild-type mouse and nullified by repopulation with AhR-deficient bone marrow cells: time after benzene treatment and recovery.	Chemosphere	73	S290-S294	2008
Baba T, Shima Y, Owaki A, Mimura J, Oshima M, <u>Fujii-Kuriyama Y</u> , Morohashi KI.	Disruption of aryl hydrocarbon receptor (AhR) induces regression of the seminal vesicle in aged male mice.	Sex Development	2	1-11	2008
Ohtake F, Baba A, <u>Fujii-Kuriyama Y</u> , Kato S.	Intrinsic AhR function underlies cross-talk of dioxins with sex hormone signalings.	Biochem Biophys Res Commun.	370	541-546	2008
Hosoya T, Harada N, Mimura J, Motohashi H, Takahashi S, Nakajima O, Morita M, Kawauchi S, Yamamoto M, <u>Fujii-Kuriyama Y</u> .	Inducibility of cytochrome P450 1A1 and chemical carcinogenesis by benzo[a]pyrene in AhR repressor-deficient mice.	Biochem Biophys Res Commun.	365	562-567	2008
Oshima M, Mimura J, Yamamoto M, <u>Fujii-Kuriyama Y</u> .	Molecular mechanism of transcriptional repression of AhR repressor involving ANKRA2, HDAC4, and HDAC5.	Biochem Biophys Res Commun.	364	276-282	2007
Matsumoto Y, Ide F, Kishi R, Akutagawa T, Sakai S, Nakamura M, Ishikawa T, <u>Fujii-Kuriyama Y</u> , Nakatsuru Y.	Aryl hydrocarbon receptor plays a significant role in mediating airborne particulate-induced carcinogenesis in mice.	Environ Sci Technol.	41	3775-3780	2007
Kawajiri K and <u>Fujii-Kuriyama Y</u> .	Cytochrome P450 Gene Regulation and Physiological Functions mediated by the Aryl Hydrocarbon Receptor.	Arch Biochem Biophys.	464	207-212	2007
Ohtake F, Baba A, Takada I, Okada M, Iwasaki K, Miki H, Takahashi S, Kouzmenko A, Nohara K, Chiba T, <u>Fujii-Kuriyama Y</u> and Kato S.	Dioxin receptor is a ligand-dependent E3 ubiquitin ligase.	Nature	446	562-566	2007
Goryo K, Suzuki A, Carpio CA, Siizaki K, Kuriyama E, Mikami Y, Kinoshita K, Yasumoto K, Rannug A, Miyamoto A, <u>Fujii-Kuriyama Y</u> , Sogawa K.	Identification of amino acid residues in the Ah receptor involved in ligand binding.	Biochem Biophys Res Commun.	354	396-402	2007
Hiromori, Y., <u>Nishikawa, J.</u> , Yoshida I, Nagase H, Nakanishi T.	Structure-dependent activation of peroxisome proliferator-activated receptor (PPAR) $\gamma$ by organotin compounds.	Chem. Biol. Interact.	180	238-244	2009

Zhen, H., Hu, J., Xiao, Y., Yang, M., Hirotsuji, J., Nishikawa, J., Nakanishi, T. and Ike, M.	Identification of retinoic acid receptor agonists in sewage treatment plants.	Environ. Sci. Technol.	43	6611-6616	2009
Takahashi M, Kanayama T, Yashiro T, Kondo H, Murase T, Hase T, Tokimitsu I, Nishikawa J, Sato R.	Effects of coumestrol on lipid and glucose metabolism as a farnesoid X receptor ligand.	Biochem Biophys Res Commun.	372	395-399	2008
Kamata R, Shiraishi F, Nishikawa J, Yonemoto J, Shiraishi H.	Screening and detection of the in vitro agonistic activity of xenobiotics on the retinoic acid receptor.	Toxicol in Vitro	22	1050-1061	2008
中西剛、西川淳一	重金属汚染による生物攪乱作用の分子基盤	細胞工学	26	1374-1379	2007
Matsumoto A, Mizukami H, Mizuno S, Umegaki K, Nishikawa J, Shudo K, Kagechika H, Inoue M.	Beta-Cryptoxanthin, a novel natural RAR ligand, induces ATP-binding cassette transporters in macrophages.	Biochem Pharmacology	74	256-264	2007
Ohta K, Ohigashi M, Naganawa A, Ikeda H, Sakai M, Nishikawa J, Imagawa M, Osada S, Nishihara T.	Histone acetyltransferase MOZ acts as a co- activator of Nrf2-MafK and induces tumour marker gene expression during hepatocarcinogenesis.	Biochem J.	402	559-566	2007
Jung J, Ishida K, Nishikawa J, Nishihara T.	Inhibition of estrogen action by 2-phenyl-chromone as AhR agonist in MCF-7 cells.	Life Sci.	81	1446-1451	2007
井上大介、松井久恵、清和成、楊敏、胡建英、荒金淳、廣辻淳二、西川淳一、池道彦	PRTR 化学物質の各種核内受容体に対する結合性	水環境学会誌	30	89-94	2007
井上 達	内分泌攪乱化学物質の低用量作用と毒性学のあたらしい課題.	科学	374	1022-1028	2009
Inoue T, and Hirabayashi, Y	Hematopoietic neoplastic diseases develop in C3H/He and C57BL/6 mice after benzene exposure: Strain differences in bone marrow tissue responses observed using microarrays.	Chem Biol Interact.	184	240-245	2010
Hirabayashi H, Inoue T.	Benzene-induced bone-marrow toxicity: A hematopoietic stem-cell-specific, aryl hydrocarbon receptor-mediated adverse effect.	Chem Biol Interact.	184	252-258	2010
Kawasaki Y, Hirabayashi Y, Kaneko T, Kanno J, Kodama Y, Matsushima Y, Ogawa Y, Saitoh M,	Benzene-induced hematopoietic neoplasms including myeloid leukemia in Trp53-deficient C57BL/6 and C3H/He mice.	Toxicol Sci.	110	293-306	2009

Sekita K, Uchida O, Umemura T, Yoon BI, and Inoue T.					
Yi JY, Hirabayashi Y, Choi YK, Kodama Y, Kanno J, Han J., Inoue T, and Yoon BI.	Benzene activates caspase-4 and -12 at the transcription level, without an association with apoptosis, in mouse bone marrow cells lacking the p53 gene.	Arch Toxicol.	83	795-803	2009
Hirabayashi T, and Inoue T.	Aryl hydro- carbon receptor biology and xenobiotic responses in hematopoietic progenitor cells.	Biochem Pharmacology	77	521-535	2009
Tohru Inoue, Yukio Kodama	Future alternatives in “3Rs”: Learning from history	AAEX.	14	257-260	2008
井上 達	内分泌かく乱化学物質研究の世界的動向	Biohilia	4	8-12	2008

IV. 研究成果の刊行物・別掲物・・・・・・・・・・・・・・・・・・217

## IV. 研究成果の刊行物・別刷り

Original Article

## Screening of toxicological properties of 4-methylbenzoic acid by oral administration to rats

Mariko Shirota, Takayuki Seki, Kazumi Tago, Hiroyasu Katoh, Hideki Marumo,  
Mami Furuya, Tomoko Shindo and Hiroshi Ono

Hatano Research Institute, Food and Drug Safety Center, 729-5 Ochiai, Hadano, Kanagawa 257-8523, Japan

(Received April 3, 2008; Accepted May 26, 2008)

**ABSTRACT** — Oral toxicity of 4-methylbenzoic acid in male and female Sprague-Dawley rats was profiled through a twenty-eight-day repeated dose toxicity study (the 28-day study) and a screening test for reproductive/developmental toxicities (the reproduction/developmental study) conducted under Organisation for Economic Co-operation and Development (OECD) test guidelines. Daily administration of 4-methylbenzoic acid, at a dose level of 0, 100, 300 or 1,000 mg/kg, did not show any adverse effect on reproductive organs of animals in the 28-day study. In the reproductive/developmental study, however, 1,000 mg/kg/day of the compound reduced epididymal weights and increased incidence of cauda epididymal oligo/azoospermia. While the compound did not affect estrous cycle or mating performances, 1,000 mg/kg of the compound reduced fertility. Furthermore, 300 mg/kg or more of the compound increased pre-implantation loss, which resulted in a decrease in the number of offspring, and reduced body weight gain of the dams during the latter period of gestation. From these results, the no-observed-effect-level (NOEL) for reproductive/developmental toxicities is considered to be 100 mg/kg, whereas 1,000 mg/kg did not show any effect on neonates. In the 28-day study, NOEL is considered to be 300 mg/kg for male and female rats, since 1,000 mg/kg of the compound caused, in both sexes, a few minor changes, such as temporal salivation, a slight increase in food consumption and a moderate increase in blood aspartate aminotransferase (AST) activity. Thus, 4-methylbenzoic acid has the potential for reproductive toxicity, with diverse adverse effects on the epididymis, after repeated administration, observed in the two studies.

**Key words:** 4-methylbenzoic acid, Epididymis, OECD test guideline, Reproductive toxicant, Repeated dose toxicity, Rats

### INTRODUCTION

Faced with an enormous number of existing chemicals that lacked hazard information, the Organisation for Economic Co-operation and Development (OECD) decided, in 1990, to undertake an investigation of such chemicals in cooperation with its member countries. They gave priority to high production volume (HPV) chemicals in collecting the data for an initial assessment of the hazard. A project was launched in 1990 to complete the dossiers of screening information data sets (SIDS) of toxicity for HPV chemicals through testing, and the work has been done cooperatively in Japan, as well as in other countries. Since 1991, we have participated in the testing under the auspices of the Ministry of Health (, Labour) and Welfare.

4-Methylbenzoic acid (*p*-toluic acid) was selected from

the OECD List of HPVs (OECD, 2004). 4-Methylbenzoic acid is produced at levels greater than 1,000 tones per year and is used in the manufacture of dye stuffs, colorants and paints; agrochemicals; and anticorrosive additives, as well as being an intermediate in the manufacture of polyethylene terephthalate. Although oral LD<sub>50</sub> values in rats and mice have been reported as 3,113 mg/kg and 2,115 mg/kg, respectively (Mineshita *et al.*, 1978), its toxicological properties are little known. The present study was performed to profile the oral toxicity of 4-methylbenzoic acid according to the standard protocols, "Repeated Dose 28-day Oral Toxicity Study" (the 28-day study) and "Reproduction/Developmental Toxicity Screening Test" (the reproduction/developmental study), in rodents (OECD, 1997a, 1997b).

Correspondence: Mariko Shirota (E-mail: shirota.m@fdsc.or.jp)

## MATERIALS AND METHODS

### Test substance

4-Methylbenzoic acid (CAS No. 99-94-5, purity 98.95%) was supplied by Toray (Tokyo, Japan), and was kept at room temperature until use. To prepare dosing samples, the compound was finely ground in a mortar, at first. Then, the ground compound for each dose was suspended in a 0.5% sodium carboxymethylcellulose solution, and each dose was adjusted to a constant volume of 5 ml/kg. The stability of the suspended compound in the vehicle and the content and uniformity of the compound of each dose were confirmed in Hatano Research Institute.

The doses used in the studies were determined based on a preliminary, 7-day, repeated dose, oral toxicity study of the compound, in which a dose level of 1,000 mg/kg did not show any toxic effects. This was considered sufficient for the highest dose, and the doses used in the present studies were set at 0, 100, 300 and 1,000 mg/kg.

### Animals

Male and female rats of the Sprague-Dawley (Crj: CD(SD)) strain were purchased from Charles River Japan (Atsugi Breeding Center, Atsugi, Kanagawa, Japan). For the 28-day study, the animals were purchased at 4 weeks of age, and after an 8-day quarantine period, the animals were divided into 4 groups of each sex according to a stratified allocation based on body weight measured on the day before initial dosing. For the reproduction/developmental study, the animals were purchased at 7 weeks of age and quarantined for 7 days. After the quarantine period, the female rats were monitored for estrous cycle by observing daily vaginal smears for 2 weeks. At 10 weeks of age, the females revolving on a regular 4-day estrous cycle and the males were divided into each 4 groups by the same method as used in the 28-day study.

These animals were kept individually in metallic cages with metal-meshed floors, except copulated females in the reproduction and developmental study, which were kept in flat-bottomed plastic cages with bedding materials (Paper Clean, Nippon SLC, Hamamatsu, Japan) from 18 days after copulation. The animal rooms were maintained with a light-dark cycle of 12-hr (lights on 7 hr), and temperature and relative humidity were controlled to 21.0-25.0°C and to 40.0-75.0%, respectively, with air ventilation at 15 complete changes per hour. The animals were supplied with solid rodent chow (CE-2, CLEA Japan, Tokyo, Japan) and tap water, *ad libitum*.

### Experimental design

All procedures described here were approved by the Committee on Animal Experiments of Hatano Research Institute, Food and Drug Safety Center.

#### The 28-day study

Groups given 0 and 1,000 mg/kg of 4-methylbenzoic acid comprised 10 animals of each sex, including 5 animals of each sex for the 2-week recovery study after the 28-day administration of the compound. The groups given 100 and 300 mg/kg comprised 5 animals of each sex.

Administration was begun at 5 weeks of age, and the initial day of the administration was designated as Day 1 of treatment. All the animals received daily administration of the compound by gavage for 28 days at a fixed time every day, and the initial day of recovery was designated as Day 1 of recovery.

Signs of toxicity were daily observed. Detailed clinical observations were made as specified in the OECD test guideline 407 (OECD, 1997a) in all animals prior to the initial administration, and once a week thereafter until the end of the recovery period. The observers were unaware of the treatment of each animal. Findings were recorded using a scoring system. In addition to the detailed observation of clinical signs, a 4-item neurobehavioral test battery assessing auditory and visual functions was administered at the last week of treatment.

Body weight was measured 3 times during the first week of the treatment, and twice a week thereafter. Food consumption was determined weekly.

Urinalysis was performed at the final week of treatment and at the final week of the recovery study. Urine of all animals was collected for 4 and 24 hr in a metabolic cage, and was examined for pH, occult blood, protein, ketone bodies, urobilinogen and bilirubin, semi-quantitatively using a urine test strip analyzer (Clinitek 200+, Bayer Medical, Tokyo, Japan), and for its color, turbidity and sediments. The volume and weight of the 24-hr urine was measured, and the specific gravity of the urine was calculated.

Necropsy of the animals was performed after 18-24 hr of fasting on the day following final treatment and on the day following the recovery period. Under anesthesia with pentobarbital sodium, blood samples were collected from the abdominal caval vein by syringe, with sodium citrate as an anticoagulant, for determination of coagulation times; with EDTA-2K potassium, for hematological examination; and with heparin, for blood chemical examination. Then, the animals were killed by exsanguination from the axillary artery. After gross observation, dissected organs, such as brain, thymus, heart, liver, kidneys, spleen, testes, adrenals and epididymides, were weighed.

## Screening of 4-methylbenzoic acid toxicities by OECD test guidelines

In addition to these organs, spinal cord, lungs, bronchi, stomach, ileum, colon, seminal vesicles, ovaries, uterus, vagina, urinary bladder, thyroid gland, femoral marrow, mesenteric lymph nodes, mandibular lymph nodes and ischiadic nerves were dissected out and fixed in buffered formalin for histopathological examination. The testes were fixed in Bouin's solution, with post fixation in buffered formalin. These organs/tissues were then processed for paraffin embedded block, and sections cut from the blocks were stained with hematoxylin-eosin (HE).

Hematological examination was carried out using automatic blood analysis apparatus (CELL-DYN3500SL, Abbot Diagnostics, IL, USA) for cell counts (erythrocytes (RBCs), leucocytes (WBCs) and platelets), hemoglobin concentration and differential WBC counts. Hematocrit, mean concentration of hemoglobin in the RBC (MCHC) and mean content of hemoglobin in the RBC were calculated. The prothrombin time (PT) and activated partial thromboplastin time were measured using a fully automatic analyzer for blood coagulation (CA-1000, Toa Medical Electronics, Saitama, Japan).

In the blood chemical examination, plasma concentrations of total protein, albumin, total cholesterol, glucose, blood urea nitrogen, creatinine, triglyceride, total bilirubin, inorganic phosphorus and calcium and activities of aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) and  $\gamma$ -glutamyl transpeptidase (GTP) were determined using a centrifugal automatic blood chemical analyzer (COBAS-FARA, Roche Diagnostics, Basel, Switzerland). Ratios of albumin to globulin were calculated, and plasma concentrations of sodium and potassium were measured using an automated electrolyte analyzer (EA05, A & T, Yokohama, Japan).

#### The reproduction/developmental study

The study consisted of 4 dosing groups, and each group comprised 13 males and 13 females. Administration was begun at 10 weeks of age, 2 weeks prior to mating in both, males and females, and was performed by gavage at a fixed time every day. The initial day of the administration was designated as Day 1 of treatment. Mating was performed by co-housing one male and one female from the same treatment group for a maximum of 2 weeks and was confirmed by observation of a copulatory plug or by the presence of sperm in a vaginal specimen. After confirmation of mating, each female was housed individually, and the day on which mating was confirmed was designated as Day 0 of gestation. Administration was continued during the mating period and gestation period, until 3 days after delivery for females. For males, the administration was continued through and after the mating period,

for a total of 42 days. For females that had copulated but did not deliver or for females that had not copulated, the administration was continued until 25 days after copulation or to Day 52 of treatment, respectively.

Clinical conditions were observed daily, and body weights and food consumptions were measured weekly. After confirming copulation, the females were weighed on Days 0, 7, 14 and 20 of gestation, and their food consumptions were determined on Days 0-1, 7-8, 14-15 and 20-21 of gestation. When the females delivered live fetuses, they were weighed on Days 0 (the day of delivery) and 4 of lactation, and their food consumptions were determined on Day 3-4 of lactation.

The estrous cycle was monitored daily until copulation. The females that had copulated were allowed to deliver spontaneously and to nurse their own pups until Day 4 of lactation. During the lactation period, the number and sex of dead and live pups were recorded for each dam, and the external morphology and general condition of the live pups were examined daily. The dead pups were examined for external and visceral abnormalities, when possible.

All the males were killed for necropsy on the day after Day 42 of treatment by exsanguination under sodium pentobarbital anesthesia. Maternal animals were similarly killed on Day 4 of lactation, while their offspring were killed for necropsy on the same day by ether inhalation. The females that had not copulated during the mating period and the females that had copulated but did not deliver fetuses were also similarly killed.

At necropsy, males were examined grossly for abnormalities, and the testes and epididymides were dissected, weighed and fixed in Bouin's solution, with post fixation with a buffered formalin solution for processing for histopathological examination. The other reproductive organs, such as ventral prostate and seminal vesicles with coagulating glands, were also dissected and fixed in a buffered formalin solution for histopathological examination. Females were examined grossly for abnormalities. Organs with lesions were dissected for histopathological examination. Ovaries, uterus and vagina from dams were dissected to determine the number of corpora lutea and implantation sites. Then, these organs were processed for histopathological examination. The absence of implantation sites in the uterus of females that had not copulated and females that had copulated but did not deliver fetuses was confirmed under a stereomicroscope. Live pups were euthanized by ether inhalation and were examined for external and visceral abnormalities.

#### **Statistical analysis**

Fisher's direct probability test was applied to analysis

of the incidence of animals in which the estrous cycle was altered, the copulation rate, the fertility index, the incidence of offspring with morphological abnormalities, and the incidence of histopathological findings. Graded findings in the histopathological examination were analyzed using the Mann-Whitney U-test. Data from the urinalysis urine quality test were analyzed using a chi-square test.

The other data were analyzed with multiple comparisons when comparing the data from more than 2 groups with those from the control. When comparing the data from a single group with those from the control, Student's t-test or Aspin-Welch's t-test was applied, following an F-test. In the multiple comparisons, an analysis of variance (ANOVA) test or Kruskal-Wallis's rank test was applied, following examination of the uniformity of variations by Bartlett's method. Significant differences from the control were determined by Dunnett's test. A *p* value of less than 0.5% was judged a significant difference.

## RESULTS

### Effects of repeated dosing in the 28-day study

Neither, death nor moribund condition, was observed in any animal, but a few male and female animals in the 1,000 mg/kg group showed temporary salivation after dosing. Except that, there were no clinical signs related to the treatment, and the scores of the detailed observation of clinical signs in the compound treated groups were comparable to those in the control (data not shown). In addition, no abnormalities were observed in the neurobehavioral test at the last week of treatment (data not shown).

Changes in body weight and food consumption are illustrated in Figs. 1 and 2, respectively. While body weights of males and females in the compound treated groups changed similarly to those in the control group at any period, food consumption of females in the 1,000 mg/kg treated group was slightly greater than in the control group, from Day 7 to 8 of treatment. Whereas the food consumption of females in this group tended to be greater thereafter, until the end of the treatment, no differences in food consumption were observed in males among any groups throughout the study.

As summarized in Table 1, urine volume measured on the Day 23 of treatment was larger in the males given 300 mg/kg or more of the compound and in females given 1,000 mg/kg when compared with those in the control. In addition, urine specific gravities were decreased in the males of these groups. These changes accompanied by an increase in water consumption, which was confirmed by an observation of water feeding bottles placed in the metabolic cages. Urinalysis showed some minor changes and

was not suggestive of any toxic effects (data not shown).

The hematology data are presented in Table 2. At the end of treatment period, platelet counts were slightly lower in the 1,000 mg/kg treated females, but were statistically insignificant when compared with those in the control. No changes were observed in any of the hematology parameters of males at the end of treatment period. At the end of recovery period, there were statistically significant differences between the males of compound treated and




Fig. 1. Body weight changes of male (A) and female (B) rats treated orally with 4-methylbenzoic acid for 28-days at dose levels of 0 ( $\circ$ ), 100 ( $\blacksquare$ ), 300 ( $\blacktriangle$ ) and 1,000 mg/kg/day ( $\blacklozenge$ ) in the repeated dose 28-day oral toxicity study. Values during the treatment period represent the average for 5 animals in the 100 or 300 mg/kg-treated group and for 10 animals in the 0 or 1,000 mg/kg-treated group. Those during the recovery period represent the average for 5 animals.



## Screening of 4-methylbenzoic acid toxicities by OECD test guidelines

control groups, and were not suggestive any toxic effects.

The blood chemistry data are presented in Table 3. At the end of treatment period, a moderate increase in AST activity and a slight decrease in total protein concentration were observed in the 1,000 mg/kg-treated females, compared with those of control. In those data from males and those at the end of the recovery period from animals of both sexes, no significant differences were observed between the compound treated and control groups.

No significant differences from the control were observed in absolute organ weights or in organ weights relative to body weight in any of the compound treated groups of males (Table 4) or females (data not shown).

No abnormalities suggestive of any toxic effects were observed in any organs or tissues on gross examination at necropsy or histopathological examination (data not shown).

#### Effects of repeated dosing in the reproduction/developmental study

As found in the 28-day study, temporary salivation was observed in a few animals given 1,000 mg/kg (data not shown). Except that, there were no clinical signs relating to the treatment.

Changes in body weight and food consumption of males are shown in Figs. 3 and 4, respectively. The compound did not affect body weight increase or food consumption, at any dose level.

Changes in body weight and food consumption of females are shown in Figs. 5 and 6, respectively. The compound did not affect body weights until Day 14 of gestation, at any dose level, while food consumption of the

1,000 mg/kg treated group was slightly higher than that of the control at the beginning of the dosing. During Days 14-20 of gestation, however, maternal body weight gain was reduced in the 300 mg/kg or more treated groups, while the food consumption that was determined during this period (Day 14-15 of gestation) and body weight on




Fig. 2. Changes in food consumption of male (A) and female (B) rats treated orally with 4-methylbenzoic acid for 28-days at dose levels of 0 (open column), 100 (dashed column), 300 (hatched column) or 1,000 mg/kg/day (closed column) in the repeated dose 28-day oral toxicity study. Each column represents average for 5 animals in the 100 or 300 mg/kg-treated group and for 10 animals in the 0 or 1,000 mg/kg-treated group. During the recovery period, it represents the average for 5 animals. \* indicates significant difference from control at  $p < 0.05$ .

Table 1. Urinalysis of rats treated orally with 4-methylbenzoic acid in repeated dose 28-day oral toxicity study

Dose (mg/kg)	On Day 23 of treatment				On Day 9 of recovery	
	0	100	300	1,000	0	1,000
<u>Number of animals (males/females)</u>						
	10/10	5/5	5/5	10/10	5/5	5/5
<u>Urinary volume (ml/24 hr)</u>						
Males	15.6 ± 2.2	15.6 ± 2.1	20.9 ± 4.7*	23.8 ± 4.7**	18.3 ± 4.7	24.3 ± 4.6
Females	11.7 ± 3.5	11.9 ± 3.4	13.0 ± 4.3	22.1 ± 7.5**	13.3 ± 2.8	18.3 ± 5.6
<u>Specific gravity</u>						
Males	1.058 ± 0.008	1.051 ± 0.007	1.043 ± 0.011**	1.045 ± 0.006**	1.056 ± 0.009	1.038 ± 0.006**
Females	1.045 ± 0.012	1.046 ± 0.008	1.043 ± 0.007	1.038 ± 0.009	1.041 ± 0.010	1.044 ± 0.011

Values represent average ± S.D.

\* and \*\*, significant difference from control at  $p < 0.05$  and 0.01, respectively.

**Table 2.** Hematology data of rats treated orally with 4-methylbenzoic acid in the repeated dose 28-day oral toxicity study

Dose (mg/kg)	End of treatment						The end of recovery					
	Males			Females			Males			Females		
	0	100	300	1,000	0	100	300	1,000	0	1,000	0	1,000
RBC ( $\times 10^6/\mu\text{l}$ )	755 $\pm$ 53	780 $\pm$ 15	770 $\pm$ 27	755 $\pm$ 18	742 $\pm$ 19	771 $\pm$ 26	750 $\pm$ 25	751 $\pm$ 30	797 $\pm$ 13	766 $\pm$ 22**	749 $\pm$ 25	766 $\pm$ 24
Hemoglobin (g/dl)	14.9 $\pm$ 0.6	15.2 $\pm$ 0.2	15.2 $\pm$ 0.3	15.0 $\pm$ 0.2	14.8 $\pm$ 0.3	15.1 $\pm$ 0.5	14.9 $\pm$ 0.3	14.7 $\pm$ 0.5	15.2 $\pm$ 0.1	14.6 $\pm$ 0.3**	14.5 $\pm$ 0.4	14.4 $\pm$ 0.7
Hematocrit (%)	44.8 $\pm$ 2.9	46.0 $\pm$ 0.7	45.5 $\pm$ 1.1	45.2 $\pm$ 0.4	44.2 $\pm$ 1.0	44.9 $\pm$ 1.3	44.4 $\pm$ 0.7	43.6 $\pm$ 1.9	45.3 $\pm$ 0.8	43.5 $\pm$ 1.0**	42.6 $\pm$ 1.3	42.7 $\pm$ 1.9
MCV (fl)	59.4 $\pm$ 2.2	58.9 $\pm$ 0.8	59.0 $\pm$ 1.3	60.0 $\pm$ 1.5	59.5 $\pm$ 1.1	58.3 $\pm$ 1.2	59.2 $\pm$ 1.4	58.0 $\pm$ 1.1	56.8 $\pm$ 1.4	56.8 $\pm$ 0.8	56.9 $\pm$ 1.2	55.8 $\pm$ 2.0
MCH (pg)	19.8 $\pm$ 0.8	19.5 $\pm$ 0.6	19.8 $\pm$ 0.6	19.9 $\pm$ 0.5	20.0 $\pm$ 0.6	19.6 $\pm$ 0.3	19.8 $\pm$ 0.6	19.6 $\pm$ 0.3	19.1 $\pm$ 0.4	19.0 $\pm$ 0.3	19.4 $\pm$ 0.4	18.8 $\pm$ 0.7
MCHC (g/dl)	33.3 $\pm$ 0.1	33.0 $\pm$ 0.7	33.6 $\pm$ 0.3	33.2 $\pm$ 0.2	33.6 $\pm$ 0.4	33.7 $\pm$ 0.3	33.5 $\pm$ 0.2	33.8 $\pm$ 0.6	33.5 $\pm$ 0.6	33.5 $\pm$ 0.4	34.0 $\pm$ 0.3	33.7 $\pm$ 0.3
Platelet ( $\times 10^4/\mu\text{l}$ )	102.9 $\pm$ 13.9	104.5 $\pm$ 16.3	106.6 $\pm$ 8.1	101.6 $\pm$ 10.4	96.3 $\pm$ 7.0	100.5 $\pm$ 9.4	93.3 $\pm$ 9.3	83.9 $\pm$ 3.1	96.1 $\pm$ 19.6	99.3 $\pm$ 8.4	101.8 $\pm$ 12.7	102.7 $\pm$ 10.2
PT (sec)	18.4 $\pm$ 3.0	18.9 $\pm$ 3.1	17.2 $\pm$ 4.0	14.9 $\pm$ 1.7	12.6 $\pm$ 0.7	12.7 $\pm$ 0.5	12.9 $\pm$ 0.6	13.2 $\pm$ 0.6	16.4 $\pm$ 2.7	15.2 $\pm$ 1.4	11.9 $\pm$ 0.7	11.5 $\pm$ 0.7
APTT (sec)	22.1 $\pm$ 0.8	22.9 $\pm$ 1.7	20.5 $\pm$ 2.4	20.1 $\pm$ 1.0	18.5 $\pm$ 1.0	18.2 $\pm$ 1.4	17.8 $\pm$ 1.2	16.7 $\pm$ 1.4	21.6 $\pm$ 1.5	21.0 $\pm$ 0.4	17.1 $\pm$ 0.8	16.9 $\pm$ 0.8
WBC ( $\times 100/\mu\text{l}$ )	90.8 $\pm$ 16.7	61.5 $\pm$ 20.5	67.7 $\pm$ 17.6	74.4 $\pm$ 21.7	61.0 $\pm$ 18.1	54.9 $\pm$ 14.8	46.7 $\pm$ 12.4	59.8 $\pm$ 7.8	69.4 $\pm$ 13.4	65.4 $\pm$ 14.2	43.9 $\pm$ 2.9	58.8 $\pm$ 14.5
Differential leukocyte counts (%)												
Neutrophil	10 $\pm$ 5	11 $\pm$ 2	10 $\pm$ 2	11 $\pm$ 3	7 $\pm$ 2	6 $\pm$ 3	8 $\pm$ 4	8 $\pm$ 4	13 $\pm$ 5	15 $\pm$ 3	10 $\pm$ 4	10 $\pm$ 3
Eosinophil	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	0 $\pm$ 0*	1 $\pm$ 0	1 $\pm$ 1	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 1	2 $\pm$ 1	1 $\pm$ 1
Basophil	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0	0 $\pm$ 0
Monocyte	3 $\pm$ 1	4 $\pm$ 1	5 $\pm$ 1	4 $\pm$ 1	3 $\pm$ 1	4 $\pm$ 1	3 $\pm$ 1	3 $\pm$ 1	4 $\pm$ 1	4 $\pm$ 2	4 $\pm$ 1	4 $\pm$ 2
Lymphocyte	85 $\pm$ 5	85 $\pm$ 3	85 $\pm$ 3	85 $\pm$ 3	89 $\pm$ 3	88 $\pm$ 3	88 $\pm$ 4	88 $\pm$ 4	81 $\pm$ 4	79 $\pm$ 4	85 $\pm$ 5	85 $\pm$ 5

Values represent average for five animals  $\pm$  S.D.

\* and \*\*, significant difference from control at  $p < 0.05$  and  $0.01$ , respectively.

## Screening of 4-methylbenzoic acid toxicities by OECD test guidelines

**Table 3.** Blood chemistry data of rats treated orally with 4-methylbenzoic acid in repeated dose 28-day oral toxicity study

Dose (mg/kg)	The end of treatment						The end of recovery					
	Males			Females			Males			Females		
	0	100	300	1,000	0	100	300	1,000	0	1,000	0	1,000
TP (g/dl)	5.0 ± 0.2	5.2 ± 0.2	5.1 ± 0.1	5.0 ± 0.0	5.4 ± 0.3	5.3 ± 0.3	5.2 ± 0.3	4.8 ± 0.3**	5.6 ± 0.4	5.4 ± 0.1	5.5 ± 0.1	5.7 ± 0.3
Albumin (g/dl)	3.0 ± 0.2	3.0 ± 0.3	3.0 ± 0.1	3.0 ± 0.1	3.2 ± 0.2	3.2 ± 0.2	3.2 ± 0.2	2.9 ± 0.2	3.0 ± 0.2	3.0 ± 0.1	3.2 ± 0.2	3.3 ± 0.4
A/G	1.48 ± 0.15	1.46 ± 0.27	1.38 ± 0.16	1.44 ± 0.08	1.51 ± 0.17	1.48 ± 0.17	1.64 ± 0.11	1.56 ± 0.09	1.14 ± 0.08	1.28 ± 0.14	1.41 ± 0.20	1.37 ± 0.24
BUN (mg/dl)	16 ± 2	17 ± 2	16 ± 3	16 ± 1	23 ± 3	21 ± 2	23 ± 1	20 ± 2	19 ± 3	15 ± 3	21 ± 1	24 ± 3
Creatinine (mg/dl)	0.6 ± 0.1	0.6 ± 0.1	0.6 ± 0.0	0.6 ± 0.1	0.6 ± 0.1	0.6 ± 0.1	0.7 ± 0.1	0.7 ± 0.1	0.7 ± 0.0	0.6 ± 0.1	0.7 ± 0.0	0.8 ± 0.1
Glucose (mg/dl)	130 ± 13	137 ± 32	143 ± 13	155 ± 24	109 ± 13	113 ± 21	118 ± 7	108 ± 10	137 ± 10	129 ± 16	125 ± 12	128 ± 10
T. Cholest.(mg/dl)	34 ± 8	33 ± 4	37 ± 5	40 ± 5	41 ± 8	48 ± 9	45 ± 10	31 ± 8	41 ± 15	38 ± 10	48 ± 9	54 ± 4
Triglyceride (mg/dl)	19 ± 5	22 ± 7	28 ± 14	35 ± 15	11 ± 5	9 ± 3	10 ± 3	9 ± 2	24 ± 13	29 ± 12	13 ± 2	22 ± 12
T. Bil.(mg/dl)	0.07 ± 0.04	0.08 ± 0.03	0.09 ± 0.05	0.07 ± 0.03	0.09 ± 0.03	0.07 ± 0.03	0.09 ± 0.04	0.09 ± 0.03	0.08 ± 0.02	0.09 ± 0.03	1.10 ± 0.04	0.08 ± 0.02
Inorg. P. (mg/dl)	9.0 ± 0.7	8.7 ± 0.8	8.4 ± 0.3	8.6 ± 0.4	7.9 ± 0.8	8.1 ± 0.7	8.2 ± 1.1	8.4 ± 0.5	7.2 ± 0.9	6.5 ± 0.6	7.4 ± 0.9	7.5 ± 0.5
Ca (mg/dl)	8.9 ± 0.1	8.9 ± 0.3	9.0 ± 0.1	9.2 ± 0.2	9.1 ± 0.1	9.1 ± 0.2	9.1 ± 0.2	8.8 ± 0.4	8.9 ± 0.2	8.8 ± 0.1	9.1 ± 0.2	9.2 ± 0.2
Na (mEq/l)	145.8 ± 0.6	145.1 ± 0.9	145.8 ± 1.3	145.5 ± 0.9	145.5 ± 0.8	144.7 ± 0.7	145.6 ± 0.8	146.0 ± 1.4	146.2 ± 0.4	145.9 ± 0.6	143.1 ± 0.8	143.4 ± 0.2
K (mEq/l)	4.48 ± 0.53	4.33 ± 0.36	4.08 ± 0.26	4.29 ± 0.24	4.18 ± 0.39	4.31 ± 0.20	4.20 ± 0.28	4.05 ± 0.38	4.03 ± 0.26	3.96 ± 0.12	4.11 ± 0.28	3.98 ± 0.32
Cl (mEq/l)	109.0 ± 1.7	108.3 ± 1.7	108.1 ± 1.2	107.8 ± 1.4	109.1 ± 1.2	107.5 ± 1.0	109.9 ± 1.1	109.3 ± 2.3	108.2 ± 1.1	108.7 ± 0.6	107.2 ± 0.6	107.8 ± 0.6
ALP (U/l)	441 ± 143	451 ± 58	511 ± 190	459 ± 56	319 ± 65	261 ± 62	299 ± 75	266 ± 46	366 ± 80	374 ± 29	219 ± 36	231 ± 39
ALT (U/l)	31 ± 6	27 ± 4	29 ± 3	37 ± 15	25 ± 2	22 ± 2	22 ± 5	29 ± 5	35 ± 5	32 ± 4	24 ± 3	28 ± 5
AST (U/l)	72 ± 10	66 ± 11	67 ± 7	88 ± 24	69 ± 3	66 ± 5	69 ± 8	94 ± 16*	83 ± 14	66 ± 2	62 ± 3	66 ± 5
γ-GTP (U/l)	0 ± 1	0 ± 1	1 ± 0	1 ± 1	1 ± 0	1 ± 0	1 ± 1	1 ± 0	0 ± 1	1 ± 1	1 ± 1	1 ± 1

Values represent average for five animals ± S.D.

\* and \*\*, significant difference from control at p &lt; 0.05 and 0.01, respectively.

TP, total protein; A/G, albumin ratio to globulin (TP - albumin); BUN, blood urea nitrogen; T. Cholest., total cholesterol; T. Bil., total bilirubin; Inorg. P., Inorganic phosphorus; ALP, alkaline phosphatase; ALT, alanine aminotransferase; AST, aspartate aminotransferase; γ -GTP, γ-glutamyl transpeptidase

**Table 4.** Organ weights of male rats treated orally with 4-methylbenzoic acid in repeated dose 28-day oral toxicity study

Dose (mg/kg)	The end of treatment				End of recovery	
	0	100	300	1,000	0	1,000
Number of animals	5	5	5	5	5	5
Body weight (g)	307.9 ± 24.7	301.4 ± 32.0	311.2 ± 4.2	311.8 ± 30.7	380.8 ± 43.4	370.2 ± 15.1
<u>Absolute weight</u>						
Brain (g)	1.92 ± 0.07	1.87 ± 0.06	1.89 ± 0.07	1.84 ± 0.07	1.92 ± 0.11	1.91 ± 0.07
Thymus (mg)	587 ± 86	493 ± 61	537 ± 95	535 ± 98	423 ± 42	481 ± 132
Heart (g)	1.10 ± 0.09	1.01 ± 0.06	1.00 ± 0.05	1.06 ± 0.11	1.21 ± 0.15	1.18 ± 0.12
Liver (g)	9.62 ± 0.99	9.97 ± 1.87	9.73 ± 0.46	10.26 ± 1.15	10.95 ± 2.00	10.88 ± 0.46
Kidneys (g)	2.39 ± 0.20	2.42 ± 0.24	2.43 ± 0.17	2.42 ± 0.28	2.72 ± 0.30	2.69 ± 0.27
Spleen (mg)	707 ± 192	611 ± 74	643 ± 85	591 ± 106	743 ± 114	694 ± 93
Testes (g)	2.69 ± 0.01	2.84 ± 0.14	3.00 ± 0.28	2.87 ± 0.35	2.94 ± 0.16	2.95 ± 0.15
Epididymides (g)	0.67 ± 0.04	0.71 ± 0.07	0.70 ± 0.02	0.69 ± 0.10	0.91 ± 0.06	0.90 ± 0.04
Adrenal glands (mg)	50.9 ± 3.6	45.6 ± 5.6	54.6 ± 4.6	47.1 ± 6.3	52.7 ± 6.0	55.8 ± 6.8
<u>Relative weight (g/100 g)</u>						
Brain	0.63 ± 0.04	0.62 ± 0.05	0.61 ± 0.03	0.60 ± 0.06	0.51 ± 0.03	0.52 ± 0.01
Thymus	0.19 ± 0.02	0.16 ± 0.02	0.17 ± 0.03	0.17 ± 0.02	0.11 ± 0.02	0.13 ± 0.04
Heart	0.36 ± 0.04	0.34 ± 0.03	0.32 ± 0.01	0.34 ± 0.02	0.32 ± 0.05	0.32 ± 0.03
Liver	3.12 ± 0.16	3.29 ± 0.36	3.13 ± 0.17	3.29 ± 0.14	2.87 ± 0.32	2.94 ± 0.08
Kidneys	0.78 ± 0.05	0.80 ± 0.05	0.78 ± 0.05	0.78 ± 0.06	0.72 ± 0.09	0.73 ± 0.07
Spleen	0.23 ± 0.04	0.20 ± 0.02	0.21 ± 0.03	0.19 ± 0.03	0.20 ± 0.02	0.19 ± 0.02
Testes	0.88 ± 0.13	0.95 ± 0.11	0.97 ± 0.09	0.92 ± 0.05	0.78 ± 0.07	0.80 ± 0.06
Epididymides	0.22 ± 0.03	0.24 ± 0.02	0.23 ± 0.00	0.22 ± 0.02	0.24 ± 0.02	0.24 ± 0.02
Adrenal glands	0.017 ± 0.002	0.015 ± 0.001	0.018 ± 0.002	0.015 ± 0.001	0.014 ± 0.001	0.015 ± 0.002

Values represent average ± S.D.

the day of delivery (Day 0 of lactation) were higher in the 1,000 mg/kg treated group and in the 300 mg/kg or more treated groups, respectively, compared with those in the control. During the lactation period, maternal body weight gain and food consumption in the 1,000 mg/kg treated group were smaller than those in the control.

Terminal body weight and the weights of testes and epididymides of the male rats treated for 42 days are shown in Table 5. The compound did not affect the terminal body weight, at any dose level. While no differences were observed in the absolute or relative testicu-

lar weights between the control and the compound treated groups, those of the epididymides were significantly lower in the 1,000 mg/kg treated group than in the control group. Histopathological examination revealed that there were lumens containing no or few spermatozoa, i.e., oligo/azoospermia, in the cauda epididymis in all the males of this group (Fig. 7), while none of their caput epididymis showed the same abnormality. In addition to this, the number of animals with cell debris in the cauda epididymal lumen was increased in this group, although it was statistically insignificant when compared with that in the