

3. Smith AG. Embryo-derived stem cells of mice and men. *Annu Rev Cell Dev Biol.* 2001;17:435-62.
4. Chambers I, Smith A. Self-renewal of teratocarcinoma and embryonic stem cells. *Oncogene.* 2004;23:7:50-60.
5. Smith AG, Heath JK, Donaldson DD, et al. Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature.* 1988;336:688-90.
6. Williams RL, Hilton DJ, Pease S, et al. Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. *Nature.* 1988;336:684-7.
7. Ying QL, Nichols J, Chambers I, et al. BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell.* 2003;115:281-92.
8. Nichols J, Zevnik B, Anastassiadis K, et al. Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct1. *Cell.* 1998;95:379-91.
9. Mitsui K, Tokuzawa Y, Itoh H, et al. The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell.* 2003;113:631-42.
10. Chambers I, Colby D, Robertson M, et al. Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell.* 2003;113:643-55.
11. Niwa H, Miyazaki J, Smith AG. Quantitative expression of Oct 3/1 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet.* 2000;21:372-6.
12. Shambhutt MJ, Axelman J, Wang S, et al. Derivation of pluripotent stem cells from cultured human primordial germ cells. *Proc Natl Acad Sci U S A.* 1998;95:13776-31.
13. Thomson JA, Itskovitz-Eldor J, Shapiro SS, et al. Embryonic stem cell lines derived from human blastocysts. *Science.* 1998;282:1145-7.
14. Rao M. Conserved and divergent paths that regulate self-renewal in mouse and human embryonic stem cells. *Dev Biol.* 2004;275:260-86.
15. Herrmann BG, Labat S, Poustka A, et al. Cloning of the T gene required in mesoderm formation in the mouse. *Nature.* 1990;343:617-22.
16. Wilkinson DG, Bhatt S, Herrmann BG. Expression pattern of the mouse T gene and its role in mesoderm formation. *Nature.* 1990;343:657-9.
17. Ramalho Santos M, Yoon S, Matsuzaki Y, et al. "Stemness": transcriptional profiling of embryonic and adult stem cells. *Science.* 2002;298:597-600.
18. Raz R, Lee CK, Cannizzaro LA, et al. Essential role of STAT3 for embryonic stem cell pluripotency. *Proc Natl Acad Sci U S A.* 1999;96:2846-51.
19. Gossen M, Freundlieb S, Bender G, et al. Transcriptional activation by tetracyclines in mammalian cells. *Science.* 1995;268:1766-9.
20. Hofmann A, Czichos S, Kaps C, et al. The T-box transcription factor Brachyury mediates cartilage development in mesenchymal stem cell line C3H10T1/2. *J Cell Sci.* 2002;115:769-81.
21. Kispert A, Herrmann BG. The Brachyury gene encodes a novel DNA binding protein. *EMBO J.* 1993;12:3211-20.
22. Haroi Y, Kudoh S, Monzen K, et al. Tbx5 associates with Nkx2-5 and synergistically promotes cardiomyocyte differentiation. *Nat Genet.* 2001;28:276-80.
23. Stennard EA, Costa MW, Elliott DA, et al. Cardiac T-box factor Tbx20 directly interacts with Nkx2-5, GATA4, and GATA5 in regulation of gene expression in the developing heart. *Dev Biol.* 2003;262:206-24.
24. Garg V, Kathuria IS, Barnes R, et al. GATA4 mutations cause human congenital heart defects and reveal an interaction with TBX5. *Nature.* 2003;424:443-7.
25. Zhu M, John S, Berg M, et al. Functional association of Nfu with Stat5 and Stat1 in IL-2- and IFN-gamma-mediated signaling. *Cell.* 1999;96:121-30.
26. Cellum RG, Brutsaert S, Lee G, et al. A Stat3-interacting protein (STIP1) regulates cytokine signal transduction. *Proc Natl Acad Sci U S A.* 2000;97:10120-5.
27. Dale L, Howes G, Price BM, et al. Bone morphogenetic protein 4: a ventralizing factor in early Xenopus development. *Development.* 1992;115:573-85.
28. Jones CM, Lyons KM, Lipan PM, et al. DVR-4 (bone morphogenetic protein-4) as a posterior-ventralizing factor in Xenopus mesoderm induction. *Development.* 1992;115:639-7.
29. Johansson BM, Wilks MV. Evidence for involvement of activin A and bone morphogenetic protein 4 in mammalian mesoderm and hematopoietic development. *Mol Cell Biol.* 1995;15:141-51.
30. Finley MF, Devata S, Haettner JF. BMP-4 inhibits neural differentiation of murine embryonic stem cells. *J Neurobiol.* 1999;40:271-87.
31. Czyz J, Wobus A. Embryonic stem cell differentiation: the role of extracellular factors. *Differentiation.* 2001;68:167-74.
32. Holley SA, Neul JL, Attisano L, et al. The Xenopus dorsalizing factor noggin ventralizes Drosophila embryos by preventing DPP from activating its receptor. *Cell.* 1996;86:607-17.
33. Zimmerman LB, De Jesus-Escobar JM, Harland RM. The Spemann organizer signal noggin binds and inactivates bone morphogenetic protein 4. *Cell.* 1996;86:599-606.
34. Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell.* 2003;113:685-700.
35. Hoffrage A, Oehlmann V, Heymer J, et al. Id genes are direct targets of bone morphogenetic protein induction in embryonic stem cells. *J Biol Chem.* 1999;274:19838-45.
36. Nakashima K, Takizawa T, Ochiai W, et al. BMP2-mediated alteration in the development of fetal mouse brain cells from neurogenesis to astrocytogenesis. *Proc Natl Acad Sci U S A.* 2001;98:5868-73.
37. Feng XII, Zhang Y, Wu RY, et al. The tumor suppressor Smad4/DPC4 and transcriptional adaptor CBP/p300 are coactivators for smad3 in TGF-beta-induced transcriptional activation. *Genes Dev.* 1998;12:2153-63.
38. Janknecht R, Wells NJ, Hunter T. TGF-beta-stimulated cooperation of smad proteins with the coactivators CBP/p300. *Genes Dev.* 1998;12:2114-9.
39. Pearson KL, Hunter T, Janknecht R. Activation of Smad1-mediated transcription by p300/CBP. *Biochim Biophys Acta.* 1999;1489:354-64.
40. Latinkic BV, Umbhauer M, Neal KA, et al. The Xenopus Brachyury promoter is activated by FGF and low concentrations of activin and suppressed by high concentrations of activin and by paired-type homeodomain proteins. *Genes Dev.* 1997;11:3265-76.
41. Kim J, Johnson K, Chen HJ, et al. Drosophila Mad binds to DNA and directly mediates activation of vestigial by Decapentaplegic. *Nature.* 1997;388:304-8.
42. Rathjen J, Lake JA, Bettess MD, et al. Formation of a primitive ectoderm like cell population, EPL cells, from ES cells in response to biologically derived factors. *J Cell Sci.* 1999;112 (Pt 5):601-12.
43. Lake J, Rathjen J, Remiszewski J, et al. Reversible programming of pluripotent cell differentiation. *J Cell Sci.* 2000;113(Pt 3):555-66.
44. Tropepe V, Hitoshi S, Sirard C, et al. Direct neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a default mechanism. *Neuron.* 2001;30:65-78.

45. Rathjen PD, Nichols J, Toth S, et al. Developmentally programmed induction of differentiation inhibiting activity and the control of stem cell populations. *Genes Dev.* 1990;4:2308-18.
46. Nichols J, Chambers I, Taga T, et al. Physiological rationale for responsiveness of mouse embryonic stem cells to gp130 cytokines. *Development.* 2001;128:2333-9.
47. Daheron L, Opitz SL, Zaehres H, et al. LIF/STAT3 signaling fails to maintain self-renewal of human embryonic stem cells. *Stem Cells.* 2004;22:770-8.
48. Suzuki A, Raya A, Kawakami Y, et al. Maintenance of embryonic stem cell pluripotency by Nanog-mediated reversal of mesoderm specification. *Nat Clin Pract Cardiovasc Med.* 2006;3 Suppl 1:S114-22.
49. Suzuki A, Raya A, Kawakami Y, et al. Nanog binds to Smad1 and blocks bone morphogenetic protein-induced differentiation of embryonic stem cells. *Proc Natl Acad Sci U S A.* 2006;103:10294-9.

X11-Like Protein Deficiency Is Associated with Impaired Conflict Resolution in Mice

Yoshitake Sano,¹ Veravej G. Ornthanalai,² Kazuyuki Yamada,³ Chihiro Homma,³ Hitomi Suzuki,¹ Toshiharu Suzuki,⁴ Niall P. Murphy,² and Shigeyoshi Itohara¹

¹Laboratory for Behavioral Genetics, ²Molecular Neuropathology Group, and ³Research Resource Center, RIKEN Brain Science Institute, Wako, Saitama 351-0198, Japan, and ⁴Laboratory of Neuroscience, Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo, Hokkaido 060-0812, Japan

Understanding how emotion is generated, how conflicting emotions are regulated, and how emotional states relate to sophisticated behaviors is a crucial challenge in brain research. Model animals showing selective emotion-related phenotypes are highly useful for examining these issues. Here, we describe a novel mouse model that withdraws in approach-avoidance conflicts. X11-like (X11L)/Mint2 is a neuronal adapter protein with multiple protein-protein interaction domains that interacts with several proteins involved in modulating neuronal activity. X11L-knock-out (KO) mice were subordinate under competitive feeding conditions. X11L-KO mice lost significantly more weight than cohoused wild-type mice without signs of decreased motivation to eat or physical weakness. In a resident-intruder test, X11L-KO mice showed decreased intruder exploration behavior. Moreover, X11L-KO mice displayed decreased marble-burying, digging and burrowing behaviors, indicating aberrant ethological responses to attractive stimuli. In contrast, X11L-KO mice were indistinguishable from wild-type mice in the open field, elevated plus maze, and light/dark transition tests, which are often used to assess anxiety-like behavior. Neurochemical analysis revealed a monoamine imbalance in several forebrain regions. The defective ethological responses and social behaviors in X11L-KO mice were rescued by the expression of X11L under a *Camk2a* promoter using the Tet-OFF system during development. These findings suggest that X11L is involved in the development of neuronal circuits that contribute to conflict resolution.

Introduction

Behavior arises from the integration of multiple emotional processes, which may themselves conflict. Approach-avoidance conflict in response to various emotional stimuli (e.g., seeking friendship, overcoming shyness or desiring success, avoiding competition) is a common everyday experience. Studies of human genetics and genetically engineered mice have gradually begun to unravel the molecular and neuronal mechanisms underlying emotion-related behavior, and cumulative evidence supports a crucial role of monoaminergic systems (Lesch et al., 1996; Zhuang et al., 1999; Gross et al., 2002; Finn et al., 2003; Holmes et al., 2003; Reif and Lesch, 2003). Behavioral paradigms designed to assess anxiety-related behaviors are relatively well established. In many of these paradigms, anxiety-like behavior is inferred by the avoidance of unpleasant stimuli under conflicting situations. However, more sophisticated paradigms and model animals that allow us to distinguish various emotional states are

required to clarify how conflicting emotions are regulated in the brain and how emotional states underlie behavior.

The neuronal adaptor protein X11L (also called X11 β and Munc18–1 interacting protein 2; Mint2) interacts with amyloid precursor protein (APP) and regulates APP metabolism (Tomita et al., 1999; Araki et al., 2003; Lee et al., 2004; Sano et al., 2006a; Saito et al., 2008). X11L suppresses amyloidogenic processing of APP via intracellular sorting machinery, and thus the lack of X11L in mice is associated with an increase in amyloid β (A β) 40 and 42 in the hippocampus (Sano et al., 2006a). X11L constitutes the X11 protein family with X11/X11 α /Mint1 and X11L2/X11 γ /Mint3, which comprises a poorly conserved N-terminal region, a conserved central phosphotyrosine interaction domain, and two C-terminal PDZ [postsynaptic density-95/*Drosophila* disks-large/zona occludens-1] domains (Tomita et al., 1999). X11L and X11 are brain-specific proteins that interact with many molecules, such as Munc18–1, calcium channels, β -neurexin, and hyperpolarization-activated cyclic nucleotide-gated potassium channel 2, involved in the regulation of neuronal/synaptic activity (Okamoto and Südhof, 1997; Maximov et al., 1999; Tomita et al., 1999; Biederer and Südhof, 2000; Kimura et al., 2004; Rogelj et al., 2006). Approximately 80% of Mint1/2 (X11/X11L) double-knock-out (KO) mice, but not single-gene KO mice, die at birth, and presynaptic release probability is decreased in the hippocampal excitatory neurons of double-KO mice, but not Mint1 (X11) single-KO mice (Ho et al., 2003, 2006), indicating substantial redundancies between X11 and X11L functions. The physiological roles of X11L, however, are not well understood.

Received Dec. 3, 2008; revised March 12, 2009; accepted April 4, 2009.

This work is supported in part by a Grant-in-Aid for Young Scientists (B) from Japan Society for the Promotion of Science 18700403 to Y.S. We thank Chadwick Boulay for preliminary neurochemical analyses, Sam Gandy for the G369 antibody, Takayuki Sassa for the HLS-LacZ plasmid, Hidenori Taru for the FLAG-hX11L plasmid, and the staff of the RIKEN Brain Science Institute animal facility for their commitment to maintaining animals and preparing the behavioral apparatuses. We also thank all of our colleagues in our laboratories for their encouragement and discussions.

Correspondence should be addressed to Shigeyoshi Itohara, Laboratory for Behavioral Genetics, RIKEN BSI, 2-1 Hirosawa, Wako 351-0198, Japan. E-mail: sitoehara@brain.riken.jp.

DOI:10.1523/JNEUROSCI.5756-08.2009

Copyright © 2009 Society for Neuroscience 0270-6474/09/295884-13\$15.00/0

Here, we performed studies to identify the neuronal mechanisms regulated by X11L and to determine whether X11L-KO mice could model Alzheimer's disease (AD). Thus, we extensively analyzed the behavioral phenotypes, and neurochemical and neurophysiological characteristics of X11L-KO mice. Although no deficits in learning, memory, or synaptic properties were detected in X11L-KO mice, we observed characteristic deficits in motivated approach behaviors associated with emotional conflict. Restoration of expression of X11L during development restored normal performance in the resident-intruder and ethological tests designed to observe species-typical behaviors to attractive stimuli, such as unfamiliar-objects/thick-bedding and a clogged-tunnel. We suggest that X11L-KO mice provide a unique opportunity to address the neuronal mechanisms that underlie emotional conflict.

Materials and Methods

Animals

All experimental protocols were approved by the Animal Care and Use Committees of the RIKEN Brain Science Institute. X11L-KO mice were previously generated as a C57BL/6 coisogenic strain (Sano et al., 2006a). A tTA transgenic mouse line, B6;CBA-Tg(Camk2a-tTA)1Mmay/J, under control of the *Camk2a* promoter (Mayford et al., 1996), was obtained from The Jackson Laboratory. The constructs of *FLAG-humanX11L* (*hX11L*) (Taru and Suzuki, 2004) and *NLS-LacZ* (Sassa et al., 2004) have been described previously. These coding sequences together with splicing signals derived from the pMSG vector (GE Healthcare) were inserted downstream from the bidirectional tet-response element (TRE)-dependent promoters of the pBI Tet vector (Clontech). The resulting *FLAG-X11L-TRE-NLS-LacZ* construct was microinjected into fertilized C57BL/6 eggs. The generated mouse lines were crossed with *Camk2a-tTA* mice. We selected a representative line that bidirectionally expressed *FLAG-X11L* and *NLS-LacZ* at the highest level depending on the presence or absence of doxycycline (DOX) (Sigma-Aldrich).

Experimental groups of wild-type (WT) and X11L-KO mice were generated by intercrossing X11L heterozygous mutant mice. To obtain *FLAG-X11L-TRE-NLS-LacZ/Camk2a-tTA* double-transgenic (DTg) mice under the X11L-KO background (DTg:X11L-KO mice), we first generated each single-transgenic mouse under the X11L-KO background and intercrossed them. X11L-KO littermates from the crosses were used for the behavioral studies. These mice theoretically carry 99% or more of the C57BL/6-derived genome. Genetic background-matched and age-matched WT mice were obtained independently from crosses between X11L-mutation heterozygotes of the same generation.

Adult male mice >3 months old were used. Mice were maintained under a 12 h light and dark cycle (lights on, 8:00 A.M.–8:00 P.M.), and *ad libitum* feeding and drinking conditions. Mice were group-housed by combining equal numbers of each genotype, unless otherwise stated. An investigator ignorant to the genotype of the animals performed all behavioral, pharmacological, and electrophysiological experiments.

Antibodies

The mouse monoclonal anti-X11L C-terminal domain (347–5) was described previously (Sumioka et al., 2003). Anti-Mint2/X11L (611033) was purchased from BD Biosciences; horseradish peroxidase (HRP)-conjugated anti- β -tubulin (s.c.-9104) was from Santa Cruz Biotechnology; anti-FLAG M2 (F1804) was from Sigma-Aldrich; anti-galactosidase (55976) was obtained from Cappel Laboratories; and HRP-conjugated anti-mouse IgG (NA931) and HRP-conjugated anti-rabbit IgG (NA934) were purchased from GE Healthcare.

Western blotting analysis

Western blotting was performed as described previously (Sano et al., 2006a,b). Details are described in the supplemental materials, available at www.jneurosci.org.

Behavioral analysis

Competitive food restriction. Groups of mice were cohoused in the same cage for at least a week before food restriction. In mixed genotype hous-

ing cages, equal numbers of WT and X11L-KO mice [(2 + 2) or (3 + 3) per cage] were cohoused. In uniform genotype housing cages, genotype-matched mice (4–6 mice/cage) were cohoused. During the food restriction period, mice were provided with 2 or 3 food pellets, equivalent to 4% of their free-feeding body weight, per day. During all test periods, mice were allowed water *ad libitum*. Weight was measured daily before feeding.

Food intake. To examine food intake of the mice under noncompetitive conditions, mice that were food-restricted as described above in mixed genotype housing cages were housed individually and then fed *ad libitum* for 90 min. The amount of food consumed and weight-gain after feeding was measured. Calculated food intake and weight-gain were normalized to the body weight measured before feeding.

Grip strength. A spring balance attached to the distal end of a wire-mesh end was used to test grip strength (O'Hara & Co.). Mice were allowed to grip the wire-mesh with their forelimbs, and were then pulled gently in a horizontal direction by their tails, away from the spring balance. The force applied to the spring balance at the moment the mice released the wire-mesh was recorded. Measurements were repeated twice with a 30 min intertrial interval. The mean of the two trials was calculated. In this test, mice were normally housed without food restriction.

Resident-intruder test. The resident-intruder test was performed according to the procedure described previously (Wada et al., 1997). Resident mice were maintained in individual housing at least for 5 weeks before experimentation. Corn cob (GreenTru; Green Products Company) was used for the bedding material to allow for easy observation of the mouse behavior. Intruder mice (male, DBA/2J strain; JCL Inc.; 7 weeks) were housed in groups of 4–5 mice per cage with standard bedding material (Tek-Fresh; Harlan) for 2 weeks before the experiment.

In the resident-intruder test, residents' cages were placed in an area with low lighting (120 lux). The intruder was transferred to the resident home cage, then social behaviors of the resident mouse toward the intruder mouse were observed for 5 min. The test was conducted twice (24 h intertrial interval) using different intruder mice for each trial. The following behavioral indices were measured. Approaching: exploratory behavior toward the intruder, sometimes involving contact with the intruder and sniffing behavior; Following: exploratory following and chasing of the intruder; and Aggression: aggressive movements against the intruder, such as fighting, biting, and other aggressive responses (aggressive-sniffing, -grooming, -following). As some indices may or may not involve aggression, behaviors were counted as aggressive only when the intruder responded with a submissive posture (Grant and Mackintosh, 1963). Behaviors of the resident mice were analyzed and counted using commercial software (VideoStudio, Corel) after the experiment.

Marble burying test and digging test. The marble burying test (MBT) was performed according to the procedure described previously (Yamada et al., 2002) with some modifications. Twenty clear blue glass marbles (17 mm diameter), white paper bedding material (Paperclean; Japan SLC), and transparent cages (24 × 45 × 21 [H] cm) were used for each trial. The cage was filled with bedding (5 cm deep). Mice were individually habituated to the cages for 30 min (habituation session) and then returned to their home cages. The 20 glass marbles were placed on the bedding surface in the habituated cage, evenly spaced at 5 cm intervals. The mice were then placed into the same cages to which they were habituated (MBT session) and allowed to remain for 20 min. After 20 min, the mice were removed from the cage and the number of marbles buried more than two-thirds in the bedding was counted. In both the habituation and MBT sessions, horizontal and vertical activity of the mice was measured using a two-level infrared beam apparatus (Scanet, Melquest). In the digging test, the mice were placed on the bedding (5 cm deep) without any marbles present. Mouse behavior was recorded from above for 30 min using a video camera, and the time spent digging was measured manually.

Burrowing test. The burrowing test was performed according to previously described procedures (Deacon et al., 2002) with some modifications. Both WT and X11L-KO mice were maintained in individual housing. Two gray vinyl chloride tubes (inside tube diameter, 56 mm; length, 150 mm; open end of the tube raised 30 mm; lower end closed with a cap)

were placed in their home cage at ~3 h before the start of dark period. One tube was filled with 200 g of the usual food pellets, each weighing ~2 g, while other tube was empty to provide a shelter. The weight of displaced food pellets from the tube was weighed after a 15 h period.

Methods for the other behavioral tasks are described in the supplemental materials, available at www.jneurosci.org.

β-Galactosidase (LacZ) staining

LacZ staining was performed according to previously described procedure (Iwasato et al., 2000; Tanaka et al., 2008) with some modifications. Mice were deeply anesthetized with 2,2,2-tribromoethanol (Avertin, Sigma-Aldrich) and transcardially perfused with physiological saline and then 4% (w/v) paraformaldehyde in 0.1 M sodium phosphate buffer (PB), pH 7.4, at 4°C for 20 min. The brains were excised, postfixed with the same fixative at 4°C for 2 h, and equilibrated in 30% (w/v) sucrose in PB as a cryoprotectant. The brains were embedded in OCT compound (Sakura Finetech), and frozen sections (20 μm) were prepared. Sections were then washed in phosphate buffered saline (PBS) on ice for 5 min and stained in 1 mg/ml X-gal, 5 mM K₃Fe(CN)₆, 5 mM K₄Fe(CN)₆, 20 mM Tris/HCl pH 7.5, and 2 mM MgCl₂ in PB at 37°C overnight. LacZ-stained sections were washed in PBS and then counter-stained with eosin.

Monoamine measurement

Circular tissue punches (1 mm diameter) were taken from 150 μm thick frozen coronal brain sections obtained from 4- to 5-month-old mice ($n = 8$ per group) and stored at -80°C until assayed. Monoamines and metabolites were extracted by sonication in 0.1 M perchloric acid containing a mixture of internal standards (200 nM isoproterenol, 200 μM L-norleucine, and 5 μM ethylhomocholine). Protein was then measured using a DC Protein Assay kit (Bio-Rad). The remainder of the sample was centrifuged for 15 min at 15,000 rpm and the supernatant filtered through polyvinylidene fluoride 0.22 μm micropore filters (Millipore Corp.) by centrifugation (14,000 rpm for 5 min). The filtrate was analyzed by high pressure liquid chromatography coupled to electrochemical detection (Eicom). Briefly, an Eicompak SC-50DS 3.0 × 150 mm column was used for separations, perfused with a mobile phase consisting of citrate (41.4 mM), sodium acetate (39.2 mM), methanol (17%), sodium 1-octanesulfonate (190 mg/L), and EDTA (5 mg/L), adjusted to pH 3.7 using glacial acetic acid and pumped at a rate of 0.5 ml/min. The working electrode (WE-3G) potential was set at +0.75 V. Column temperature was maintained at 25°C. Data were collected and analyzed by Ezchrom Elite software (Scientific Software Inc.). All analyte information, including the retention times, peak heights, concentrations, and recovery rate of internal standards, was calculated by comparison to standard curves generated for known concentrations of external standards that were run daily. Turnover rates were calculated as follows: noradrenaline (NA) turnover rate = [3-methoxy-4-hydroxyphenylglycol (MHPG)]/[NE]; dopamine (DA) turnover rate = [3,4-dihydroxyphenylacetic acid (DOPAC)] + [homovanillic acid (HVA)]/[DA]; serotonin (5-HT) turnover rate = [5-hydroxyindoleacetic acid (5-HIAA)]/[5-HT]. Correlation maps were constructed using iVici software (Michnick Laboratory, <http://michnick.bcm.umontreal.ca/ivici/>) using correlation coefficients calculated by pairwise comparisons (Tarassov and Michnick, 2005).

8-Hydroxy-2-dipropylaminotetralin-induced hypothermia

(±)-8-Hydroxy-2-dipropylaminotetralin (8-OH-DPAT) hydrobromide was purchased from Tocris Bioscience. The 8-OH-DPAT induced hypothermia experiment was performed according to the procedure described previously (Li et al., 1999) with minor modifications. Mice were handled every day for 1 week before the experiment. Mouse body temperature was measured using a digital thermometer with a temperature probe (Physitemp BAT-12R and RET-3) inserted 2–2.5 cm into the rectum, with the mouse slightly restrained by the tail. WT and X11L-KO mice were subcutaneously injected with saline or 8-OH-DPAT at doses of 0.1 or 0.5 mg/kg. Body temperature was monitored every 10 min for 60 min before the injection and every 15 min for 90 min after the injection. Mean temperature before injection was used as the basal temperature.

Statistics

Statistical analyses were performed using SPSS 14.0J or Excel statistics 2008 (Social Survey Research Information Co.) software. For comparison of two groups in most experiments, we applied an unpaired two-tailed *t* test. A repeated measures ANOVA was used to analyze learning data through training from the water maze and radial maze tests, and the 2 d of data from the resident-intruder test. We used a single-factor ANOVA followed by Scheffe's *post hoc* analysis to test for differences among three or more independent groups. Nonparametric tests were used to analyze data from the marble burying, digging and burrowing tests (Mann-Whitney *U* test for comparison of two groups or Steel test for rescue experiments). Differences were considered to be statistically significant when the probability value was <0.05.

Results

X11L-KO mice do not exhibit an AD-model like phenotype in terms of learning and memory

Amyloid β40 and 42 levels are increased in X11L-KO mice (Sano et al., 2006a). Patients with AD as well as AD mouse models that exhibit Aβ accumulation are cognitively impaired (Chen et al., 2000; Janus et al., 2000; Morgan et al., 2000). To determine whether X11L-KO mice have an AD-like phenotype, we analyzed their learning and memory ability. The Morris water maze task is a hippocampus-dependent task that is most frequently used to detect dysfunction of learning and memory performance in mouse models of AD. Despite the increase in Aβ in X11L-KO hippocampi, the learning and memory performance of X11L-KO and WT mice was comparable (supplemental Fig. 1A–C, available at www.jneurosci.org as supplemental material). Working memory in the 8-arm radial maze test and associative memory in context-dependent and cue-dependent fear conditioning tests were not impaired in X11L-KO mice compared with WT mice (supplemental Fig. 1D,E, available at www.jneurosci.org as supplemental material). These data suggest that learning and memory functions in X11L-KO mice are largely intact.

An accumulation of Aβ may depress neuronal transmission (Chapman et al., 1999; Kamenetz et al., 2003). Additionally, X11L interacts with the synaptic machinery that regulates neurotransmitter release such as Munc 18-1 and β-neurexin (Okamoto and Südhof, 1997; Biederer and Südhof, 2000; Ho et al., 2003, 2006). Expression of X11L is most abundant in the CA3 pyramidal neurons (Nakajima et al., 2001). Thus, we next examined the synaptic properties of the CA3-CA1 synapse in hippocampal slices from WT and X11L-KO mice. Basal synaptic transmission, paired-pulse facilitation (a form of short-term plasticity), and long-term potentiation in slices from X11L-KO mice were similar to those of WT littermates (supplemental Fig. 2A–C, available at www.jneurosci.org as supplemental material). In the hippocampus, no differences were detected between X11L-KO and WT mice in the amounts of X11/X11L binding, or of the levels of presynaptic and postsynaptic molecules examined (supplemental Table 1, available at www.jneurosci.org as supplemental material).

Therefore, the behavioral and physiological data do not support the tenet that X11L-KO mice are AD-like animal model associated with dementia.

X11L-KO mice are subordinated by their WT peers under competitive conditions

We noticed that X11L-KO mice showed a unique behavioral phenotype during a set of repeated radial maze tests. When food restriction was implemented by limiting the amount of time during which food was available (90 min free feeding/d), group housed X11L-KO and WT mice had a similar rate of body weight loss. When the food restriction regimen was transiently shifted to




Figure 1. X11L-KO mice are subordinated in competitive feeding conditions. *A–C*, The percentages of weight reduction by food deprivation for the mixed genotype housing condition (WT mice, black square; X11L-KO mice, red circle). Free-feeding body weight was defined as 100%. *A*, From day 1 to day 7, mice could freely feed for 90 min per day. At day 8 (arrow), only an amount of food equivalent to 8% of the sum of the mouse body weight per cage was provided. *B*, Food pellets (2 pellets/4 mice; amount of food = 4% of the sum of the mouse body weights per cage under free-feeding conditions) were placed into the cages from day 1 to 7. *C*, Food was restricted by a time-limited manner of free feeding for 90 min/d. *D*, The percentages of weight reduction by food deprivation are shown for 10 d under different housing conditions (WT mice in mixed genotype housing, black square; X11L-KO mice in mixed genotype housing, red circle; WT mice in uniform genotype housing, blue square; X11L-KO mice in uniform genotype housing, pink circle). Body weights in the *ad libitum* feeding condition were defined as 100%. *E*, Percentage body weight at day 10 is shown. In only the mixed genotype housing, X11L-KO mice lost significantly more weight than did WT mice. *F*, The percentage of food intake and weight gain during the 90 min free feeding after 11 d of competitive food restriction was the same between genotypes. Data represent mean \pm SE. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. (WT mice, black and blue columns; X11L-KO mice, red, and pink columns).

restricting the amount of food available on day 8 (amount of food = 8% of the summed body weight per cage), WT mice had a transiently larger weight gain than cohoused X11L-KO mice (7–8 months old, $n = 12$ per group; at day 9, $p < 0.01$) (Fig. 1*A*). This differential weight loss in X11L-KO mice was reproduced in another experiment. X11L-KO mice lost significantly more weight than cohoused WT mice under the amount-limited (competitive) feeding regimen (amount food = 4% of the summed body weight per cage per day; 3–4 months old, $n = 12$ per group; at day 7, $p < 0.001$) (Fig. 1*B*). Under the time-limited (no or very low-competitive) condition (90 min free feeding per day), there were no differences in body weight between genotypes (8–9 months old, $n = 12$ per group, $p = 0.68$) (Fig. 1*C*).

Based on these observations, we hypothesized that X11L-KO mice were subordinated under competitive feeding conditions. To test this hypothesis, we restricted feeding using two types of group housing conditions: equal numbers of WT and X11L-KO mice per cage (referred to as “mixed genotype

housing”) or groups of a single genotype mice per cage (referred to as “uniform genotype housing”). To avoid the influence of body weight at the start, the WT and X11L-KO mice in the mixed genotype housing were weight matched. During 10 d of feeding restriction using the limited-amount regimen, the X11L-KO mice in the mixed genotype housing cages lost significantly more weight than the WT mice (5–6 months old, $n = 6$ per group; weight ratio at day 10; WT, $82.9 \pm 1.14\%$; X11L-KO, $77.3 \pm 1.56\%$). No differences between genotypes were induced by the same treatment in uniform genotype housing (weight ratio at day 10; WT, $80.2 \pm 0.68\%$, $n = 5$; X11L-KO, $80.2 \pm 0.97\%$, $n = 6$). ANOVA and *Post hoc* Scheffe’s test revealed significant differences between genotypes only in the mixed genotype housing groups ($F_{(3,19)} = 4.05$, $p < 0.05$; Scheffe’s test, WT vs X11L-KO in mixed genotype housing, $p < 0.05$; in uniform genotype housing, $p = 1.00$) (Fig. 1*D,E*).

To rule out the possibility that the larger weight loss in X11L-KO mice was caused by a loss of appetite, the amount of food ingested and of weight gain after eating were measured under noncompetitive conditions in which each food-deprived mouse were fed individually. Food intake and weight gain in X11L-KO mice were indistinguishable from those of WT mice (3–4 months old, $n = 12$ per group; food intake, $p = 0.85$; weight gain, $p = 0.33$) (Fig. 1*F*). Lever-pressing behavior to obtain food pellets in an operant conditioning test, in which mice received a 20 mg pellet when they pressed the lever 1, 10, or 50 times, was not different between genotypes (data not shown). These data indicate that the weight loss in X11L-KO mice in the competitive condition was not likely caused by a decreased motivation to eat, or a reduction in the incentive value of the food.

We considered the possibility that X11L-KO mice were less able to access food when competing with WT mice due to a physical limitation, such as weaker muscle strength. We assessed muscle (grip) strength of the forelimbs of WT and X11L-KO mice, and observed no differences between genotypes (7–8 months old, $n = 24$ per group; WT, 0.71 ± 0.03 Newtons (N); X11L-KO, 0.80 ± 0.04 N; grip strength, $p = 0.08$). Moreover, neither X11L-KO nor WT mice had wounds consistent with aggressive behavior in the home cage. Together, these findings suggested that X11L-KO mice are subordinated only under highly competitive conditions, and that this response is not due to a physical factor.

X11L-KO mice exhibit reduced social interaction

Because the subordinate feeding behaviors of the X11L-KO mice were observed under competitive conditions, we hypothesized that X11L-KO might show aberrant social interaction. To assess social behavior in X11L-KO mice, we used the resident-intruder test. We measured nonaggressive approach and following, and aggressive behaviors against intruder mice (see experimental procedure for details). In our experiment, fighting behavior was rarely observed. Approaching and following behaviors were lower in resident X11L-KO mice compared with WT mice (5–6 months old, $n = 10$ per group; approach, $F_{(1,18)} = 4.27$, $p = 0.05$; following, $F_{(1,18)} = 4.43$, $p < 0.05$; aggression, $F_{(1,18)} = 1.59$, $p = 0.22$; Figure 2*A–C*). The total number of resident social interactions with intruder mice was significantly decreased in X11L-KO mice compared with WT mice ($F_{(1,18)} = 9.14$, $p < 0.01$) (Fig. 2*D*). These data indicate that social interaction is suppressed in X11L-KO mice.




Figure 2. Social interaction against an intruder is significantly decreased in X11L-KO mice. *A–C*, The resident-intruder test (WT mice, black square; X11L-KO mice, red circle) was performed for 2 consecutive days using different intruder mice for each trial. The numbers of approaches (*A*) and following behaviors (*B*) of resident mice against intruder mice were lower in X11L-KO mice. *C*, Differences in the amount of time spent by resident mice exhibiting aggressive behavior in which the intruder mice showed a typical submissive pose was not statistically significant between genotypes. *D*, The summed number of social interactions of resident mice toward intruder mice was significantly decreased in X11L-KO mice compared with WT mice. Data represent mean \pm SE. * $p < 0.05$; ** $p < 0.01$.

X11L-KO mice do not show altered anxiety-like behavior in conventional behavioral tests

Social interaction is the result of an inherent conflict in approach-avoidance factors, and is decreased under anxiogenic conditions (File, 1992; Crawley, 2007). Even if anxiety in the social interaction test qualitatively differs from that assessed on the elevated plus maze test, anxiolytic drugs are effective in both models (File, 1992; Nakamura and Kurasawa, 2001; Crawley, 2007). To gain insight into anxiety-related behaviors in X11L-KO mice, we subjected X11L-KO mice to open field, elevated plus maze, and light-dark transition tests. The behavior of the X11L-KO mice was comparable to that of WT mice in all of these tests (supplemental

Table 2, available at www.jneurosci.org as supplemental material). Furthermore, X11L-KO mice did not show abnormalities in model behavioral tests for mental disorders such as the tail suspension test (a model of depression-like behavior) and prepulse inhibition test (a model for sensory motor gating activities associated with schizophrenia; supplemental Table 2, available at www.jneurosci.org as supplemental material). X11L-KO and WT mice showed comparable locomotor activity and circadian rhythms (supplemental Table 2, available at www.jneurosci.org as supplemental material). These results suggest that suppressed social interaction is not caused by enhanced anxiety-like, depressive behavior, or decreased locomotor activity.

X11L-KO mice show attenuated marble burying, digging, and burrowing behavior

The behavioral paradigms described above identify anxiety-related phenotypes based on avoidance behavior elicited by aversive stimuli. We examined whether behaviors that reflect approach motivation to some stimuli are selectively altered in X11L-KO mice. Behavior of X11L-KO mice and WT mice was therefore evaluated in the MBT. The MBT is proposed to be a model of defensive behavior, as mice actively bury glass marbles under a layer of bedding material (Broekkamp et al., 1986; Nicolas et al., 2006) (see also below). X11L-KO mice buried significantly fewer marbles than did WT mice (4–5 months old, $n = 12$ per group; WT, 11.2 ± 1.1 ; X11L-KO mice, 1.7 ± 0.76 ; number of buried marbles, $p < 0.001$) (Fig. 3*A*). Horizontal and vertical activity of X11L-KO mice was indistinguishable from that of WT mice in both the habituation and MBT sessions (locomotor activity in habituation session, $p = 0.75$; locomotor activity in MBT session, $p = 0.97$; rearing in habituating session, $p = 0.40$; rearing in MBT session, $p = 0.88$) (Fig. 3*B, C*). These data indicate that impaired marble burying behavior was not due to a reduction in general activity.

Marble burying behavior is argued to be the result of innate digging behavior (Njung'e and Handley, 1991a; Gyertyán, 1995; Deacon, 2006b). Thick bedding, which is used for the MBT, strongly motivates mice to dig the bedding material. This innate behavior is not habituated, is evoked in the absence of glass marbles, and continues even after all marbles are buried (Webster et al., 1981; Dudek et al., 1983; Gyertyán, 1995; Masuda et al., 2000; Deacon, 2006b). To assess whether impaired marble burying is associated with digging behavior, we measured the time spent digging during the habituation session. X11L-KO mice spent significantly less time digging (4–5 months old, $n = 12$ per group; WT, 288.9 ± 34.3 s; X11L-KO, 90.2 ± 15.3 s; $p < 0.001$) (Fig. 3*D*). Digging behavior was initiated within the first 10 min under conditions of a novel environment with thick bedding materials (supplemental Fig. 3*A*, available at www.jneurosci.org as supplemental material). In an independent experiment, strong and compulsive digging behavior was induced by bedding materials in the absence of marbles, depending on the bedding thickness (supplemental Fig. 3*B*, available at www.jneurosci.org as supplemental material). In this condition, X11L-KO mice again exhibited significantly reduced digging behavior. These data suggest that marble burying and digging behaviors are closely associated, and that the approach behavior motivated by incentives, such as marbles and bedding materials, is attenuated in X11L-KO mice.

Burying and digging behaviors are species-typical behaviors to some rodents. The burrowing test is conceptually related to the above ethological tests, as mice typically burrow




Figure 3. Marble burying and digging behaviors are impaired in X11L-KO mice. *A*, Typical images of the cage after the marble burying test are shown. *B*, *C*, Horizontal (*B*) and vertical (*C*) activity in the habituation and MBT sessions was not significantly different in X11L-KO mice compared with WT mice. *D*, The number of marbles buried and time spent digging during the habituation session (30 min) are plotted for each mouse. Data represent mean \pm SE. *** $p < 0.001$ (WT mice, black column; X11L-KO mice, red column).

into tunnel-like containers filled with small-solid objects such as soils, small clay balls and food pellets (Deacon, 2006a). X11L-KO mice displaced significantly less food pellets from a tube (4–5 months old, $n = 14$ per group; WT, 80.6 ± 15.4 g; X11L-KO mice, 10.1 ± 1.9 g; weight of displaced pellets, $p < 0.001$). These data indicate that X11L plays a crucial role in these species-typical behaviors, which likely reflect defensive behaviors, seeking safety, exploration and hoarding food seen in the wild (Dudek et al., 1983; Deacon, 2006a,b), and further suggest X11L-KO mice are less attracted to what are usually attractive stimuli.

Impaired species-typical behaviors and reduced social interaction in X11L-KO mice are rescued by the expression of X11L during development

To confirm the causal link between X11L and a specific behavior impaired in X11L-KO mice, we performed a rescue experiment using the Tet-regulated system, which is used for conditional expression in mice (Mayford et al., 1996). In normal WT mice, X11L is highly expressed in the forebrain (Nakajima et al., 2001). Therefore, *Camk2a-tTA* mice (Mayford et al., 1996; Gross et al., 2002) were used to express X11L selectively in the forebrain after birth. In the *TRE-hX11L:Camk2tTA* DTg mice (Fig. 4*A*), FLAG-hX11L and β -galactosidase are expected to be coexpressed in the same cells. Western blot analyses using antibodies against the C-terminal epitope of X11L and FLAG successfully detected the expression of FLAG-hX11L in adult DTg brains (Fig. 4*B*). LacZ staining revealed that β -galactosidase was strongly expressed in the forebrain region, as well as in some other regions, such as the substantia nigra, ventral tegmental area, and raphe (Fig. 4*C*). Western blot analysis revealed that X11L protein expression was completely depleted by administering 2 mg/ml DOX in the drinking water for 3 weeks, and then recovered to $\sim 80\%$ of the initial expression level within 1 week after the withdrawal of DOX (Fig. 4*D*).

To perform a rescue experiment using this system, we generated DTg mice under the X11L-KO background (DTg:X11L-KO mice). We first focused on the MBT. DTg:X11L-KO mice buried significantly more marbles than X11L-KO mice (4–5 months old, WT, 8.1 ± 1.2 , $n = 21$; X11L-KO, 2.7 ± 0.7 , $n = 27$; DTg:X11L-KO, 5.6 ± 1.0 , $n = 29$; X11L-KO vs DTg:X11L-KO, $p < 0.05$) (Fig. 4*G*). The increased burying behavior in DTg:X11L-KO mice was not likely due to an increase in general activity as the horizontal activity of DTg:X11L-KO mice was slightly but significantly decreased compared with X11L-KO mice in the habituation session and indistinguishable from X11L-KO mice in the MBT session (habituation session, X11L-KO vs DTg:X11L-KO, $p < 0.05$; MBT session, X11L-KO vs DTg:X11L-KO; $p = 0.53$) (Fig. 4*J*).

To test whether X11L protein regulates marble burying behavior during information processing, the expression of FLAG-hX11L was acutely inhibited in DTg:X11L-KO mice in the adult stage by administering DOX (DTg:X11L-KO/DOXad). FLAG-hX11L proteins were not detected in DTg:X11L-KO/DOXad mice administered DOX in their drinking water (2 mg/ml) for 3 weeks (Fig. 4*E*). The DTg:X11L-KO/DOXad mice, however, still buried significantly more marbles compared with X11L-KO mice (4–5 months old, WT/DOXad, 9.4 ± 1.4 , $n = 21$, X11L-KO/DOXad, 3.0 ± 0.7 , $n = 27$; DTg:X11L-KO/DOXad, 6.8 ± 1.0 , $n = 30$; X11L-KO/DOXad vs DTg:X11L-KO/DOXad, $p < 0.01$) (Fig. 4*H*). The recovery of burying behavior was not associated with an increase in general activity (Fig. 4*K*). These data suggest that the X11L protein is required during the developmental stages but not in adults, for normal marble burying behavior. The partial recovery may indicate incomplete expression and/or a cell type specificity of X11L in DTg mice.

To further confirm the role of X11L in developmental stages, we administered DOX (25 μ g/ml) in the drinking water of the parents from the mating period through postnatal day 28. This concentration of DOX was sufficient to completely suppress the expression of FLAG-hX11L in DTg:X11L-KO mice (DTg:X11L-KO/DOXdev) (Fig. 4*F*). The expression of FLAG-hX11L did not recover after DOX withdrawal, even after 5 months, in DTg:X11L-KO/DOXdev mice, unlike the case of DOX administration in adults (Fig. 4*F*). The transgenic locus silenced during the developmental stage likely acquires irreversible modifications. The numbers of marbles buried by DTg:X11L-KO/DOXdev mice and




Figure 4. Impairment in marble burying behavior in X11L-KO mice was rescued by the expression of X11L during development, but not in adults. *A*, The tTA protein is driven by *Camk2a* promoter. The active form of tTA binds to TRE and bidirectionally drives the expression of the *FLAG-hX11L* and *MLS-LacZ* genes. DOX inhibits the expression of both transgenes by binding to tTA and converting it into an inactive conformation. *B*, Immunoblot of whole brain lysates obtained from transgenic and DTg mice. Anti-X11L N antibody only recognizes the N-terminal region of the mouse X11L protein. Anti-X11L C antibody recognizes the C-terminal region of both human and mouse X11L proteins. *C*, Expression of the transgenes was detected by LacZ staining. Scale bars, 500 μ m. HyTh, hypothalamus; SNC, substantia nigra pars compacta; VTA, ventral tegmental area; DR, dorsal raphe; MR, median raphe; LC, locus ceruleus. *D*, Immunoblot of whole brain lysates obtained from DTg mice with administration and withdrawal of 2 mg/ml DOX in the drinking water. *E*, Immunoblot of whole brain lysates obtained from DTg:X11L-KO and control mice. Complete depletion of X11L in DTg:X11L-KO mice by 2 mg/ml DOX administration was confirmed. *F*, Immunoblot of whole brain lysate obtained from DTg:X11L-KO mice after withdrawal from 25 μ g/ml DOX administration during the developmental period. *D–F*, Experimental design and results are schematically represented at the bottom part of each panel. Blue lines indicate X11L protein level. *G–I*, MBT and concurrent measurement of locomotor activity were performed using DTg:X11L-KO mice with the various DOX administration paradigms. *G–I*, MBT. The number of marbles buried was counted for each genotype of experimental group: without DOX administration (*G*), with DOX administration during the adult period (*H*), and with DOX administration in the developmental period (*I*) with subsequent withdrawal of DOX administration. *J–L*, Locomotor activity in MBT and habituating sessions for each experimental group without DOX administration (*J*), the same as *G*, with DOX in the adult period (*K*), the same as *H*, and with DOX in the developmental period (*L*), the same as *I*. Data represent mean \pm SE. * p < 0.05, ** p < 0.01, *** p < 0.001 (WT mice, black column; DTg:X11L-KO mice, brown column; X11L-KO mice, red column).

X11L-KO/DOXdev mice were indistinguishable (4–5 months old, WT/DOXdev, 10.0 ± 1.4 , $n = 16$; X11L-KO/DOXdev, 2.3 ± 1.0 , $n = 16$; DTg;X11L-KO/DOXdev, 2.5 ± 1.1 , $n = 15$; X11L-KO/DOXdev vs DTg;X11L-KO/DOXdev, $p = 0.77$) (Fig. 4I). These results further strengthened the notion that X11L has an important role during development.

Locomotor activity in DTg;X11L-KO mice treated with or without DOX was significantly lower than that in X11L-KO mice (Fig. 4J–L). Decreased locomotor activity in *Camk2a-tTA* mice was reported by another group (McKinney et al., 2008). This decreased locomotor activity might lead to an underestimation of the degree of the rescued phenotype in our experiments.

Next, we examined the impact of X11L expression on burrowing behavior. We also observed that DTg;X11L-KO mice displaced significantly more food pellets from a tube than X11L-KO mice (5–6 months old, DTg;X11L-KO, 144.7 ± 11.6 g, $n = 14$; X11L-KO, 30.5 ± 9.6 g, $n = 16$; weight of displaced pellets, $p < 0.001$).

Finally, we performed the resident-intruder test using DTg;X11L-KO and X11L-KO mice to confirm whether the expression of X11L can also restore the suppressed social interaction in X11L-KO mice as is the case in the MBT. Resident social interactions with intruder mice were significantly increased in DTg;X11L-KO mice compared with X11L-KO mice (4–5 months old, DTg;X11L-KO $n = 14$, X11L-KO $n = 16$; approach, $F_{(1,28)} = 5.86$, $p < 0.05$; following, $F_{(1,28)} = 4.32$, $p < 0.05$; aggression, $F_{(1,28)} = 0.22$, $p = 0.64$; social interaction, $F_{(1,28)} = 6.86$, $p < 0.05$; Figure 5A–D). These data indicate that X11L protein is required not only for normal burying/burrowing behavior but also for normal social behavior, and suggest that social behavior and species-typical behavior may share some common neuronal mechanisms associated with X11L protein.

Monoamine systems are disrupted in X11L-KO mice

Marble burying behavior is affected by anxiolytic drugs, particularly 5-HT agents. Drugs acting on the 5-HT system, such as 5-HT reuptake inhibitors, which increase 5-HT levels at release sites, and some 5-HT agonists, inhibit marble burying behavior without affecting locomotor activity (Njung'e and Handley, 1991b; Shinomiya et al., 2005; Nicolas et al., 2006; Bruins Slot et al., 2008). Some antipsychotics also inhibit marble burying behavior (Broekkamp et al., 1986; Nicolas et al., 2006; Bruins Slot et al., 2008). These reports suggest that monoaminergic activity underlie marble burying behavior. Therefore, we measured the amounts of monoamines and their metabolites in selected brain regions of X11L-KO mice.

Biopsies of the following seven regions were obtained from frozen brain sections: anterior cingulate cortex, striatum, hypothalamus (mainly containing the paraventricular nucleus), somatosensory cortex (mainly containing the barrel field), amygdala, hippocampus (mainly containing the CA3 region), and dorsal and median raphe (Fig. 6A). These areas were selected on the basis of high X11L expression, or their known involvement in the behaviors studied here. Although the levels of most monoamines and their metabolites in the X11L-KO mice brains were not significantly different from those in WT mice, 5-HT levels were significantly higher in the hypothalamus and somatosensory cortex (Table 1). Because monoamines are synthesized and broken down in sophisticated biochemical networks, we sought evidence of an altered flux through these networks by analyzing the correlations among monoamines and their metabolites. The correlational relationships in the striatum, hippocampus, and somatosensory cortex of X11L-KO mice were substantially different from those in WT mice, which were reflected in an asym-




Figure 5. Reduced social interaction in X11L-KO mice was restored by the expression of the X11L protein. The resident-intruder test (DTg;X11L-KO mice, brown square; X11L-KO mice, red circle) was performed for 2 consecutive days using different intruder mice for each trial. *A, B*, The numbers of approaches (*A*) and following behaviors (*B*) of resident mice against intruder mice were significantly higher in DTg;X11L-KO mice. *C*, Differences in the amount of time spent by resident mice exhibiting aggressive behavior was not statistically significant between genotypes. *D*, The summed number of social interactions of resident mice toward intruder mice was significantly increased in DTg;X11L-KO mice compared with X11L-KO mice. Data represent mean \pm SE. * $p < 0.05$.

metrical pattern in the correlation maps (Fig. 6B). In contrast, these relationships were highly comparable in the raphe and amygdala (Fig. 6B). These results suggest that X11L has an active role in regulating the synthesis and/or breakdown of monoamine neurotransmitters.

To gain insight into the function of the serotonergic system in X11L-KO mice, we analyzed the hypothermic responses induced by the selective 5-HT_{1A} receptor agonist 8-OH-DPAT, which reflects somatodendritic autoreceptor function in raphe neurons (Goodwin et al., 1985; Bill et al., 1991; Martin et al., 1992; Gross et




Figure 6. Balance of monoamine systems is disturbed in X11L-KO mice. *A*, Representative photomicrographs of Nissl-stained brain sections showing locations of biopsies used for neurochemical analyses. *B*, Correlation maps showing relationships between tissue contents of monoamines, metabolites, and calculated turnover values. Red indicates positive correlation. Green indicates negative correlation. *C*, Time course and dose–response of 8-OH-DPAT-induced hypothermia in X11L-KO and WT mice. Data represent mean \pm SE. * p < 0.05 (WT mice, black square; X11L-KO mice, red circle).

al., 2002). Stimulation of 5-HT_{1A} autoreceptors attenuates 5-HT release and induces a hypothermic response. The body temperature of WT mice decreased after 8-OH-DPAT administration (Fig. 6C). This hypothermic response was significantly attenuated in X11L-KO mice 15 and 30 min after injection with 0.5 mg/kg 8-OH-DPAT (6–7 months old, WT and X11L-KO mice, n = 8 per group; at 15 and 30 min of 0.5 mg/kg dose, p < 0.05)

(Fig. 6C), suggesting that 5-HT_{1A} receptor function or its downstream signaling is attenuated in X11L-KO mice.

Discussion

The main finding of the present study is that X11L-KO mice show a selective deficit in motivated approach behavior, but not in motivated avoidance behavior. That is, X11L-KO mice showed

Table 1. The amount of monoamines and metabolites

Monoamines and metabolites	Genotypes	Anterior cingulate cortex	Striatum	Hypothalamus	Somatosensory cortex	Amygdala	Hippocampus	Raphe
5-HT	WT	1681 ± 144	2444 ± 165	4039 ± 170	917 ± 45	5403 ± 342	3566 ± 318	9779 ± 697
	KO	1491 ± 97	2709 ± 61	5021 ± 245**	1130 ± 78*	4819 ± 287	3464 ± 188	9069 ± 908
5-HIAA	WT	2300 ± 219	2500 ± 221	3939 ± 221	758 ± 23	3663 ± 235	4661 ± 381	11,438 ± 933
	KO	2148 ± 64	2620 ± 79	3964 ± 212	759 ± 54	3247 ± 209	4121 ± 211	10,006 ± 855
NA	WT	3420 ± 214	73 ± 11	13,750 ± 607	2275 ± 78	3539 ± 188	2713 ± 217	5644 ± 493
	KO	3129 ± 134	117 ± 27	15,455 ± 654	2562 ± 150	3142 ± 141	2584 ± 93	5988 ± 619
MHPG	WT	513 ± 40	29 ± 7	577 ± 67	169 ± 7	573 ± 39	277 ± 28	1042 ± 78
	KO	449 ± 37	42 ± 8	553 ± 51	168 ± 8	452 ± 22*	245 ± 18	990 ± 118
DA	WT	460 ± 37	10,9843 ± 8939	2431 ± 133	90 ± 4	5345 ± 451	219 ± 30	1156 ± 99
	KO	394 ± 21	12,0274 ± 3273	2605 ± 157	94 ± 9	4227 ± 401	189 ± 12	1138 ± 117
DOPAC	WT	502 ± 46	18,005 ± 2298	854 ± 49	146 ± 12	2040 ± 152	265 ± 29	639 ± 41
	KO	420 ± 22	21,580 ± 1520	826 ± 28	141 ± 11	1654 ± 186	237 ± 14	594 ± 52
HVA	WT	1113 ± 148	14,317 ± 1178	1293 ± 97	192 ± 17	2145 ± 169	289 ± 20	1300 ± 70
	KO	889 ± 63	14,513 ± 312	1215 ± 65	153 ± 17	1647 ± 198	273 ± 21	1233 ± 111
5-HT Turnover	WT	1.27 ± 0.07	0.94 ± 0.05	0.90 ± 0.05	0.77 ± 0.04	0.63 ± 0.03	1.22 ± 0.07	1.08 ± 0.04
	KO	1.37 ± 0.10	0.89 ± 0.03	0.74 ± 0.05*	0.63 ± 0.06	0.62 ± 0.03	1.12 ± 0.08	1.03 ± 0.04
NE Turnover	WT	0.14 ± 0.01	0.36 ± 0.06	0.04 ± 0.004	0.07 ± 0.003	0.15 ± 0.004	0.09 ± 0.003	0.17 ± 0.01
	KO	0.13 ± 0.01	0.36 ± 0.04	0.03 ± 0.003	0.06 ± 0.001*	0.13 ± 0.01	0.09 ± 0.005	0.16 ± 0.01
DA Turnover	WT	3.16 ± 0.53	0.26 ± 0.02	0.77 ± 0.03	3.30 ± 0.29	0.69 ± 0.02	2.36 ± 0.18	1.47 ± 0.05
	KO	2.92 ± 0.20	0.27 ± 0.02	0.69 ± 0.04	2.82 ± 0.20	0.69 ± 0.06	2.44 ± 0.24	1.44 ± 0.10

Data shown are mean ± SE (pg/mg protein).

significantly attenuated ethological responses to attractive stimuli and withdrew during approach-avoidance conflict, such as that encountered during competitive feeding conditions and social interaction. Additionally, attenuated marble-burying, burrowing and reduced social behavior in X11L-KO mice was rescued by expression of X11L protein during development. Together, these findings suggest that X11L is involved in the maturation of neuronal circuits regulating conflict resolution-related behaviors. We propose that X11L-KO mice provide a novel model for analyzing emotional behavior, which differs from the existing models of generalized anxiety-like and depression-like behaviors (Finn et al., 2003; Cryan and Mombereau, 2004; Crawley, 2007).

X11L-KO mice lost more body weight than WT mice, specifically under a competitive feeding regimen with WT mice. This was unlikely due to physical defects or changes in general activity, but is likely related to the suppression of a social interaction-induced behavioral response. The results from the resident-intruder test support this notion: That is, X11L-KO mice exhibited reduced exploratory behavior toward intruder mice. The social behavior of one mouse toward another mouse probably reflects the net result of conflicts from coexisting incentives, including those that might induce approach (e.g., investigation of intruder mice and approach to housemates that have a food pellet) and those that might induce avoidance (e.g., threat from novel intruders, avoiding competition with housemates, and generalized anxiety). Anxiety-like behaviors, which are often expressed as avoidance, are thought to be closely associated with decreased social interaction (File, 1992; Sankoorikal et al., 2006; Crawley, 2007; Moy et al., 2007). Interestingly, X11L-KO mice did not show anxiety-like behavior in standard behavioral tests, such as the elevated plus maze, light and dark transition, and open field tests. Thus, low levels of social interaction in X11L-KO mice appear to be independent of “traditional” anxiety-like behaviors that are expressed as avoidance of aversive stimuli.

X11L-KO mice showed prominent defective ethological stimuli-responses, such as reduced burying, digging and burrowing behaviors, whereas they showed no phenotypes in many of the most common behavioral tests (e.g., locomotor activity, cir-

cadian rhythm, open field, elevated plus maze, light and dark transition, tail suspension, prepulse inhibition and some learning/memory tests) commonly applied in the study of behavioral genetics. These data highlight the importance of testing species-typical behaviors to better understand brain functions. So-called ethological tests likely have strong potential to extract brain function and mechanisms. Burrowing behavior is highly sensitive to dysfunctions of the hippocampus induced by cytotoxic lesion, scrapie infection and in AD model mice (Deacon et al., 2001, 2002, 2008). Marble burying behavior is correlated very well with activity in the serotonin system and is proposed to have predictive validity as a model of obsessive-compulsive disorder (OCD)-like behavior as drugs used to treat OCD inhibit marble burying behavior (Njung'e and Handley, 1991b; Shinomiya et al., 2005; Nicolas et al., 2006; Bruins Slot et al., 2008; Kobayashi et al., 2008). Thus, species-typical ethological responses may provide the most sensitive parameters to estimate the therapeutic effect using animal models.

Transgene rescue experiments using the *Camk2a-tTA* mouse line suggested that X11L is involved in the development of neuronal circuits regulating emotional and conflict-related behaviors. We found that monoamine concentrations in forebrain regions of X11L-KO mice were imbalanced. Monoamine signaling is critically involved in developing neuronal circuits that modulate emotion in adults (Reif and Lesch, 2003). The development of 5-HT function in the postnatal period determines adult emotional behavior (Andersen et al., 2002; Gross et al., 2002; Ansorge et al., 2004, 2008; Lo Iacono and Gross, 2008). Catecholamine levels during early life also influence adult emotional behavior (Estelles et al., 2007; Gray et al., 2007). Maturation and/or refinement of forebrain circuits during postnatal weeks 2 through 5 via somatodendritic 5HT1A receptor interactions is considered critical for programming anxiety-related behaviors (Gross et al., 2002; Lo Iacono and Gross, 2008). X11L expression begins during development, is strongly localized to the somatodendritic compartment of forebrain region (Nakajima et al., 2001; Ho et al., 2003), and binds many molecules involved in regulating neuronal activity. Therefore, X11L may function as a neuronal media-

tor in the development of emotional circuits via molecules relating to monoaminergic systems.

Alternatively, or additionally, the behavioral deficits observed in X11L-KO mice in adulthood may reflect ongoing neurochemical perturbations. In the present study, the statistically strongest neurochemical difference was increased 5-HT content in the hypothalamus. Previous studies show that decreased hypothalamic 5-HT levels are associated with increased aggression in rats [Vergnes et al., (1988); see Popova (2006) for discussion of the role of 5-HT in aggression]. By the same token, the subordinate behavior of X11L-KO mice reported here may relate to increased 5-HT levels and/or altered 5-HT metabolism. It should be noted, however, that monoamine turnover rates were marginally lower in X11L-KO mice in some regions studied, in addition to monoamine-metabolite imbalances being evident. Thus, the possibility that other neurochemical changes underlie the behavioral deficits observed here cannot be ruled out. There is undoubtedly a close relationship between activity in monoaminergic systems, as exemplified by studies of 5-HT_{1A} and 5-HT_{1B} receptor KO mice, which show increases in both 5-HT and dopamine turnover (Ase et al., 2000). Indeed, all three monoaminergic systems are targeted in the treatment of anxiety-related disorders and social phobias in the clinical setting, and so much further research is necessary to elucidate the detailed interactions between these systems that bring about the sophisticated behaviors studied here.

How X11L affects monoaminergic systems remains to be determined. 8-OH-DPAT (5-HT_{1A} receptor specific antagonist)-induced hypothermia was attenuated in the X11L-KO mice. The 5-HT_{1A} receptor is a G_i-type G-protein coupled receptor. Its activation opens G-protein-coupled inwardly rectifying K⁺ channels (GIRKs) by binding liberated G_{βγ} proteins from 5-HT_{1A} receptor, resulting in cell hyperpolarization and inhibition of 5-HT release (Andrade and Nicoll, 1987; Williams et al., 1988; Lüscher et al., 1997). Functional coupling of 5-HT_{1A} receptors to GIRKs does not occur until postnatal day 14 (Béique et al., 2004). GIRKs are also downstream regulators of several other G-protein coupled receptors such as dopaminergic (D₂), adrenergic (α₂), and muscarinic (M₂) autoreceptors, and they act to regulate cellular excitability and coupling (North, 1989; Hille, 1992; Yamada et al., 1998). GIRKs containing the GIRK2 subunit are control hubs for the hypothermic response induced by various G-protein coupled receptor agonists (Costa et al., 2005). The GIRK2 subunit is widely distributed throughout the brain and strongly localized to the somatodendritic compartment (Murer et al., 1997; Schein et al., 1998; Inanobe et al., 1999; Saenz del Burgo et al., 2008). Additionally, GIRK3 and GIRK2c, which is a splice variant of GIRK2, subunits have a PDZ-binding motif (-ESKV) at their C-terminal and their trafficking is regulated by PDZ-containing adaptor proteins (Inanobe et al., 1999; Ma et al., 2002; Lunn et al., 2007). In contrast, X11L regulates the intracellular localization of its binding protein and also has coat protein properties (Hill et al., 2003; Sano et al., 2006a). Therefore, it is possible that X11L regulates the intracellular localization of GIRK2, and the functional attenuation of GIRKs, by ultimately altering neuronal excitability, may disturb monoamine balance and reduce the hypothermic response in X11L-KO mice. Indeed, we have observed specific binding between X11L and GIRK2c by coimmunoprecipitation analyses using a cell culture system (YS, HS, and SI, unpublished data).

Other mechanisms may underlie the role of X11L in neurochemical activity and behavior. X11L regulates APP metabolism (Sano et al., 2006a) and interacts with many other types of neu-

ronal/synaptic machinery (Okamoto and Südhof, 1997; Tomita et al., 1999; Biederer and Südhof, 2000; Gotthardt et al., 2000; Lau et al., 2000; Kitano et al., 2002; Araki et al., 2003; Hill et al., 2003; Kimura et al., 2004). The unique behavioral abnormalities in X11L-KO mice may be due to the combined dysfunction of interacting molecules and the compensatory effect of X11, which has binding partners and cellular functions similar to those of X11L (Ho et al., 2006; Miller et al., 2006; Rogelj et al., 2006; Sano et al., 2006a). Interestingly, compared with WT mice, X11-KO mice exhibit a vigorous and rapid escape response when the sides and body of the animal are stroked (Mori et al., 2002). Comparative studies using these mice and mice with conditional double mutations in limited subsets of neurons might help to better understand the mechanisms that underlie emotional control.

Finally, the X11L gene, namely *APBA2*, is located at the 15q11-q12 locus. The close association between abnormalities in the 15q11-q13 region and autism is frequently reported (Cook et al., 1997, 1998; Gillberg, 1998; Schroer et al., 1998; Shao et al., 2002; Muhle et al., 2004). Autism is a neurodevelopmental disorder characterized by impaired social interaction and communication. It is possible that X11L dysfunction during the developmental stage relates to the pathogenesis of developmental disorders with mental retardation such as autism, and X11L may have an important role in the development of emotionality and sociability in humans.

References

- Andersen SL, Dumont NL, Teicher MH (2002) Differences in behavior and monoamine laterality following neonatal clomipramine treatment. *Dev Psychobiol* 41:50–57.
- Andrade R, Nicoll RA (1987) Pharmacologically distinct actions of serotonin on single pyramidal neurones of the rat hippocampus recorded in vitro. *J Physiol* 394:99–124.
- Ansorge MS, Zhou M, Lira A, Hen R, Gingrich JA (2004) Early-life blockade of the 5-HT transporter alters emotional behavior in adult mice. *Science* 306:879–881.
- Ansorge MS, Morelli E, Gingrich JA (2008) Inhibition of serotonin but not norepinephrine transport during development produces delayed, persistent perturbations of emotional behaviors in mice. *J Neurosci* 28:199–207.
- Araki Y, Tomita S, Yamaguchi H, Miyagi N, Sumioka A, Kirino Y, Suzuki T (2003) Novel cadherin-related membrane proteins, Alcadeins, enhance the X11-like protein-mediated stabilization of amyloid beta-protein precursor metabolism. *J Biol Chem* 278:49448–49458.
- Ase AR, Reader TA, Hen R, Riad M, Descarries L (2000) Altered serotonin and dopamine metabolism in the CNS of serotonin 5-HT_{1A} or 5-HT_{1B} receptor knockout mice. *J Neurochem* 75:2415–2426.
- Béique JC, Campbell B, Perring P, Hamblin MW, Walker P, Mladenovic L, Andrade R (2004) Serotonergic regulation of membrane potential in developing rat prefrontal cortex: coordinated expression of 5-hydroxytryptamine (5-HT_{1A}), 5-HT_{2A}, and 5-HT₇ receptors. *J Neurosci* 24:4807–4817.
- Biederer T, Südhof TC (2000) Mints as adaptors. Direct binding to neuroligins and recruitment of munc18. *J Biol Chem* 275:39803–39806.
- Bill DJ, Knight M, Forster EA, Fletcher A (1991) Direct evidence for an important species difference in the mechanism of 8-OH-DPAT-induced hypothermia. *Br J Pharmacol* 103:1857–1864.
- Broekkamp CL, Rijk HW, Joly-Gelouin D, Lloyd KL (1986) Major tranquilizers can be distinguished from minor tranquilizers on the basis of effects on marble burying and swim-induced grooming in mice. *Eur J Pharmacol* 126:223–229.
- Bruins Slot LA, Bardin L, Auclair AL, Depoortere R, Newman-Tancredi A (2008) Effects of antipsychotics and reference monoaminergic ligands on marble burying behavior in mice. *Behav Pharmacol* 19:145–152.
- Chapman PF, White GL, Jones MW, Cooper-Blacketer D, Marshall VJ, Irizarry M, Younkin L, Good MA, Bliss TV, Hyman BT, Younkin SG, Hsiao KK (1999) Impaired synaptic plasticity and learning in aged amyloid precursor protein transgenic mice. *Nat Neurosci* 2:271–276.
- Chen G, Chen KS, Knox J, Inglis J, Bernard A, Martin SJ, Justice A, McCon-

- logue L, Games D, Freedman SB, Morris RG (2000) A learning deficit related to age and beta-amyloid plaques in a mouse model of Alzheimer's disease. *Nature* 408:975–979.
- Cook EH Jr, Lindgren V, Leventhal BL, Courchesne R, Lincoln A, Shulman C, Lord C, Courchesne E (1997) Autism or atypical autism in maternally but not paternally derived proximal 15q duplication. *Am J Hum Genet* 60:928–934.
- Cook EH Jr, Courchesne RY, Cox NJ, Lord C, Gonen D, Guter SJ, Lincoln A, Nix K, Haas R, Leventhal BL, Courchesne E (1998) Linkage-disequilibrium mapping of autistic disorder, with 15q11–13 markers. *Am J Hum Genet* 62:1077–1083.
- Costa AC, Stasko MR, Stoffel M, Scott-McKean JJ (2005) G-protein-gated potassium (GIRK) channels containing the GIRK2 subunit are control hubs for pharmacologically induced hypothermic responses. *J Neurosci* 25:7801–7804.
- Crawley JN (2007) Emotional behaviors: animal models of psychiatric disease. In: *What's wrong with my mouse?: Behavioral phenotyping of transgenic and knockout mice*, Ed 2, pp 227–265. Hoboken, NJ: Wiley.
- Cryan JF, Mombereau C (2004) In search of a depressed mouse: utility of models for studying depression-related behavior in genetically modified mice. *Mol Psychiatry* 9:326–357.
- Deacon RM (2006a) Burrowing in rodents: a sensitive method for detecting behavioral dysfunction. *Nat Protoc* 1:118–121.
- Deacon RM (2006b) Digging and marble burying in mice: simple methods for *in vivo* identification of biological impacts. *Nat Protoc* 1:122–124.
- Deacon RM, Raley JM, Perry VH, Rawlins JN (2001) Burrowing into prion disease. *Neuroreport* 12:2053–2057.
- Deacon RM, Croucher A, Rawlins JN (2002) Hippocampal cytotoxic lesion effects on species-typical behaviours in mice. *Behav Brain Res* 132:203–213.
- Deacon RM, Cholerton LL, Talbot K, Nair-Roberts RG, Sanderson DJ, Romberg C, Koros E, Bornemann KD, Rawlins JN (2008) Age-dependent and -independent behavioral deficits in Tg2576 mice. *Behav Brain Res* 189:126–138.
- Dudek BC, Adams N, Boice R, Abbott ME (1983) Genetic influences on digging behaviors in mice (*Mus musculus*) in laboratory and seminatural settings. *J Comp Psychol* 97:249–259.
- Estelles J, Lluich J, Rodriguez-Arias M, Aguilar MA, Miñarro J (2007) Cocaine exposure during adolescence affects anxiety in adult mice. *Brain Res Bull* 71:393–403.
- File SE (1992) Behavioral detection of anxiolytic action. In: *Experimental approaches to anxiety and depression* (Elliott JM, Heal DJ, Marsden CA, eds), pp 25–44. New York: Wiley.
- Finn DA, Rutledge-Gorman MT, Crabbe JC (2003) Genetic animal models of anxiety. *Neurogenetics* 4:109–135.
- Gillberg C (1998) Chromosomal disorders and autism. *J Autism Dev Disord* 28:415–425.
- Goodwin GM, De Souza RJ, Green AR (1985) The pharmacology of the hypothermic response in mice to 8-hydroxy-2-(di-*n*-propylamino)tetralin (8-OH-DPAT). A model of presynaptic 5-HT1 function. *Neuropharmacology* 24:1187–1194.
- Gotthardt M, Trommsdorff M, Nevitt MF, Shelton J, Richardson JA, Stockinger W, Nimpf J, Herz J (2000) Interactions of the low density lipoprotein receptor gene family with cytosolic adaptor and scaffold proteins suggest diverse biological functions in cellular communication and signal transduction. *J Biol Chem* 275:25616–25624.
- Grant EC, Mackintosh JH (1963) A comparison of the social postures of some laboratory rodents. *Behavior* 21:246–259.
- Gray JD, Punsoni M, Tabori NE, Melton JT, Fanslow V, Ward MJ, Zupan B, Menzer D, Rice J, Drake CT, Romeo RD, Brake WG, Torres-Reveron A, Milner TA (2007) Methylphenidate administration to juvenile rats alters brain areas involved in cognition, motivated behaviors, appetite, and stress. *J Neurosci* 27:7196–7207.
- Gross C, Zhuang X, Stark K, Ramboz S, Oosting R, Kirby L, Santarelli L, Beck S, Hen R (2002) Serotonin1A receptor acts during development to establish normal anxiety-like behaviour in the adult. *Nature* 416:396–400.
- Gyertyán I (1995) Analysis of the marble burying response: marbles serve to measure digging rather than evoke burying. *Behav Pharmacol* 6:24–31.
- Hill K, Li Y, Bennett M, McKay M, Zhu X, Shern J, Torre E, Lah JJ, Levey AI, Kahn RA (2003) Munc18 interacting proteins: ADP-ribosylation factor-dependent coat proteins that regulate the traffic of beta-Alzheimer's precursor protein. *J Biol Chem* 278:36032–36040.
- Hille B (1992) G protein-coupled mechanisms and nervous signaling. *Neuron* 9:187–195.
- Ho A, Morishita W, Hammer RE, Malenka RC, Südhof TC (2003) A role for Mints in transmitter release: Mint 1 knockout mice exhibit impaired GABAergic synaptic transmission. *Proc Natl Acad Sci U S A* 100:1409–1414.
- Ho A, Morishita W, Atasoy D, Liu X, Tabuchi K, Hammer RE, Malenka RC, Südhof TC (2006) Genetic analysis of Mint/X11 proteins: essential presynaptic functions of a neuronal adaptor protein family. *J Neurosci* 26:13089–13101.
- Holmes A, Yang RJ, Lesch KP, Crawley JN, Murphy DL (2003) Mice lacking the serotonin transporter exhibit 5-HT(1A) receptor-mediated abnormalities in tests for anxiety-like behavior. *Neuropsychopharmacology* 28:2077–2088.
- Inanobe A, Yoshimoto Y, Horio Y, Morishige KI, Hibino H, Matsumoto S, Tokunaga Y, Maeda T, Hata Y, Takai Y, Kurachi Y (1999) Characterization of G-protein-gated K⁺ channels composed of Kir3.2 subunits in dopaminergic neurons of the substantia nigra. *J Neurosci* 19:1006–1017.
- Iwasato T, Datwani A, Wolf AM, Nishiyama H, Taguchi Y, Tonegawa S, Knöpfel T, Erzurumlu RS, Itoharu S (2000) Cortex-restricted disruption of NMDAR1 impairs neuronal patterns in the barrel cortex. *Nature* 406:726–731.
- Janus C, Pearson J, McLaurin J, Mathews PM, Jiang Y, Schmidt SD, Chishti MA, Horne P, Heslin D, French J, Mount HT, Nixon RA, Mercken M, Bergeron C, Fraser PE, St George-Hyslop P, Westaway D (2000) A beta peptide immunization reduces behavioural impairment and plaques in a model of Alzheimer's disease. *Nature* 408:979–982.
- Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S, Malinow R (2003) APP processing and synaptic function. *Neuron* 37:925–937.
- Kimura K, Kitano J, Nakajima Y, Nakanishi S (2004) Hyperpolarization-activated, cyclic nucleotide-gated HCN2 cation channel forms a protein assembly with multiple neuronal scaffold proteins in distinct modes of protein-protein interaction. *Genes Cells* 9:631–640.
- Kitano J, Kimura K, Yamazaki Y, Soda T, Shigemoto R, Nakajima Y, Nakanishi S (2002) Tamalin, a PDZ domain-containing protein, links a protein complex formation of group 1 metabotropic glutamate receptors and the guanine nucleotide exchange factor cytohesins. *J Neurosci* 22:1280–1289.
- Kobayashi T, Hayashi E, Shimamura M, Kinoshita M, Murphy NP (2008) Neurochemical responses to antidepressants in the prefrontal cortex of mice and their efficacy in preclinical models of anxiety-like and depression-like behavior: a comparative and correlational study. *Psychopharmacology (Berl)* 197:567–580.
- Lau KF, McLoughlin DM, Standen C, Miller CC (2000) X11 alpha and x11 beta interact with presenilin-1 via their PDZ domains. *Mol Cell Neurosci* 16:557–565.
- Lee JH, Lau KF, Perkinton MS, Standen CL, Rogelj B, Falinska A, McLoughlin DM, Miller CC (2004) The neuronal adaptor protein X11beta reduces amyloid beta-protein levels and amyloid plaque formation in the brains of transgenic mice. *J Biol Chem* 279:49099–49104.
- Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Benjamin J, Müller CR, Hamer DH, Murphy DL (1996) Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 274:1527–1531.
- Li Q, Wichems C, Heils A, Van De Kar LD, Lesch KP, Murphy DL (1999) Reduction of 5-hydroxytryptamine (5-HT)(1A)-mediated temperature and neuroendocrine responses and 5-HT(1A) binding sites in 5-HT transporter knockout mice. *J Pharmacol Exp Ther* 291:999–1007.
- Lo Iacono L, Gross C (2008) Alpha-Ca²⁺/calmodulin-dependent protein kinase II contributes to the developmental programming of anxiety in serotonin receptor 1A knock-out mice. *J Neurosci* 28:6250–6257.
- Lunn ML, Nassirpour R, Arrabit C, Tan J, McLeod I, Arias CM, Sawchenko PE, Yates JR 3rd, Slesinger PA (2007) A unique sorting nexin regulates trafficking of potassium channels via a PDZ domain interaction. *Nat Neurosci* 10:1249–1259.
- Lüscher C, Jan LY, Stoffel M, Malenka RC, Nicoll RA (1997) G protein-coupled inwardly rectifying K⁺ channels (GIRKs) mediate postsynaptic but not presynaptic transmitter actions in hippocampal neurons. *Neuron* 19:687–695.
- Ma D, Zerangue N, Raab-Graham K, Fried SR, Jan YN, Jan LY (2002) Diverse trafficking patterns due to multiple traffic motifs in G protein-

- activated inwardly rectifying potassium channels from brain and heart. *Neuron* 33:715–729.
- Martin KF, Phillips I, Hearson M, Prow MR, Heal DJ (1992) Characterization of 8-OH-DPAT-induced hypothermia in mice as a 5-HT_{1A} autoreceptor response and its evaluation as a model to selectively identify antidepressants. *Br J Pharmacol* 107:15–21.
- Masuda Y, Ishigooka S, Matsuda Y (2000) Digging behavior of ddY mouse. *Exp Anim* 49:235–237.
- Maximov A, Südhof TC, Bezprozvanny I (1999) Association of neuronal calcium channels with modular adaptor proteins. *J Biol Chem* 274:24453–24456.
- Mayford M, Bach ME, Huang YY, Wang L, Hawkins RD, Kandel ER (1996) Control of memory formation through regulated expression of a CaMKII transgene. *Science* 274:1678–1683.
- McKinney BC, Schneider JS, Schafer GL, Lowing JL, Mohan S, Zhao MX, Heng MY, Albin RL, Seasholtz AF, Akil H, Murphy GG (2008) Decreased locomotor activity in mice expressing tTA under control of the CaMKII alpha promoter. *Genes Brain Behav* 7:203–213.
- Miller CC, McLoughlin DM, Lau KF, Tennant ME, Rogelj B (2006) The X11 proteins, Abeta production and Alzheimer's disease. *Trends Neurosci* 29:280–285.
- Morgan D, Diamond DM, Gottschall PE, Ugen KE, Dickey C, Hardy J, Duff K, Jantzen P, DiCarlo G, Wilcock D, Connor K, Hatcher J, Hope C, Gordon M, Arendash GW (2000) A beta peptide vaccination prevents memory loss in an animal model of Alzheimer's disease. *Nature* 408:982–985.
- Mori A, Okuyama K, Horie M, Taniguchi Y, Wadatsu T, Nishino N, Shimada Y, Miyazawa N, Takeda S, Niimi M, Kyushiki H, Kondo M, Mitsumoto Y (2002) Alteration of methamphetamine-induced striatal dopamine release in *mint-1* knockout mice. *Neurosci Res* 43:251–257.
- Moy SS, Nadler JJ, Young NB, Perez A, Holloway LP, Barbaro RP, Barbaro JR, Wilson LM, Threagill DW, Lauder JM, Magnuson TR, Crawley JN (2007) Mouse behavioral tasks relevant to autism: phenotypes of 10 inbred strains. *Behav Brain Res* 176:4–20.
- Muhle R, Trentacoste SV, Rapin I (2004) The genetics of autism. *Pediatrics* 113:e472–e486.
- Murer G, Adelbrecht C, Lauritzen I, Lesage F, Lazdunski M, Agid Y, Raisman-Vozari R (1997) An immunocytochemical study on the distribution of two G-protein-gated inward rectifier potassium channels (GIRK2 and GIRK4) in the adult rat brain. *Neuroscience* 80:345–357.
- Nakajima Y, Okamoto M, Nishimura H, Obata K, Kitano H, Sugita M, Matsuyama T (2001) Neuronal expression of *mint1* and *mint2*, novel multimodular proteins, in adult murine brain. *Brain Res Mol Brain Res* 92:27–42.
- Nakamura K, Kurasawa M (2001) Anxiolytic effects of aniracetam in three different mouse models of anxiety and the underlying mechanism. *Eur J Pharmacol* 420:33–43.
- Nicolas LB, Kolb Y, Prinszen EP (2006) A combined marble burying-locomotor activity test in mice: a practical screening test with sensitivity to different classes of anxiolytics and antidepressants. *Eur J Pharmacol* 547:106–115.
- Njung'e K, Handley SL (1991a) Evaluation of marble-burying behavior as a model of anxiety. *Pharmacol Biochem Behav* 38:63–67.
- Njung'e K, Handley SL (1991b) Effects of 5-HT uptake inhibitors, agonists and antagonists on the burying of harmless objects by mice; a putative test for anxiolytic agents. *Br J Pharmacol* 104:105–112.
- North RA (1989) Twelfth Gaddum memorial lecture. Drug receptors and the inhibition of nerve cells. *Br J Pharmacol* 98:13–28.
- Okamoto M, Südhof TC (1997) *Mint1*, *Munc18*-interacting proteins in synaptic vesicle exocytosis. *J Biol Chem* 272:31459–31464.
- Popova NK (2006) From genes to aggressive behavior: the role of serotonergic system. *Bioessays* 28:495–503.
- Reif A, Lesch KP (2003) Toward a molecular architecture of personality. *Behav Brain Res* 139:1–20.
- Rogelj B, Mitchell JC, Miller CC, McLoughlin DM (2006) The X11/Mint family of adaptor proteins. *Brain Res Rev* 52:305–315.
- Saenz del Burgo L, Cortes R, Mengod G, Zarate J, Echevarria E, Salles J (2008) Distribution and neurochemical characterization of neurons expressing GIRK channels in the rat brain. *J Comp Neurol* 510:581–606.
- Saito Y, Sano Y, Vassar R, Gandy S, Nakaya T, Yamamoto T, Suzuki T (2008) X11 proteins regulate the translocation of amyloid beta-protein precursor (APP) into detergent-resistant membrane and suppress the amyloidogenic cleavage of APP by beta-site-cleaving enzyme in brain. *J Biol Chem* 283:35763–35771.
- Sankoorikal GM, Kaercher KA, Boon CJ, Lee JK, Brodtkin ES (2006) A mouse model system for genetic analysis of sociability: C57BL/6J versus BALB/cJ inbred mouse strains. *Biol Psychiatry* 59:415–423.
- Sano Y, Syuzo-Takabatake A, Nakaya T, Saito Y, Tomita S, Itohara S, Suzuki T (2006a) Enhanced amyloidogenic metabolism of the amyloid beta-protein precursor in the X11L-deficient mouse brain. *J Biol Chem* 281:37853–37860.
- Sano Y, Nakaya T, Pedrini S, Takeda S, Iijima-Ando K, Iijima K, Mathews PM, Itohara S, Gandy S, Suzuki T (2006b) Physiological mouse brain Abeta levels are not related to the phosphorylation state of threonine-668 of Alzheimer's APP. *PLoS ONE* 1:e51.
- Sassa T, Gomi H, Itohara S (2004) Postnatal expression of *Cdkl2* in mouse brain revealed by LacZ inserted into the *Cdkl2* locus. *Cell Tissue Res* 315:147–156.
- Schein JC, Hunter DD, Roffler-Tarlov S (1998) *Girk2* expression in the ventral midbrain, cerebellum, and olfactory bulb and its relationship to the murine mutation weaver. *Dev Biol* 204:432–450.
- Schroer RJ, Phelan MC, Michaelis RC, Crawford EC, Skinner SA, Cuccaro M, Simensen RJ, Bishop J, Skinner C, Fender D, Stevenson RE (1998) Autism and maternally derived aberrations of chromosome 15q. *Am J Med Genet* 76:327–336.
- Shao Y, Wolpert CM, Raiford KL, Menold MM, Donnelly SL, Ravan SA, Bass MP, McClain C, von Wendt L, Vance JM, Abramson RH, Wright HH, Ashley-Koch A, Gilbert JR, DeLong RG, Cuccaro ML, Pericak-Vance MA (2002) Genomic screen and follow-up analysis for autistic disorder. *Am J Med Genet* 114:99–105.
- Shinomiyama K, Fujii Y, Sugimoto Y, Azuma N, Tokunaga S, Kitazumi K, Kamei C (2005) Effect of paroxetine on marble-burying behavior in mice. *Methods Find Exp Clin Pharmacol* 27:685–687.
- Sumioka A, Imoto S, Martins RN, Kirino Y, Suzuki T (2003) X11L (X11-like protein)-dependent and -independent mechanisms. *Biochem J* 374:261–268.
- Tanaka M, Yamaguchi K, Tatsukawa T, Nishioka C, Nishiyama H, Theis M, Willecke K, Itohara S (2008) Lack of connexin43-mediated Bergmann glial gap junctional coupling does not affect cerebellar long-term depression, motor coordination, or eyeblink conditioning. *Front Behav Neurosci* 2:1–14.
- Tarassov K, Michnick SW (2005) iVici: interrelational visualization and correlation interface. *Genome Biol* 6:R115.
- Taru H, Suzuki T (2004) Facilitation of stress-induced phosphorylation of beta-amyloid precursor protein family members by X11-like/Mint2 protein. *J Biol Chem* 279:21628–21636.
- Tomita S, Ozaki T, Taru H, Oguchi S, Takeda S, Yagi Y, Sakiyama S, Kirino Y, Suzuki T (1999) Interaction of a neuron-specific protein containing PDZ domains with Alzheimer's amyloid precursor protein. *J Biol Chem* 274:2243–2254.
- Vergnes M, Depaulis A, Boehrer A, Kempf E (1988) Selective increase of offensive behavior in the rat following intrahypothalamic 5,7-DHT-induced serotonin depletion. *Behav Brain Res* 29:85–91.
- Wada E, Watase K, Yamada K, Ogura H, Yamano M, Inomata Y, Eguchi J, Yamamoto K, Sunday ME, Maeno H, Mikoshiba K, Ohki-Hamazaki H, Wada K (1997) Generation and characterization of mice lacking gastrin-releasing peptide receptor. *Biochem Biophys Res Commun* 239:28–33.
- Webster DG, Williams MH, Owens RD, Geiger VB, Dewsbury DA (1981) Digging behavior in 12 taxa of muroid rodents. *Anim Learn Behav* 9:173–177.
- Williams JT, Colmers WF, Pan ZZ (1988) Voltage- and ligand-activated inwardly rectifying currents in dorsal raphe neurons *in vitro*. *J Neurosci* 8:3499–3506.
- Yamada M, Inanobe A, Kurachi Y (1998) G protein regulation of potassium ion channels. *Pharmacol Rev* 50:723–760.
- Yamada K, Wada E, Yamano M, Sun YJ, Ohara-Imazumi M, Nagamatsu S, Wada K (2002) Decreased marble burying behavior in female mice lacking neuromedin-B receptor (NMB-R) implies the involvement of NMB/NMB-R in 5-HT neuron function. *Brain Res* 942:71–78.
- Zhuang X, Gross C, Santarelli L, Compan V, Trillat AC, Hen R (1999) Altered emotional states in knockout mice lacking 5-HT_{1A} or 5-HT_{1B} receptors. *Neuropsychopharmacology* 21:525–605.

Physical Properties of Bedding Materials Determine the Marble Burying Behavior of Mice (C57BL/6J)

Chihiro Homma and Kazuyuki Yamada*

Support Unit for Animal Experiments, Research Resources Center, Brain Science Institute, RIKEN, 2-1 Hirosawa, Wako-city, Saitama, 351-0198, Japan

Abstract: The marble burying test is a simple behavioral psychopharmacology task in which rodents, especially rats and mice, are exposed to glass marbles laying on thick bedding materials. This test has been widely used to assess the effects of a variety of psychoactive drugs. Although the marble burying test clearly reflects drug efficacy, many aspects of the behavior itself remain unclear. In the present study, we assessed the effects of different bedding materials and the number of glass marbles presented on the marble burying behavior of mice (C57BL/6J). Six (bedding materials) × three (number of glass marbles) factorial experiments revealed that the number of marbles buried correlates closely with the quality of the bedding material. Our results provide a basic understanding of certain factors that govern marble burying behavior, which may be applicable to behavioral sciences including psychopharmacology.

Keywords: Marble burying, defensive burying, digging, bedding materials, mice.

INTRODUCTION

Defensive burying is a widely observed behavior in rodents [1-8]. Rats and mice eagerly bury noxious things, such as bad tasting and/or smelling food [1-3], shock prod [4], and unusual objects like glass marbles [2], under their bedding materials. This behavior is thought to be an expression of defensiveness, some forms of anxiety, or compulsiveness [5]. When rats and/or mice are placed in a cage in which glass marbles are laying on sawdust, they immediately and eagerly begin burying the marbles [2]. The marble burying behavior is quite easy to observe and is sensitive to anxiolytic and/or anxiogenic compounds. Therefore, this behavior is often used to assess the properties of psychoactive drugs [6-11].

Certain unique properties of marble burying behavior have been revealed. Animals bury glass marbles even though they are not harmful. Marble burying behavior does not show any habituation after repeated trials [6,12-14], indicating that this behavior is not likely to be affected by learning processes or cognitive function. Furthermore, marble burying behavior does not differ by gender [12]. Although marble burying behavior has been well characterized, the factors governing the burying behavior itself remain unclear. It was recently reported that the marble burying test simply reflects digging and other related behaviors [14-17], calling into question the validity of this behavior for assessing anxiety, defensiveness, compulsiveness, and so on. However, marble burying behavior correlates well with both serotonin (5-HT) transporter (SERT) binding of selective 5-HT reuptake inhibitors (SSRIs) and extracellular 5-HT levels in brains treated with SSRIs [18,19], and thus the behavior may reflect a relation between 5-HT neuronal

activity and anxiety, defensiveness, compulsiveness. Therefore, the marble burying test is often used to assess the efficacy of drugs such as SSRIs.

The number of marbles an animal buries is determined by certain factors such as the type of bedding material and the number of marbles used. However, these aspects have not been well studied. For example, many studies have used "saw dust" as the bedding material to study burying behavior because it is lightweight, allowing animals to easily bury the marbles. This feature of "saw dust" might prompt digging, however, leading to a misunderstanding of the emotional state of the animals and/or drug effects. Thus, knowledge of the effects of different bedding materials is required to properly assess burying behavior.

The aim of this study was to assess the effects of bedding material type and number of glass marbles on marble burying test results. Two-factorial within-subject analysis revealed significant effects of bedding materials on the marble burying behavior of C57BL/6J mice.

MATERIALS AND METHODOLOGY

Animals

Seventeen male C57BL/6J mice were used as subjects. Mice were bred in the animal facility of RIKEN BSI (originally purchased from JCL, Inc. Tokyo, Japan, F2 generation) and were about 10 weeks of age at the beginning of the experiment. The breeding room was air-conditioned (22 ± 1°C, 50-60% humidity) with a 12h:12h light-dark cycle (lights on at 0800). Mice were individually housed beginning one week prior to the experiment, and food and water were freely available except during experimentation. All experiments were conducted in the breeding room during the light cycle between 1300 and 1700.

All animal experiments in this study were performed in strict accordance with the guidelines of The Institute of Physical and Chemical Research (RIKEN), Brain Science

*Address correspondence to this author at the Support Unit for Animal Experiments, Research Resources Center, Brain Science Institute, RIKEN, 2-1 Hirosawa, Wako-city, Saitama, 351-0198, Japan; Tel: +81-48-467-7659; Fax: +81-48-467-6287; E-mail: kaz-yamada@brain.riken.jp

Institute (BSI), and were approved by the Animal Investigation Committee of the Institute.

Apparatus

All experiments were conducted in clear plastic caging with filter tops (Allentown, PA, USA: $29 \times 19.5 \times 13.5$ cm (H)). Glass marbles were clear, green in color and 1.5 cm in diameter. A spontaneous activity monitor (AB System: Neuroscience, Tokyo, Japan) was used to measure activity in the cage during both habituation and test periods.

Bedding Materials

Six types of commercially available bedding materials were used: Soft-chip and Paper-clean (SLC, Shizuoka, Japan), TEK-Fresh (Harlan, IN, USA), Cellu-dri and ALPHA-dri (Shepherd Specialty Papers, TN, USA), and GreenTru (GREENPRODUCTS, IA, USA) (Fig. 1). All bedding materials were autoclaved before use.

Spontaneous Activity Monitoring

The ceiling of each compartment of the breeding rack was equipped with a pyroelectric sensor to monitor the movements of mice throughout the experiment. Data were collected and analyzed with a PC equipped with Windows and other commercially available software (AB System). Activity counts represent the number of active time bin (approximately 0.20 - 0.25 sec each) in which spontaneous activity including locomotor activity, rearing and other activities such as stereotypic movements were detected.

Marble Burying Test

A modified procedure based on Yamada *et al.* [20] was employed. Mice were placed individually in plastic cages with the designated bedding material for 30 min (habituation period) and then placed into waiting cages. Ten, twenty or forty glass marbles were then evenly spaced 3 to 7 cm apart on a 4-cm layer of bedding material in the habituation cages. Mice were then reintroduced into the same cage in which they had been habituated. After 30 min, the marble burying period was terminated by removing the mice, and the number of marbles that were more than two-thirds covered with bedding material was counted. Whether a marble was buried was established by two experimenters and was

confirmed by manual post-experimental assessment of digital photographs on a PC. After each trial, cages and bedding materials were replaced with fresh ones, and glass marbles were washed with water, dried with a paper towel, and left to return to room temperature.

The marble burying test was repeated twenty times. The first, tenth and the last tests were calibration trials, in which mice were exposed to twenty glass marbles on TEK-Fresh, the bedding material used in our breeding facility. In the second to ninth and eleventh to nineteenth trials each mouse was assigned to one of 17 designated conditions—number of glass marbles (10, 20, or 40) and bedding materials (Soft-chip, TEK-Fresh, Paper-clean, Cellu-dri, ALPHA-dri, or GreenTru). The order of condition for each mouse was counterbalanced using *Latin square design*. Each trial was separated by two to three days. On days between trials, mice were individually housed in a cage with a 2-cm layer of TEK-Fresh.

Statistical Analysis

Statistical analyses were performed by the SPSS™ statistical package (ver 16.0). The repeated measure ANOVA and one-way within-subject ANOVA were used. *Post-hoc* analysis was conducted using Dunnett's *t*-test or Tukey's HSD test. The correlation between the density of the bedding materials and the number of buried marbles was analyzed using Pearson's correlation coefficient. Statistical significance was set at $p < 0.05$.

RESULTS


Effects of Repeated Trials on Spontaneous Activity and Marble Burying Behavior

Because many trials were conducted in this study, we assessed the effects of repeated trials both on spontaneous activity and marble burying behavior (Fig. 2). Spontaneous activity increased linearly over the trials, and one-way ANOVA revealed a statistically significant main effect of trial number ($F(2,32) = 10.2$, $p < 0.001$, Fig. 2A). In contrast, the number of buried marbles did not differ significantly with trial number ($F(2,32) = 2.16$, Fig. 2B), consistent with previous studies [6,12,13].



Fig. (1). Bedding materials used in this study. Numbers under the bedding material names indicate the density (g/L).

A Spontaneous activity



B Marble burying




Fig. (2). Spontaneous activity increases with repeated trials, but the number of marbles buried does not. (A) spontaneous activity, (B) number of marbles buried. Number of marbles used was twenty in each trial. Data represent mean ± SEM. **: $p < 0.01$.


Spontaneous Activity is Affected by Bedding Material

It is possible that the spontaneous activity level of mice could affect marble burying behavior and might be related to properties of the bedding material. Therefore, we assessed the spontaneous activity of mice during both habituation and the marble burying period (Fig. 3). Activity was clearly affected by the type of bedding material. During the habituation period (Fig. 3A), one-way ANOVA revealed a significant main effect of bedding material in 10-, 20-, and 40-marble trials ($F(5,80) = 6.03, p < 0.0001$; $F(5,80) = 7.01, p < 0.0001$; $F(5,80) = 4.37, p < 0.01$, respectively). During the marble burying period (Fig. 3B), one-way ANOVA revealed a significant main effect of bedding material in the 20- and 40-marble trials ($F(5,80) = 2.52, p < 0.05$; $F(5,80) = 3.11, p < 0.05$, respectively). Throughout the experiment, mice exhibited the highest activity in the TEK-Fresh trials and showed the lowest activity in GreenTru trials. These results strongly suggest that spontaneous activity of mice is affected by bedding material properties.

Low-Density Bedding Increase Marble Burying Behavior in Mice

Results of the marble burying test are summarized in Fig. (4). Mice buried many glass marbles when the density of

A Habituation period



B Marble burying period




Fig. (3). Mean spontaneous activity during habituation and the marble burying period. (A) habituation period; (B) marble burying period. The terms ‘marble 10’, ‘marble 20’, and ‘marble 40’ indicate marble burying test trials with 10, 20, and 40 marbles, respectively. TEK: TEK-Fresh, ALP: ALPHA-dri, Cellu: Cellu-dri, Sof: Soft-chip, Pap: Paper-clean, Gre: GreenTru. Statistics are presented in the text.

bedding material was low, for example in Soft-chip (20 g/L) and TEK-Fresh (80.3 g/L) bedding. In contrast, they buried only a few marbles when the density of the bedding material was high, for example in ALPHA-dri (194.7 g/L) and GreenTru (277.4 g/L) bedding. In Paper-clean (124.2 g/L) and Cellu-dri (173.2 g/L) bedding, which are moderately dense, the marble burying behavior was moderate. Furthermore, the effect of the number of glass marbles on the burying behavior was salient. In Soft-chip, TEK-Fresh, Cellu-dri and Paper-clean (partially) trials, the number of buried marbles increased linearly with the number of marbles in the cage. The number of buried marbles did not show a linear increment in the ALPHA-dri and GreenTru trials, however. Fig. (5) shows the negative correlation between the number of buried marbles and the density of bedding materials (10 marbles (Fig. 5A): $r = -0.973, p < 0.01$; 20 marbles (Fig. 5B): $r = -0.962, p < 0.01$; 40 marbles (Fig. 5C): $r = -0.941, p < 0.01$, respectively). These results indicate that bedding material density strongly affects the number of marbles buried by mice. In addition, bedding material density may also affect how the mice bury the marbles. Representative photos of burying behaviors in each of the bedding materials are shown in Fig. (6). In Soft-chip and TEK-Fresh, mice completely buried the marbles under

the low-density bedding material. In Cellu-dri, ALPHA-dri and GreenTru, however, the mice only partially buried the marbles. The texture, as well as density, of the bedding material may also affect burying behavior. Mice buried more marbles in bedding materials with a soft texture (Soft-chip, TEK-Fresh, Paper-clean, Cellu-dri) than in bedding with a hard texture (ALPHA-dri, GreenTru). We replicated experiments several times with a somewhat smaller design and obtained almost identical results (data not shown). These results suggest that bedding material properties may affect how mice bury marbles.




Fig. (4). Marble burying changes with the type of bedding and number of marbles presented. Sof: Soft-chip, TEK: TEK-Fresh, Pap: Paper-clean, Cellu: Cellu-dri, ALP: ALPHA-dri, Gre: GreenTru. 10, 20, and 40 represent trials with 10, 20, and 40 marbles, respectively. Data represent mean ± SEM. Statistics are presented in the text.


Correlation Between Spontaneous Activity and Number of Marbles Buried

As described above, spontaneous activity may affect marble burying behavior. Therefore, we examined the correlation between the spontaneous activity score and the number of marbles buried (Fig. 7). In both 10- and 20-marble trials, correlation coefficients were not statistically significant, contrary, in the 40-marble trials, the correlation coefficient was statistically significant ($r = 0.20, p < 0.05$). Although there is a significant correlation between spontaneous activity and number of marbles, the small effect size for this measure suggests that spontaneous activity does not play a crucial role for marble burying behavior.


DISCUSSION

In this study, we demonstrated that the physical properties of bedding material and the number of glass marbles in the cage significantly affect the marble burying behavior of mice. The number of marbles buried depended on bedding material density and texture. Mice buried more marbles in low-density, soft bedding than in high-density, hard bedding (Figs. 1, 4). Furthermore, the number of marbles buried correlated with the number of marbles used for the test when the bedding was not high-density and/or hard in texture (Fig. 4). Thus, the properties of bedding, rather than the number of marbles used, seems to be the principal determining factor governing marble burying behavior.

A 10 marbles



B 20 marbles



C 40 marbles




Fig. (5). Correlation between the number of marbles buried and bedding material density. (A) trials with 10 marbles; (B) trials with 20 marbles; (C) trials with 40 marbles. Red lines represent the linear regression.




Fig. (6). Representative marble burying behavior in each bedding material.

Bedding material also affected spontaneous activity of mice both quantitatively and qualitatively. Animals were most active in the TEK-Fresh trial and were least active in the GreenTru trial (Fig. 3). The highest activity in the TEK-Fresh trial might be attributed to the 'familiar environment' not to the physical property of the bedding, because TEK-Fresh is used in our breeding facility. However, this possibility might be excluded in this study because mice show high activity when they are exposed to fresh bedding, showing it is not 'familiar'. Activity also differed qualitatively with bedding type. Mice eagerly dug the layer of bedding in TEK-Fresh and Soft-chip trials, but rarely dug in GreenTru trials (data not shown). Certain studies have suggested that marble burying behavior correlates strongly with digging behavior [14-16], and some marbles might be covered with bedding as an indirect consequence of digging. Thus, marble burying behavior, at least in trials using low-density bedding materials, consists of two qualitatively different behaviors, burying and digging. These mixed behaviors may confound the results of studies assessing drugs effects.

Because animals did not dig much in ALPHA-dri and GreenTru trials, the small number of marbles buried in these trials (Fig. 4) might reflect "real burying behavior", raising a fundamental question about the marble burying test. In TEK-Fresh and Soft-chip trials, a majority of marbles might have been buried indirectly by the digging behavior. Previous work has suggested that the marble burying test simply measures digging behavior [14-15]. Is marble burying

behavior equal to the digging behavior? Or is the contribution of burying to marble burying behavior so small? In this way, marble burying test has some theoretical confusion. Thus, to elucidate this confusion we added some experiments in which the digging behavior in the habituation trial was measured using TEK-Fresh, Soft-chip and GreenTru. The correlation coefficient between the digging duration and the number of buried marbles was statistically significant in Soft-chip experiment ($n=5$, $r=0.897$, $p<0.05$), contrary, not significant in TEK-Fresh and GreenTru experiments ($n=6$, $r=-0.01$, n.s.; $n=7$, $r=0.208$, n.s., respectively). These results suggest that a selection of bedding materials (e.g. TEK-Fresh and/or GreenTru) may be able to dissect out marble burying behavior.

Regardless, the marble burying test is still quite useful for assessing psychotropic drugs [6-11]. For example, in this study, we repeated the marble burying test 20 times for each set of mice, and the number of buried marbles did not change (Fig. 2B). This lack of habituation provides reliability and validity to experiments having a within-subject design. The marble burying test may have other advantages in psychopharmacological studies, but further study is required to elucidate these advantages.

ACKNOWLEDGEMENTS

This study was supported by the Ministry of Education, Culture, Sports, Science and Technology (MEXT): Grants-in-Aid for Scientific Research (C) (19530670, K.Y.).