

53

今後の方針

- RSort GeneListと一致率比較をベースにした毒性推測サポートプログラムを作成する。
- 新たに追加される化学物質データについては、全ての既存データとの一致率計算をルーチン化する。
- 各種アノテーション情報(フェノタイプ情報など)の取り込みを進める。
- データベース公開用Webアプリケーションに組み込み、一般公開する。

54

別添 5

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書籍名	出版社名	出版地	出版年	ページ
Kitano, H.	The Theory of Biological Robustness and its Implication to Cancer.	Pengcheng Fu and Sven Panke	SYSTEMS BIOLOGY AND SYNTHETIC BIOLOGY	John Wiley & Sons, Inc	USA	2009	529-547
北嶋 聡	食品, 食品添加物, 食品汚染物質, 飼料添加物	上野光一 ら	新版トキシコロジー	朝倉書店	東京	2009	118-126

雑誌

発表者氏名	論文タイトル名	発表誌名	巻名	ページ	出版年
Matsunaga N, Kanno J, Hamada C, Yoshimura I.	An experimental design for judging synergism on consideration to endocrine disruptor animal experiments.	Environmetrics	20	1 - 13	2009
Woo GH, Takahashi M, Inoue K, Fujimoto H, Igarashi K, Kanno J, Hirose M, Nishikawa A, Shibutani M.	Cellular distributions of molecules with altered expression specific to thyroid proliferative lesions developing in a rat thyroid carcinogenesis model.	Cancer Sci.	100 (4)	617 - 625	2009
Ishimaru N, Takagi A, Kohashi M, Yamada A, Arakaki R, Kanno J, Hayashi Y.	Neonatal exposure to low-dose 2,3,7,8-tetrachlorodibenzo-p-dioxin causes autoimmunity due to the disruption of T cell tolerance.	J Immunol.	182 (10)	6576 - 6586	2009
Kanno J.	Overview: "Children's toxicology", a renovating study field of irreversible "early exposure-delayed effects".	J Toxicol Sci.	34 Suppl 2	SP199 - 200	2009
Tanemura K, Igarashi K, Matsugami TR, Aisaki K, Kitajima S, Kanno J.	Intrauterine environment-genome interaction and children's development (2): Brain structure impairment and behavioral disturbance induced in male mice offspring by a single intraperitoneal administration of domoic acid (DA) to their dams.	J Toxicol Sci.	34 Suppl 2	SP279 - 286	2009
Sekine H, Mimura J, Oshima M, Okawa H, Kanno J, Igarashi K, Gonzalez FJ, Ikuta T, Kawajiri K, Fujii-Kuriyama Y.	Hypersensitivity of aryl hydrocarbon receptor-deficient mice to lipopolysaccharide-induced septic shock.	Mol Cell Biol.	29 (24)	6391 - 6400	2009

<p>Saegusa Y, Woo GH, Fujimoto H, Inoue K, Takahashi M, Hirose M, Igarashi K, Kanno J, Mitsumori K, Nishikawa A, Shibutani M.</p>	<p>Gene Expression Profiling and Cellular Distribution of Molecules with Altered Expression in the Hippocampal CA1 Region after Developmental Exposure to Anti-Thyroid Agents in Rats.</p>	<p>J Vet Med Sci.</p>	<p>72 (2)</p>	<p>187 - 195</p>	<p>2010</p>
<p>Suzuki A, Igarashi K, Aisaki KI, Kanno J, Saga Y.</p>	<p>NANOS2 interacts with the CCR4-NOT deadenylation complex and leads to suppression of specific RNAs.</p>	<p>Proc Natl Acad Sci</p>	<p>23 107 (8)</p>	<p>3594 - 3599</p>	<p>2010</p>
<p>Hase, T.; Tanaka, H.; Suzuki, Y.; Nakagawa, S.; Kitano, H</p>	<p>Structure of Protein Interaction Networks and Their Implications on Drug Design.</p>	<p>PLoS Computational Biology.</p>	<p>5, 10</p>	<p>e100 0550</p>	<p>2009</p>
<p>Le Novere, N.; Hucka, M.; Mi, H.; Moodie, S.; Schreiber, F.; Sorokin, A.; Demir, E.; Wegner, K.; Aladjem, M.; Wimalaratne, S.; Bergman, F.; Gauges, R.; Ghazal, P.; Kawaji, H.; Li, L.; Matsuoka, Y.; Villeger, A.; Boyd, S.; Calzone, L.; Courtot, M.; Dogrusoz, U.; Freeman, T.; Funahashi, A.; Ghosh, S.; Jouraku, A.; Kim, S.; Kolpakov, F.; Luna, A.; Sahle, S.; Schmidt, E.; Watterson, S.; Wu, G.; Goryanin, I.; Kell, D.; Sander, C.; Sauro, H.; Snoep, J.; Kohn, K.; & Kitano, H</p>	<p>The Systems Biology Graphical Notation.</p>	<p>Nature biotechnolog</p>	<p>27, 8</p>	<p>735 - 741</p>	<p>2009</p>
<p>Ktantz, M.; Ahmadpour, D.; Ottosson, L-G.; Warringer, J.; Waltermann, C.; Nordlander, B.; Klipp, E.; Blomberg, A.; Hohmann, S.; Kitano, H</p>	<p>Robustness and fragility in the yeast high osmolarity glycerol (HOG) signal-transduction pathway.</p>	<p>Molecular Systems Biology.</p>	<p>5:281</p>	<p>1 - 7</p>	<p>2009</p>
<p>Matsuoka, Y.; Ghosh, S.; Kitano, H. Consistent design schematics for biological systems</p>	<p>Standardization of representation in biological engineering.</p>	<p>Journal of the Royal Society Interface</p>	<p>6, Suppl 4</p>	<p>S393 - 404</p>	<p>2009</p>
<p>Nils Gehlenborg; Seán I. O'Donoghue; Nitin S Baliga; Alexander Goemann; Matthew A Hibbs; Hiroaki Kitano; Oliver Kohlbacher; Heiko Neuweger; Reinhard Schneider; Dan Tenenbaum and Anne-Claude Gavin</p>	<p>Visualization of omics data for systems biology.</p>	<p>Nature Methods.</p>	<p>7 3s</p>	<p>S56 - S68</p>	<p>2010</p>

92. Arkin A, Ross J, McAdams HH. Stochastic kinetic analysis of developmental pathway bifurcation in phage lambda-infected *Escherichia coli* cells. *Genetics* 1998;149:1633–1648.
93. Arkin AP. Synthetic cell biology. *Curr Opin Biotechnol* 2001;12:638–644.
94. Nimfa AJ, Magasanik B. Covalent modification of the *glnG* product, NRI, by the *glnL* product, NRII, regulates the transcription of the *ghrALG* operon in *Escherichia coli*. *Proc Natl Acad Sci USA* 1986;83:5909–5913.
95. Fuqua WC, Winans SC, Greenberg EP. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J Bacteriol* 1994;176:269–275.
96. Hu MC, Davidson N. The inducible lac operator-repressor system is functional in mammalian cells. *Cell* 1987;48:555–566.
97. Weber W, Schoenmakers R, Spielmann M, El-Baba MD, Foicher M, Keller B, Weber CC, Link N, van de Wetering P, Heinzen C, et al. Streptomyces-derived quorum-sensing systems engineered for adjustable transgene expression in mammalian cells and mice. *Nucleic Acids Res* 2003;31:e71.
98. Semenza GL, Wang GL. A nuclear factor induced by hypoxia via de novo protein synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional activation. *Mol Cell Biol* 1992;12:5447–5454.

16

THE THEORY OF BIOLOGICAL ROBUSTNESS AND ITS IMPLICATION TO CANCER

Hiroaki Kitano

Sony Computer Science Laboratories, Inc., 3-14-13 Higashi-Gotanda,
Shinagawa, Tokyo 141-0022, Japan

The Systems Biology Institute, 6-31-15 Jingumae,
Shibuya, Tokyo 150-0001, Japan

Department of Cancer Systems Biology, Cancer Institute of Japanese
Foundation for Cancer Research, Tokyo, Japan

16.1 INTRODUCTION

Systems biology aims at system-level understanding of biological systems [1,2]. Investigations of biological systems at system level are not a new concept and can be traced back to homeostasis by Walter Cannon, Cybernetics by Norbert Wiener [3], and the general systems theory by von Bertalanffy [4]. Numbers of approaches in physiology have also taken a systemic view of the biological subjects. Systems biology is gaining renewed interest today because of progress in genomics, molecular biology, nonlinear dynamics, computational science, and other related fields.

However, “system-level understanding” is a rather vague notion and is often hard to define. This is due to the fact that the system is not a tangible object. Genes and

proteins are more tangible because they are identifiable matters. Although the system is composed of these matters and they are components of the system, the system itself cannot be made tangible. Often, diagrams of the gene regulatory networks and the protein interaction networks are shown as representations of systems. It is certainly true that such diagrams capture any one aspect of the structures of the system, but they are still only static slices of the system. The heart of the system lies within the dynamics it creates and the logic behind it. It is science on the dynamical state of affairs.

There are four distinct phases that lead us to system-level understanding at various levels. First, system structure identification enables us to understand the structure of the system. While this may be only a static view of the system, it is an essential first step. The structure shall then be identified, ultimately, in both physical and interaction structures. Interaction structures are represented as gene regulatory networks and biochemical networks that indicate how components interact within and in between cells. Physical details of specific regions of the cell, overall structure of cells, and organisms are also important because such physical structure imposes constraints on possible interactions and the outcome of interactions impacts the formation of physical structures. Nature of interaction could be different if proteins involved in interaction move by simple diffusion or under specific guidance from cytoskeleton.

Second, system dynamics needs to be understood. Understanding the dynamics of the system is an essential aspect of the study in systems biology. This requires integrative efforts of experiments, measurement technology development, computational model development, and theoretical analysis. Various methods, such as bifurcation analysis, have been used, but further investigations are necessary to handle the dynamics of systems with very high dimensional space.

Third, methods to control the system shall be investigated. One of the implications is to find a therapeutic approach based on system-level understanding. Many drugs have been developed through extensive effect-oriented screening. It is only recently that specific molecular targets have been identified and lead compounds are designed accordingly. Success in control methods of cellular dynamics may enable us to exploit intrinsic dynamics of the cell so that its effects can be precisely predicted and controlled.

Finally, designing the system that is to modify and construct biological system with designed features. Bacteria and yeast may be redesigned to yield desired properties for drug production and alcohol production. Artificially created gene regulatory logic could be introduced and linked to innate genetic circuits to attain desired functions [5].

Several different approaches can be taken within systems biology field. One may decide to carry out a large-scale, high-throughput experiment and try to find out the overall picture of the system at coarse-grain resolution [6-9]. Alternatively, working on precise details of specific signal transduction [10,11], cell cycle [12,13], and other biological issues to find out the logic behind them are a viable research approach. Both approaches are essentially complementary, and, together, can reshape our understanding of biological systems.

16.2 ROBUSTNESS IS THE FUNDAMENTAL ORGANIZATIONAL PRINCIPLE OF BIOLOGICAL SYSTEMS

Robustness is a property of the system that maintains a certain function despite external and internal perturbations that are ubiquitously observed in various aspects of biological systems [14]. It is distinctively a system-level property that cannot be observed by just looking at components. Specific aspects of the system, the functions to be maintained, and the types of perturbations that the system is robust against must be well defined to make solid arguments. For example, a modern airplane (system) has a function to maintain its flight path (function) against atmospheric turbulences (perturbations).

Bacteria chemotaxis is one of the most well-documented examples in which chemotaxis is a function maintained against the perturbations that are changes in ligand concentration and rate constants for the interactions involved [15-17]. The network for segmental polarity formation during *Drosophila* embryogenesis robustly produces repetitive stripes of differential gene expressions despite variations in initial concentration of substances involved, as well as kinetic parameters of interactions [18,19]. Various aspects of robustness of biological systems have been studied extensively, but more remains to be explored and formalized to create solid theoretical foundations.

Why is robustness so important? First of all, it is a feature that is observed to be so ubiquitous in biological systems; from such a fundamental process like phage fate decision switch [20] and bacteria chemotaxis [15-17] to developmental plasticity [18] and tumor resistance against therapies [21,22]. This implies that it may be a basis for principles that are universal in biological systems, as well as being opportunistic toward finding cures for cancer and other complicated diseases.

Second, robustness is a system-level property of the system in which interactions of components give rise to this feature. Robustness in this context refers to a feature of the system to maintain its function instead of structures or specific states. Structures or states can be dynamically changed if they lead to maintenance of the function of the system.

Third, robustness against environmental and genetic perturbation is essential for evolvability [23-25]. Evolvability requires generation of variety of nonlethal phenotype and genetic buffering [26,27]. Mechanisms that attain robustness against environmental perturbation may be used also for attaining robustness against mutations, developmental stability, and other features that facilitate evolvability [14,23-25].

Fourth, it is one of the features that distinguish biological systems and man-made engineering systems. Although some man-made systems, such as airplanes, are designed to be robust against the range of perturbations, most man-made systems are not as robust as biological systems. Some engineering systems that are designed to be highly robust entail mechanisms that are also present in life forms, which imply existence of the universal principle.

16.3 UNDERLYING MECHANISMS FOR ROBUSTNESS

16.3.1 System Control

First, extensive systems control is used, mostly saliently negative feedback loops but also feedforward and positive feedback controls, to make a system dynamically stable around the specific state of the system. An integral feedback is used in bacteria with chemotaxis as a typical example [15–17]. Due to integral feedback, bacteria can sense changes of chemoattractant and chemorepellant independent of absolute concentration so that proper chemotaxis behavior is maintained over a wide range of ligand concentration. In addition, the same mechanism makes it insensitive to changes in rate constants involved in the circuit. Positive feedbacks are often used to create bistability in signal transduction and cell cycle, so that the system is tolerant to minor perturbation in the stimuli [10,12,13].

16.3.2 Fault Tolerance (Redundancy and Diversity)

Second, fault tolerance mechanisms increase tolerance against components failure and environmental changes by providing alternative components or methods to ultimately maintain a function of the system. Sometimes there are multiple components that are similar to each other and are redundant. Other cases are different means that they are used to cope with perturbations that cannot be handled by the other means. This is often called phenotypic plasticity [28,29] or diversity. Redundancy and phenotypic plasticity are often considered as opposite things, but it is more consistent to view them as different ways to meet an alternative fail-safe mechanism.

16.3.3 Modularity

Third, modularity provides isolation of perturbation from the rest of the system. The cell is the most significant example. More subtle and less obvious examples are modules of biochemical and gene regulatory networks. Modules also play an important role during developmental processes that buffer perturbations so that proper pattern formation can be accomplished [18,30,31]. The definition of the module and the methods of how to detect such modules are still controversial, but the general consensus is that the module does exist and play an important role [32].

16.3.4 Decoupling (Buffering)

Fourth, decoupling isolates low-level noise and fluctuations from functional-level structures and dynamics. One example here is genetic buffering by Hsp90 in which misfolding of proteins due to environmental stresses is fixed, and thus effects of such perturbations are isolated from the functions of the circuits. This mechanism also applies to genetic variations where genetic changes in coding region that may affect protein structures are masked because protein folding is fixed by Hsp90, unless such masking is removed by extreme stress [24,33,34]. Emergent behaviors of complex

networks also exhibit such buffering properties [35]. These effects may constitute canalization proposed by Waddington [36]. A recent discovery by Uri Alon's group on oscillatory expression of p53 upon DNA damage may exemplify decoupling at signal-encoding level [37], because stimuli invoked pulses of p53 activation level, instead of gradual changes, effectively converting analogue into digital signal. Digital pulse encoding may indicate robust information transmission, although further investigations are clearly warranted to draw any conclusion at this moment.

An example of a sophisticated engineering system clearly illustrates how these mechanisms work as a whole system. An airplane is supposed to maintain a flight path following the command of the pilot against atmospheric perturbations and various internal perturbations, including changes in the center of gravity due to fuel consumption and movement of passengers, as well as mechanical inaccuracies. This function is carried out by controlling flight control surfaces (rudder, flaps, elevators, etc.) and a propulsion system (engines) by an automatic flight control system (AFCS). Extensive negative feedback control is used to correct deviations of flight path. The reliability of the AFCS is critically important for stable flight. To increase reliability, the AFCS is composed of three independently implemented modules (a triple redundancy system) all of which meet the same functional specification. Most parts of the AFCS are digitalized, so that low-level noise of voltage fluctuations is effectively decoupled from digital signals that define the function of the system. Due to these mechanisms, modern airplanes are highly robust against various perturbations.

16.4 INTRINSIC FEATURES OF ROBUST SYSTEMS: EVOLVABILITY AND TRADE-OFFS

For the system to be evolvable, it must be able to produce variety of nonlethal phenotypes [27]. At the same time, genetic variations need to be accumulated as a neutral network so that pools of genetic variants are exposed when the environment suddenly changes. Systems that are robust against environmental perturbations entail mechanisms such as system control, alternative, modularity, and decoupling that also support, by congruence, generation of nonlethal phenotype and genetic buffering. In addition, the capability to generate flexible phenotype and robustness requires the emergence of the bow tie structure as an architectural motif [38]. One of the reasons why robustness in biological systems is so ubiquitous is that it facilitates evolution, and evolution tends to select traits that are robust against environmental perturbations. This leads to successive addition of system controls.

Systems that acquire robustness against certain perturbations through design or evolution have intrinsic trade-offs between robustness, fragility, performance, and resource demands. Carlson and Doyle argued, using simple examples from physics and forest fire, that systems that are optimized for specific perturbations are extremely fragile against unexpected perturbations [39,40]. A system that has been designed, or evolved, optimally (either globally optimal or suboptimal) against certain perturbations is called a high optimized tolerance (HOT) system. Ceste and Doyle further

argued that robustness is a conserved quantity [41]. This means when robustness is enhanced against a range of perturbations, it must then be paid off by fragility elsewhere as well as compromised performance and increased resource demands.

Robust-yet-fragile trade-offs can be understood intuitively using the airplane example yet again. When comparing modern commercial airplanes with the Wright Flyer, modern commercial airplanes are, by a great magnitude, more robust against atmospheric perturbations than the Wright flyer, and are thus attributed to a sophisticated flight control system. However, such a flight control system fully relies on electricity. In a very unthinkable event of total power failure in which all electricity is lost in the airplane, the airplane cannot be controlled at all. Obviously, airplane manufacturers are well aware of this issue and take all possible counter measures to minimize such a risk. On the other hand, despite its vulnerability against atmospheric perturbations, the Wright flyer will never be affected by the power failure because there is no reliance on electricity. This extreme example illustrates that systems that are optimized for certain perturbations could be extremely fragile against unusual perturbations.

HOT model systems are successively optimized/designed (not necessarily globally optimized, though) against perturbations in contrast to self-organized criticality (SOC) [42] or scale-free networks [43] that are unconstrained stochastic additions of components without design or optimization involved. Such differences actually affect failure patterns of the system, and thus have direct implications on understanding the nature of disease and therapy design.

Unlike scale-free networks, HOT systems are robust against perturbations like removal of hubs as far as systems are optimized against such perturbations. However, systems are generally fragile against "Fail-on" type failure in which component's failure results in continuous malfunction, instead of cease to function "Fail-off," so that incorrect signals are kept transmitted. This type of failure is known in the engineering field as the Byzantine Generals Problem [44], named after the problem in the Byzantine army composed of numbers of generals dispersed in the field, some of them traitors who sent incorrect messages to confuse the army.

Disease often reflects the systemic failure of the system triggered by the fragility of the system. Diabetes mellitus is an excellent example of how systems that are optimized for near-starving, intermittent food supply, high energy utilization lifestyle, and highly infectious conditions are fragile against unusual perturbations such as high energy containing foods, and a low energy utilization lifestyle [45]. Due to optimization toward a near-starving condition, the extensive control to maintain a minimum blood glucose level is acquired so that activities of central neural systems and innate immunity are maintained. However, no effective regulatory loop has been developed against excessive energy intake and feedback regulations work to reduce glucose uptake by adipocyte and skeletal muscle cells because it may reduce plasma glucose level below the acceptable level. These mechanisms lead to a state where blood glucose level is chronically maintained higher than the desired level, from the longer time scale that has not been optimized for, further leading to cardiovascular complications. Similar observations have been made for autoimmune disorders where the

evolution of robust immunity also entails proinflammatory and hyperactive immune system [46].

16.5 SELF-EXTENDING SYMBIOSIS

So far, robustness and its relationship with evolution have been argued within the framework of Mendel's genetics in a sense that mutation and crossover through mating has been considered as a mechanism for evolutionary innovations. Emergence of specific mechanisms for increasing robustness and enrichment of bow-tie structure has been discussed within this paradigm. I have previously proposed that there may be other means of enhancing robustness through evolution, but by extending "self" with foreign biologic substances, a notation that I termed "self-extending symbiosis" [47]. Self-extending symbiosis is a phenomenon where evolvable robust systems continue to extend their system boundary by incorporating foreign biologic forms (genes, microorganisms, etc.) to enhance their adaptive capability against environmental perturbations, hence improving their survivability and reproduction potential. In other words, robust evolvable systems have consistently extended themselves by incorporating nonself into tightly coupled symbiotic states.

Looking at the history of evolutionary innovations, it has become clear that some of the major innovations are the result of acquisition of "nonself" into "self" at various levels. Horizontal gene transfer (HGT) facilitates evolution by exchanging genes of different species that have evolved for different optimization contexts, and was shown to be a frequently observed phenomenon in prokaryotes, archaea, and unicellular eukaryotes [48,49]. Microorganisms acquire novel functions, mostly to enhance their robustness against environmental challenges, through horizontal exchange of genes. For example, it has been argued that global emergence of antibiotic-resistant bacteria may be caused by horizontal transfer of antibiotic genes [50-52]. In metazoan species, HGT has not been reported (at best, reported highly controversially) except in some rare instances on insect-bacteria symbiosis between the adzuki bean beetle *Callosobruchus chinensis* and *Wolbachia* [53].

The serial endosymbiosis theory by Lynn Margulis [54,55] argues that eukaryotic cells have been created by acquiring bacteria as their organelles. This resulted in greater functionalities of eukaryotic cells, hence more robust against environmental challenges. Here, symbiosis resulted in incorporation of foreign biologic entity into cytoplasm as well as into its own genome.

While HGT and endosymbiosis resulted in incorporation of foreign biologic entity into genome and cellular structure, there are forms of symbiosis that do not directly alter genome but essential to the survival of the species. There are species that allow certain bacteria to be vertically inherited through the host's oocytes as observed in sponges, clams [56], and aphids [57]. Aphids, for example, are infected with the genus *Buchnera*, resulting in an endosymbiotic relationship and acquired dramatically improved energy utilization and terrain exploration capability. It was shown that aphids and *Buchnera* undergo parallel evolution where the phylogeny trees of the host (aphids) and symbionts (genus *Buchnera*) are consistent [57]. A case

of parallel evolution has also been observed in endosymbiosis of *Psyllid* and *Candidatus* [58].

Apart from such tight coupling of host and symbiont, horizontal (environmental) acquisition of symbionts [59] is yet another approach in extending the self by incorporating a broader range of microbes, thereby allowing the host to be able to adapt to a broader range of environments and nutrients. Commensal bacterial flora are ubiquitously observed in various metazoan species, including termites [60], cockroaches [61], prawns [62], and mammals, and have established inseparable relationships with the host organisms, and are even considered to have coevolved [63]. In human beings, the commensal bacterial flora in the gut consists of diverse microorganisms up to 500–1000 species, amounting to about 10^{14} bacteria weighing a total of 1.5 kg [64]. The human being as a symbiotic system consists of approximately 90 percent prokaryotes and 10 percent eukaryotes [65], and a random shotgun sequencing of the whole human symbiotic system would result in predominantly bacterial genome readouts of about 2 million genes with sporadic mammalian genes [66]. Such commensal intestinal bacteria play a critical role in various aspects of the host physiology. Mammalian bacterial flora has been considered to constitute an integral part of host protection by mutually beneficial symbiosis with the host immune system.

The line of observations point to the characteristic property of biological systems that the greater levels of robustness and functionalities is gained by incorporating foreign biologic entities into their own system in the form of different degree of symbiosis. HGT and endosymbiosis incorporate foreign entities into genome and cellular structures, where vertical inheritance based endosymbiosis do not directly alter the genome. Bacterial flora simply adds a layer of adaptive system that is symbiotically interacting with mucosal immune system of the host. A general tendency observed here is the continuous addition of external layers by symbiotic incorporation of foreign entities, and increased level of robustness against environmental perturbation is gained in this process.

16.6 CANCER AS A ROBUST SYSTEM

Cancer is a heterogeneous and highly robust disease that represents worse case scenario of system failure; a fail-on fault where malfunction components are protected by mechanisms that support robustness in normal physiology [21,22]. It is a robustness hijack. Survival and proliferation capability of tumor cells are robustly maintained against a range of therapies due to intratumoral genetic diversity, feedback loops for multidrug resistance, tumor–host interactions, and so on.

Intratumoral genetic heterogeneity is a major source of robustness in cancer cells. Chromosome instability facilitates generation of intratumoral genetic heterogeneity through gene amplification, chromosomal translocation, point mutations, aneuploidy, and so on [67–70]. Intratumoral genetic heterogeneity is one of the most important features of cancer that provides alternative, or fail-safe mechanisms for tumor to survive and grow again despite various therapies, because some tumor cells may have

genetic profile that are resistant to the therapies carried out. Although there are only a few studies on intratumoral genetic heterogeneity, available observations in certain types of solid tumors indicate that there are multiple subclusters of tumor cells within one tumor cluster in which each subcluster has different chromosomal aberrations [71–75]. This implies that each subcluster is developed as clonal expansion of a single mutant cell, and creation of a new subcluster depends upon the emergence of a new mutant that is viable for clonal expansion. A computational study demonstrates that spatial distribution within a tumor cluster enables the coexistence of multiple subclusters [76].

Multidrug resistance is a cellular-level mechanism that provides robustness of viable tumor cell against toxic anticancer drugs. In general, this mechanism involves overexpression of genes such as MDR1 that encodes ATP-dependent efflux pump, P-glycoprotein (P-gp) that effectively pumps out broad range of cytotoxins [77,78]. Trials to mitigate function of P-gp using verapamil, cyclosporine its derivative PSC833 have been disappointing [79].


Tumor–host interactions play major roles in tumor growth and metastasis [80]. When tumor growth is not balanced by vascular growth, hypoxic condition emerges in a tumor cluster [81]. This triggers HIF-1 upregulation that induces a series of reactions that normally function to maintain normal physiological conditions [82]. Upregulation of HIF-1 induces upregulation of VEGF that facilitates angiogenesis, and uPAR and other genes that enhance cell motility [81]. These responses solve hypoxia of tumor cells either by providing oxygen to tumor cluster or by moving tumor cells to a new environment—resulting in further tumor growth or metastasis. Interestingly, macrophages are found to chemotaxis into tumor cluster. Such macrophages are called tumor-associated macrophage (TAM), and found to over-express HIF-1[83]. This means that the macrophage that is supposed to remove tumor cells may be built-in to feedback loops to facilitate tumor growth and metastasis.

In addition, it can be considered that tumor cells may evolve through self-extending symbiosis. If this is the case, tumor cells shall enhance their robustness against various perturbations through horizontal gene transfer, symbiosis with other cells in the form of cell fusion, and formation of symbiotic relationship with surrounding environments. Interestingly, recent reports indicate that tumor cells may be actively involved in cell fusion and uptake of chromosomes of other cells [84–87]. In addition, artificially produced hybridomas between antibody-producing plasma cell and tumor cell are used for monoclonal antibody production indicating stable maintenance of cellular function upon hybridization. These series of observations imply that tumor cells may be considered as a group of cells that have become somewhat detached from the host system and have begun evolving independently, so that a wide range of phenomena, such as self-extending symbiosis, also occur on tumor cells and thereby their robustness against perturbation is enhanced (Fig. 16-1).

So far, such phenomena have only been reported independently, and not been placed in the perspective. Reorganizing these findings under the coherent view of cancer robustness will provide us a guideline for further research.

Self-extending symbiosis

(a)



Cancer self-extending symbiosis

(b)




Figure 16-1 Self-extending symbiosis and cancer evolution. Self-extending symbiosis is a path that multicellular organism might have gone through in the course of evolution. Acquisition of oneself into self at various levels of flexibility enhances robustness of organisms against various perturbations. Cancer may also evolve through self-extending symbiosis. Assume cancer as an independent species diverted from somatic cell, it may rapidly evolve through bacteria-like horizontal gene transfer, cell fusion, and microenvironment remodeling to enhance robustness against environmental perturbation. In self-extending symbiosis, there is clear evidence of oocyte-mediated vertical infection. There is no conclusive report if any bacterial infection is observed in any type of cancer that affects robustness of cancer against perturbation. Such phenomena might have been simply unnoticed waiting for future discovery.

16.7 THEORETICALLY MOTIVATED THERAPY STRATEGIES

Given the highly complex control and heterogeneity of tumor, random trial of potential targets is not as effective as one wish it to be. There is a need for the theoretically motivated approach that guides us to identify a set of therapies to best counter the disease. The implication of the theory of cancer robustness is that there are specific patterns of behaviors and weakness in robust systems as well as rational way of controlling and fixing the system, and such general principles also apply to cancer. Thus, there must be theoretically motivated approach for the prevention and treatment of cancer. This section discusses therapeutic implications of the theory.

Strategy for cancer therapy may depend upon the level of robustness that the tumor of a specific patient has. When robustness is low, and genetic heterogeneity is low, then there is a good chance that the use of drugs with specific molecular targets may effectively cure cancer by causing the common mode failure: a type of failure in which all redundant subsystems fail for the same reason. An example of CML (chronic myeloid leukemia) therapy by imatinib methylate (Glivec; Novartis) may provide us some insights [88,89]. Dramatic effects of imatinib methylate for early stage CML stem from the fact that it selectively target BCR-ABL protein that is specifically expressed in tumor cells and tumor growth depends on BCR-ABL [90]. Thus, it causes the common mode failure in tumor cells that have similar fragility. However, it is resistant in advanced stage due to heterogeneity of mutations so that the drug cannot inhibit diverse emergent mutant proteins [91]. For this strategy to be effective, there must be proper means to diagnose the degree of intratumoral genetic variations. Then, the most effective molecule as a target needs to be recognized that directs the lead identification and optimization processes.

However, for patients with an advanced stage cancer, intratumoral genetic heterogeneity may be already high and various feedback controls may be significantly upregulated. In these cases, drugs that are effective in the early stage may not work as expected, due to heterogeneous response of tumor cells and feedbacks to compensate for perturbations. For these cases, therapy and drug design need a drastic shift from molecule-oriented approach to a system-oriented approach. Then, the question is which approach shall be taken to target the system, instead of the molecule. I would consider that there are three theoretically motivated countermeasures.

First, robustness/fragility trade-off implies that the cancer cells that have gained increased robustness against various therapies may have a point of extreme fragility. Targeting such a point of fragility may bring dramatic effects for the disease. The major challenge is to find such a point of fragility. Since this trade-off emerged due to successive modifications of the system design to optimally cope with specific perturbations, it is essential to identify the perturbations that the system is optimized against and the underlying mechanisms that enable such optimization. For example, one mechanism for tumor robustness is enhanced genetic heterogeneity that is generated by chromosomal instability, so that some cells may have genetic profile suitable for survival under the specific pressure from the therapy. Then, a method to enhance chromosomal instability selectively in cells that already have unstable chromosome could be one candidate. The point here is whether such

effects can be done with sufficient selectivity. Nonselective approach to increase chromosomal instability has been proposed [92], but it may enhance chromosome instability of the cells that are relatively stable now and thus potentially promotes malignancy.

Second, approaches that avoid the increase of robustness constitute the other possibility. Since genetic heterogeneity is enhanced, at least in part, by somatic recombination, selectively inducing cell cycle arrest to tumor cells can effectively control the robustness. There is a theoretical possibility that such subtle control can be done by careful combination of multiple drugs that specifically perturb biochemical interactions. A computational study indicates that the removal or attenuation of specific feedback loops involved in cell cycle reduces the robustness of cell cycle against changes in rate constant [93]. The challenge is to find appropriate combination of drugs that can effectively induce cell cycle arrest only in tumor cells, but not in other cells. Although this approach uses combination of multiple drugs, there is hope to find a set of drugs that can be administered at minimum dosage and toxicity. This approach results in the dormancy of the tumor. Cancer dormancy has already been proposed [94,95] and many report that induced dormancy has been found in mouse [96,97]. However, these studies report cases where tumor cell proliferation is offset by increased apoptosis. Since heterogeneity may increase by cell proliferation, this type of dormancy, which I call "pseudo dormancy" does not prevent increase in heterogeneity, hence robustness is not controlled. Genuine dormancy needs to induce selective cell cycle arrest.

Third, an approach to actively reduce intratumoral genetic heterogeneity followed by a therapy by molecular targeted drugs may be a viable option. If we can design an initial therapy to impose a specific selection pressure on the tumor in which there are only cells with specific genetic variations to survive the therapy, then reduction of genetic heterogeneity may be achieved. Then, if a tumor cell population is sufficiently homogeneous, a drug that specifically targets a certain molecule may have significant impact on the remaining tumor cell population. An important point here is that the drugs used shall not enhance mutation and chromosomal instability. If mutations and chromosomal instability are enhanced, particularly by the initial therapy, heterogeneity may quickly increase so that the second line therapy will be ineffective.

Fourth, one may wish to retake control of the feedback loops that give rise to robustness in an epidemic state. Since the robustness of tumor is often caused by host tumor feedback controls, robustness of tumor can be seriously mitigated if such feedback loops can be controlled. One possible approach is to introduce a decoy that effectively disrupts feedback control or invasive mechanisms of the epidemic. Such an approach is proposed in AIDS therapy where conditionally replicating HIV-1 (or HIV-1) vector that has only *cis* region but no *trans* is introduced [98,99]. This decoy virus dominates the replication machinery, so that HIV-1 virus is pushed into latency, instead of eradication. In solid tumor, an interesting idea has been expressed to use TAM as delivery vehicle of the vector [83,100]. TAM migrates into solid tumor cluster and upregulates HIF-1 that facilitates angiogenesis and metastasis. If TAM can be used to retake a control, robustness may be well controlled and self-extending symbiosis in cancer evolution may be aborted.

Finally, multicomponent drugs may be designed where each component targets molecule in which perturbation differentially affects tumor and normal cells. A certain perturbation affects more the tumor cells but less the normal cells. Even if each of such perturbations does not eliminate tumor cells or is not able to stop their proliferation, there may be specific combination of such drugs that in synergy affects drastically and selectively the tumor cells. One extreme of such approach is the "long-tail drug," recently proposed by the author, that uses large numbers of weakly interacting compounds to affect the tumor cells [101].

16.8 A PROPER INDEX OF TREATMENT EFFICACY

It is important to recognize that, in the light of cancer robustness theory, tumor mass reduction is not an appropriate index for therapy and drug efficacy judgment. As discussed already, reduction of tumor mass does not mean that proliferation potential of tumor has generally decreased. It merely means that subpopulation of tumor cells that are respondent of the therapy were eradicated, or significantly reduced. The problem is that the remaining tumor cells may be more malignant and aggressive, so that therapies for relapsed tumor could be extremely ineffective. This is particularly the case, drugs used to reduce tumor mass are toxic and potentially promote mutations and chromosomal instability in nonspecific ways. It may even enhance malignancy but imposing selective pressures to select resistant phenotype, enhance genetic diversity, as well as providing niche for growth by eradicating fragile subpopulation of tumor cells.

The proper index shall be based on control of robustness: either minimizes the increase of robustness or reduces robustness. This can be achieved by inducing dormancy, actively imposing selective pressure to reduce heterogeneity or exposing fragility that can be the target of therapies to follow, and retaking control of the feedback regulations. The outcome of controlling the robustness may vary from moderate growth of tumor, dormancy that is no tumor mass growth or significant reduction in tumor mass. It should be noted that robustness control does not exclude the possibility of significant tumor mass reduction. If we can target a point of fragility of tumor, it may trigger a common mode failure and may result in significant tumor mass reduction. However, this is a result of controlling robustness, and should not be confused as therapy aimed that tumor mass reduction because robustness has to be controlled to the first to actively exploit a point of fragility. Except for the fragility attack, other options seek for dormancy that results in no tumor growth.

However, this criterion poses a problem for drug design, because current efficacy index of antitumor drugs is measured on the basis of tumor mass reduction. Drugs that induce dormancy will not satisfy this efficacy criterion; thus, they are most likely to be rejected in Phase-II stage. On the other hand, this means that many compounds that have been rejected in Phase-II could be effective from the point of robustness control. Whether such approach can be taken depends on perception change in practitioners, drug industries, and regulatory authorities.

16.9 CONCLUSION

This chapter discussed basic ideas behind the theory of biological robustness and its implications for cancer research and treatment. Biological robustness is one of the essential features of living systems that argued to be tightly coupled with evolution. It may also shape the basic architectural feature of biological systems that are robust and evolvable. One of major consequences is trade-offs between robustness, fragility, resource demands, and performance. Fragility is particularly relevant to diseases. At the same time, cancer established its own robustness. It may be the result of hijacking the robustness intrinsic to the host system. Understanding of this complex nature of biological systems may have profound implications for biomedical research in future.

ACKNOWLEDGMENTS

This study is supported by ERATO-SORST Program (Japan Science and Technology Agency) and the Genome Network Project (Ministry of Education, Culture, Sports, Science, and Technology) of the Systems Biology Institute, the Center of Excellence Program, and the special coordination funds (Ministry of Education, Culture, Sports, Science, and Technology) to Keio University.

REFERENCES

1. Kitano H. Systems biology: a brief overview. *Science* 2002;295(5560):1662–1664.
2. Kitano H. Computational systems biology. *Nature* 2002;420(6912):206–210.
3. Wiener N. *Cybernetics: or Control and Communication in the Animal and the Machine*. Cambridge, MA: The MIT Press, 1948.
4. Bertalanffy, LV. *General System Theory*, New York: George Braziller, 1968.
5. Hasty J, McMillen D, Collins JJ. Engineered gene circuits. *Nature* 2002;420(6912):224–230.
6. Guelzim N, Bottani S, Bourgoin P, Kepes F. Topological and causal structure of the yeast transcriptional regulatory network. *Nat Genet* 2002;31(1):60–63.
7. Ideker T, Ozier O, Schwikowski B, Siegel AF. Discovering regulatory and signaling circuits in molecular interaction networks. *Bioinformatics* 2002;18(Suppl 1): S233–S240.
8. Ideker T, Thorsson V, Ranish JA, Christmas R, Buhler J, Eng JK, Bumgarner R, Goodlett DR, Aebersold R, Hood L. Integrated genomic and proteomic analyses of a systematically perturbed metabolic network. *Science* 2001;292(5518):929–934.
9. Ihmels J, Friedlander G, Bergmann S, Szig O, Ziv Y, Barkai N. Revealing modular organization in the yeast transcriptional network. *Nat Genet* 2002;31(4):370–377.
10. Ferrell JE Jr. Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr Opin Cell Biol* 2002;14(2):140–8.
11. Bhalla US, Iyengar R. Emergent properties of networks of biological signaling pathways. *Science* 1999;283(5400):381–387.

12. Tyson JJ, Chen K, Novak B. Network dynamics and cell physiology. *Nat Rev Mol Cell Biol* 2001;2(12):908–916.
13. Chen KC, Calzone L, Csikasz-Nagy A, Cross FR, Novak B, Tyson JJ. Integrative analysis of cell cycle control in budding yeast. *Mol Biol Cell* 2004;15(8):3841–3862.
14. Kitano H. Biological robustness. *Nat Rev Genet* 2004;5:826–837.
15. Alon U, Surette MG, Barkai N, Leibler S. Robustness in bacterial chemotaxis. *Nature* 1999;397(6715):168–171.
16. Barkai N, Leibler S. Robustness in simple biochemical networks. *Nature* 1997;387(6636):913–917.
17. Yi TM, Huang Y, Simon MI, Doyle J. Robust perfect adaptation in bacterial chemotaxis through integral feedback control. *Proc Natl Acad Sci USA* 2000;97(9):4649–4653.
18. von Dassow G, Meir E, Munro EM, Odell GM. The segment polarity network is a robust developmental module. *Nature* 2000;406(6792):188–192.
19. Ingolia NT. Topology and robustness in the *Drosophila* segment polarity network. *PLoS Biol* 2004;2(6):E123.
20. Little JW, Shepley DP, Wert DW. Robustness of a gene regulatory circuit. *Embo J* 1999;18(15):4299–4307.
21. Kitano H. Cancer as a robust system: implications for anticancer therapy. *Nat Rev Cancer* 2004;4(3):227–235.
22. Kitano H. Cancer robustness: tumour tactics. *Nature* 2003;426(6963):125.
23. Wagner GP, Altenberg L. Complex adaptations and the evolution of evolvability. *Evolution* 1996;50(3):967–976.
24. Rutherford SL. Between genotype and phenotype: protein chaperones and evolvability. *Nat Rev Genet* 2003;4(4):263–274.
25. de Visser J, Hermissin J, Wagner GP, Meyers L, Bagheri-Chaichian H, Blanchard J, Chao L, Cheverud J, Elena S, Fontana W, et al. Evolution and detection of genetics robustness. *Evolution* 2003;57(9):1959–1972.
26. Gerthart J, Kirschner M. *Cells, Embryos, and Evolution: Toward a Cellular and Developmental Understanding of Phenotypic Variation and Evolutionary Adaptability*. Malden Massachusetts: Blackwell Science, 1997.
27. Kirschner M, Gerhart J. Evolvability. *Proc Natl Acad Sci USA* 1998;95(15):8420–8427.
28. Agrawal AA. Phenotypic plasticity in the interactions and evolution of species. *Science* 2001;294(5541):321–326.
29. Schlichting C, Pigliucci M. *Phenotypic Evolution: A Reaction Norm Perspective*. Sunderland, MA: Sinauer Associates, Inc., 1998.
30. Eldar A, Dorfman R, Weiss D, Ashe H, Shilo BZ, Barkai N. Robustness of the BMP morphogen gradient in *Drosophila* embryonic patterning. *Nature* 2002;419(6904):304–308.
31. Meir E, von Dassow G, Munro E, Odell GM. Robustness, flexibility, and the role of lateral inhibition in the neurogenic network. *Curr Biol* 2002;12(10):778–786.
32. Schlosser G, Wagner G, editors. *Modularity in Development and Evolution*. Chicago: The University of Chicago Press, 2004.
33. Rutherford SL, Lindquist S. Hsp90 as a capacitor for morphological evolution. *Nature* 1998;396(6709):336–342.

34. Queitsch C, Sangster TA, Lindquist S. Hsp90 as a capacitor of phenotypic variation. *Nature* 2002;417(6889):618–624.
35. Siegal ML, Bergman A. Waddington's canalization revisited: developmental stability and evolution. *Proc Natl Acad Sci USA* 2002;99(16):10528–10532.
36. Waddington CH. *The Strategy of the Genes: A Discussion of Some Aspects of Theoretical Biology*. New York: Macmillan, 1957.
37. Lahav G, Rosenfeld N, Sigal A, Geva-Zatorsky N, Levine AJ, Elowitz MB, Alon U. Dynamics of the p53-Mdm2 feedback loop in individual cells. *Nat Genet* 2004;36(2):147–150.
38. Cséte ME, Doyle J. Bow ties, metabolism and disease. *Trends Biotechnol* 2004;22(9):446–450.
39. Caulson JM, Doyle J. Highly optimized tolerance: a mechanism for power laws in designed systems. *Phys Rev E Stat Phys Plasmas Fluids Relat Interdiscip Topics* 1999;60(2 Part A):1412–1427.
40. Carlson JM, Doyle J. Complexity and robustness. *Proc Natl Acad Sci USA* 2002;99(Suppl 1): 2538–2545.
41. Cséte ME, Doyle JC. Reverse engineering of biological complexity. *Science* 2002;295(5560):1664–1669.
42. Bak P, Tang C, Wiesenfeld K. Self-organized criticality. *Physical Review A* 1988;38(1):364–374.
43. Barabasi AL, Oltvai ZN. Network biology: understanding the cell's functional organization. *Nat Rev Genet* 2004;5(2):101–113.
44. Lamport L, Shostak R, Pease M. The Byzantine generals problem. *ACM Trans on Program Lang Syst* 1982;4(3):382–401.
45. Kitano H, Kimura T, Oda K, Matsuka Y, Cséte ME, Doyle J, Muramatsu M. Metabolic syndrome and robustness trade-offs. *Diabetes* 2004;53(Suppl 3): S1–S10.
46. Kitano H, Oda K. Robustness trade-offs and host? Microbial symbiosis in the immune system. *Mol Syst Biol* 2006;doi:10. 1038/msb4100039.
47. Kitano H, Oda K. Self-extending symbiosis: a mechanism for increasing robustness through evolution. *Biol Theory* 2006;1(1):61–66.
48. Gogarten JP, Townsend JP. Horizontal gene transfer: genome innovation and evolution. *Nat Rev Microbiol* 2005;3(9):679–687.
49. Brown JR. Ancient horizontal gene transfer. *Nat Rev Genet* 2003;4(2):121–132.
50. Monroe S, Polk R. Antimicrobial use and bacterial resistance. *Curr Opin Microbiol* 2000;3(5):496–501.
51. Levy SB, Marshall B. Antibacterial resistance worldwide: causes, challenges and responses. *Nat Med* 2004;10(Suppl 12):S122–S129.
52. Smeets BF, Barkay T. Horizontal gene transfer: perspectives at a crossroads of scientific disciplines. *Nat Rev Microbiol* 2005;3(9):675–678.
53. Kondo N, Nikoh N, Ijichi N, Shimada M, Fukatsu T. Genome fragment of *Wolbachia* endosymbiont transferred to X chromosome of host insect. *Proc Natl Acad Sci USA* 2002;99(22):14280–14285.
54. Margulis L. Symbiosis and evolution. *Sci Am* 1971;225(2):48–57.
55. Margulis L, Bermudes D. Symbiosis as a mechanism of evolution: status of cell symbiosis theory. *Symbiosis* 1985;1:101–24.
56. Cary SC, Giovannoni SJ. Transovarial inheritance of endosymbiotic bacteria in clams inhabiting deep-sea hydrothermal vents and cold seeps. *Proc Natl Acad Sci USA* 1993; 90(12):5695–5699.
57. Baumann P, Baumann L, Lai CY, Roubbaksh D, Moran NA, Clark MA. Genetics, physiology, and evolutionary relationships of the genus *Buchnera*: intracellular symbionts of aphids. *Annu Rev Microbiol* 1995;49:55–94.
58. Thao ML, Clark MA, Burkhardt DH, Moran NA, Baumann P. Phylogenetic analysis of vertically transmitted psyllid endosymbionts (*Candidatus Carsonella riddii*) based on apAGD and rpoC: comparisons with 16S-23S rDNA-derived phylogeny. *Curr Microbiol* 2001;42(6):419–421.
59. Nyholm SV, McFall-Ngai MJ. The winnowing: establishing the squid–vibrio symbiosis. *Nat Rev Microbiol* 2004;2(8):632–642.
60. Schmitt-Wagner D, Friedrich MW, Wagner B, Brune A. Phylogenetic diversity, abundance, and axial distribution of bacteria in the intestinal tract of two soil-feeding termites (*Cubitermes* spp.). *Appl Environ Microbiol* 2003;69(10):6007–6017.
61. Bracke JW, Cruden DL, Markovetz AJ. Intestinal microbial flora of the American cockroach. *Periplaneta americana* L. *Appl Environ Microbiol* 1979;38(5):945–955.
62. Oxley AP, Shipton W, Owens L, McKay D. Bacterial flora from the gut of the wild and cultured banana prawn. *Pentaeus merguensis*. *J Appl Microbiol* 2002;93(2):214–223.
63. Backhed F, Ley R, Sonnenburg J, Peterson D, Gordon JI. Host-bacterial mutualism in the human intestine. *Science* 2005;307:1915–1920.
64. Xu J, Gordon JI. Inaugural article: honor thy symbionts. *Proc Natl Acad Sci USA* 2003; 100(18):10452–10459.
65. Savage DC. Microbial ecology of the gastrointestinal tract. *Annu Rev Microbiol* 1977;31:107–133.
66. Hooper LV, Midtvedt T, Gordon JI. How host–microbial interactions shape the nutrient environment of the mammalian intestine. *Annu Rev Nutr* 2002;22:283–307.
67. Lengauer C, Kinzler KW, Vogelstein B. Genetic instabilities in human cancers. *Nature* 1998;396(6712):643–649.
68. Li R, Sonik A, Stindl R, Rasnick D, Duesberg P. Aneuploidy vs. gene mutation hypothesis of cancer: recent study claims mutation but is found to support aneuploidy. *Proc Natl Acad Sci USA* 2000;97(7):3236–3241.
69. Tischfield JA, Shao C. Somatic recombination redux. *Nat Genet* 2003;33(1):5–6.
70. Rasnick D. Aneuploidy theory explains tumor formation, the absence of immune surveillance, and the failure of chemotherapy. *Cancer Genet Cytogenet* 2002;136(1):66–72.
71. Baisse B, Bouzouene H, Saraga EP, Bosmao FT, Benhattar J. Intra-tumor genetic heterogeneity in advanced human colorectal adenocarcinoma. *Int J Cancer* 2001;93(3):346–352.
72. Fujii H, Yoshida M, Gong ZX, Matsumoto T, Hainano Y, Fukunaga M, Hruban RH, Gabrielson E, Shirai T. Frequent genetic heterogeneity in the clonal evolution of gynecological carcinomas and its influence on phenotypic diversity. *Cancer Res* 2000;60(1):114–120.
73. Gorunova L, Hoglund M, Andren-Sandberg A, Dawicki S, Jin Y, Mitelman F, Johansson B. Cytogenetic analysis of pancreatic carcinomas: intra-tumor heterogeneity and nonrandom pattern of chromosome aberrations. *Genes Chromosomes Cancer* 1998;23(2):81–99.

74. Gorunova L, Dawiskiba S, Andren-Sandberg A, Høglund M, Johansson B. Extensive cytogenetic heterogeneity in a benign retroperitoneal schwannoma. *Cancer Genet Cytogenet* 2001;127(2):148–154.
75. Frigyesi A, Gisselsson D, Mittelman F, Høglund M. Power law distribution of chromosome aberrations in cancer. *Cancer Res* 2003;63(21):7094–7097.
76. Gonzalez-Garcia I, Sole RV, Costa J. Metapopulation dynamics and spatial heterogeneity in cancer. *Proc Natl Acad Sci USA* 2002;99(20):13085–13089.
77. Juliano RL, Ling V. A surface glycoprotein modulating drug permeability in Chinese hamster ovary cell mutants. *Biochim Biophys Acta* 1976;455(1):152–162.
78. Nooter K, Herweijer H, Multidrug resistance (mdr) genes in human cancer. *Br J Cancer* 1991;63(5):663–669.
79. Tsuruo T, Iida H, Tsukagoshi S, Sakurai Y. Overcoming of vincristine resistance in P388 leukemia *in vivo* and *in vitro* through enhanced cytotoxicity of vincristine and vinblastine by verapamil. *Cancer Res* 1981;41(5):1967–1972.
80. Bissell MJ, Radisky D. Putting tumours in context. *Nat Rev Cancer* 2001;1(1):46–54.
81. Harris AL. Hypoxia: a key regulatory factor in tumour growth. *Nat Rev Cancer* 2002;2(1):38–47.
82. Sharp FR, Bernaudin M. HIF1 and oxygen sensing in the brain. *Nat Rev Neurosci* 2004;5(6):437–448.
83. Bingle L, Brown NJ, Lewis CE. The role of tumour-associated macrophages in tumour progression: implications for new anticancer therapies. *J Pathol* 2002;196(3):254–265.
84. Pawelek JM. Tumour-cell fusion as a source of myeloid traits in cancer. *Lancet Oncol* 2005;6(12):988–993.
85. Pawelek J, Chakraborty A, Lazova R, Yilmaz Y, Cooper D, Brash D, Handerson T. Co-opting macrophage traits in cancer progression: a consequence of tumor cell fusion? *Contrib Microbiol* 2006;13:138–155.
86. Vignery A. Macrophage fusion: Are somatic and cancer cells possible partners? *Trends Cell Biol* 2005;15(4):188–193.
87. Ogle BM, Cascalho M, Platt JL. Biological implications of cell fusion. *Nat Rev Mol Cell Biol* 2005;6(7):567–575.
88. Hochhaus A. Cytogenetic and molecular mechanisms of resistance to imatinib. *Semin Hematol* 2003;40(2 Suppl 3): 69–79.
89. Hochhaus A, Kreil S, Corbin A, La Rusec P, Lahaye T, Berger U, Cross NC, Linkesch W, Druker BJ, Hehlmann R, et al. Roots of clinical resistance to STI-571 cancer therapy. *Science* 2001;293(5538):2163.
90. Druker BJ, Talpaz M, Resta DJ, Peng B, Buchdunger E, Ford JM, Lydon NB, Kantarjian H, Capdeville R, Ohno-Jones S, et al. Efficacy and safety of a specific inhibitor of the BCR-ABL tyrosine kinase in chronic myeloid leukemia. *N Engl J Med* 2001;344(14):1031–1037.
91. Druker BJ. Overcoming resistance to imatinib by combining targeted agents. *Mol Cancer Ther* 2003;2(3):225–226.
92. Sole RV. Phase transitions in unstable cancer cell populations. *Eur Phys J B* 2003;35:117–123.
93. Morohashi M, Winn AE, Borisuk MT, Bolouri H, Doyle J, Kitano H. Robustness as a measure of plausibility in models of biochemical networks. *J Theor Biol* 2002;216(1):19–30.
94. Takahashi Y, Nishioka K. Survival without tumor shrinkage: re-evaluation of survival gain by cytostatic effect of chemotherapy. *J Natl Cancer Inst* 1995;87(16):1262–1263.
95. Uhr JW, Scheuermann RH, Street NE, Vittra ES. Cancer dormancy: opportunities for new therapeutic approaches. *Nat Med* 1997;3(5):505–509.
96. Holmgren L, O'Reilly MS, Folkman J. Dormancy of micrometastases: balanced proliferation and apoptosis in the presence of angiogenesis suppression. *Nat Med* 1995;1(2):149–153.
97. Murray C. Tumour dormancy: not so sleepy after all. *Nat Med* 1995;1(2):117–118.
98. Dropic B, Hermankova M, Pitha PM. A conditionally replicating HIV-1 vector interferes with wild-type HIV-1 replication and spread. *Proc Natl Acad Sci USA* 1996;93(20):11103–11108.
99. Weinberger LS, Schaffer DV, Arkin AP. Theoretical design of a gene therapy to prevent AIDS but not human immunodeficiency virus type 1 infection. *J Virol* 2003;77(18):10028–10036.
100. Owen MR, Byrne HM, Lewis CE. Mathematical modelling of the use of macrophages as vehicles for drug delivery to hypoxic tumour sites. *J Theor Biol* 2004;226(4):377–391.
101. Kitano H. A robustness-based approach to systems-oriented drug design. *Nat Rev Drug Discov* 2007;6(3):202–210.

SYSTEMS BIOLOGY AND SYNTHETIC BIOLOGY

Edited by

Pengcheng Fu

China University of Petroleum Beijing, China

Sven Panke

ETH Zurich, Institute of Process Engineering Zurich, Switzerland



WILEY

A JOHN WILEY & SONS, INC., PUBLICATION

CONTENTS

Foreword	ix
Contributors	xi
1 Introduction <i>Pengcheng Fu</i>	1
2 Basics of Molecular Biology, Genetic Engineering, and Metabolic Engineering <i>Michael X. Wang and Huidong Shi</i>	9
3 High-Throughput Technologies and Functional Genomics <i>Xiu-Feng Wan and Dorothea K. Thompson</i>	67
4 Genomic Signal Processing of DNA Microarray Data for the Enhanced Prediction of Axillary Lymph Node Status of Breast Cancer Tumors <i>Gordon S. Okamoto</i>	115
5 Recombinant Genomes: Novel Resources for Systems Biology and Synthetic Biology <i>Mitsuhiko Itaya</i>	155
6 In silico Genome-Scale Metabolic Models: The Constraint-Based Approach and its Applications <i>Andrew R. Joyce and Bernhard Ø. Palsson</i>	193
7 Mathematical Modeling of Genetic Regulatory Networks: Stress Responses in <i>Escherichia coli</i> <i>Delphine Ropers, Hidde de Jong, and Johannes Geiselmann</i>	235
8 Synthetic Life: Ethobricks for a New Biology <i>Jordi Valverde and Claes Gustafsson</i>	273

9	Yeast as a Prototype for Systems Biology <i>Goutham Vemuri and Jens Nielsen</i>	287
10	Construction and Applications of Genome-Scale in silico Metabolic Models for Strain Improvement <i>Sang Yup Lee, Jin Sik Kim, Hongseok Yun, Tae Yong Kim, Seung Bum Sohn, and Hyun Uk Kim</i>	355
11	Synthetic Biology: Putting Engineering into Bioengineering <i>Matthias Heinemann and Sven Panke</i>	387
12	Rationales of Gene Design and de novo Gene Construction <i>Marcus Graf, Thomas Schoebl, and Ralf Wagner</i>	411
13	Self-Replication in Chemistry and Biology <i>Philipp Holliger and David Lookes</i>	439
14	The Synthetic Approach for Regulatory and Metabolic Circuits <i>Wilson W. Wong and James C. Liao</i>	467
15	Synthetic Gene Networks <i>David Greber and Martin Fussenegger</i>	489
16	The Theory of Biological Robustness and its Implication to Cancer <i>Hiroaki Kitano</i>	529
17	Nucleic Acid Engineering <i>WenLong Cheng, Liang Ding, Hisakage Funabashi, Nokyong Park, Soong Ho Um, Jianfeng Xu, and Dan Luo</i>	549
18	Potential Applications of Synthetic Biology in Marine Microbial Functional Ecology and Biotechnology <i>Guangyi Wang and Juanita Mathews</i>	577
19	On Fundamental Implications of Systems and Synthetic Biology <i>Cliff Hooker</i>	593
20	Outstanding Issues in Systems and Synthetic Biology <i>Pengcheng Fu and Cliff Hooker</i>	615
	Index	647

FOREWORD

The popular use of the term “systems biology” arose following the appearance of the first full genome sequences. These genome sequences suggested that we would have a full delineation of the molecular components of an organism. Expression profiling and proteomic data then could tell us when these components were actually used in a context-specific manner.

The need to track the interrelationship of all such components created the need to develop networks of the interactions of such components. Protein–protein interaction maps are one manifestation of this need, stoichiometric models are another; they are, however, amenable to rigorous mathematical analysis and prospective uses. Network reconstruction took center stage in systems biology, as networks describe the interactions between the gene products and the chemical compounds they make, provide context for high-throughput data mapping, and give the basis for mechanistic models that can compute phenotypic functions.

Having molecular manipulation tools and mathematical models in turn provides tools that allow the synthesis of biological components and biological functions. We thus witnessed the emergence of “synthetic biology.” It is practiced on multiple scales, from component design, that is akin to classical molecular biology, to design of whole cell functions, such as metabolic engineering.

Thus, in retrospect we can state that genomics gave rise to systems, and systems biology in turn gave rise to synthetic biology. This of course is a simplified view, but provides a first-order approximation to the historical origin and appearance of these popularly used terms. This volume contains a series of chapters that highlight the development and status of the various aspects of systems and synthetic biology.

BERNHARD Ø. PALSSON

University of California, San Diego, CA

④スダン I : 赤色 219号に含有されていた不純物であるスダン I が原因で、女性の顔面に、初期には発赤、かゆみなどの炎症症状をきたし、やがて紫灰色から紫褐色の色素沈着へと進展するリール黒皮症(女子顔面黒皮症)が1970年代中頃を中心に発生した。その後メーカーの安全性保証の強化に伴い、発生率は減少し、現在ではほとんど認められなくなった。

5.2.7 化粧品の安全管理^{1, 25)}

GVP (Good Vigilance Practice; 製造販売後安全管理基準)に関する省令に基づき、医薬品、医薬部外品、化粧品および医療機器の各々に携わる製造販売業者は、市販後の安全性を担保するために、定められた基準にしたがって、情報の収集、作成、保管および報告を実行することが求められている。とくに、化粧品においては、成分のほとんどを自己責任のもとで配合することができるため、GVPは非常に重要なものといえる。 [森 眞輝]

文 献

- 1) 薬事日報社 化粧品・医薬部外品製造販売ガイドブック検討会 (2006) : 化粧品・医薬部外品 製造販売ガイドブック 2006, 薬事日報社.
- 2) 日本化粧品工業連合会 編 (2008) : 化粧品の安全性評価に関する指針 2008, 薬事日報社.
- 3) 板垣 宏, 萩野滋延 (2008) : 動物実験代替法への化粧品企業における取り組み, フェルマシア, 44, 863-868.
- 4) The international agency for research on cancer (IARC) : IARC monographs on the evaluation of carcinogenic risks to humans, IARC, Lyon.
- 5) Rietschel, R.L. and Fowler, J.F., Jr., Eds. (2001) : Fisher's contact dermatitis 5th ed., Lippincott Williams and Wilkins, Pennsylvania.
- 6) 荒田次郎監修 (2004) : 標準皮膚科学 第7版, 医学書院.
- 7) 独立行政法人 科学技術振興機構 : 日本化学物質辞書 Web, http://nikkajiweb.jst.go.jp/nikkaji_web/pages/top.html
- 8) The national center for biotechnology information (NCBI) : PubMed, <http://www.ncbi.nlm.nih.gov/pubmed/>
- 9) Klaassen, C.D., Ed. (2007) : Casarett and Doull's toxicology - The basic science of poisons - 7th ed., McGraw-Hill, New York.
- 10) Champion, R.H., Burton, J.L. and Ebling, F.J.G., Eds. (1992) : Textbook of dermatology 5th ed., Blackwell Scientific Publications, Oxford.
- 11) 国際化学物質安全性計画 (IPCS) : 国際化学物質安全性カード (ICSC) - 日本語版 -, <http://www.nihs.go.jp/ICSC/>
- 12) Baran, R. and Maibach, H.I., Eds. (1998) : Textbook of cosmetic dermatology 2nd ed., Martin Dunitz, London.
- 13) 日本ビタミン学会編 (1996) : ビタミンの事典, 朝倉書店.
- 14) 田村建夫, 廣田 博 (1999) : 化粧品科学—理論と実際 第3版, フレグランスジャーナル社.
- 15) The U.S. food and drug administration (FDA) : <http://www.fda.gov/>
- 16) 中村 淳, 高野勝弘 (2006) : 化粧品の国際法規制の現状とハーモナイゼーションの課題 国際法規制の現状とグローバルゼイション, フレグランスジャーナル, 1月号, 17-20.
- 17) 高橋理佳 (2006) : 米国の化粧品の法規制の現状と課題 国際法規制の現状とグローバルゼイション, フレグランスジャーナル, 1月号, 21-24.
- 18) The cosmetic ingredient review (CIR) (2007) : 2007 CIR Compendium, CIR, Washington, D.C.
- 19) The national institute for occupational safety and health (NIOSH) : Search the NIOSH Website, <http://www.cdc.gov/niosh/srchpage.html>
- 20) 伊東和子 (2006) : EUにおける化粧品規制とその課題 国際法規制の現状とグローバルゼイション, フレグランスジャーナル, 1月号, 25-30.
- 21) European Commission-Enterprise and Industry : http://ec.europa.eu/enterprise/cosmetics/html/consolidated_dir.htm
- 22) The international fragrance association (IFRA) : <http://www.ifraorg.org/>
- 23) 光井武夫編 (2001) : 新化粧品学 第2版, 南山堂.
- 24) Sato, Y., Katsumura, Y., Ichikawa, H., Kobayashi, T., Kozuka, T., Morikawa, F. and Ohta, S. (1981) : A modified technique of guinea pig testing to identify delayed hypersensitivity allergens, Contact Dermatitis, 7, 225-237.
- 25) 厚生労働省 : 医薬品, 医薬部外品, 化粧品及び医療機器の製造販売後安全管理の基準に関する省令, 平成16年9月22日, 厚生労働省令第135号.

5.3 食品, 食品添加物, 食品汚染物質, 飼料添加物

5.3.1 食 品

食品とは、食品衛生法第4条に「食品とは、すべての飲食物をいう。ただし、薬事法に規定する医薬品および医薬部外品は、これを含まない」と定義されている。飲食によって起こる健康障害は食性病害 (food borne disease) とも呼ばれている。食品衛生上、ヒトの健康維持に有害な因子が食品中に含まれる可能性は、その生産、採取、製造、輸入、加工、処理、保存、輸送、陳列、販売、調理、から摂取までの各段階で考慮しなければならない。

a. 食品による健康障害 この有害因子としては、食中毒や経口感染症を引き起こすコレラ、赤痢などの細菌、A型肝炎ウイルスなどのウイルス、トキソプラズマ、アニサキスなどの寄生虫のほかに、マイコトキシン (かび毒)、環境汚染物質 (ダイオ

キシシキ類など), 残留農薬, 容器からの溶出物といったいわゆる食品汚染物質, テトロドトキシシキ (フグ毒), ドーモイ酸 (記憶喪失性貝毒), ムスカリシキ (キノコ毒), サイカシキ (ソテツ実に含まれる毒) といった天然毒性物質などもあり, きわめて多種多用である. 本来トキシコロジー領域では, 細菌, ウィルス, 寄生虫などによる感染症ではなく, 細菌毒素, 天然毒性物質などの生物由来の化学物質と, 産業化学物質, 残留農薬や環境に由来する物質等人工的な化学物質が対象となるが, 食中毒時などでは感染と毒素, 双方が関与してくることに注意を払う必要がある. 以下に, 飲食によって起こる健康障害の代表例を原因により分類し記載する.

1) 食品自身によるもの 植物性・動物性の天然毒性物質 (フグ毒, 貝毒, キノコ毒, 青酸化合物など), 過剰摂取による被害 (イシナギ肝臓中のビタミン A) などがある. 急性毒性だけでなく, 発がんのおそれのある変異原性物質 (ソテツ実に含まれるサイカシキが腸内細菌により代謝を受けて生じるメチルアゾキシメタノール, ワラビの中のプタキロサイドなど) にも注意が必要である. また, 食品由来の成分には, その作用・機能が未知の化学物質が存在するため, いわゆる健康食品を代表例として, 過去の食経験が活かされないような抽出や濃縮が行われると, この未知成分による健康被害が引き起こされる可能性がある. 実例として, トウダイグサ科の植物アマメシバによる閉塞性細気管支炎を挙げることができる¹⁾.

2) 食品の変質によるもの²⁾ 食品成分が微生物, 酵素, 酸素, 光などにより質的に変化する (変質) ことがある. タンパク質が微生物により分解され, 悪臭物質 (アンモニア, メルカプタン, 硫化水素, インドール, アミン類など) を生成し変質することを一般に腐敗という. 腐敗の進行は, 温度, pH, 水分含量, 成分により著しく異なる. 糖質, 脂質等の成分も変質するがこの場合は一般に変敗といい, アルコールなど有用な化合物を生成する場合は発酵といい, 区別されている. 変敗の中では, 自動酸化による油脂の変質が問題となることが多いが, これは熱, 光, 金属などにより加速される. 高度不飽和脂肪酸 (リノール酸, リノレン酸, アラキドン酸など) の油脂過酸化物および二次生成物であるマロンアルデヒドやアクロレインなどのアルデヒ

ド類などが生成する.

3) 細菌, かび, ウィルス, 寄生虫などによる付着・汚染によるもの 食中毒菌 (腸炎ビブリオ菌, サルモネラ菌, ブドウ球菌など), 経口感染症の病原菌 (赤痢菌, 腸チフス菌, コレラ菌, 腸管出血性大腸菌 O 157 など) や, ウィルス (ノロウィルス, A 型肝炎ウィルスなど), 原虫 (アメーバ赤痢など), 寄生虫 (アニサキスなど) が原因で起きる.

かびの場合, マイコトキシシキが問題となる. これは, かびなどの真菌類が産生する低分子の二次代謝産物で急性あるいは慢性毒性を現す. かびは熱に弱い, 耐熱性のマイコトキシシキが存在する (アフラトキシシキは 300℃ でも安定). マイコトキシシキの例として, アフラトキシシキ, オクラトキシシキ, フモニシキ, パツリン, デオキシニバレノール, 麦角アルカロイド (エルゴタミンなど) が挙げられる.

ヘリコバクター・ピロリ菌 (*Helicobacter pylori*) は, 胃および十二指腸潰瘍の原因として注目されているらせん型の細菌である. 経口感染が疑われている. ウレアーゼにより胃粘液中の尿素をアンモニアと二酸化炭素に分解し, 生じたアンモニアで局所的に胃酸を中和し胃へ定着する.

牛海綿状脳症 (bovine spongiform encephalopathy, BSE) は, 異常プリオンタンパクによって媒介されると考えられている病気の 1 つで, 1990 年代になって BSE に感染した牛の特定危険部位 (脳, せき髄など) をヒトが食べたことによる発症が疑われる変異型クロイツフェルト・ヤコブ病の症例が見つかり世界的に大問題となった. 異常プリオンは, 通常の調理による加熱では失活しない.

4) 加工過程で生成する化学物質によるもの²⁾

発酵, 燻蒸, 加熱などの加工過程で実験動物において発がん性を有する, ニトロソアミンやヘテロサイクリックアミン (複素環アミン) などが生成することが示されている. ニトロソアミンは食品中にも微量存在しているが, それ以上の量が, 食品どうしあるいは添加物等との組み合わせにより, 生体内で容易に生じる. たとえば, 野菜などに多量に含まれている硝酸塩が口腔や消化管内の細菌によって還元されて亜硝酸塩になり, 胃内の酸性条件下で魚などに多く含まれる第二級アミンと反応しニトロソアミンが生成する. ヘテロサイクリックアミンは, アミノ酸やタンパク質の加熱により生成するもので, ト

リプトファンからは Trp-P-1 や Trp-P-2 が、グルタミン酸からは Glu-P-1 や Glu-P-2 などが生じる。魚干物を加熱調理すると MeIQ などが生成する。

フェオホルバイド (pheophorbide) は、クロロフィル分解物のひとつで、摂取したヒトに光過敏症を誘発することがある。1997年に粗悪なクロレラを摂取し光過敏症が起き問題となった。アワビの中腸腺の摂取時でも、含まれる葉緑素由来のフェオホルバイドと考えられる光過敏症例が報告されている。

不飽和脂肪酸は天然ではほとんどの場合シス型であるが、最近になって、植物油等の製造時の水素添加などの過程や加熱調理過程等でシス型の不飽和脂肪酸から生じるトランス脂肪酸による心血管系障害が懸念されている。なお、トランス脂肪酸は牛など反すう動物の胃内細菌により生成することも知られている。

5) 外来物質の混入、残留、汚染によるもの
環境汚染物質(ダイオキシン類, PCB, メチル水銀, カドミウム, 鉛, ヒ素, クロムなど), 過量の残留農薬・食品添加物・飼料添加物(ホルモン剤など), 食器や包装物からの溶出物(フタル酸エステルやビスフェノール A など), 放射性物質などが挙げられる。また水の汚染, たとえば, 塩素消毒時の消毒副生成物(トリハロメタンなど)や湖沼の富栄養化により大量発生したアオコ等らん藻由来の毒(ミクロシスチンなど)などにも注意が必要である。

また, 加工過程での生成や容器包装からの溶出による混入ではなく, 製品の保存中に新たな化学物質が生じるという事例もある。清涼飲料水中にベンゼンが低濃度検出されることが欧米等諸外国で公表され(2006年), 製品中の安息香酸(保存料)とアスコルビン酸(酸味料, 酸化防止剤)が, 製品の保存中に特定の条件下で反応し低濃度のベンゼンが生成することが判明した。その後ベンゼン生成を低減する措置がとられている³⁾。

6) 食物アレルギーによるもの ヒトによっては, 特定の食物の摂取によってアレルギーが引き起こされる。その多くは即時型のアレルギー反応であり, 重篤な場合は, 呼吸困難等のアナフラキシー症状を呈する。多くの場合, 抗原はタンパク質である。発生頻度が高い食物としては, えび, かに, 小麦, そば, 卵, 乳および落花生が知られている。これら7品目は特定原材料として, 表示が義務づけら

れている。また特定原材料に準ずるものとして, あわび, いくら, さば等の18品目についても表示が推奨されている。

7) 飲食物と医薬品の相互作用によるもの この例として, 1) 納豆(ビタミンK合成能が高い納豆菌)の摂取により, ワルファリンの作用が減弱, 2) 牛乳や高脂肪食摂取により, 難水溶性薬物(インドメタシン, ニトロフラントインなど)の吸収率が増加, 3) 食餌による胃酸分泌亢進により, 酸に不安定な薬物(ペニシリン, エリスロマイシンなど)の吸収率低下, 4) モノアミン酵素阻害薬(イソニアジドなど)投与を受けているヒトでの, ヒスチジンを含む魚の摂取によるヒスタミンによる中毒あるいはチーズやワインに含まれるチラミンの血圧上昇作用の促進, 5) グレープフルーツジュース中のフラノクマリン類に起因する薬物代謝酵素(CYP3A4)阻害による薬物(ニフェジピン, シクロスポリンなど)血中濃度の増大, あるいは酵素誘導による血中濃度の低下(セントジョーンズワートなどの摂取によるCYP3A4誘導, 芽キャベツによるCYP1A2誘導), などを挙げることができる。

このような食性病害を防ぐには食品の衛生管理と品質管理も重要となるが, この管理法として, 検査結果が明らかとなったときには出荷済みとなり手遅れとなる最終製品の抜き取り検査のようなものではなく, アメリカで開発されたHACCP(hazard analysis and critical control point)という食品品質管理システムがある。食品の製造工程全般を通じて危害の発生原因を分析し重要管理事項を定めるもので, 日本の食品衛生法でも採用され日本の企業でも導入されつつある²⁾。

b. 食品衛生に関する法規²⁾ 日本での食品衛生行政は「飲食物その他の物品取締に関する法律」の制定にはじまり(1900年), 当時は内務省が行政を担当し警察官により取締りが行なわれた。第二次大戦後, 新憲法のもと食品衛生行政が厚生省の所管となり, これまでの食品衛生に関する法律, 規則が見直され「食品衛生法」が公布(1947年), 翌年施行された。この法律の目的は, 「食品の安全性の確保のために公衆衛生の見地から必要な規制その他の措置を講ずることにより, 飲食に起因する衛生上の危害の発生を防止し, もって国民の健康の保護を図ること」と定められている。食品衛生に関する監視,

指導のため, 国, 都道府県および政令指定都市に「食品衛生監視員」が設けられている。なお, 2003年に残留農薬の規制に関して食品衛生法が改正となり, ポジティブリスト制が導入され, 基準値が設定されていない農薬を一律基準である0.01ppm以上含む食品は, 国内では流通できないこととなった。厚生労働省が管轄する調査・研究機関としては, 国立医薬品食品衛生研究所, (独) 国立健康・栄養研究所などがある。また厚生労働大臣の諮問機関として厚生労働省内に「薬事・食品衛生審議会」が設置されている。他方, 厚生労働省が食品衛生法を管轄するのに対し, 農林水産省は, 『農林物質の規格化および品質表示に関する法律 (JAS法: Japanese Agricultural Standard)』(1950年制定) を所管している。

なお, 食品衛生法及びJAS法において, 同じ期限表示項目に「品質保持期限」及び「賞味期限」という2つの用語が使用され, わかりにくい表現となっていたが, 両省による「食品の表示に関する共同会議」の協議を経て, 2003年より「賞味期限」に統一された。またこれと同時に食品衛生法とJAS法において別々に定義されていた「消費期限」の定義が統一された。

食品の安全を確保するための法律が厚生労働省と農林水産省に分かれていたが, 2003年に『食品安全基本法』が公布・施行され両省から独立した『食品安全委員会』が設立された。食品安全委員会は, 7名の委員から構成される。食品の安全性の確保に関する施策の策定に当たってはリスク評価とリスク管理を分離し, 1) 食品健康影響評価(リスク評価)の実施, 2) リスク評価に基づいた施策の策定(リスク管理), 3) 関係者相互間の情報および意見の交換(リスクコミュニケーション)を行うこととなった。リスク評価は新設の食品安全委員会が担当し, リスク管理はこれまで通り厚生労働省および農林水産省などが担当することとなった。また食品安全委員会は, 関係者相互の情報と意見の交換によって社会的合意を得るという「リスクコミュニケーション」を推進することとなった。

c. 保健機能食品⁴⁾ 食品のなかには, 国が定めた安全性や有効性に関する基準等を満たした「保健機能食品」がある(健康増進法・食品衛生法)。他方, いわゆる健康食品(栄養補助食品, 健康補助

食品, サプリメントなどの名称も含む)は, 法律上の定義はなく, 広く健康の保持増進に資する食品として販売・利用されるもの全般を指している。保健機能食品は, 特定保健用食品および栄養機能食品の2種類の類型からなり, それぞれ独自の表示をすることができる。

特定保健用食品は, 身体の生理学的機能などに影響を与える保健機能成分を含み, 厚生労働大臣により許可または承認された食品であり約850品目がある(2009年4月現在)(個別許可型)。栄養機能食品は, 不足しがちな栄養成分(ビタミン・ミネラル)の補給のために利用される食品で, 規格基準に適合すれば許可申請や届け出は不要である(規格基準型)。その他, 健康や栄養に関する表示が行える食品として, 病者用, 乳児用, 妊産婦用などの特別の用途に適するという表示ができる特別用途食品がある(病者用食品, 高齢者用食品など)。特別用途食品として食品を販売するには, その表示について国の許可を受ける必要がある。これらを除いては, 食品のもつ効果や機能を根拠なく表示することは, 医薬品と誤認されるような効能・効果表示と見なされ, 薬事法により禁止されている。

d. 遺伝子組換え食品(遺伝子組換え食品表示)²⁾

遺伝子組換え技術によって改良した生物から得られた食品を遺伝子組換え食品と呼ぶ。この技術により, 食品生産を量的・質的に向上させるだけでなく, 害虫や病気に強い農作物の改良や, 加工特性などの品質向上に利用されることが期待されている。具体的には, 除草剤グリホサートに抵抗性をもたせた大豆などの除草剤に耐性がある農作物, パチルス属の細菌由来の殺虫タンパク遺伝子を発現させたトウモロコシなどの害虫に抵抗性がある農作物, 遺伝子操作でオレイン酸含量を増加させた大豆などの高栄養価の農作物などである。遺伝子組換えサケなども食用に検討されている。

遺伝子組換え食品のうち, 大豆, とうもろこし, ばれいしょ, 菜種, 綿実, アルファルファ, てん菜を原材料とする加工食品については, 食品衛生法施行規則で組換えDNA技術応用作物の食品に関する表示に関する規定が設けられており, 農産物およびこれを原材料とする加工食品であって, 加工後も組み換えられたDNAまたはこれによって生じたタンパク質が残存するものについては, 「遺伝子組換え