

**DIVISION OF AIDS TABLE FOR GRADING THE SEVERITY OF
ADULT AND PEDIATRIC ADVERSE EVENTS
PUBLISH DATE: DECEMBER, 2004**

LABORATORY				
PARAMETER	GRADE 1 MILD	GRADE 2 MODERATE	GRADE 3 SEVERE	GRADE 4 POTENTIALLY LIFE-THREATENING
Glucose, serum, low				
Adult and Pediatric ≥ 1 month	55 – 84 mg/dL 3.05 – 3.55 mmol/L	40 – 54 mg/dL 2.22 – 3.06 mmol/L	30 – 39 mg/dL 1.67 – 2.23 mmol/L	< 30 mg/dL < 1.67 mmol/L
Infant [†] , < 1 month	50 – 54 mg/dL 2.78 – 3.00 mmol/L	40 – 49 mg/dL 2.22 – 2.77 mmol/L	30 – 39 mg/dL 1.67 – 2.21 mmol/L	< 30 mg/dL < 1.67 mmol/L
Lactate	< 2.0 x ULN without acidosis	≥ 2.0 x ULN without acidosis	Increased lactate with pH < 7.3 without life- threatening consequences	Increased lactate with pH < 7.3 with life- threatening consequences
LDL cholesterol (fasting)				
Adult ≥ 18 years	130 – 159 mg/dL 3.37 – 4.12 mmol/L	160 – 190 mg/dL 4.13 – 4.90 mmol/L	≥ 190 mg/dL ≥ 4.91 mmol/L	NA
Pediatric > 2 - < 18 years	110 – 129 mg/dL 2.85 – 3.34 mmol/L	130 – 159 mg/dL 3.35 – 4.20 mmol/L	≥ 190 mg/dL ≥ 4.91 mmol/L	NA
Lipase	1.1 – 1.5 x ULN	1.5 – 3.0 x ULN	3.1 – 5.0 x ULN	> 5.0 x ULN
Magnesium, serum, low	1.2 – 1.4 mEq/L 0.60 – 0.70 mmol/L	0.9 – 1.1 mEq/L 0.45 – 0.55 mmol/L	0.6 – 0.8 mEq/L 0.30 – 0.41 mmol/L	< 0.60 mEq/L < 0.30 mmol/L
Pancreatic amylase	1.1 – 1.5 x ULN	1.5 – 2.0 x ULN	2.1 – 5.0 x ULN	> 5.0 x ULN
Phosphate, serum, low				
Adult and Pediatric > 14 years	2.5 mg/dL – < LLN 0.81 mmol/L – < LLN	2.0 – 2.4 mg/dL 0.65 – 0.80 mmol/L	1.0 – 1.9 mg/dL 0.32 – 0.64 mmol/L	< 1.00 mg/dL < 0.32 mmol/L
Pediatric 1 year – 14 years	3.0 – 3.5 mg/dL 0.97 – 1.13 mmol/L	2.5 – 2.9 mg/dL 0.81 – 0.95 mmol/L	1.5 – 2.4 mg/dL 0.48 – 0.80 mmol/L	< 1.50 mg/dL < 0.48 mmol/L
Pediatric < 1 year	3.5 – 4.5 mg/dL 1.13 – 1.45 mmol/L	2.5 – 3.4 mg/dL 0.81 – 1.12 mmol/L	1.5 – 2.4 mg/dL 0.48 – 0.80 mmol/L	< 1.50 mg/dL < 0.48 mmol/L
Potassium, serum, high	5.6 – 6.0 mEq/L 5.6 – 6.0 mmol/L	6.1 – 6.5 mEq/L 6.1 – 6.5 mmol/L	6.6 – 7.0 mEq/L 6.6 – 7.0 mmol/L	> 7.0 mEq/L > 7.0 mmol/L
Potassium, serum, low	3.0 – 3.4 mEq/L 3.0 – 3.4 mmol/L	2.5 – 2.9 mEq/L 2.5 – 2.9 mmol/L	2.0 – 2.4 mEq/L 2.0 – 2.4 mmol/L	< 2.0 mEq/L < 2.0 mmol/L
Sodium, serum, high	146 – 150 mEq/L 146 – 150 mmol/L	151 – 154 mEq/L 151 – 154 mmol/L	155 – 159 mEq/L 155 – 159 mmol/L	≥ 160 mEq/L ≥ 160 mmol/L
Sodium, serum, low	130 – 135 mEq/L 130 – 135 mmol/L	125 – 129 mEq/L 125 – 129 mmol/L	121 – 124 mEq/L 121 – 124 mmol/L	≤ 120 mEq/L ≤ 120 mmol/L
Triglycerides (fasting)	NA	500 – 750 mg/dL 5.65 – 8.48 mmol/L	751 – 1,200 mg/dL 8.49 – 13.56 mmol/L	> 1,200 mg/dL > 13.56 mmol/L
Uric acid	7.5 – 10.0 mg/dL 0.45 – 0.59 mmol/L	10.1 – 12.0 mg/dL 0.60 – 0.71 mmol/L	12.1 – 15.0 mg/dL 0.72 – 0.89 mmol/L	> 15.0 mg/dL > 0.89 mmol/L

* Values are for term infants.

† Use age and sex appropriate values (e.g., bilirubin), including preterm infants.

**DIVISION OF AIDS TABLE FOR GRADING THE SEVERITY OF
ADULT AND PEDIATRIC ADVERSE EVENTS
PUBLISH DATE: DECEMBER, 2004**

LABORATORY				
PARAMETER	GRADE 1 MILD	GRADE 2 MODERATE	GRADE 3 SEVERE	GRADE 4 POTENTIALLY LIFE-THREATENING
URINALYSIS <i>Standard International Units are listed in italics</i>				
Hematuria (microscopic)	0 – 10 RBC/HPF	> 10 RBC/HPF	Gross, with or without clots OR with RBC casts	Transfusion indicated
Proteinuria, random collection	1 +	2 – 3 +	4 +	NA
Proteinuria, 24 hour collection				
Adult and Pediatric ≥ 10 years	200 – 999 mg/24 h <i>0.200 – 0.999 g/d</i>	1,000 – 1,999 mg/24 h <i>1.000 – 1.999 g/d</i>	2,000 – 3,500 mg/24 h <i>2.000 – 3.500 g/d</i>	> 3,500 mg/24 h <i>> 3.500 g/d</i>
Pediatric > 3 mo - < 10 years	201 – 499 mg/m ² /24 h <i>0.201 – 0.499 g/d</i>	500 – 799 mg/m ² /24 h <i>0.500 – 0.799 g/d</i>	800 – 1,000 mg/m ² /24 h <i>0.800 – 1.000 g/d</i>	> 1,000 mg/m ² /24 h <i>> 1.000 g/d</i>

*Values are for term infants.

† Use age and sex appropriate values (e.g., bilirubin), including preterm infants.

付録 2)

重篤な有害事象発現時に、製造販売元または KFDA へ連絡すべき内容一覧

1. 報告医師名
2. 施設名
3. 試験薬 (一般名: メロキシカム)
4. 被験者の識別番号
5. 被験者の性別 男性
6. 被験者の年齢
7. 有害事象 1 ※有害事象が複数ある場合は、8 以下の項目をコピーして繰り返して下さい。

8. 有害事象名
9. 重篤と判断した理由
 - ※該当するもののみ記載
 - ※複数選択可 (死に至るもの/生命を脅かすもの/入院または入所期間の延長/永続的または顕著な障害・機能不全/先天異常を来たすもの/その他、医学的に重大なもの)
10. 試験薬 (メロキシカム) との因果関係 関連あり/多分関連あり/関連不明/多分関連なし/関連なし
11. 判定理由 ※「判定理由」は必須
12. 発現日 2010/MM/DD
13. 転帰 ※該当するもののみ記載
(死亡/未回復/回復したが後遺症あり/回復/不明/その他)
14. 転帰日 2010/MM/DD
15. 試験薬 (メロキシカム) の投与 7.5 mg 錠の単回投与
16. 投与開始日 2010/MM/DD
17. 有害事象発現後の措置
 - ※副作用に対する治療薬や処置
18. 有害事象の概要 (発現状況、症状、場所、処置等の経過)
 - ※可能な限り、経時的に記載

