

LETTERS




Figure 1 HMGGA2 is an essential positive regulator of cardiomyocyte differentiation. (a) Expression of *HMGGA2* mRNA during P19CL6 differentiation into cardiomyocytes was assessed by northern blot analysis. (b–f) The effect of *HMGGA2* overexpression or *HMGGA2* knockdown on cardiomyocyte differentiation was assessed with MF20 staining in control (b), empty vector-transfected (c), *HMGGA2* overexpressing (d), *HMGGA2* siRNA-transfected (e) and control siRNA-transfected (f) P19CL6 cells. Scale bar is

100 μ m. (g) Quantification of MF20-positive area. The results are shown as mean \pm s. d. * $P < 0.05$, compared with control or vector-transfected cells, ** $P < 0.01$ compared with control or irrelevant siRNA-transfected cells (one-way ANOVA, followed by Fisher's PLSD test; $n = 6$) (h) The effect of *HMGGA2* overexpression or *HMGGA2* knockdown on cardiac marker gene expression was assessed by RT-PCR (for *Nkx2.5* and *atrial natriuretic peptide (ANP)*) and northern blot (for *GATA-4*, *MEF2C* and *myosin light chain 2v (MLC2v)*).

generated by alternative splicing of *HMGGA1* gene transcripts, and HMGGA2, encoded by the *HMGGA2* gene. All HMGGA proteins except HMGGA1c contain three short basic repeats called AT-hooks, which bind to the minor groove of AT-rich DNA stretches¹. HMGGA proteins are also able to associate with multiple transcription factors and regulate the expression of their target genes. For example, expression of the interferon- β gene is regulated by a multiprotein complex

containing NF- κ B, interferon regulatory factor, activating transcription factor-2/c-Jun and HMGGA1a⁹. HMGGA proteins also participate in the regulation of the genes for interleukin-2 receptor α and the insulin receptor^{10,11}. HMGGA proteins are expressed ubiquitously and abundantly during embryogenesis, whereas their expression is low or undetectable in fully differentiated adult tissues¹². This suggests that HMGGA proteins regulate normal cell growth and differentiation.




Figure 2 cDNA sequence and expression pattern of *XHMGA2*. (a) Amino acid sequences of *XHMGA2* protein were compared with those of human (NM_001094371), mouse (NM_010441), rat (NM_032070), chicken (NM_205001) and zebrafish (NM_212680) *HMGA2*. Dots indicate identities and dashed lines are introduced to maximize the alignment. Boxes represent the three AT-hooks. (b) Expression of *XHMGA2* mRNA during embryogenesis analysed by RT-PCR. *Ornithine decarboxylase* (*ODC*) represents internal controls. (c–h) Expression of *XHMGA2* mRNA during embryogenesis analysed by *in situ*

hybridization. Lateral (c) and ventral (d) view of stage-15 embryo. *XHMGA2* expression was detected in neural crest (NC) and weakly in precardiac region (arrowhead). Lateral (e) and ventral (f) view of stage-23 embryo. *XHMGA2* was expressed in anterior neural tissue (AN), tailbud (TB) and weakly in precardiac region (arrowhead). Lateral (g) and ventral (h) view of stage-32 embryo. *XHMGA2* was detected in brain (BR), spinal cord (SC), eye anlage (EA), otic vesicle (OV), branchial arch (BA), pronephric duct (PD) and heart anlage (arrowhead). Anterior is left and dorsal is top for c, e, g, and anterior is left for d, f, h.

To examine whether *HMGA2* regulates cardiomyocyte differentiation, we analysed the ability of P19CL6 clones stably overexpressing wild-type *HMGA2* (P19CL6-*HMGA2*) to differentiate into cardiomyocytes. The extent of cardiomyocyte differentiation was assessed by the area positive for an anti- α -myosin heavy chain antibody (MF20). In response to DMSO, P19CL6-*HMGA2* differentiated into beating cardiomyocytes more efficiently than the parental P19CL6 cells (Fig. 1b–d, g), and there was a positive correlation between the extent of cardiomyocyte differentiation and the level of *HMGA2* expression in multiple P19CL6-*HMGA2* clones (Supplementary Information, Fig. S1a). On the other hand, siRNA-mediated knockdown of *HMGA2* blocked cardiomyocyte

differentiation of P19CL6 cells without altering the expression of mesodermal marker genes such as *Brachyury* and *Flk-1*, whose expressions precede those of early cardiac marker genes such as *Nkx2.5* (Fig. 1e–g; Supplementary Information, Fig. S1b). The expression levels of cardiac marker genes was upregulated in response to *HMGA2* overexpression, whereas they were downregulated by knockdown of *HMGA2* expression (Fig. 1h). Together, these results suggest that *HMGA2* is an essential positive regulator of cardiomyocyte differentiation.

To investigate the role of *HMGA2* in cardiogenesis *in vivo*, we performed experiments using *Xenopus* embryos. We first cloned *XHMGA2*, a *Xenopus* orthologue of *HMGA2*. *XHMGA2* encodes a protein of 105

LETTERS




Figure 3 XHMGGA2 is essential for cardiogenesis. (a–d) Overexpression of a dominant-negative XHMGGA2 mutant in *Xenopus* embryos blocked *XNkx2.5* expression. Ventral view of stage-23 (a, b) and lateral view of stage-34 (c, d) embryos. Expression of *XNkx2.5* mRNA was decreased in XHMGGA2-EnR mRNA-injected embryos (b, d, arrowheads), compared with that of control embryos (a, c, arrowheads). (e–p) MO-mediated XHMGGA2 knockdown results in impaired cardiogenesis. Whole-mount *in situ* hybridization analysis was

performed sequentially for *Xenopus* embryos at stage 15 (e–g), stage 23 (h–j) and stage 34 (k–m) for *XNkx2.5*, and at stage 41 for *cardiac troponin I (cTnl)* (n–p). Expression of *XNkx2.5* mRNA was detected in uninjected control embryos (e, h, k, arrows), whereas it was attenuated in both XHMGGA2-MO1 (f, i, l) and XHMGGA2-MO2 (g, j, m)-injected embryos (arrowheads). *In situ* hybridization analysis for *cardiac troponin I* revealed that the heart size in MO-injected embryos (o, p, arrowheads) was smaller than that of control embryos (n, arrows).

amino acids that contains three highly conserved AT-hooks and is 60–70% homologous with HMGGA2 proteins in other species (Fig. 2a). Reverse transcriptase-polymerase chain reaction (RT-PCR) analysis revealed that XHMGGA2 was maternally transcribed in the early stage of embryogenesis and its expression levels increased at stage 30 (Fig. 2b). *In situ* hybridization analysis revealed that XHMGGA2 was

expressed in neural crest cells and the precardiac region at the neurula stage (Fig. 2c, d), and in anterior neural tissue, tailbud and precardiac region at the early tailbud stage (Fig. 2e, f). At the late tailbud stage, a strong signal was detected in brain, spinal code, eye anlage, otic vesicle, branchial arch, pronephric duct and heart anlage (Fig. 2g, h). To examine whether XHMGGA2 regulates cardiogenesis *in vivo*, we injected

LETTERS




Figure 5 HMGGA2 binding is required for full activation of *Nkx2.5* promoter. (a) Sequence comparison of SBE-2/HBE-1 and SBE-4/HBE-2 elements in the -3059/-2554 *Nkx2.5* cardiac enhancer. These elements are evolutionarily conserved in human, mouse and chick, and SBE-4/HBE-2 sites are also conserved in *Xenopus*. Nucleotides that are not conserved are shown by small letters: m, mouse; h, human; c, chick; x, *Xenopus*. (b) Mutations introduced into conserved SBE-2/HBE-1 and SBE-4/HBE-2 elements attenuated *Nkx2.5* promoter activity during cardiomyocyte differentiation of P19CL6 cells. *Nkx2.5* promoter activity was assessed in P19CL6 cells at day 6 of differentiation. GM, growth medium (without DMSO); DM, differentiation medium (containing 1% DMSO); WT, wild-type *Nkx2.5* promoter; mS2, mH1, mS4 and mH2, *Nkx2.5* promoter containing mutations at SBE-2, HBE-1, SBE-4 and HBE-2,

respectively ($n = 3$). (c) Interaction between Smad1/4 and HMGGA2 on the SBE-4/HBE-2 element, as revealed by EMSA. Smad1/4 and HMGGA2, either alone or in combination, bound to a 24-bp probe containing the SBE-4/HBE-2 element. (d) Sequences used for the EMSA probe. Mutations introduced into the HBE-2 site are shown in red. WT, wild-type probe; MT, mutated probe. (e) Binding of HMGGA2 to SBE-4/HBE-2 element was reduced by mutations introduced into the HBE-2 site. (f) *GFP* reporter genes used to generate transgenic *Xenopus* embryos are shown on the left. pXNkx2.5(mt)-GFP plasmid contains mutations in the conserved HBE-2 site, which are indicated by red. *GFP* expression as assessed by *in situ* hybridization is shown on the right panels. The number of embryos showing *GFP* expression (arrowhead) per total number of surviving embryos for both plasmids are indicated.

in *Xenopus*¹⁸, suggesting that this cardiac enhancer may be a target of HMGGA2 and Smads. Indeed, there were two potential AT-rich HMGGA2 binding elements (HBEs) adjacent to SBE-2 and SBE-4 (Fig. 5a). These HBEs were conserved in chick, mouse and human, and HBE-2 was also conserved in *Xenopus*^{15,16,18}. Mutations introduced into SBE-2 or HBE-1 reduced the promoter activity of *Nkx2.5* by 50–60%, and more than 80% reduction in the promoter activity was observed by mutations introduced into SBE-4 or HBE-2 (Fig. 5b), suggesting that HMGGA2 and Smads regulate *Nkx2.5* promoter activity cooperatively through these elements. As mutations introduced into the proximal element containing SBE-4/HBE-2 had a profound effect on *Nkx2.5* promoter activity, a 24-bp sequence containing SBE-4/HBE-2 (Fig. 5a, d) was used as a probe to examine protein–DNA interactions. Electrophoretic mobility shift assay (EMSA) showed that Smad1/4, HMGGA2 and a complex containing Smad1/4 and HMGGA2 bound to the 24-bp sequence (Fig. 5c). Mutation of the A/T-rich HBE-2 markedly attenuated the ability of HMGGA2 to bind to this sequence (Fig. 5d, e). We also analysed *XNkx2.5* promoter

activity *in vivo* in transgenic *Xenopus* embryos. Transgenic embryos expressing *green fluorescent protein* (*GFP*) gene under the control of the -4295 bp *XNkx2.5* promoter, which contains conserved SBE-4 and HBE-2, were assayed for their *GFP* mRNA expression by *in situ* hybridization. The *GFP* transgene containing the -4295 bp *XNkx2.5* promoter induced detectable *GFP* expression in the heart, and mutations introduced into HBE-2 abolished it (Fig. 5f). These findings suggest that HMGGA2 and BMP-responsive Smads upregulate *Nkx2.5* promoter activity cooperatively through Smad- and HMGGA2-binding elements, and that the conserved HMGGA2 binding site is essential for *Nkx2.5* expression both *in vitro* and *in vivo*.

In this study we have demonstrated that HMGGA2 promotes cardiomyocyte differentiation *in vitro* and is essential for cardiogenesis both *in vitro* and *in vivo*. Expression of HMGGA2 is high during embryogenesis but low or undetectable in the adult tissues, suggesting that HMGGA2 regulates normal cell growth and differentiation in general. Of note, transient expression of HMGGA2 was observed in the heart after myocardial

infarction (Supplementary Information, Fig. S4), suggesting that HMGA2 is one of the fetal genes re-expressed in the myocardium in response to biomechanical stress. HMGA2 has been implicated in both benign and malignant tumours. Rearrangement of *HMGA2*, which results in the generation of a chimaeric or a truncated HMGA2 protein that contains three AT-hooks but lacks its carboxy-terminus, is frequently observed in benign human tumours of mesenchymal origin¹⁹. Overexpression of wild-type HMGA2 has also been reported in several malignant tumours¹⁹. Moreover, transgenic mice overexpressing the C-terminal-truncated form of HMGA2 are large and obese with lipomas^{20,21}. Targeted disruption of the *HMGA2* gene in mice causes general growth retardation and impaired adipocyte differentiation^{22,23}, consistent with the suggestion that HMGA2 regulates cell growth and differentiation. In the experiments shown in Fig. 3, MOs were injected into the dorsal region of two dorsal-vegetal blastomeres fated to be heart and liver anlage. Examination of other lineage markers under these experimental conditions revealed that the expression of *XHex*, a marker gene for liver and thyroid gland, was attenuated in the liver and absent in the thyroid gland (Supplementary Information, Fig. S5d–f). Similarly, *Xmsr*, an endothelial-cell marker gene, showed a perturbed pattern of expression (Supplementary Information, Fig. S5g–i). Furthermore, when *XHMGA2*-MOs were injected into other regions at the 8-cell stage (Supplementary Information, Fig. S6a), different phenotypes of morphants were observed, depending on the site of MO injection and the area of MO distribution (Supplementary Information, Fig. S6b–f). Thus, the phenotype of *XHMGA2* knockdown is not necessarily restricted to the heart.

In this study we have also demonstrated that HMGA2 forms a protein complex with BMP-responsive Smad transcription factors that coordinately upregulate the promoter activity of *Nkx2.5* through evolutionarily conserved Smad- and HMGA2-binding elements. As HMGA2 has been shown to regulate the proliferation and/or differentiation of multiple cell types, there is a possibility that the regulation of cardiogenesis by HMGA2 is indirect and mediated by its effects on other cell types. However, we favour the idea that HMGA2 regulates cardiomyocyte differentiation directly, as our data suggest that HMGA2 promotes cardiogenesis through the transcriptional activation of *Nkx2.5*. Specifically, reduced expression of *Nkx2.5* by inhibition of HMGA2 and downregulation of *Nkx2.5* promoter activity by mutations in the HBE-2 site both *in vitro* and *in vivo* strongly suggest that *Nkx2.5* is a direct target of HMGA2. In this regard it is noteworthy that HMGA2 regulates the expression of an organ-specific transcription factor in collaboration with a growth factor-mediated signalling system and consequently contributes to organogenesis. In contrast to the *Nkx2.5* promoter, there was no cooperative upregulation of the BMP-responsive reporter BRE-luc or the TGF- β -responsive reporter p3TP-luc by the co-expression of Smad1/4/HMGA2 or Smad2/4/HMGA2, (Supplementary Information, Fig. S7). Thus, HMGA2 is not generally involved in transcriptional regulation mediated by Smads, but rather, is required for a specific subset of Smad-responsive transcriptional regulation in a context-dependent manner. Simultaneous interaction of HMGA2 with DNA and Smads with DNA may be necessary for synergistic transactivation by HMGA2 and Smads.

We speculate that the defect in cardiogenesis induced by HMGA2 inactivation is caused by downregulation of BMP-mediated *Nkx2.5*, as inactivation of *Nkx2.5* is sufficient to disrupt normal heart formation

*in vivo*⁴. However, the possibility that HMGA2 is required for the proliferation of embryonic cardiomyocytes cannot be excluded. *HMGA2* knockdown in *Xenopus* embryos resulted in cardiac defects, whereas no cardiac abnormalities have been reported in *HMGA2* knockout mice. Although the exact reason for this apparent discrepancy is not clear at this time, the lack of cardiac phenotype in *HMGA2* mutant mice may be due to the genetic redundancies between *HMGA1* and *HMGA2* or among other HMG family members. As HMGA2 seems to regulate the early stage of cardiac development, further studies on this molecule may provide insights into myocardial regenerative medicine and the pathophysiology of congenital heart diseases. □

METHODS

Plasmids and reagents. pcDNA3-HMGA2 was provided by G. Manfioletti (University of Trieste, Italy). FLAG-tagged Smad1/2/4, GST-Smad1/4 and p3TP-luc have been described previously²⁴. BRE-luc was provided by P. ten Dijke (Leiden University Medical Center, Netherlands)²⁵. *Nkx2.5*(-3059)-luc and *Nkx2.5*(-959)-luc were provided by K. E. Yutzey (Cincinnati Children's Medical Center, Cincinnati, OH)¹⁹. pCS-Fast-EnR was provided by M. Whitman (Harvard Medical School, Boston, MA)¹². XCarGFP3 was provided by E. Amaya (University of Manchester, UK)²⁶. Natural bovine BMP cocktail was purchased from Sangi.

Differential mRNA display. Differential mRNA display and subcloning of re-amplified cDNA fragments were performed as described previously²⁷.

P19CL6 cell culture and stable transformants. P19CL6 cells were cultured and induced to differentiate into cardiomyocytes as described previously⁸. To obtain P19CL6 clones stably overexpressing *HMGA2*, pcDNA3-HMGA2 was transfected into P19CL6 cells and neomycin-resistant clones were selected.

siRNA-mediated knockdown in P19CL6 cells. For *HMGA2* knockdown, 2-For-Silencing siRNA kit (Qiagen) was used. siRNAs were transfected at day 2 of differentiation. Sequences of *HMGA2* siRNA were as follows. Duplex 1: r(AGU AUA AGU UAA UAC UGA A)dTdT for sense, r(UUC AGU AUU AAC UUA UAC U)dGdA for antisense. Duplex 2: r(GGA AAU CUA CAC AGC CAA A)dTdT for sense, r(UUU GGC UGU GUA GAU UUC C)dCdG for antisense. MARK1 siRNA included in the kit was used as an irrelevant control siRNA.

Immunocytochemistry. Immunostaining with an anti-MF20 antibody was performed as described previously⁸. MF20-positive areas were measured at day 14 of differentiation.

RNA analysis. Northern blot, RT-PCR and quantitative real-time PCR for RNA analysis in P19CL6 cells were performed as described previously^{8,28}. In *Xenopus* embryos, RT-PCR was performed as described previously²⁹. PCR primers and PCR conditions are available in the Supplementary Information.

IP-western blot analysis. Total cell lysate was prepared from COS7 cells transfected with expression vectors for HA-HMGA2 and FLAG-Smad1 from P19CL6 cells induced to differentiate for 6 days. IP-western blot analysis was performed essentially as described previously³⁰ using anti-HA monoclonal antibody 12CA5 (Roche) and anti-FLAG monoclonal antibody M2 (Kodak) for COS7 cell lysate, and anti-Smad1/5/8 and anti-HMGA2 (HMG1-C) rabbit polyclonal antibodies (Santa Cruz) for P19CL6 cell lysate.

Luciferase reporter gene assay. *Nkx2.5*-luc, pRL-CMV (an internal control) and effector constructs were transfected into COS7 cells or P19CL6 cells, and luciferase activity was measured with a luminometer (Berthold Lumat LB9507) 48 h after transfection. BMP cocktail (100 ng ml⁻¹) was added to COS7 medium. Mutations were introduced into *Nkx2.5*-luc plasmid using QuikChange II Site-Directed Mutagenesis Kit (Stratagene).

EMSA. Probes for EMSA were labelled using Biotin 3' End DNA Labeling Kit (Pierce Biotechnology) and EMSA was performed using the LightShift Chemiluminescent EMSA Kit (Pierce Biotechnology). The HMGA2 protein

LETTERS

(HMGIC(48–109)-NH₂) was purchased from Phoenix Pharmaceuticals. GST-Smad1 and GST-Smad4 fusion proteins were purified with B-PER GST Spin Purification Kit (Pierce Biotechnology).

Isolation of *XHMG2* cDNA. The entire coding region of *XHMG2* was amplified by RT-PCR using mRNA obtained from embryos at the early tailbud stage, based on the sequence of a *Xenopus* EST (NM_001094371), which encodes a full-length protein similar to human HMGA2. The following primers were used: *XHMG2*-U (5'-ATG AGC TCA AGG GAA GGA GCC-3'), *XHMG2*-D (5'-CTA GTC GTC TTC AGA TTC CTG GG-3'). We cloned the PCR product into a pCS2+ vector (pCS2+*XHMG2*).

Microinjection of *XHMG2-EnR* mRNA. An expression vector for *XHMG2-EnR* was constructed by a PCR-based cloning strategy. A cDNA fragment encoding Engrailed repressor domain (EnR) was amplified from *FAST-EnR* plasmid by PCR. mRNA for microinjection was synthesized with mMESSAGE mMACHINE kit (Ambion) from plasmids encoding *XHMG2-EnR*. Synthesized mRNAs were microinjected as described previously²⁹.

MO experiments. The sequences of *XHMG2* MOs were: *XHMG2*-MO1, 5'-AGC TCA TGG TAG AGA GTG TGT GTG C-3'; *XHMG2*-MO2, 5'-GCC CGG CGA TCC TGG AGC ACC TTA A-3'. MO activities and specificities were checked by co-injection of 5' *XHMG2-EGFP* or *XHMG2-EGFP* mRNA (Supplementary Information, Fig. S2). For this, the *XHMG2* coding region with 73 bp 5' untranslated region (UTR) and *XHMG2* coding region without 5' UTR (Supplementary Information, Fig. S2a) were inserted into the *Clal* site of the EGFP-CS2 vector²⁹ to construct expression vectors for 5' *XHMG2-EGFP* and *XHMG2-EGFP*, respectively. We injected the MOs into two dorsal-vegetal blastomeres at the 8-cell stage. Rescue experiments were performed by injecting optimal-effect doses of *XHMG2*-MO1 or *XHMG2*-MO2 in conjunction with pCS2+*XHMG2* plasmid DNA, which lacks 5' UTR and therefore is MO-resistant (100 pg per embryo). For tracing of injected MOs, β -gal mRNA was co-injected and embryos were pre-stained with Red-gal (Research Organics) before whole-mount *in situ* hybridization.

Whole-mount *in situ* hybridization. The following plasmid templates were linearized, and digoxigenin-substituted antisense RNA probes were transcribed with T7 or SP6 RNA polymerase: *XNkx2.5*, *XHex*, *Xmsr* and *XHMG2* (a PCR-amplified coding region subcloned into pBluescript II SK+); *Xenopus cardiac troponin I* (a PCR amplified coding region subcloned into pGEM-T Easy). Embryos were processed for whole-mount *in situ* hybridization using BM purple substrate (Roche) and then the processed pigmented embryos were bleached by 9% H₂O₂, 21% H₂O and 70% methanol.

Generation and analysis of transgenic *Xenopus* embryos. pXNkx2.5-GFP3 was generated by replacing the *cardiac actin* promoter in XCarGFP3 (ref. 18) with 4295 bp promoter sequence of *XNkx2.5*. This fragment was amplified by PCR from *Xenopus* tailbud genomic DNA using the XNkx2.5p-U primer (5'-ACC TGA GCT CGG GGG GAA TAT ACA CAA GGC C-3') and XNkx2.5p-D primer (5'-GCA CCG GTG ACG GTA TCA GGT AAA CCC CAC A-3'). pXNkx2.5(mt)-GFP3 was created by site-directed mutagenesis. Both plasmids were digested by *SacI* and injected. Generation of transgenic *Xenopus* embryos was carried out as described previously³¹.

Accession codes. USCD-Nature Signaling Gateway (<http://www.signaling-gateway.org>): A001667, A001029 and A001503

Note: Supplementary Information is available on the Nature Cell Biology website.

ACKNOWLEDGEMENTS

We thank G. Manfioletti, P. ten Dijke, K. E. Yutzey, M. Whitman and E. Amaya for providing plasmids, and C. Masuo and Y. Itoh for their excellent technical assistance. This work was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology, and Health and Labor Sciences Research Grants; an Academic Award of the Mochida Memorial Foundation and Uehara Memorial Foundation (to I. K.); and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to K. M.).

AUTHOR CONTRIBUTIONS

K. Monzen, Y. I. and A. T. M. contributed equally to this work; I. K. designed and supervised the research; K. Monzen, Y. I., A. T. M., H. K., Y. H. and D. H. performed

experiments; I. S., T. Y., K. Miyazono, M. A. and R. N. contributed new reagents/analytical tools; K. Monzen, Y. I. and A. T. M. analysed the data; K. Monzen, Y. I., I. S. and I. K. prepared the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Published online at <http://www.nature.com/naturecellbiology/>

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>

1. Reeves, R. Molecular biology of HMGA proteins: hubs of nuclear function. *Gene* **277**, 63–81 (2001).
2. Sgarra, R. *et al.* Nuclear phosphoproteins HMGA and their relationship with chromatin structure and cancer. *FEBS Lett.* **574**, 1–8 (2004).
3. Hock, R., Furusawa, T., Ueda, T. & Bustin, M. HMG chromosomal proteins in development and disease. *Trends Cell. Biol.* **17**, 72–79 (2007).
4. Srivastava, D. Genetic assembly of the heart: implications for congenital heart disease. *Annu. Rev. Physiol.* **63**, 451–469 (2001).
5. Olson, E. N. & Schneider, M. D. Sizing up the heart: development redux in disease. *Genes Dev.* **17**, 1937–1956 (2003).
6. Hoffman, J. I. & Kaplan, S. The incidence of congenital heart disease. *J. Am. Coll. Cardiol.* **39**, 1890–1900 (2002).
7. Foley, A. & Mercola, M. Heart induction: embryology to cardiomyocyte regeneration. *Trends Cardiovasc. Med.* **14**, 121–125 (2004).
8. Monzen, K. *et al.* Bone morphogenetic proteins induce cardiomyocyte differentiation through the mitogen-activated protein kinase kinase TAK1 and cardiac transcription factors Csx/Nkx-2.5 and GATA-4. *Mol. Cell. Biol.* **19**, 7096–105 (1999).
9. Yie, J., Merika, M., Munshi, N., Chen, G. & Thanos, D. The role of HMGI(Y) in the assembly and function of the IFN- β enhanceosome. *EMBO J.* **18**, 3074–3089 (1999).
10. Reeves, R., Leonard, W. J. & Nissen, M. S. Binding of HMGI(Y) imparts architectural specificity to a positioned nucleosome on the promoter of the human interleukin-2 receptor α gene. *Mol. Cell. Biol.* **20**, 4666–4679 (2000).
11. Brunetti, A., Manfioletti, G., Chiefari, E., Goldfine, I. D. & Foti, D. Transcriptional regulation of human insulin receptor gene by the high-mobility group protein HMGI(Y). *FASEB J.* **15**, 492–500 (2001).
12. Watanabe, M. & Whitman, M. FAST-1 is a key maternal effector of mesoderm inducers in the early *Xenopus* embryo. *Development* **126**, 5621–5634 (1999).
13. Schultheiss, T. M., Burch, J. B. & Lassar, A. B. A role for bone morphogenetic proteins in the induction of cardiac myogenesis. *Genes Dev.* **11**, 451–462 (1997).
14. Heldin, C. H., Miyazono, K. & ten Dijke, P. TGF β signalling from cell membrane to nucleus through SMAD proteins. *Nature* **390**, 465–471 (1997).
15. Liberatore, C. M., Searcy-Schrick, R. D., Vincent, E. B. & Yutzey, K. E. *Nkx-2.5* gene induction in mice is mediated by a Smad consensus regulatory region. *Dev. Biol.* **244**, 243–256 (2002).
16. Lien, C. L., McAnally, J., Richardson, J. A. & Olson, E. N. Cardiac-specific activity of an Nkx2-5 enhancer requires an evolutionarily conserved Smad binding site. *Dev. Biol.* **244**, 257–266 (2002).
17. Schwartz, R. J. & Olson, E. N. Building the heart piece by piece: modularity of cis-elements regulating Nkx2-5 transcription. *Development* **126**, 4187–4192 (1999).
18. Sparrow, D. B. *et al.* Regulation of the tinman homologues in *Xenopus* embryos. *Dev. Biol.* **227**, 65–79 (2000).
19. Fusco, A. & Fedele, M. Roles of HMGA proteins in cancer. *Nature Rev. Cancer* **7**, 899–910 (2007).
20. Battista, S. *et al.* The expression of a truncated HMGI-C gene induces gigantism associated with lipomatosis. *Cancer Res.* **59**, 4793–4797 (1999).
21. Ariotta, P. *et al.* Transgenic mice expressing a truncated form of the high mobility group I-C protein develop adiposity and an abnormally high prevalence of lipomas. *J. Biol. Chem.* **275**, 14394–14400 (2000).
22. Zhou, X., Benson, K. F., Ashar, H. R. & Chada, K. Mutation responsible for the mouse pygmy phenotype in the developmentally regulated factor HMGI-C. *Nature* **376**, 771–774 (1995).
23. Anand, A. & Chada, K. *In vivo* modulation of Hmgic reduces obesity. *Nature Genet.* **24**, 377–380 (2000).
24. Imamura, T. *et al.* Smad6 inhibits signalling by the TGF- β superfamily. *Nature* **389**, 622–626 (1997).
25. Korchynskiy, O. & ten Dijke, P. Identification and functional characterization of distinct critically important bone morphogenetic protein-specific response elements in the Id1 promoter. *J. Biol. Chem.* **277**, 4883–4891 (2002).
26. Breckenridge, R. A., Mohun, T. J. & Amaya, E. A role for BMP signalling in heart looping morphogenesis in *Xenopus*. *Dev. Biol.* **232**, 191–203 (2001).
27. Hosoda, T. *et al.* A novel myocyte-specific gene *Midori* promotes the differentiation of P19CL6 cells into cardiomyocytes. *J. Biol. Chem.* **276**, 35978–35989 (2001).
28. Naito, A. T. *et al.* Developmental stage-specific biphasic roles of Wnt/ β -catenin signalling in cardiomyogenesis and hematopoiesis. *Proc. Natl Acad. Sci. USA* **103**, 19812–19817 (2006).
29. Michiue, T. *et al.* Xldax, an inhibitor of the canonical Wnt pathway, is required for anterior neural structure formation in *Xenopus*. *Dev. Dyn.* **230**, 79–90 (2004).
30. Hiroi, Y. *et al.* Tbx5 associates with Nkx2-5 and synergistically promotes cardiomyocyte differentiation. *Nature Genet.* **28**, 276–280 (2001).
31. Kroll, K. L. & Amaya, E. Transgenic *Xenopus* embryos from sperm nuclear transplantations reveal FGF signaling requirements during gastrulation. *Development* **122**, 3173–3183 (1996).

Vascular Endothelial Growth Factor Receptor-1 Regulates Postnatal Angiogenesis Through Inhibition of the Excessive Activation of Akt

Jun-ichiro Nishi,* Tohru Minamino,* Hideyuki Miyauchi, Aika Nojima, Kaoru Tateno, Sho Okada, Masayuki Orimo, Junji Moriya, Guo-Hua Fong, Kenji Sunagawa, Masabumi Shibuya, Issei Komuro

Abstract—Vascular endothelial growth factor (VEGF) binds both VEGF receptor-1 (VEGFR-1) and VEGF receptor-2 (VEGFR-2). Activation of VEGFR-2 is thought to play a major role in the regulation of endothelial function by VEGF. Recently, specific ligands for VEGFR-1 have been reported to have beneficial effects when used to treat ischemic diseases. However, the role of VEGFR-1 in angiogenesis is not fully understood. In this study, we showed that VEGFR-1 performs “fine tuning” of VEGF signaling to induce neovascularization. We examined the effects of retroviral vectors expressing a small interference RNA that targeted either the VEGFR-1 gene or the VEGFR-2 gene. Deletion of either VEGFR-1 or VEGFR-2 reduced the ability of endothelial cells to form capillaries. Deletion of VEGFR-1 markedly reduced endothelial cell proliferation and induced premature senescence of endothelial cells. In contrast, deletion of VEGFR-2 significantly impaired endothelial cell survival. When VEGFR-1 expression was blocked, VEGF constitutively activated Akt signals and thus induced endothelial cell senescence via a p53-dependent pathway. VEGFR-1^{+/-} mice exhibited an increase of endothelial Akt activity and showed an impaired neovascularization in response to ischemia, and this impairment was ameliorated in VEGFR-1^{+/-} Akt1^{+/-} mice. These results suggest that VEGFR-1 plays a critical role in the maintenance of endothelial integrity by modulating the VEGF/Akt signaling pathway. (*Circ Res.* 2008;103:261-268.)

Key Words: VEGF ■ Akt ■ senescence ■ p53

Angiogenesis involves the differentiation, proliferation, and migration of endothelial cells, leading to tubulogenesis and the formation of vessels.¹ One of the most important receptors for angiogenesis is the vascular endothelial growth factor (VEGF) receptor, which is a member of the receptor tyrosine kinase family.^{2,3} VEGF receptor (VEGFR)-1 and VEGFR-2 are closely related receptor tyrosine kinases and have both common and specific ligands. VEGFR-1 has weaker kinase activity, whereas VEGFR-2 is a highly active kinase that stimulates a variety of signaling pathways and induces a broad range of biological responses. Despite its weak kinase activity, VEGFR-1 is essential for normal development and angiogenesis.⁴ VEGFR-1 null mutant mice die in utero because of the overgrowth of endothelial cells and vascular disorganization.^{5,6} In contrast, mice expressing the VEGFR-1 that lacks the tyrosine kinase domain develop a normal cardiovascular system,⁷ suggesting that VEGFR-1 kinase activity might not be required for

vascular development during embryogenesis and that VEGFR-1 may act as a decoy receptor. Consistent with this concept, selective activation of chimeric VEGFR-1 (in the absence of chimeric VEGFR-2)⁸ or a VEGF mutant that binds to VEGFR-1 does not influence cell proliferation, migration, or survival in vitro.⁹⁻¹¹

However, recent studies have demonstrated that the role of VEGFR-1 in postnatal angiogenesis is more complicated than was initially recognized. For example, treatment with placenta growth factor (PlGF), a specific ligand for VEGFR-1, was reported to promote angiogenesis in vitro^{11,12} and in vivo.¹³ Overexpression of PlGF also induced angiogenesis in tumors¹⁴ and the skin.¹⁵ It has been suggested that stimulation by PlGF induces the heterodimerization of VEGFR-1 with VEGFR-2, leading to transactivation of VEGFR-2 and the promotion of angiogenesis.^{8,16,17} Another possible explanation for the positive effect of PlGF on angiogenesis is that it prevents VEGF from binding to VEGFR-1, thereby

Original received July 3, 2007; resubmission received February 18, 2008; revised resubmission received June 11, 2008; accepted June 16, 2008.

From the Department of Cardiovascular Science and Medicine (J.N., T.M., H.M., A.N., K.T., S.O., M.O., J.M., I.K.), Chiba University Graduate School of Medicine, Japan; PRESTO (T.M.), Japan Science and Technology Agency, Saitama, Japan; the Department of Physiology (G.-H.F.), University of Connecticut Health Center, Farmington; the Department of Cardiovascular Medicine (J.N., K.S.), Kyushu University Graduate School of Medical Sciences, Fukuoka, Japan; and the Department of Molecular Oncology (M.S.), Graduate School of Medicine and Dentistry, Tokyo Medical and Dental University, Japan.

*These authors contributed equally to this study.

Correspondence to Issei Komuro, MD, PhD, Department of Cardiovascular Science and Medicine, Chiba University Graduate School of Medicine, 1-8-1 Inohana, Chuo-ku, Chiba 260-8670, Japan. E-mail komuro-ty@umin.ac.jp

© 2008 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.108.174128

Downloaded from circres.ahajournals.org at SWETS SUBS SERV-#25476254 on March 30, 2011

increasing the binding and activation of VEGFR-2. In other studies, PIGF was shown to protect against hyperoxic vascular damage in the retina without provoking retinal neovascularization.¹⁸ These results suggest that VEGFR-1 can either positively or negatively regulate angiogenesis depending on the circumstances, but further studies are required to better understand the role of this receptor in postnatal angiogenesis.

In the present study, we examined the effects of VEGFR-1 deletion on angiogenesis by using the retroviral vector expressing a small interference RNA that targeted the VEGFR-1 gene. Deletion of VEGFR-1 markedly reduced endothelial cell proliferation and thus impaired angiogenesis. Likewise, VEGFR-1^{+/-} mice exhibited an impaired neovascularization in response to ischemia. This impairment was restored by inhibiting the excessive activation of Akt by VEGF. These results suggest that VEGFR-1 plays a critical role in the maintenance of endothelial integrity by modulating the VEGF/Akt signaling pathway.

Materials and Methods

Short Hairpin Interference RNA Vectors

The mammalian retrovirus expression vector pSIREN-RetroQ (Clontech) was used to achieve the expression of short hairpin interference RNA (shRNA) in human endothelial cells.

Statistical Analysis

Data are shown as mean ± SEM. Differences between groups were examined by Student *t* test or ANOVA followed by the Bonferroni procedure for comparison of means. Values of *P* < 0.05 were considered statistically significant.

Results

Effect of VEGF Receptor Gene Silencing on Endothelial Cell Function

To elucidate the role of VEGFR-1 in angiogenesis, we constructed mammalian retroviral vectors expressing a short hairpin interference RNA that targeted either the VEGFR-1 gene (shVEGFR-1) or the VEGFR-2 gene (shVEGFR-2). Northern blot and Western blot analyses revealed that introduction of each construct into human umbilical vein endothelial cells caused effective and stable downregulation of the expression of the target molecule (Figure 1A and 1B, and supplemental Figure 1A [available online at <http://circres.ahajournals.org>]). It is noted that either shVEGFR-1 or shVEGFR-2 did not affect VEGFR-2 or VEGFR-1 expression, respectively (Figure 1B, and supplemental Figure 1A). We used two kinds of constructs for the following experiments and both of them achieved similar results. The nonsilencing control vector (shNega) was used as a control. After infected endothelial cells were purified by incubation with antibiotics, we performed the tube formation assay. Deletion of VEGFR-1 or VEGFR-2 significantly impaired tube formation compared with control cells (Figure 1C). We next examined the proliferative activity of infected cells. We seeded 2 × 10⁵ infected cells into 100-mm dishes with VEGF-A on day 0 and counted cell number on day 3. Compared with shNega-infected control endothelial cells, both shVEGFR-1- and

shVEGFR-2-infected cells showed significantly lower proliferation (Figure 1D). Deletion of VEGFR-1 caused more marked impairment of cell proliferation than deletion of VEGFR-2 (Figure 1D). This inhibitory effect of VEGFR-1 deletion was more evident when infected endothelial cells were subjected to long-term culture. Although VEGFR-2 deletion slightly reduced the lifespan of cells compared with that of control cells, VEGFR-1 deletion significantly shortened the lifespan of endothelial cells (Figure 1E). As a result, shVEGFR-1-infected cells underwent irreversible growth arrest earlier than shVEGFR-2-infected cells (Figure 1E). After growth arrest, the cells exhibited characteristics of senescence, becoming flatter and larger and showing an increase of senescence-associated β-galactosidase activity (Figure 1F). These findings suggest that VEGFR-1 deletion induces premature endothelial cell senescence. We next examined the effect of VEGFR-1 deletion on endothelial survival. We cultured infected cells in regular growth medium for 24 hours and subsequently cultured the cells under serum-free conditions with VEGF-A. After 24 hours, the number of viable cells was counted. As compared with the viability of control cells, deletion of VEGFR-2, but not VEGFR-1, markedly decreased cell viability (Figure 1G). Consistent with these findings, activation of caspase 3 was detected in cells with VEGFR-2 deletion, but not VEGFR-1 deletion (Figure 1H). These results suggest that VEGFR-1 is involved in the regulation of angiogenesis by regulating endothelial cell proliferation and senescence, whereas VEGFR-2 may be crucial for endothelial survival as well as cell proliferation.

VEGFR-1 Deletion Induces Endothelial Dysfunction by Activating Akt

To investigate the molecular mechanisms of premature senescence induced by VEGFR-1 deletion, we examined the transcriptional activity of p53 and its target gene p21. We transfected VEGFR-1-deleted endothelial cells with the luciferase reporter gene containing 13 copies of the p53-binding consensus sequence (pPG13-Luc). Deletion of VEGFR-1 significantly induced p53 transcriptional activity compared with that in shNega-infected cells, whereas VEGFR-2 deletion had no effect (Figure 2A). Likewise, p21 expression was significantly higher in VEGFR-1-deleted endothelial cells than in control cells or VEGFR-2-deleted cells (Figure 2B). However, expression of bax, another target molecule regulated by p53, was not altered in VEGFR-1-deleted endothelial cells compared with control cells (supplemental Figure 1B). Ablation of p53 by the introduction of HPV16 E6 oncoprotein abolished the inhibitory effect of VEGFR-1 deletion on cell proliferation (Figure 2C). These results suggest that VEGFR-1 deletion induces endothelial cell senescence via a p53-dependent pathway.

We have previously demonstrated that Akt negatively regulates the endothelial cell lifespan by activating the p53/p21 pathway.¹⁹ It has also been shown that Akt plays a central role in the regulation of angiogenesis by VEGF.²⁰ Thus, we examined the level of phosphorylated Akt in VEGFR-1-deleted endothelial cells. Western blot analysis




Figure 1. Effect of VEGF receptor gene silencing on endothelial cell function. A, Human umbilical vein endothelial cells were infected with retroviral vectors expressing a short hairpin interference RNA that targeted either the VEGFR-1 gene (shVEGFR-1a and shVEGFR-1b) or the VEGFR-2 gene (shVEGFR-2a and shVEGFR-2b), and then the cells were purified by culture with antibiotics. shNega served as the control vector. Total RNA (30 μ g) was extracted from infected cells and analyzed to assess the expression of VEGFR-1 or VEGFR-2 by Northern blot analysis. * P <0.05, ** P <0.01, and *** P <0.001 vs shNega (n =3). B, Total RNA (30 μ g) was extracted from endothelial cells infected with shVEGFR-1 or shVEGFR-2 and simultaneously analyzed the expression of VEGFR-1 and VEGFR-2 by Northern blot analysis. C, Infected endothelial cells were seeded into 96-well plates in serum-free basic medium with VEGF-A (50 ng/mL). After 16 hours, capillary-like tube formation was

estimated by using an angiogenesis image analyzer. * P <0.01, ** P <0.0001 vs shNega (n =4 to 6). Scale bar: 300 μ m. D, Infected endothelial cells were seeded at a density of 2×10^5 cells per 100-mm dish and cultured with VEGF-A (day 0). Then cell number was counted on day 3. * P <0.001, ** P <0.0001 vs shNega, # P <0.001 vs shVEGFR-2 (n =13 to 14). E, Infected cell populations were passaged until cells underwent senescence, and the total number of population doublings was determined. * P <0.01 vs shNega, # P <0.05 vs shVEGFR-2 (n =4 to 6). F, Morphology and senescence-associated β -galactosidase staining (arrow) of endothelial cells infected with shNega, shVEGFR-1, or shVEGFR-2. Scale bar: 100 μ m. G, Infected endothelial cells were seeded at the density of 1×10^5 cells per 60-mm dish and cultured for 24 hours in growth medium. After washing twice with PBS, the cells were cultured in serum-free DMEM with VEGF-A (10 ng/mL). After 24 hours of serum starvation, the number of viable cells and the total number of cells were counted by a hemocytometer. * P <0.0001 vs shNega (n =4 to 6). H, The lysates were extracted from cells, which are prepared as described in legend to G, and analyzed for cleaved caspase-3 expression by Western blotting.

showed that VEGFR-1 deletion led to a marked increase of the phosphorylated Akt level compared with that in control cells or cells with VEGFR-2 deletion, even under serum-free conditions (Figure 3A). VEGFR-1 deletion increased pAkt levels even in the absence of VEGF, presumably attributable to autocrine VEGF signaling (Figure 3B). Treatment with VEGF markedly increased pAkt levels within 5 to 15 minutes in VEGFR-1–deleted cells but not in VEGFR-2–deleted cells (Figure 3B). Treatment with a neutralizing anti-VEGF antibody reduced the phosphorylated Akt level in VEGFR-1–deleted cells (Figure 3C), suggesting that VEGFR-1 inhibits the activation of Akt by VEGF. To further investigate the relationship between constitutive Akt activation and endothelial cell dysfunction induced by VEGFR-1 deletion, we examined the effect of inhibition of Akt. We infected human endothelial cells with a retroviral vector encoding a dominant-negative form of Akt (DN-Akt)¹⁹ or the empty vector encoding resistance to neomycin alone (Mock). Both cell populations were then infected with shNega or

shVEGFR-1. We found that VEGFR-1 deletion markedly inhibited the proliferation of mock-infected endothelial cells (Figure 3D, Mock), whereas this inhibitory effect was significantly ameliorated in DN-Akt–infected cells (Figure 3D, DN-Akt). Consequently, VEGFR-1 deletion significantly impaired tube formation by mock-infected cells, but not DN-Akt–infected cells (Figure 3E). Likewise, inhibition of Akt activation prevented the induction of p21 expression by VEGFR-1 deletion (supplemental Figure II). These results suggest that VEGFR-1 deletion causes dysregulation of activation of the VEGFR-2/Akt signaling pathway by VEGF-A, and that constitutive activation of Akt is related to the impaired ability of VEGFR-1–deleted endothelial cells to proliferate and form capillary-like structures. VEGF-induced phosphorylation of eNOS was enhanced, but production of cGMP was significantly reduced by VEGFR-1 deletion, presumably because constitutive activation of Akt increases cellular reactive oxygen species¹⁹ that inactivate this enzyme (supplemental Figure IC and ID).




Figure 2. VEGFR-1 deletion induces activation of the p53/p21 signal pathway. **A**, A luciferase reporter gene plasmid (pPG13-Luc) containing the p53-binding sequence was transfected into endothelial cells infected with shNega, shVEGFR-1, or shVEGFR-2. Luciferase activity was measured at 48 hours after transfection in the presence of VEGF-A (10 ng/mL) as described in Methods. * $P < 0.05$ vs shNega (n=5). **B**, Whole cell lysates (30 μ g) were prepared from infected endothelial cells and p21 expression was assessed by Western blot analysis. * $P < 0.05$ vs shNega, # $P < 0.01$ vs shVEGFR-2 (n=4). **C**, Human endothelial cells were infected with pLNCX (Mock) or pLNCX E6 (E6). Infected cell populations were then transduced with shNega or shVEGFR-1. After purification, double-infected cells were seeded at a density of 2×10^5 cells per 100-mm dish in the presence of VEGF-A (day 0), and cell number was counted on day 3. * $P < 0.05$ vs Mock/shNega (n=3). Western blot analysis revealed that introduction of E6 effectively ablated p53 expression (right panel).

Influence of VEGFR-1 Deletion on Neovascularization In Vivo

To examine the influence of VEGFR-1 deletion on neovascularization in vivo, we produced a hindlimb ischemia model in VEGFR-1^{+/-} mice and assessed blood flow recovery and the capillary density of ischemic tissue. VEGFR-1 mRNA levels were significantly lower in VEGFR-1^{+/-} mice than in wild-type mice (Figure 4A). Aortic expression of VEGFR-1 protein was decreased in VEGFR-1^{+/-} mice compared with wild-type mice (Figure 4B). Consistent with the in vitro data, phospho-Akt levels were significantly higher in VEGFR-1^{+/-} mice than in wild-type mice (Figure 4C and supplemental Figure III). There was no significant difference in plasma VEGF levels between the two groups (data not shown). Laser Doppler image analysis revealed that blood flow recovery

was significantly impaired in VEGFR-1^{+/-} mice compared with their wild-type littermates (Figure 4D). Likewise, VEGFR-1^{+/-} mice exhibited significantly fewer CD31-positive cells in the ischemic tissues than their wild-type littermates (Figure 4E), suggesting that decreased expression of VEGFR-1 led to reduced neovascularization of ischemic tissue.

There are several reports indicating that VEGFR-1 kinase activity is required for VEGF-induced migration of hematopoietic cells including macrophages,²¹⁻²⁶ and it was reported that infiltration of macrophages plays a critical role in pathological angiogenesis during ischemia, inflammation, and tumor development.²⁷⁻²⁹ Therefore, we examined the number of infiltrating macrophages in ischemic tissue, but we found no significant difference in the number of Mac3-




Figure 3. VEGFR-1 deletion induces endothelial dysfunction by activating Akt. **A**, Human endothelial cells were infected with shVEGFR-1 or shVEGFR-2. Infected cells were cultured in the presence of VEGF-A (10 ng/mL). Whole cell lysates (30 μ g) were prepared and the expression of phosphorylated Akt (pAkt) was detected by Western blot analysis. * $P < 0.05$ vs shNega, # $P < 0.05$ vs shVEGFR-2 (n=5). **B**, Infected cells were cultured in serum-free basal medium (without VEGF-A) for 8 hours and subsequently treated with VEGF-A (10 ng/mL) for 5 to 15 minutes. Whole cell lysates were extracted at indicated times and phospho-Akt (pAkt) expression was investigated by Western blot analysis. **C**, Infected cells were treated with a neutralizing antibody for VEGF (500 ng/mL) (+) or a control antibody (-) for 24 hours. Whole cell lysates were extracted and phospho-Akt expression was assessed by Western blot analysis. **D**, Human endothelial cells were

infected with pLNCX (Mock) or pLNCX DN-Akt (DN-Akt). Infected cell populations were then transduced with shNega or shVEGFR-1 and were subjected to the proliferation assay as described in legend for Figure 2C. * $P < 0.005$ vs Mock/shNega, # $P < 0.005$ vs Mock/shVEGFR-1 (n=6 to 8). Expression of c-Myc-tagged DN-Akt was confirmed by Western blot analysis (right panel). **E**, Double-infected endothelial cells (prepared as in Figure 3C) were subjected to the tube-forming assay. * $P < 0.05$ vs Mock/shNega (n=3).




Figure 4. Impaired neovascularization after ischemia in VEGFR-1^{+/-} mice. A, Total RNA (30 μg) was extracted from the lung of VEGFR-1^{+/-} mice and wild-type littermates to investigate VEGFR-1 expression by Northern blot analysis. *P<0.001 vs wild-type littermates (n=5). B, Whole cell lysates (30 μg) were prepared from the aorta of VEGFR-1^{+/-} mice and wild-type littermates to investigate VEGFR-1 expression by Western blot analysis. *P<0.05 vs wild-type littermates (n=3). C, Whole cell lysates (30 μg) were prepared as described in Figure 4B to investigate phospho-Akt (pAkt) expression by Western blot analysis. *P<0.05 vs wild-type littermates (n=3). D, Limb perfusion was measured by a laser Doppler analyzer at 1 to 3 weeks after ischemia. The graph shows the ratio of ischemic (right) to nonischemic limb (left) blood flow. *P<0.05 vs wild-type littermates (n=16). E, Immunohistochemistry for CD31 (brown) in ischemic limbs. Scale bar: 50 μm. The number of CD31-positive cells per square millimeter is shown in the graph. *P<0.05 vs wild-type littermates (n=4).

nonischemic limb (left) blood flow. *P<0.05 vs wild-type littermates (n=16). E, Immunohistochemistry for CD31 (brown) in ischemic limbs. Scale bar: 50 μm. The number of CD31-positive cells per square millimeter is shown in the graph. *P<0.05 vs wild-type littermates (n=4).

positive cells between VEGFR-1^{+/-} mice and their wild-type littermates (Figure 5A). To further test the possible involvement of bone marrow-derived cells, we transplanted wild-type bone marrow cells into VEGFR-1^{+/-} mice or their wild-type littermates. We then produced a hindlimb ischemia model and assessed blood flow recovery and the capillary density of ischemic tissue. Despite the transplantation of wild-type bone marrow, blood flow recovery was still significantly impaired in VEGFR-1^{+/-} mice (Figure 5B). The number of CD31-positive cells was also lower in VEGFR-1^{+/-} mice than in their wild-type littermates (Figure 5C). Thus, it is unlikely that impaired neovascularization in VEGFR-1^{+/-} mice is attributed to reduced migration of bone marrow-derived cells. We could not detect VEGFR-1 expression in muscle cells (supplemental Figure IV). It was noted that the number of endothelial cells double positive for phospho-Akt and CD31 was significantly higher in VEGFR-1^{+/-} mice than in their wild-type littermates (Figure 5D).

Inhibition of Akt Signaling Ameliorates the Impairment of Neovascularization in VEGFR-1^{+/-} Mice

Next, we examined whether an increase of endothelial Akt activity contributed to impaired neovascularization in VEGFR-1^{+/-} mice. Akt1 is the predominant isoform of Akt in endothelial cells and is thought to play an important role in postnatal angiogenesis.³⁰ It has been reported that the angiogenic response of Akt1^{-/-} mice was enhanced in a tumor angiogenesis model, but was decreased in a hindlimb ischemia

model.^{30,31} so we thus used Akt1^{+/-} mice for our in vivo experiments. Consistent with the previous reports,³² phospho-Akt levels were lower in the aorta of Akt1^{+/-} mice compared with wild-type littermates (supplemental Figure V). After creating hindlimb ischemia in VEGFR-1^{+/-} Akt1^{+/-} mice, we examined the extent of blood flow recovery and the capillary density 1 week later. We found that there were no significant differences of blood flow recovery and capillary density between Akt1^{+/-} mice and Akt1^{+/+} mice (Figure 6A and 6B). Decreased VEGFR-1 expression significantly reduced blood flow recovery in Akt1^{+/+} mice, but not in Akt1^{+/-} mice (Figure 6A). Likewise, the capillary density of ischemic tissue was significantly reduced in VEGFR-1^{+/-} Akt1^{+/+} mice compared with wild-type mice, but VEGFR-1^{+/-} Akt1^{+/-} mice had a similar capillary density to that of VEGFR-1^{+/+} Akt1^{+/-} mice (Figure 6B). These results suggest that an increase of endothelial Akt activity may be responsible for impaired neovascularization in VEGFR-1^{+/-} mice.

Discussion

In the present study, we demonstrated that VEGFR-1 modulates postnatal angiogenesis through inhibition of the excessive activation of Akt by VEGF. It has been reported that VEGF and VEGFR-1 can be simultaneously induced by various stimuli, including hypoxia.³³ Thus, the role of VEGFR-1 may vary, depending on the extent of activation of Akt. For example, when overproduction of growth factors such as VEGF and insulin leads to excessive activation of Akt and impairs normal regulation of endothelial proliferation,




Figure 5. Role of bone marrow–derived cells in impaired neovascularization in VEGFR-1^{+/-} mice. A, Immunohistochemistry for Mac3 (brown) in ischemic limbs. Scale bar: 50 μ m. The number of Mac3-positive cells per square millimeter is shown (n=4). B, Wild-type bone marrow cells were transplanted into VEGFR-1^{+/-} mice or their wild-type littermates. Limb perfusion was measured by a laser Doppler analyzer at 1 week after ischemia. *P<0.05 vs wild-type littermates (n=6). C, Immunohistochemistry for CD31 (brown) in ischemic limbs of bone marrow–transplanted mice. Scale bar: 50 μ m. *P<0.05 vs wild-type littermates (n=6). D, Activation of Akt in endothelial cells of ischemic limbs from VEGFR-1^{+/-} mice. Representative immunostainings for phospho-Akt (red) and CD31 (green) were shown. Arrows indicate phospho-Akt/CD31-positive cells (yellow). Scale bar: 50 μ m. The graph shows the ratio of phospho-Akt/CD31-positive cell number to all CD31-positive cell number. *P<0.05 vs wild-type littermates (n=5).

VEGFR-1 may act as a positive regulator of angiogenesis by inhibiting activation of VEGFR-2. Conversely, VEGFR-1 may exert a negative effect on angiogenesis when growth factors appropriately activate the Akt signaling pathway to induce endothelial cell proliferation. These mechanisms may provide an explanation as to why the effects of PIGF on angiogenesis were reported to differ.

Although there is evidence to suggest that VEGFR-1 interacts with the p85 subunit of phosphatidylinositol-3 ki-

nase (PI3K) to regulate its activity,^{34–36} VEGFR-1 appears to exert its inhibitory effect on angiogenesis mainly by blocking the activation of Akt mediated by VEGF via VEGFR-2 for the following reasons. First, treatment with VEGF-A increased Akt activity in VEGFR-1–deleted cells, but not in VEGFR-2–deleted cells (Figure 3A and 3B). Second, treatment with a neutralizing anti-VEGF antibody reduced the enhanced activation of Akt in VEGFR-1–deleted cells (Figure 3C). Finally, treatment with PIGF did not provoke any




Figure 6. Inhibition of Akt signaling ameliorates the impairment of neovascularization in VEGFR-1^{+/-} mice. A, Limb perfusion was measured by a laser Doppler analyzer at 1 week after creation of ischemia. *P<0.01 vs wild-type littermates (n=14 to 18). B, Immunohistochemistry for CD31 (brown) in ischemic limbs. Scale bar: 50 μ m. *P<0.05 vs wild-type littermates (n=6 to 7).

biological response in the presence of anti-VEGF antibody (J. Nishi, T. Minamino, unpublished data, 2007). Our results are consistent with previous studies^{37,38} demonstrating that tyrosine phosphorylation of VEGFR-2 was elevated in VEGFR-1-deficient embryonic stem cells, whereas loss of VEGFR-1 led to decreased sprout formation and migration, which resulted in reduced vascular branching. This reduction was restored by blockade of the VEGFR-2 signaling pathway as well as by treatment with soluble VEGFR-1. Although Bussolati et al demonstrated that VEGFR-1 but not VEGFR-2 increases endothelial production of NO, thereby promoting tube formation,³⁹ cGMP production was significantly decreased in VEGFR-1-deleted endothelial cells (supplemental Figure ID). Moreover, VEGF treatment failed to activate Akt in VEGFR-2-deleted endothelial cells (Figure 3B) and introduction of mutant VEGFR-1 lacking the sites for interaction with PI3K did not mimic the effects of shVEGFR-1 (J. Nishi, T. Minamino, unpublished data, 2007). Taken together, these results suggest that VEGFR-1 acts to provide “fine tuning” of VEGF signaling to achieve the proper formation of blood vessels. The biological consequences of VEGFR-1 deletion appears to be related to loss of its decoy effect, but other mechanisms might be involved such as “cross talk” between VEGFR-1 and VEGFR-2.^{8,16,17} direct regulation of the VEGFR-2 signaling pathway by VEGFR-1,^{39,40} and some undefined effect of the extracellular domain of membrane-bound VEGFR-1.⁴¹

We have previously demonstrated that constitutive activation of Akt induced by insulin promotes senescence-like arrest of endothelial cell growth via a p53/p21-dependent pathway.¹⁹ Moreover, tube formation was significantly reduced by overactivation of Akt. Likewise, constitutive activation of Akt has been reported to promote the senescence in other types of cells such as endothelial progenitors and mouse embryonic fibroblasts.^{42,43} The study using conditional transgenic mice has demonstrated that sustained activation of Akt in endothelial cells causes increased blood vessel size and generalized edema within 2 weeks and that these changes are reversible.⁴⁴ Using the same mouse model, it has been reported that chronic activation of Akt over 8 weeks leads to endothelial cell senescence and loss of endothelium-dependent stroke protection.⁴⁵ Recent studies by several groups demonstrated that diabetic state induces activation of the Akt pathway, thereby contributing to the pathology of diabetic complications.^{42,46–48} We also detected increased Akt activity in endothelial cells on the surface of coronary atherosclerotic lesions in patients with diabetes.¹⁹ Moreover, accumulating evidence suggests that vascular cell senescence contributes to the pathogenesis of age-associated vascular diseases including diabetic vasculopathy.⁴⁹ Thus, these results suggest the potential of the treatment for vascular dysfunction associated with diabetes and aging by modulating Akt activity with a soluble form of VEGFR-1.

Acknowledgments

We thank Dr B. Vogelstein and Dr T Zioncheck for reagents, Dr M. Birnbaum for mice, and E. Fujita, Y. Ishiyama, R. Kobayashi, and Y. Ishikawa for their excellent technical assistance.

Sources of Funding

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture, and Health and Labor Sciences Research Grants (to I.K.) and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan, and the grants from the Suzuken Memorial Foundation, the Japan Diabetes Foundation, the Ichiro Kanehara Foundation, the Tokyo Biochemical Research Foundation, the Takeda Science Foundation, the Cell Science Research Foundation, and the Japan Foundation of Applied Enzymology (to T.M.).

Disclosures

None.

References

- Carmeliet P. Angiogenesis in life, disease and medicine. *Nature*. 2005; 438:932–936.
- Ferrara N, Gerber HP, LeCouter J. The biology of VEGF and its receptors. *Nat Med*. 2003;9:669–676.
- Coults L, Chawengsaksophak K, Rossant J. Endothelial cells and VEGF in vascular development. *Nature*. 2005;438:937–945.
- Fong GH, Rossant J, Gertsenstein M, Breitman ML. Role of the Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium. *Nature*. 1995;376:66–70.
- Fong GH, Zhang L, Bryce DM, Peng J. Increased hemangioblast commitment, not vascular disorganization, is the primary defect in flt-1 knock-out mice. *Development*. 1999;126:3015–3025.
- Kearney JB, Ambler CA, Monaco KA, Johnson N, Rapoport RG, Bautch VL. Vascular endothelial growth factor receptor Flt-1 negatively regulates developmental blood vessel formation by modulating endothelial cell division. *Blood*. 2002;99:2397–2407.
- Hiratsuka S, Minowa O, Kuno J, Noda T, Shibuya M. Flt-1 lacking the tyrosine kinase domain is sufficient for normal development and angiogenesis in mice. *Proc Natl Acad Sci U S A*. 1998;95:9349–9354.
- Rahimi N, Dayanir V, Lashkari K. Receptor chimeras indicate that the vascular endothelial growth factor receptor-1 (VEGFR-1) modulates mitogenic activity of VEGFR-2 in endothelial cells. *J Biol Chem*. 2000;275: 16986–16992.
- Keyt BA, Nguyen HV, Berleau LT, Duarte CM, Park J, Chen H, Ferrara N. Identification of vascular endothelial growth factor determinants for binding KDR and FLT-1 receptors. Generation of receptor-selective VEGF variants by site-directed mutagenesis. *J Biol Chem*. 1996;271: 5638–5646.
- Yang S, Xin X, Zlot C, Ingle G, Fuh G, Li B, Moffat B, de Vos AM, Gerritsen ME. Vascular endothelial cell growth factor-driven endothelial tube formation is mediated by vascular endothelial cell growth factor receptor-2, a kinase insert domain-containing receptor. *Arterioscler Thromb Vasc Biol*. 2001;21:1934–1940.
- Gille H, Kowalski J, Li B, LeCouter J, Moffat B, Zioncheck TF, Pelletier N, Ferrara N. Analysis of biological effects and signaling properties of Flt-1 (VEGFR-1) and KDR (VEGFR-2). A reassessment using novel receptor-specific vascular endothelial growth factor mutants. *J Biol Chem*. 2001;276:3222–3230.
- Errico M, Riccioni T, Iyer S, Pisano C, Acharya KR, Persico MG, De Falco S. Identification of placenta growth factor determinants for binding and activation of Flt-1 receptor. *J Biol Chem*. 2004;279:43929–43939.
- Luttun A, Tjwa M, Moons L, Wu Y, Angelillo-Scherrer A, Liao F, Nagy JA, Hooper A, Priller J, De Klerck B, Compemolle V, Daci E, Bohlen P, Dewerchin M, Herbert JM, Fava R, Matthys P, Carmeliet G, Collen D, Dvorak HF, Hicklin DJ, Carmeliet P. Revascularization of ischemic tissues by PIGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nat Med*. 2002;8:831–840.
- Adini A, Kornaga T, Firoozbakht F, Benjamin LE. Placental growth factor is a survival factor for tumor endothelial cells and macrophages. *Cancer Res*. 2002;62:2749–2752.
- Odorisio T, Schietroma C, Zaccaria ML, Cianfarani F, Tiverson C, Tatangelo L, Failla CM, Zambruno G. Mice overexpressing placenta growth factor exhibit increased vascularization and vessel permeability. *J Cell Sci*. 2002;115:2559–2567.
- Autiero M, Waltenberger J, Communi D, Kranz A, Moons L, Lambrechts D, Kroll J, Plaisance S, De Mol M, Bono F, Kliche S, Fellbrich G, Ballmer-Hofer K, Maglione D, Mayr-Beyrle U, Dewerchin M, Dom-

- browski S, Stanimirovic D, Van Hummelen P, Dehio C, Hicklin DJ, Persico G, Herbert JM, Communi D, Shibuya M, Collen D, Conway EM, Carmeliet P. Role of PlGF in the intra- and intermolecular cross talk between the VEGF receptors Flt1 and Flk1. *Nat Med*. 2003;9:936–943.
17. Neaogoe PE, Lemieux C, Sirois MG. Vascular endothelial growth factor (VEGF)-A165-induced prostacyclin synthesis requires the activation of VEGF receptor-1 and -2 heterodimer. *J Biol Chem*. 2005;280:9904–9912.
 18. Shih SC, Ju M, Liu N, Smith LE. Selective stimulation of VEGFR-1 prevents oxygen-induced retinal vascular degeneration in retinopathy of prematurity. *J Clin Invest*. 2003;112:50–57.
 19. Miyauchi H, Minamoto T, Tateno K, Kunieda T, Toko H, Komuro I. Akt negatively regulates the in vitro lifespan of human endothelial cells via a p53/p21-dependent pathway. *Embo J*. 2004;23:2112–220.
 20. O'Neill BT, Abel ED. Akt1 in the cardiovascular system: friend or foe? *J Clin Invest*. 2005;115:2059–2064.
 21. Clauss M, Weich H, Breier G, Knies U, Rockl W, Waltenberger J, Risau W. The vascular endothelial growth factor receptor Flt-1 mediates biological activities. Implications for a functional role of placenta growth factor in monocyte activation and chemotaxis. *J Biol Chem*. 1996;271:17629–17634.
 22. Barleon B, Sozzani S, Zhou D, Weich HA, Mantovani A, Marme D. Migration of human monocytes in response to vascular endothelial growth factor (VEGF) is mediated via the VEGF receptor flt-1. *Blood*. 1996;87:3336–3343.
 23. Sawano A, Iwai S, Sakurai Y, Ito M, Shitara K, Nakahata T, Shibuya M. Flt-1, vascular endothelial growth factor receptor 1, is a novel cell surface marker for the lineage of monocyte-macrophages in humans. *Blood*. 2001;97:785–791.
 24. Lyden D, Hattori K, Dias S, Costa C, Blaikie P, Butros L, Chadburn A, Heissig B, Marks W, Witte L, Wu Y, Hicklin D, Zhu Z, Hackett NR, Crystal RG, Moore MA, Hajjar KA, Manova K, Benezra R, Rafii S. Impaired recruitment of bone-marrow-derived endothelial and hematopoietic precursor cells blocks tumor angiogenesis and growth. *Nat Med*. 2001;7:1194–1201.
 25. Hattori K, Heissig B, Wu Y, Dias S, Tejada R, Ferris B, Hicklin DJ, Zhu Z, Bohlen P, Witte L, Hendrikx J, Hackett NR, Crystal RG, Moore MA, Werb Z, Lyden D, Rafii S. Placental growth factor reconstitutes hematopoiesis by recruiting VEGFR1(+) stem cells from bone-marrow micro-environment. *Nat Med*. 2002;8:841–849.
 26. Jin DK, Shido K, Kopp HG, Petit I, Shmelkov SV, Young LM, Hooper AT, Amano H, Avicilla ST, Heissig B, Hattori K, Zhang F, Hicklin DJ, Wu Y, Zhu Z, Dunn A, Salari H, Werb Z, Hackett NR, Crystal RG, Lyden D, Rafii S. Cytokine-mediated deployment of SDF-1 induces revascularization through recruitment of CXCR4+ hemangiocytes. *Nat Med*. 2006;12:557–567.
 27. Carmeliet P. Mechanisms of angiogenesis and arteriogenesis. *Nat Med*. 2000;6:389–395.
 28. Hiratsuka S, Maru Y, Okada A, Seiki M, Noda T, Shibuya M. Involvement of Flt-1 tyrosine kinase (vascular endothelial growth factor receptor-1) in pathological angiogenesis. *Cancer Res*. 2001;61:1207–1213.
 29. Murakami M, Iwai S, Hiratsuka S, Yamauchi M, Nakamura K, Iwakura Y, Shibuya M. Signaling of vascular endothelial growth factor receptor-1 tyrosine kinase promotes rheumatoid arthritis through activation of monocytes/macrophages. *Blood*. 2006;108:1849–1856.
 30. Chen J, Somanath PR, Razorenova O, Chen WS, Hay N, Bornstein P, Byzova TV. Akt1 regulates pathological angiogenesis, vascular maturation and permeability in vivo. *Nat Med*. 2005;11:1188–1196.
 31. Ackah E, Yu J, Zoellner S, Iwakiri Y, Skurk C, Shibata R, Ouchi N, Easton RM, Galasso G, Birnbaum MJ, Walsh K, Sessa WC. Akt1/protein kinase B α is critical for ischemic and VEGF-mediated angiogenesis. *J Clin Invest*. 2005;115:2119–2127.
 32. Chen WS, Xu PZ, Gottlob K, Chen ML, Sokol K, Shiyanova T, Roninson I, Weng W, Suzuki R, Tobe K, Kadowaki T, Hay N. Growth retardation and increased apoptosis in mice with homozygous disruption of the Akt1 gene. *Genes Dev*. 2001;15:2203–2208.
 33. Pugh CW, Ratcliffe PJ. Regulation of angiogenesis by hypoxia: role of the HIF system. *Nat Med*. 2003;9:677–684.
 34. Cunningham SA, Waxham MN, Arrate PM, Brock TA. Interaction of the Flt-1 tyrosine kinase receptor with the p85 subunit of phosphatidylinositol 3-kinase. Mapping of a novel site involved in binding. *J Biol Chem*. 1995;270:20254–20257.
 35. Igarashi K, Isohara T, Kato T, Shigeta K, Yamano T, Uno I. Tyrosine 1213 of Flt-1 is a major binding site of Nck and SHP-2. *Biochem Biophys Res Commun*. 1998;246:95–99.
 36. Yu Y, Hulmes JD, Herley MT, Whitney RG, Crabb JW, Sato JD. Direct identification of a major autophosphorylation site on vascular endothelial growth factor receptor Flt-1 that mediates phosphatidylinositol 3'-kinase binding. *Biochem J*. 2001;358:465–472.
 37. Roberts DM, Kearney JB, Johnson JH, Rosenberg MP, Kumar R, Bautch VL. The vascular endothelial growth factor (VEGF) receptor Flt-1 (VEGFR-1) modulates Flk-1 (VEGFR-2) signaling during blood vessel formation. *Am J Pathol*. 2004;164:1531–1535.
 38. Kearney JB, Kappas NC, Ellerstrom C, DiPaola FW, Bautch VL. The VEGF receptor flt-1 (VEGFR-1) is a positive modulator of vascular sprout formation and branching morphogenesis. *Blood*. 2004;103:4527–4535.
 39. Bussolati B, Dunk C, Grohman M, Kontos CD, Mason J, Ahmed A. Vascular endothelial growth factor receptor-1 modulates vascular endothelial growth factor-mediated angiogenesis via nitric oxide. *Am J Pathol*. 2001;159:993–1008.
 40. Zeng H, Dvorak HF, Mukhopadhyay D. Vascular permeability factor (VPF)/vascular endothelial growth factor (VEGF) receptor-1 down-modulates VPF/VEGF receptor-2-mediated endothelial cell proliferation, but not migration, through phosphatidylinositol 3-kinase-dependent pathways. *J Biol Chem*. 2001;276:26969–26979.
 41. Hiratsuka S, Nakao K, Nakamura K, Katsuki M, Maru Y, Shibuya M. Membrane fixation of vascular endothelial growth factor receptor 1 ligand-binding domain is important for vasculogenesis and angiogenesis in mice. *Mol Cell Biol*. 2005;25:346–354.
 42. Rosso A, Balsamo A, Gambino R, Dentelli P, Falcioni R, Cassader M, Pegoraro L, Pagano G, Brizzi MF. p53 Mediates the accelerated onset of senescence of endothelial progenitor cells in diabetes. *J Biol Chem*. 2006;281:4339–4347.
 43. Chen Z, Trotman LC, Shaffer D, Lin HK, Dotan ZA, Niki M, Koutcher JA, Scher HI, Ludwig T, Gerald W, Cordon-Cardo C, Pandolfi PP. Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature*. 2005;436:725–730.
 44. Phung TL, Ziv K, Dabydeen D, Eyyah-Mensah G, Riveros M, Perruzzi C, Sun J, Monahan-Earley RA, Shiojima I, Nagy JA, Lin MI, Walsh K, Dvorak AM, Briscoe DM, Neeman M, Sessa WC, Dvorak HF, Benjamin LE. Pathological angiogenesis is induced by sustained Akt signaling and inhibited by rapamycin. *Cancer Cell*. 2006;10:159–170.
 45. Wang C, Kim H, Hiroi Y, Mukai Y, Satoh M, Liao JK. Increase cellular senescence and cerebral infarct size in mice with chronic activation of endothelial protein kinase Akt. *Circulation*. 2006;114:111–116.
 46. Hojlund K, Staehr P, Hansen BF, Green KA, Hardie DG, Richter EA, Beck-Nielsen H, Wojtaszewski JF. Increased phosphorylation of skeletal muscle glycogen synthase at NH₂-terminal sites during physiological hyperinsulinemia in type 2 diabetes. *Diabetes*. 2003;52:1393–1402.
 47. Sheu ML, Ho FM, Yang RS, Chao KF, Lin WW, Lin-Shiau SY, Liu SH. High glucose induces human endothelial cell apoptosis through a phosphoinositide 3-kinase-regulated cyclooxygenase-2 pathway. *Arterioscler Thromb Vasc Biol*. 2005;25:539–545.
 48. Clodfelder-Miller B, De Samo P, Zmijewska AA, Song L, Jope RS. Physiological and pathological changes in glucose regulate brain Akt and glycogen synthase kinase-3. *J Biol Chem*. 2005;280:39723–39731.
 49. Minamoto T, Komuro I. Vascular cell senescence: contribution to atherosclerosis. *Circ Res*. 2007;100:15–26.

Peroxisome Proliferator-Activated Receptor γ and Cardiovascular Diseases

Hiroyuki Takano, MD; Issei Komuro, MD

Peroxisome proliferator-activated receptors (PPARs) are members of the nuclear receptor superfamily and form heterodimers with retinoid X receptor. Three PPAR isoforms have been isolated and termed α , β (or δ) and γ . Although PPAR γ is expressed predominantly in adipose tissue and associated with adipocyte differentiation and glucose homeostasis, PPAR γ is also present in a variety of cell types. Synthetic antidiabetic thiazolidinediones (TZDs) are well known as ligands and activators for PPAR γ . After it was reported that activation of PPAR γ suppressed production of pro-inflammatory cytokines in activated macrophages, medical interest in PPAR γ has grown and there has been a huge research effort. PPAR γ is currently known to be implicated in various human chronic diseases such as diabetes mellitus, atherosclerosis, rheumatoid arthritis, inflammatory bowel disease, and Alzheimer's disease. Many studies suggest that TZDs not only ameliorate insulin sensitivity, but also have pleiotropic effects on many tissues and cell types. Although activation of PPAR γ seems to have beneficial effects on cardiovascular diseases, the mechanisms by which PPAR γ ligands prevent their development are not fully understood. Recent data about the actions and its mechanisms of PPAR γ -dependent pathway in cardiovascular diseases are discussed here. (Circ J 2009; 73: 214–220)

Key Words: Atherosclerosis; Cardiac hypertrophy; Heart failure; PPAR γ ; Thiazolidinedione

Peroxisome proliferator-activated receptors (PPARs) are transcription factors belonging to the nuclear receptor superfamily that heterodimerize with the retinoid X receptor (RXR) and bind to specific response elements termed PPAR responsive elements (PPREs) in target gene promoters. The PPREs are direct repeats of the hexameric consensus sequence AGGTCA, separated by 1 nucleotide. These nuclear receptors are ligand-dependent transcription factors, and activation of target gene transcription depends on the binding of the ligand to the receptor. PPARs have 3 isoforms, α , β (or δ) and γ . PPAR α regulates genes involved in fatty acid oxidation, whereas PPAR γ promotes adipocyte differentiation and glucose homeostasis. The main function of PPAR β/δ has yet to be ascertained, but involvement in the regulation of fatty acid oxidation seems likely. PPAR α is present mainly in the liver, kidney, and muscle, whereas PPAR γ is expressed predominantly in adipose tissue. PPAR β/δ is almost ubiquitously expressed. It was recently demonstrated that PPAR γ is also expressed in a variety of cell types. After it was reported that activation of PPAR γ suppresses production of inflammatory cytokines in activated macrophages, medical interest in PPAR γ has grown, along with a huge research effort.

PPAR γ

Peroxisome is a subcellular organelle that plays a crucial role in cellular metabolism. Peroxisome enzymes are implicated in a broad range of catabolic and anabolic enzymatic pathways, such as fatty acid oxidation, biosynthesis of both glycerolipids and cholesterol, and metabolism of reactive oxygen species. Peroxisome proliferation induced in rodents is associated with cellular responses to a range of chemical compounds. In 1990, Issemann and Green reported that peroxisome proliferators activate a member of the steroid hormone receptor superfamily in mouse liver! This nuclear receptor was named PPAR. Soon after, 3 major types of PPAR (α , β/δ , and γ) were recognized. PPAR γ is associated with adipocyte differentiation and glucose homeostasis. PPAR γ is expressed in a variety of cell types, including adipocytes, macrophages, vascular smooth muscle cells (VSMCs), endothelial cells (ECs), and cardiomyocytes^{2–7}. Several lines of evidence have demonstrated the functional significance of PPAR γ in atherosclerotic lesions^{8,9}.

Activity of PPAR γ is depressed by phosphorylation of a serine residue (Ser¹¹²) in the N-terminal domain, mediated by a member of the mitogen-activated protein (MAP) kinase family, extracellular signal-regulated protein kinase (ERK). In addition, another member of MAP kinase family, c-Jun N-terminal kinase (JNK) also phosphorylates PPAR γ at Ser⁸² and reduces the transcriptional activity of PPAR γ . The association of PPAR γ polymorphism with metabolic syndrome has also been examined^{10,11}. In the presence of ligand, PPAR γ binds to coactivator complexes, resulting in the activation of target genes. In the absence of ligand, PPAR γ binds to the promoters of several target genes and associates with a corepressor complex, leading to active repression of target genes. This process is referred to as active repression (Fig 1). The corepressor complex constitutes corepressor proteins, such as nuclear receptor corepressor

(Received November 17, 2008; accepted December 8, 2008; released online January 8, 2009)

Department of Cardiovascular Science and Medicine, Chiba University Graduate School of Medicine, Chiba, Japan

Mailing address: Hiroyuki Takano, MD, Department of Cardiovascular Science and Medicine, Chiba University Graduate School of Medicine, 1-8-1 Inohana, Chuo-ku, Chiba 260-8670, Japan. E-mail: htakano-cib@umin.ac.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp




Fig 1. Transactivation and active repression. PPAR γ functions as a heterodimer with RXR. (A) In the presence of ligand, PPAR γ binds to coactivator complexes, resulting in the activation of target genes. (B) In the absence of ligand, PPAR γ binds to the promoters of several target genes and associates with corepressor complexes, leading to active repression of target genes. HDAC, histone deacetylase; PPAR, peroxisome proliferator-activated receptor; PPRE, PPAR responsive element; RXR, retinoid X receptor.

(NCoR) and silencing mediator of retinoid and thyroid hormone receptors, histone deacetylases (HDACs) and transducin β -like protein 1 (TBL1). HDACs are essential in maintaining repressed chromatin structure and TBL1 exchanges a corepressor complex for a coactivator complex in the presence of ligand.¹²

Many nuclear receptors are proposed to sequester inflammatory transcription factors, such as nuclear factor- κ B (NF- κ B) and AP-1, by inhibiting their DNA-binding activities, resulting in inhibition of inflammatory target genes. In the presence of ligand, PPAR γ also interacts with inflammatory transcription factors and inhibits their DNA-binding activities. PPAR γ blocks clearance of the corepressor complex in a ligand-dependent manner, and PPAR γ stabilizes the corepressor complex bound to the promoter of inflammatory genes.¹³ It was demonstrated that PPAR γ associates with the protein inhibitor of activated STAT1 (PIAS1), which is a small ubiquitin-like modifier (SUMO)-E3 ligase, in a ligand-dependent manner. PIAS1-induced SUMOylation of the ligand-binding domain of PPAR γ enables the receptor to maintain NCoR on the promoter of inflammatory genes.¹⁴ These are the suggested mechanisms of PPAR γ transrepression.

PPAR γ Ligands

Natural and synthetic ligands bind to PPAR γ , resulting in conformational change and activation of PPAR γ . The PGD₂ metabolite, 15d-PGJ₂, was the first endogenous ligand for PPAR γ to be discovered. Although 15d-PGJ₂ is the most potent natural ligand of PPAR γ , the extent to which its effects are mediated through PPAR γ *in vivo* remains to be determined. Two components of oxidized low density lipoprotein (ox-LDL), the 9-hydroxy and 13-hydroxy octadecadienoic acids (HODE), are also potent endogenous activators of PPAR γ .^{15,16} Activation of 12/15-lipoxygenase induced by interleukin (IL)-4 also produced endogenous ligands for PPAR γ ;¹⁷ however, whether these natural ligands act as physiological PPAR γ ligands *in vivo* remains unknown. The antidiabetic thiazolidinediones (TZDs), such as troglitazone, pioglitazone, ciglitazone and rosiglitazone, which are used to control glucose concentration in patients with diabetes mellitus (DM), are pharmacological ligands of PPAR γ . They bind PPAR γ with various affinities and it is conceiv-

able that their insulin-sensitizing and hypoglycemic effects are exerted by activating PPAR γ . However, the molecular mechanisms by which TZDs affect insulin resistance and glucose homeostasis are not fully understood. They seem to mediate their effects primarily through adipose tissue, because TZDs alter the expression level of genes that are involved in lipid uptake, lipid metabolism and insulin action in adipocytes. TZDs enhance adipocyte insulin signaling and reduce the release of free fatty acids. TZDs also decrease the inflammation of adipose tissue that is induced by obesity and contributes to increased insulin resistance. There is a possibility that TZDs improve insulin sensitivity in skeletal muscle and liver, the main insulin-sensitive organs, through these multiple adipocentric actions. PPAR γ has been demonstrated to have an antiinflammatory effect, leading to initiation of treatment trials for patients with inflammatory diseases. RXR, which interacts with the PPARs, is activated by 9-*cis* retinoic acid. When combined as a PPAR:RXR heterodimer, the PPAR ligands and 9-*cis* retinoic acid act synergistically on PPAR responses.

PPAR γ and Atherosclerosis

Atherosclerosis is a complex process to which many different factors contribute. Injury of the endothelium, proliferation of VSMCs, migration of monocytes/macrophages, and the regulatory network of growth factors and cytokines are important in the development of atherosclerosis. In addition, chronic inflammation of the vascular wall is also involved. As mentioned earlier, PPAR γ has antiinflammatory effect. PPAR γ ligands have been shown to reduce production of inflammatory cytokines, such as IL-1 β , IL-6, inducible nitric oxide synthase and tumor necrosis factor- α (TNF- α), by inhibiting the activity of transcription factors such as activator protein-1 (AP-1), signal transducers and activators of transcription (STAT), and NF- κ B in monocytes/macrophages.^{2,3} Those findings suggest that PPAR γ activation may have beneficial effects in modulating inflammatory responses in atherosclerosis. Interestingly, expression of PPAR γ has been demonstrated in atherosclerotic plaques.⁸ Macrophages affect the vulnerability of plaque to rupture and they are implicated in the secretion of matrix metalloproteinases (MMPs), enzymes that are important in the degradation of extracellular matrix. In macrophages and VSMCs, PPAR γ




Fig2. In atherosclerosis, PPAR γ inhibits progression of the atherosclerotic lesion. PPAR, peroxisome proliferator-activated receptor.

ligands have been shown to reduce the expression of MMP-9, resulting in the inhibition of migration of VSMCs, and plaque destabilization^{3,4} Although activation of T lymphocytes represents a critical step in atherosclerosis, PPAR γ ligands also reduce the activation T lymphocytes.¹⁸ Recently, it was reported that PPAR γ is a key regulator of M1/M2 polarization.¹⁹ Classically activated macrophages (M1) express a high level of pro-inflammatory cytokines and reactive oxygen species, whereas alternatively activated macrophages (M2) play an anti-inflammatory role in atherosclerosis. PPAR γ agonists prime monocytes into M2 and PPAR γ expression is enhanced by M2 differentiation.²⁰

VSMC proliferation and migration are also critical events in atherosclerosis and vascular-intervention-induced restenosis. TZDs inhibit both these changes in the VSMCs and neointimal thickening after vascular injury^{21–24} Furthermore, TZDs induce apoptosis of VSMCs via p53 and Gadd45^{25,26} Angiotensin II (AngII) plays an important role in vascular remodeling via the AngII type 1 receptor (AT₁R) and accelerates atherosclerosis. Although AngII induces transcriptional suppression of PPAR γ , activation of PPAR γ inhibits AT₁R gene expression at a transcriptional level in VSMCs^{27–29} Expression of adhesion molecule by ECs, leading to adhesion of leukocytes, is a critical early step in atherosclerosis. PPAR γ ligands inhibit the expression of vascular cell adhesion molecule-1 (VCAM-1) and intercellular adhesion molecule-1 and decreased production of chemokines, such as IL-8 and monocyte chemoattractant protein-1 (MCP-1) via suppressions of AP-1 and NF- κ B activities in ECs.^{30–32} PPAR γ ligands also inhibit MCP-1-induced monocytes migration.³³ Endothelin-1 (ET-1) is involved in the regulation of vascular tone and endothelial functions, and induces proliferation of VSMCs. In bovine aortic ECs, PPAR γ ligands suppressed transcription of the ET-1 promoter by interfering with AP-1.³⁴

PPAR γ activation by major oxidized lipid components of ox-LDL, 9-HODE and 13-HODE has an important role in the development of lipid-accumulating macrophages through transcriptional induction of CD36, a scavenger receptor.³⁵ These findings suggest that atherogenic ox-LDL particles could induce their own uptake through activation of PPAR γ and expression of CD36, leading to atherosclerosis. How-

ever, several studies have demonstrated that activation of PPAR γ does not promote lipid accumulation in either mouse or human macrophages.^{36–38} Liver X receptor α (LXR α) is an oxysterol receptor that promotes cholesterol excretion and efflux by modulating expression of ATP-binding cassette transporter 1 (ABCA1).^{37,38} LXR α was recently identified as a direct target of PPAR γ in mouse and human macrophages.^{39,40} Although the PPAR γ -induced increase in CD36 expression might accelerate lipid uptake in macrophages, subsequent activation of LXR α and upregulation of ABCA1 appear to induce lipid efflux.

Diep et al have demonstrated that rosiglitazone and pioglitazone attenuate the development of hypertension and structural abnormalities, and improve endothelial dysfunction in AngII-infused rats.⁴¹ These TZDs also prevented upregulation of AT₁R, cell cycle proteins, and inflammatory mediators. Rosiglitazone, but not the PPAR α ligand fenofibrate, prevented hypertension and endothelial dysfunction in DOCA-salt hypertensive rats.⁴² It has been reported that serum levels of the soluble CD40 ligand are elevated in acute coronary syndrome and associated with increased cardiovascular risk. Treatment with rosiglitazone decreased the serum levels of soluble CD40 and MMP-9 in type 2 diabetic patients with coronary artery disease.⁴³ Taking all the evidence together, PPAR γ ligands may prevent the progression of atherosclerotic lesions, particularly in patients with DM (Fig2).

PPAR γ and Ischemic Heart Disease

As the effects of PPAR γ on the heart are not fully understood, we and others have examined whether PPAR γ is involved in various heart diseases. Although the expression of PPAR γ in cardiac myocytes is low compared with adipocytes, PPAR γ ligands seem to act on cardiac myocytes.^{7,44} We demonstrated that PPAR γ ligands inhibited the cardiac expression of TNF- α at the transcriptional level, in part by antagonizing NF- κ B activity.⁷ Because TNF- α expression is elevated in the failing heart and has a negative inotropic effect on cardiac myocytes, treatment with PPAR γ ligands may prevent the development of congestive heart failure. Diabetic cardiomyopathy, which is characterized by




Fig 3. Actions of PPAR γ in heart diseases. PPAR γ inhibits the progression of heart failure following cardiac hypertrophy, myocardial infarction, ischemia–reperfusion injury, and myocarditis. PPAR, peroxisome proliferator-activated receptor.

systolic and diastolic dysfunction, is a major complication of DM, and therefore TZDs seem to be beneficial for the impaired cardiac function in patients with DM. Following our study, the role of PPAR γ in myocardial ischemia–reperfusion (IR) injury has been elucidated^{45–48}. In animal models, PPAR γ ligands reduced the size of the myocardial infarct and improved contractile dysfunction after IR through inhibition of the inflammatory response. IR injury activates JNK, and subsequently JNK induces increases in both AP-1 DNA-binding activity and apoptotic cells. It has been shown in rats that rosiglitazone inhibits the activation of JNK and AP-1 after myocardial IR⁴⁶. Furthermore, pioglitazone has been reported to attenuate left ventricular remodeling and heart failure after myocardial infarction (MI) in mice⁴⁹. Both of these effects of TZDs ligands were associated with decreases in inflammatory cytokines and chemokines^{49,50}.

PPAR γ and Cardiac Hypertrophy

The PPAR γ ligands, troglitazone, pioglitazone and rosiglitazone, inhibited AngII-induced hypertrophy of neonatal rat cardiac myocytes^{51–53}. Because generalized PPAR γ gene deletion causes embryonic lethality, we examined the role of PPAR γ in the development of cardiac hypertrophy *in vivo* using heterozygous PPAR γ -deficient (PPAR $\gamma^{+/-}$) mice⁵³. Pressure overload-induced cardiac hypertrophy was more prominent in heterozygous PPAR $\gamma^{+/-}$ mice than in wild-type (WT) mice. Treatment with pioglitazone strongly inhibited the pressure overload-induced cardiac hypertrophy in WT mice and moderately in PPAR $\gamma^{+/-}$ mice⁵³. Thereafter, 2 other groups examined the role of PPAR γ in the heart by using cardiomyocyte-specific PPAR γ knockout mice^{54,55}. Duan et al reported that these mice develop cardiac hypertrophy through elevated NF- κ B activity⁵⁴ and unexpectedly, rosiglitazone induced cardiac hypertrophy in both the WT mice and cardiomyocyte-specific PPAR γ knockout mice through activation of p38 MAP kinase independent of PPAR γ . Ding et al reported that cardiomyocyte-specific PPAR γ knockout mice displayed cardiac hypertrophy from approximately 3 months of age and then progress to dilated cardiomyopathy⁵⁵; most mice died from heart failure within 1 year after birth. Mitochondrial oxidative damage and reduced expression of manganese superoxide dismutase were recognized in the cardiomyocyte-specific PPAR γ knockout mice⁵⁵. These mice models demonstrate that PPAR γ is essential for protecting cardiomyocytes from

stress and oxidative damage, although the expression level of PPAR γ in cardiomyocytes is low. On the other hand, Son et al demonstrated that cardiomyocyte-specific PPAR γ transgenic mice develop dilated cardiomyopathy associated with increased uptake of both fatty acid and glucose⁵⁶. Rosiglitazone increased this glucolipotoxicity in cardiomyocyte-specific PPAR γ transgenic mice. If PPAR γ in the heart is expressed at a high level, rosiglitazone may cause cardiotoxic effects; however, as noted earlier the expression level of PPAR γ in the heart is quite low.

PPAR γ and Myocarditis

Experimental autoimmune myocarditis (EAM) is a T-cell-mediated disease characterized by infiltration of T cells and macrophages, leading to massive myocarditis necrosis, which develops into heart failure in the chronic phase⁵⁷. The onset of EAM in rats occurs approximately 2 weeks after the first immunization with porcine cardiac myosin. At this time, small numbers of CD4⁺ T cells and macrophages start to infiltrate into the myocardium and various cytokines are expressed. Macrophage inflammatory protein-1 α (MIP-1 α) is a C-C chemokine that induces leukocyte accumulation in tissue sites of inflammation. We previously demonstrated that MIP-1 α mRNA and protein are highly expressed in the hearts of rats with EAM from day 11 after first immunization⁵⁷. Th1 cells produce interferon- γ (IFN- γ), which is mainly involved in cell-mediated immune responses, whereas Th2 cells produce IL-4, IL-5, IL-6, IL-10 and IL-13, which participate in humoral responses. Immune dysfunction associated with autoimmune disease is known to involve an imbalance between Th1 and Th2 cells.

It has been reported that pioglitazone treatment markedly reduces the severity of myocarditis in a rat model of EAM^{58,59}. Pioglitazone suppressed expression of inflammatory cytokines and activation of myocardiogenic T cells in the myocardium of EAM rats⁵⁸. The mRNA levels of MIP-1 γ were upregulated in the hearts of EAM rats, but not in the hearts of those in the pioglitazone group. Furthermore, treatment with pioglitazone decreased the expression levels of pro-inflammatory cytokine (TNF- α and IL-1 β) genes and Th1 cytokine (IFN- γ) genes, and increased the expression levels of Th2 cytokine (IL-4) gene⁵⁹. These results suggest that PPAR γ ligands may have beneficial effects on myocarditis by inhibiting MIP-1 α expression and modulating the Th1/Th2 balance (Fig 3).

Efficacy and Safety of TZD Treatment in the Clinical Setting

Despite the beneficial effects of TZDs in the basic experiments, their propensity to cause fluid retention is a serious side-effect. Clinical studies report TZD-induced peripheral fluid retention, and an increase in plasma volume in 2–5% of patients on monotherapy.⁶⁰ Fluid retention was more likely to occur with concomitant insulin use, and in patients with underlying cardiac dysfunction or renal insufficiency. The exact mechanisms for TZD-induced fluid retention are not well understood, and it remains unclear whether TZDs directly cause the development of de novo congestive heart failure. It is known that the level of vascular endothelial growth factor is increased in the patients who develop fluid retention with TZD therapy⁶¹ and this may lead to peripheral edema through increased vascular permeability. The insulin-sensitizing action of TZDs also induces water and salt retention. PPAR γ is highly expressed in the kidney and collecting-duct-specific PPAR γ knockout mice demonstrated no effects of TZD on fluid retention or the expression level of sodium channel ENaC- γ .^{62,63} These findings suggest that activation of the sodium channel in the collecting duct cells expressing PPAR γ may be a mechanism of fluid retention. In patients without evidence of heart failure, careful examination did not reveal any worsening of left ventricular function by TZDs.⁶⁴ There are very few studies investigating the safety of TZDs in patients with preexisting heart failure. Although a recent study demonstrated that there is not a direct association between the risk of fluid retention and the baseline degree of severity of heart failure in diabetic patients treated with TZDs, the prescription of TZDs for patients with established heart failure should be avoided at present.^{60,65}

The PROactive (Prospective Pioglitazone Clinical Trial in Macrovascular Events) study has shown that pioglitazone significantly decreases the occurrence of all-cause mortality, nonfatal MI, and nonfatal stroke in patients with type 2 DM and macrovascular diseases.⁶⁶ Pioglitazone significantly reduced the occurrence of fatal and nonfatal MI by 28% in the PROactive study.⁶⁶ Although there was a 1.6% absolute increase in heart failure hospitalizations in the pioglitazone group compared with the placebo group, the number of heart-failure-related deaths was almost identical. In contrast to the PROactive study, it has been recently reported that rosiglitazone treatment is associated with increased incidence of MI by meta-analysis.^{67,68} Although meta-analysis has a number of limitations and the increased risk in MI is still controversial, those results attracted the attention of many clinicians. There are some differences in the actions of pioglitazone and rosiglitazone. Pioglitazone has more beneficial effects on the lipid profile than rosiglitazone.⁶⁹ As mentioned earlier, rosiglitazone, but not pioglitazone, induced cardiac hypertrophy by a non-PPAR γ -mediated pathway.⁵⁴ Pioglitazone represses NF- κ B activation and VCAM-1 expression in a PPAR α -dependent manner.⁷⁰ Pioglitazone was recently reported to increase the number and function of endothelial progenitor cells (EPCs) in patients with stable coronary artery disease and normal glucose tolerance.⁷¹ Pioglitazone may induce angiogenesis by modulating EPC mobilization and function. In the future, more mechanistic studies are required to investigate the differences in action between pioglitazone and rosiglitazone.

Conclusions

The American Heart Association (AHA) and American Diabetes Association (ADA) have released a consensus statement that advises caution regarding the use of TZDs in patients with known or suspected heart failure.⁷² Because there is a possibility that TZDs may unmask asymptomatic cardiac dysfunction by increasing plasma volume, they should be avoided in patients with congestive heart failure of New York Heart Association (NYHA) class III or IV. The data from in vitro studies suggest that TZDs exert direct actions on vascular cells and cardiomyocytes, independent of their glucose-mediated mechanisms. Further studies using tissue-specific gene targeting mice are necessary to address in vivo the pleiotropic effects of PPAR γ on the cardiovascular system. If the beneficial roles of PPAR γ can be solved, modulation of PPAR γ may become a promising therapeutic strategy for cardiovascular diseases. Because cardiac hypertrophy can be seen even in normotensive diabetic patients, and diabetic cardiomyopathy is a major complication of DM, antidiabetic agents such as the TZDs would be expected to have beneficial effects on cardiac hypertrophy and dysfunction in patients with DM. It has been already clarified that 3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors, statins, have pleiotropic effects in cardiovascular diseases. The effects of PPAR γ ligands are similar to those of statins in many respects. A recent study demonstrated that statins activate PPAR γ through ERK and p38 MAP-kinase-dependent cyclooxygenase-2 expression in macrophages.⁷³ Further studies are needed to elucidate the molecular mechanisms of the pleiotropic effects of PPAR γ ligands in cardiovascular disease.

Acknowledgments

This work was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology in Japan, Takeda Science Foundation, and Mitsui Life Social Welfare Foundation.

References

1. Issemann I, Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 1990; **347**: 645–650.
2. Jiang C, Ting AT, Seed B. PPAR- γ agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998; **391**: 82–86.
3. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor- γ is a negative regulator of macrophage activation. *Nature* 1998; **391**: 79–82.
4. Marx N, Schonbeck U, Lazar MA, Libby P, Plutzky J. Peroxisome proliferator-activated receptor γ activators inhibit gene expression and migration in human vascular smooth muscle cells. *Circ Res* 1998; **83**: 1097–1103.
5. Iijima K, Yoshizumi M, Ako J, Eto M, Kim S, Hashimoto M, et al. Expression of peroxisome proliferator-activated receptor γ (PPAR γ) in rat aortic smooth muscle cells. *Biochem Biophys Res Commun* 1998; **247**: 353–356.
6. Benson S, Wu J, Padmanabhan S, Kurtz TW, Pershadsingh HA. Peroxisome proliferator-activated receptor (PPAR)- γ expression in human vascular smooth muscle cells: Inhibition of growth, migration, and c-fos expression by the peroxisome proliferator-activated receptor (PPAR)- γ activator troglitazone. *Am J Hypertens* 2000; **13**: 74–82.
7. Takano H, Nagai T, Asakawa M, Toyozaki T, Oka T, Komuro I, et al. Peroxisome proliferator-activated receptor activators inhibit lipopolysaccharide-induced tumor necrosis factor- α expression in neonatal rat cardiac myocytes. *Circ Res* 2000; **87**: 596–602.
8. Marx N, Sukhova G, Murphy C, Libby P, Plutzky J. Macrophage in human atheroma contain PPAR γ : Differentiation-dependent peroxisomal proliferator-activated receptor γ (PPAR γ) expression and reduction of MMP-9 activity through PPAR γ activation in mononuclear phagocytes in vitro. *Am J Pathol* 1998; **153**: 17–23.