

seconds, and extension at 72°C for 20 seconds. Gene expression levels were measured by real time-PCR with TaqMan[®] Gene Expression Master Mix and StepOnePlus[™] real time PCR (Life Technologies). All probes (*Flt3-L*, *TPO*, *EPO*, *SCF*, *IL-3*, *IL-6*, *IL-11*, *G-CSF* and *GM-CSF*) were from TaqMan[®] Gene Expression Assays (Life Technologies). All samples were assayed in triplicate wells. mRNA levels were normalized to *β-actin* and the relative quantity (RQ) of expression was compared with a reference sample.

Enzyme-Linked Immunosorbent Assay (ELISA). Lysates of sorted cells were obtained using a cell lysis buffer (M-Per[®] Mammalian Protein Extraction Reagent, Thermo Fisher Scientific, Waltham, MA) containing protease inhibitors (Protease Inhibitor Cocktail, Sigma Aldrich, St. Louis, MO). The sample was centrifuged at 14000xg at 4°C for 15 minutes. Supernatants containing soluble protein were collected and protein concentration was estimated by measuring absorbance at 280 nm (NANODROP 2000C, Thermo Fisher Scientific). SCF and EPO in sorted cells were assayed using an ELISA kit (Mouse SCF Immunoassay and Mouse EPO Immunoassay, R&D systems) according to the manufacturer's instructions. Experiments were performed in duplicate. The O.D. was measured using a Thermo Multiskan EX plate reader (Thermo Fisher Scientific).

Immunohistochemistry. Dissected ICR mouse embryos were fixed in 2% paraformaldehyde in PBS overnight at 4°C and washed in PBS three times. After 27% sucrose infusion, embryos were embedded in OCT compound (Sakura Finetek, Tokyo, Japan) and frozen in liquid nitrogen (vapor phase). Frozen embryos were sectioned at 20µm, transferred onto glass slides (Matsunami, Osaka, Japan), and dried. After blocking in 1% BSA in PBS, sections were incubated with primary antibodies overnight at 4°C. Anti-mouse Dlk-1 Ab (MBL), anti-mouse Lyve-1 Ab (MBL), anti-mouse c-Kit Ab (R&D Systems), anti-mouse SCF Ab (Santa Cruz Biotechnology, Santa Cruz, CA), anti-mouse EPO Ab (Santa Cruz Biotechnology, Santa Cruz, CA) and anti-mouse Ki-67 Ab (Dako Corporation, Carpinteria, CA) served as primary antibodies. After washing in PBS three times, sections stained with Dlk-1, Lyve-1, c-Kit or Ki-67 antibodies were incubated with secondary antibodies and TOTO-3 (Life Technologies) for nuclear staining. In samples stained with SCF or EPO antibodies, the TSA Biotin System (PerkinElmer, Covina, CA) was used to amplify the signal. Donkey anti-rabbit IgG-Alexa555, Donkey anti-goat IgG-Alexa488, Donkey anti-rat IgG-Alexa488 and Alexa546 and Streptavidin Alexa546 (all from Life Technologies) served as secondary antibodies. Coverslips were mounted using fluorescent mounting medium (Dako Corporation, Carpinteria, CA). Slides were observed using a FV-1000 confocal microscope (Olympus, Tokyo, Japan).

Single cell preparation and cell counting. To prepare single cell suspensions of HCs from *Map2k4*^{-/-} and wild-type FL at 12.5 dpc, dissected FL was crushed on the 40 μ m nylon cell strainer (BD Falcon, Bedford, MA) with the inner of 2.5 mL syringe. Cells were washed in PBS and collected into the tube. The number of living cells was counted after Trypan Blue staining.

Results

Cytokine expression in hepatoblasts

We examined expression of several cytokine genes in FL tissue samples at 12.5 and 14.5 dpc using real-time PCR. Significant expression of *FMS-like tyrosine kinase 3 ligand (Flt3l)*, *thrombopoietin (TPO)*, *erythropoietin (EPO)* and *interleukin-6 (IL-6)* was seen in FL (Fig. 1A). *Dlk* (Delta-like 1 homolog) is a marker of hepatoblasts (HBs) in FL [16], while Lyve-1 (lymphatic vessel endothelial hyaluronan receptor 1) marks sinusoid endothelial cells (SECs) [17]. As shown in Figure 1B, fractions of HBs, SECs and hematopoietic cells (HCs) were isolated from mouse FL at both 12.5 and 14.5 dpc by flow cytometry based on expression of the following surface markers: (1) HBs, CD45⁻/Ter119⁻/Dlk-1⁺; (2) SECs, CD45⁻/Ter119⁻/Lyve-1⁺/CD31⁺; and (3) HCs, CD45⁺/Ter119⁺. To determine which component contributes to cytokine production, we used real-time PCR to examine *EPO*, *Flt3l*, *IL-6*, *SCF* and *TPO* expression in isolated HBs, SECs and HCs (Fig. 1C). *EPO* and *TPO* were expressed predominantly in HBs both at 12.5 dpc and 14.5 dpc, suggesting that erythropoiesis and megakaryopoiesis are activated by HBs in FL. Levels of *SCF* mRNA were higher in HBs than in SECs. Expression of *Flt3l* and *IL-6* was predominantly detected in HCs, suggesting that HSCs and HPCs expand via an autocrine mechanism. To investigate expression of the cytokine proteins EPO and SCF, we undertook Enzyme-Linked ImmunoSorbent Assay (ELISA) and found that EPO protein was predominantly detected in HBs (63.8 pg/mL/10000 cells) but was not detected in SECs and HCs (Fig. 2A). SCF protein, however, was detected in all fractions (Fig. 2B). In agreement with

SCF mRNA expression, HBs expressed SCF protein at a higher level (7.01 pg/mL/10000 cells) than did SECs (2.38 pg/mL/10000 cells) or HCs (0.025 pg/mL/10000 cells). When we compared expression levels of EPO and SCF proteins in each fraction, EPO expression was highest in HBs, suggesting that HBs secrete EPO to regulate FL erythropoiesis (Fig. 2C). To confirm EPO and SCF protein localization in FL, we performed immunohistochemistry and found that EPO protein was expressed primarily in Dlk-1-expressing HBs (Fig. 2D), in agreement with ELISA analysis. Staining of FL with anti-SCF antibody revealed SCF protein in both SECs expressing Lyve-1 and HBs expressing Dlk-1 (Fig. 2E-F).

Fetal liver of Map2k4^{-/-} mouse embryos

To characterize HB function in FL, we analyzed *Map2k4* (*mitogen-activated protein kinase kinase 4*, formerly known as *Sek1* and *MKK4*)^{-/-} mouse embryos, which lack FL HBs [18-20]. Real-time PCR analysis of whole FL from mutant and wild-type embryos showed that among cytokine-encoding genes, *EPO* and *SCF* mRNAs were down-regulated in the mutant mice, implying an impairment in hematopoiesis, particularly erythropoiesis, in *Map2k4* mutant mice (Fig. 3A). To evaluate potential alterations in hematopoiesis, we performed immunohistochemistry and found that the number of c-Kit expressing cells which represent HSCs and HPCs in the FL at this stage, decreased in a FL field from *Map2k4^{-/-}* mouse embryo relative to wild-type embryo (Fig. 3B; green) [23]. In addition, the number of cells expressing Ki-67, a marker of cell proliferation, decreased in a FL field from *Map2k4^{-/-}* mouse embryos

compared to wild-type embryos (Fig. 3B; red). In agreement with the Ki-67 staining, there were fewer FL cells seen in *Map2k4*^{-/-} versus wild-type embryos (Fig. 3C).

Discussion

In FL, HSCs differentiate into mature HCs, particularly erythroid cells [9,10]. SCF and EPO are representative cytokines that regulate erythropoiesis [14,15]. Expression levels of *SCF* and *EPO* genes were higher in whole FL tissue than in adult BM and kidney, suggesting that FL primarily functions in erythropoiesis [24]. Recently, using a competitive repopulating assay, Chou and Lodish reported that FL stromal cells expressing both SCF and Dlk-1 support HSC maintenance [25]. However, it remained unclear which cells secreted EPO in FL. HBs are regarded as common progenitors of hepatocytes and biliary epithelial cells and thought to support liver construction through formation of a mesh-like structure [16,26]. Our results are strongly indicative of an additional role of HBs in producing erythropoietic cytokines SCF and EPO. Although gene expression of *IL-6* was observed in FL at 12.5 dpc, that was not detected in cell fractions of HBs, SECs and HCs (Fig. 1). As shown in Fig. 1B, we could observe unclassified cell fraction (CD45⁻/Ter119⁻/Dlk-1⁻/Lyve-1⁻/CD31⁻), based on surface marker expression, implying that this unclassified cell fraction may express *IL-6* gene. It will be further necessary to clarify roles of the unclassified cells in FL hematopoiesis in the future.

The c-Jun NH₂-terminal protein kinases (JNKs) are members of the mitogen-activated protein kinase (MAPK) family. Map2k4 is a MAPK kinase that directly activates JNKs in response to extracellular and intracellular stresses, and its deficiency leads to abnormal hepatogenesis, resulting in loss of HBs [20,27-29]. In situ hybridization shows that expression of *Map2k4* transcripts is detected in FL by 12.5 dpc and then up-regulated until 16.5 dpc, at which time expression is down-regulated [30]. When *Map2k4*^{-/-} FL cells were transplanted into *rag1*^{-/-} adult mice, both B cells and T cells were generated, demonstrating that *Map2k4*^{-/-} FL cells contain HSCs and/or HPCs [28]. In addition, normal hematopoiesis reportedly occurs in the *Map2k4*^{-/-} YS at both 9.5 and 10.5 dpcs, based on erythroid, myeloid and mixed colony formation [28]. Taken together, hematopoietic potential of *Map2k4*^{-/-} FL looked normal compared to wild-type mice. Therefore, the decrease of hematopoietic cells were likely due to lack of HBs, but not due to alteration of hematopoietic potential in *Map2k4*^{-/-} FL. Recently, Hikita et al. reported a novel mouse model lacking both HBs and hepatocytes using an *Alb Cre* driver [31]. This mutant mouse, generated by crossing *Alb Cre* mice with both *bcl-x1*^{flox/flox} mice and *mcl-1*^{flox/flox} mice, showed a decreased number of hepatocytes at 18.5 dpc. Future studies will be required to address the function of HBs and hepatocytes in FL erythropoiesis using this model.

Conclusion

Hepatoblasts comprise a niche for erythropoiesis through cytokine secretion.

Acknowledgements

The authors acknowledge grant support from the Special Coordination Funds for Promoting Science and Technology, a Grant-in-Aid for Young Scientists (B) from The Ministry of Education, Culture, Sports, Science and Technology, Bilateral Joint Projects for Japan Society for the Promotion of Science, the Ajinomoto Scholarship Foundation and the Tokyo Biochemical Research Foundation. We thank the Research Support Center, the Graduate School of Medical Sciences, Kyushu University, for technical support, Drs. K. Akashi and K. Tani for helpful discussion, Dr. T. Inoue for technical support, and Dr. Elise Lamar for critical reading of the manuscript.

Authorship contributions

D. Sugiyama designed the research, performed research, analyzed data and wrote the paper. K. Kulkeaw performed research and analyzed data. C. Mizuochi, Y. Horio and S. Okayama performed research.

Conflict of Interest Disclosures

The authors have no conflict of interest to declare.

References

- [1] I.L. Weissman, Stem cells: units of development, units of regeneration, and units in evolution, *Cell* 100 (2000) 157-168.
- [2] E. Dzierzak, A. Medvinsky, M. de Bruijn, Qualitative and quantitative aspects of hematopoietic cell development in the mammalian embryo, *Immunol. Today*, 19 (1998) 228-236.
- [3] S. Matsuoka, K. Tsuji, H. Hisakawa, et al., Generation of definitive hematopoietic stem cells from murine early yolk sac and paraaortic splanchnopleures by aorta-gonad-mesonephros region-derived stromal cells, *Blood* 98 (2001) 6-12.
- [4] D. Sugiyama, K. Tsuji, Definitive hematopoiesis from endothelial cells in the mouse embryo; a simple guide, *Trends Cardiovasc. Med.* 16 (2006) 45-49.
- [5] B.M. Zeigler, D. Sugiyama, M. Chen, Y. Guo, K.M. Downs, N.A. Speck, The allantois and chorion, when isolated before circulation or chorio-allantoic fusion, have hematopoietic potential, *Development* 133 (2006) 4183-4192.
- [6] I.M. Samokhvalov, N.I. Samokhvalova, S. Nishikawa, Cell tracing shows the contribution of the yolk sac to adult haematopoiesis, *Nature* 446 (2007) 1056-1061.
- [7] K.E. Rhodes, C. Gekas, Y. Wang, et al., The emergence of hematopoietic stem cells is initiated in the placental vasculature in the absence of circulation, *Cell Stem Cell* 2 (2008) 252-263.

- [8] T. Sasaki, C. Mizuochi, Y. Horio, K. Nakao, K. Akashi, D. Sugiyama, Regulation of hematopoietic cell clusters in the placental niche through SCF/Kit signaling in embryonic mouse, *Development* 137 (2010) 3941-3952.
- [9] G.R. Johnson, R.O. Jones, Differentiation of the mammalian hepatic primordium in vitro. I. Morphogenesis and the onset of haematopoiesis, *J. Embryol. Exp. Morphol.* 30 (1973) 83-96.
- [10] G.R. Johnson, M.A. Moore, Role of stem cell migration in the initiation of mouse foetal liver haemopoiesis, *Nature* 258 (1975) 726-728.
- [11] J. Palis, S. Robertson, M. Kennedy, C. Wall, G. Keller, Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse, *Development* 126 (1999) 5073-5084.
- [12] Kurata H, Mancini GC, Alespeiti G, A.R. Miqliaccio, G. Miqliaccio, Stem cell factor induces proliferation and differentiation of fetal progenitor cell in the mouse, *Br. J. Haematol.* 101 (1998) 676-687.
- [13] P.D. Kingsley, J. Malik, K.A. Fantauzzo, J. Palis, Yolk sac-derived primitive erythroblasts enucleate during mammalian embryogenesis, *Blood* 104 (2004) 19-25.
- [14] H. Lodish, J. Flygare, S. Chou, From stem regulation of red cell production at multiple levels by multiple hormones, *IUBMB Life* 62 (2010) 492-496.
- [15] G. Molineux, M.A. Foote, S. Elliott, *Erythropoietins and Erythropoiesis*, Birkhauser, Basel, Switzerland, 2009.

- [16] N. Tanimizu, M. Nishikawa, H. Saito, T. Tsujimura, A. Miyajima, Isolation of hepatoblasts based on the expression of Dlk/Pref-1, *J. Cell Sci.* 116 (2003) 1775-1786.
- [17] C. Mouta Carreira, S.M. Nasser, E. di Toamaso, T.P. Padera, Y. Boucher, S.I. Tomarev, R.K. Jain, LYVE-1 is not restricted to the lymph vessels: expression in normal liver blood sinusoids and down-regulation in human liver cancer and cirrhosis, *Cancer Res.* 61 (2001) 8079-8084.
- [18] H. Nishina, K.D. Fischer, L. Radvanyi, et al., Stress-signalling kinase Sek1 protects thymocytes from apoptosis mediated by CD95 and CD3, *Nature* 385 (1997) 350-353.
- [19] H. Nishina, M. Bachmann, A.J. Oliveira-dos-Santos, et al., Impaired CD28-mediated interleukin 2 production and proliferation in stress kinase SAPK/ERK1 kinase (SEK1)/mitogen-activated protein kinase kinase 4 (MKK4)-deficient T lymphocytes, *J. Exp. Med.* 186 (1997) 941-953.
- [20] T. Watanabe, K. Nakagawa, S. Ohata, et al., SEK1/MKK4-mediated SAPK/JNK signaling participates in embryonic hepatoblast proliferation via a pathway different from NF- κ B-induced anti-apoptosis, *Dev. Biol.* 250 (2002) 332-347.
- [21] N. Suzuki, N. Suwabe, O. Ohneda, N. Obara, S. Imagawa, X. Pan, H. Motohashi, M. Yamamoto, Identification and characterization of 2 types of erythroid progenitors that express GATA-1 at distinct levels, *Blood* 102 (2003) 3575-3583.

- [22] U. Testa, Apoptotic mechanisms in the control of erythropoiesis, *Leukemia* 18 (2004) 1176-1199.
- [23] T. Inoue, D. Sugiyama, R. Kurita et al., APOA-1 is a novel marker of erythroid cell maturation from hematopoietic stem cells in mice and humans, *Stem Cell Rev and Rep* 7 (2011) 43-52.
- [24] S.J. England, K.E. McGrath, J.M. Frame, J. Palis, Immature erythroblasts with extensive ex vivo self-renewal capacity emerge from the early mammalian fetus. *Blood* 117 (2010) 2708-2717.
- [25] S. Chou, H.F. Lodish, Fetal liver hepatic progenitors are supportive stromal cells for hematopoietic stem cells, *Proc. Natl. Acad. Sci. USA.* 107 (2010) 7799-804.
- [26] K. Sasaki, Y. Sonoda, Histometrical and three-dimensional analyses of liver hematopoiesis in the mouse embryo, *Arch. Histol. Cytol.* 63 (2000) 137-146.
- [27] Ganiatsas S, Kwee L, Fujiwara Y, A. Perkins, T. Ikeda, M.A. Labow, L.I. Zon, SEK1 deficiency reveals mitogen-activated protein kinase cascade crossregulation and leads to abnormal hepatogenesis, *Proc. Natl. Acad. Sci. USA.* 95 (1998) 6881-6886.
- [28] Nishina H, Vaz C, Billia P, et al., Defective liver formation and liver cell apoptosis in mice lacking the stress signaling kinase SEK1/MKK4, *Development* 126 (1999) 505-516.
- [29] X. Wang, A. Destrumont, C. Tournier, Physiological roles of MKK4 and

MKK7: insights from animal models, *Biochim. Biophys. Acta.* 1773 (2007) 1349-1357.

- [30] J.K. Lee, W.S. Hwang, Y.D. Lee, P.L. Han, Dynamic expression of SEK1 suggests multiple roles of the gene during embryogenesis and in adult brain of mice, *Brain Res. Mol. Brain Res.* 66 (1999) 133-140.
- [31] H. Hikita, T. Takehara, S. Shimizu, Mcl-1 and Bcl-xL cooperatively maintain integrity of hepatocytes in developing and adult murine liver, *Hepatology* 50 (2009), 1217-1226.

Figure Legends

Fig. 1. Cytokine gene expression in fetal liver. (A) *FMS-like tyrosine kinase 3 ligand (Flt3l)*, *thrombopoietin (TPO)*, *erythropoietin (EPO)*, *stem cell factor (SCF)*, *interleukin-3 (IL-3)*, *interleukin-6 (IL-6)*, *interleukin-11 (IL-11)*, *granulocyte-colony stimulating factor (G-CSF)* and *granulocyte/macrophage-colony stimulating factor (GM-CSF)* mRNAs were examined in FL samples at 12.5 and 14.5 dpc by real-time PCR. Note high expression of *Flt3l*, *TPO*, *EPO* and *IL-6* in FL. (B) A single cell suspension was obtained from FL at 12.5 dpc and expression of CD45/Ter119, Dlk-1, Lyve-1 and CD31 was analyzed by flow cytometry. (1) CD45⁻/Ter119⁻/Dlk-1⁺ defines HBs; (2) CD45⁻/Ter119⁻/Lyve-1⁺/CD31⁺ defines SECs; and (3) CD45⁺/Ter119⁺ defines HCs. (C) Expression of *Flt3l*, *TPO*, *EPO* and *IL-6* was examined by real-time PCR in HBs, SECs and HCs sorted by flow cytometry, according to gates defined in Figure 1B. *EPO* and *TPO* expression was high in HBs at both 12.5 dpc and 14.5 dpc. *SCF* expression was higher in HBs than in SECs. Expression of *Flt3l* was high in HCs. Expression of *IL-6* was detected only in HCs.

Fig. 2. Expression of EPO and SCF protein in fetal liver. (A) EPO protein in sorted cells was assayed by ELISA. EPO was detected in HBs (63.8 pg/mL/10000 cells) but not in SECs and HCs at 12.5 dpc. (B) SCF protein in sorted cells was assayed by ELISA. SCF was detected in HBs (7.01 pg/mL/10000 cells), SECs (2.38 pg/mL/10000

cells) and HCs (0.025 pg/mL/10000 cells) at 12.5 dpc. (C) The data from Figure 2A with 2B were combined to compare EPO and SFC protein expression in each fraction sorted from fetal liver. EPO protein expression was highest in HBs. (D-F) Liver sections were prepared from ICR mouse embryos at 12.5 dpc and stained with Dlk-1 (green), EPO (red) and TOTO-3 (blue) (D); Lyve-1 (green), SCF (red) and TOTO-3 (blue) (E); and Dlk-1 (green), SCF (red) and TOTO-3 (blue) (F). Arrowheads show co-localization of the antigens, respectively. Samples were observed under confocal microscopy. EPO was primarily expressed in HBs expressing Dlk-1, whereas SCF was more widely expressed in FL. Original magnification is 40x. (D-F)

Fig. 3. Phenotypic analysis of FL in *Map2k4*^{-/-} mouse embryos. (A) Expression of *EPO* and *SCF* was examined by real-time PCR in *Map2k4*^{-/-} fetal liver (FL). Expression levels of both *EPO* and *SCF* were down-regulated in *Map2k4*^{-/-} FL relative to wild-type FL. (B) Liver sections were made from both *Map2k4*^{-/-} and wild-type mouse embryos at 12.5 dpc, stained with c-Kit (green), Ki-67 (red) and TOTO-3 (blue), and observed under confocal microscopy. The number of Ki-67 positive cells in a field decreased in *Map2k4*^{-/-} FL compared to wild-type FL. Original magnification is 40x. (C) Single cell suspensions were obtained from *Map2k4*^{-/-} and wild-type FL at 12.5 dpc and the number of living cells was counted after Trypan Blue staining. Significant decreases in the number of *Map2k4*^{-/-} FL cells were observed compared to wild-type FL cells.

Figure 1A




Figure 1B

